

$$\mu_0 (1 - p^2 - 1 - p^2 + 2p) = 2$$

$$\mu_0 (2p - 2p^2) = 2$$

$$\mu_0 = \frac{1}{p - p^2}$$

$$\mu_0 = \frac{1}{p(1-p)}$$

OSSERVAZIONI

Calcoliamo per completezza μ_1 e μ_2 :

$$\mu_1 = 1 + p\mu_0 = 1 + \frac{1}{p(1-p)} = \frac{1-p+1}{1-p} = \frac{2-p}{1-p}$$

$$\mu_2 = 1 + (1-p)\mu_0 = 1 + \frac{1}{p} = \frac{p+1}{p}$$


Ci si aspetta che $\mu_1 > \mu_0$ e $\mu_2 > \mu_0$. In effetti:

$$\frac{2-p}{1-p} > \frac{1}{p(1-p)}$$

$$p(2-p) > 1$$

errore qui!

SI




$$\frac{p+1}{p} > \frac{1}{p(1-p)}$$

$$(p+1)(1-p) > 1$$

errore qui!

SI



La parte finale va sostituita come segue.

Ci si aspetta che

$$\mu_1 < \mu_0 \text{ e } \mu_2 < \mu_0.$$

In effetti si verificano così:

$$\frac{2-p}{1-p} < \frac{1}{p(1-p)}, \quad p(2-p) < 1,$$

$$-p^2 + 2p - 1 < 0, \quad -(p-1)^2 < 0 \quad \text{(SI)}$$

$$\frac{p+1}{p} < \frac{1}{p(1-p)}, \quad (p+1)(1-p) < 1,$$

$$-p + p - p^2 < 1, \quad -p^2 < 0 \quad \text{(SI)}$$