

# VV186 RC Part IV

## Integration

“When learning integration, integrate all knowledge you have...”

Pingbang Hu

University of Michigan-Shanghai Jiao Tong University Joint Institute

July 14, 2022



JOINT INSTITUTE

交大密西根学院

1. Motivation
2. step Function
3. Regulated Integral
4. Darboux Integral
5. Riemann Integral
6. Exercise

The motivation for integration is simple:

- ▶ Find area under curves.
- ▶ Find signed area for real life application.
- ▶ Do different kinds of *transformation*.
- ▶ Solving equations.

- ▶ Partition
- ▶ Properties of step functions
  1. If  $\varphi$  is a step function on  $[a, b]$ , then  $\forall \alpha \in \mathbb{R}$ ,  $\alpha\varphi$  is also a step function on  $[a, b]$ .
  2. If  $\varphi, \Psi$  are two step functions on  $[a, b]$ , then  $\varphi + \Psi$  is also a step function on  $[a, b]$ .
  3. The set consisting of step functions on  $[a, b]$  where  $a, b \in \mathbb{R}$  and  $a < b$  is a **vector space**. (Why?)

## Properties of step function integral

- ▶ Given a step function  $\varphi : [a, b] \rightarrow \mathbb{R}$ , its integral exists and doesn't depend on the choice of partition.
- ▶ Let  $T : \text{Step}([a, b]) \rightarrow \mathbb{R}$  be a function (functional) and  $T(\varphi) = \int_a^b \varphi$ , then  $T$  is a linear function that maps non-negative step functions to non-negative values.
- ▶ Let  $\varphi \in [a, b]$ , then  $|\int_a^b \varphi| \leq \int_a^b |\varphi|$

( $T$  is sometimes called "positive linear functional")

Let  $f \in \text{Reg}([a, b])$  and  $(\varphi_n)$  a sequence in  $\text{Step}([a, b])$  converging uniformly to  $f$ . Then the regulated integral of  $f$ , defined by

$$\int_a^b f := \lim \int_a^b \varphi_n$$

exists and does not depend on the choice of  $(\varphi_n)$ .

Let  $[a, b] \subseteq \mathbb{R}$  be a closed interval and  $f$  a bounded real function on  $[a, b]$ . Let  $u_f$  denote the set of all step functions  $u$  on  $[a, b]$  such that  $u \geq f$  and denote  $l_f$  the set of all step functions  $l$  on  $[a, b]$  such that  $f \geq l$ . The function  $f$  is then said to be Darboux-integrable if

$$\mathbb{I}(f) = \sup_{l \in l_f} \int_a^b l = \inf_{u \in u_f} \int_a^b u = \bar{\mathbb{I}}(f)$$

We denote the integral of  $f$  by  $\mathbb{I}(f)$  to distinguish with the lower step functions  $l$ .

Let  $[a, b] \subseteq \mathbb{R}$  be a closed interval and  $f$  a bounded real function on  $[a, b]$ . Then  $f$  is Riemann-integrable with integral  $\int_a^b f \in \mathbb{R}$  if for every  $\epsilon > 0$  there exists a  $\delta > 0$  such that for any tagged partition on  $[a, b]$  with mesh size  $m(P) < \delta$ ,

$$\left| \sum_{k=1}^n f(\xi_k)(x_k - x_{k-1}) - \int_a^b f \right| < \epsilon$$

Comment. It is also a usual way to define Riemann integral by Darboux integral. In fact, Riemann integral is usually *just the Darboux integral*.



The following are some results / Theorems & comments for integrals. They apply to all the three kinds of integrals that we learn in Vv186.

1. The integral is a linear map that maps non-negative functions to non-negative values.
2. Let  $f, g$  be two regulated functions on  $[a, b]$ . Moreover, if  $f \leq g$ , then  $\int_a^b f \leq \int_a^b g$ .  
Comment. This can be proved by taking the limit of the step functions.

3. The integral of  $f$  doesn't change if  $f$  changes its value on a **finite** set and is still integrable.
4. The integral of  $f$  doesn't change if  $f$  changes its value on a **countable** set and is still integrable. If we take the usual understanding of "length of interval" to integrate.  
Comment. It is important that  $f$  is *integrable*.  
Comment. This set is usually called *measured zero set*.
5. The Riemann integral, Darboux integral and the regulated integral of  $f$  exist and coincide if  $f$  is regulated.

1. This exercise gives an example of using step function sequence to find the integral of a function. Calculate

$$\int_0^1 x^4.$$

2.

- ▶ Check whether the function  $J$  given by

$$J : [0, 1] \rightarrow \mathbb{R}, \quad J(x) = \begin{cases} 1, & x = \frac{1}{n}, \\ 0, & \text{otherwise;} \end{cases}$$

is regulated.

- ▶ Check whether the function  $J$  we defined in i) is Riemann integrable.

3. Please judge whether the following statements are true or false.

- ▶  $f(x) = x^3 + e^x$  on  $[-1, 1]$  is regulated.
- ▶ Since  $f(x) = x^2$  is regulated on each  $[-n, n]$ , where  $n \in \mathbb{N}$ ,  $f$  is regulated on  $\mathbb{R}$ , because we can let  $n \rightarrow \infty$ .
- ▶ A continuous function is piecewise continuous.
- ▶ Let  $f, g$  be two real-valued functions defined on  $[0, 1]$ . Furthermore, assume  $f - g = x$ , then the equation

$$\int_0^1 f - \int_0^1 g = \frac{1}{2}$$

holds.

- ▶ Let  $f \in \text{Reg}([0, 1])$ , let  $g$  be a real-valued function. The  $f \circ g$  is regulated.

4. Now let's prove the assertion on Slide 543. Let  $f$  be a bounded real function on  $[a, b]$  and  $f$  is Darboux integrable. Please show that it is Riemann integrable on  $[a, b]$ .

5. Let  $f$  be a piecewise continuous function on  $[a, b]$ . Prove that  $f$  is regulated, i.e., for any  $\epsilon > 0$ , there exists a step function  $\varphi$  such that

$$\sup_{x \in [a, b]} |f(x) - \varphi(x)| < \epsilon$$

6. This exercise aims at showing two more characteristics of  $\text{Reg}([a, b])$ . Let  $f, g \in \text{Reg}([a, b])$ ,

1. Show that the  $f \cdot g \in \text{Reg}([a, b])$
2. Show that  $f^2 \in \text{Reg}([a, b])$
3. Suppose  $f \geq p > 0$  for some  $p \in \mathbb{R}$ . Show that  $\frac{1}{f} \in \text{Reg}([a, b])$ .



7. Give an example such that  $\int_a^b |f|$  is Riemann integrable but  $\int_a^b f$  is not Riemann integrable. Is the converse true?

8. Let  $f : [a, b] \rightarrow \mathbb{R}$ ,  $f$  is monotonic on  $[a, b]$ . Prove that  $f$  is regulated. Is  $f$  integrable?

9. Given  $\epsilon > 0$ . For any regulated function  $f : [0, 1] \rightarrow \mathbb{R}$ . There exists a continuous function  $g : [0, 1] \rightarrow \mathbb{R}$  such that

$$\left| \int_0^1 f - \int_0^1 g \right| < \epsilon \quad \text{and} \quad |f(x) - g(x)| < \epsilon \text{ on } [0, 1]$$

except for a finite number of intervals with total length less than  $\epsilon$ .

## Any Question?

# Have Fun and Learn Well!