




The figure displays four diagnostic plots for a linear regression model, arranged in a 2x2 grid. The model is defined as $\text{stat} \sim \text{scale}(\text{tempanom_DW_tstat}) + \text{scale}(\text{prcp_DW_totastat})$.

- Top Left: Standardized Residuals vs Fitted**
 - Y-axis:** Standardized residuals (ranging from -2 to 2).
 - X-axis:** Fitted values (ranging from -1.0 to 1.0).
 - Description:** A scatter plot showing the distribution of standardized residuals against fitted values. A horizontal dotted line is drawn at y=0. Most points are clustered between y=-1 and y=1, with a few outliers at higher values.
- Top Right: Normal Q-Q**
 - Y-axis:** Standardized residuals (ranging from -2 to 2).
 - X-axis:** Theoretical Quantiles (ranging from -2 to 2).
 - Description:** A Q-Q plot showing standardized residuals against theoretical quantiles. The points closely follow a solid diagonal line, indicating that the residuals are approximately normally distributed.
- Bottom Left: Scale-Location**
 - Y-axis:** $\sqrt{|\text{Standardized residuals}|}$ (ranging from 0.4 to 1.2).
 - X-axis:** Fitted values (ranging from -1.0 to 1.0).
 - Description:** A plot showing the square root of the absolute value of standardized residuals against fitted values. The points are scattered around a value of 0.8, with no clear trend, suggesting constant variance.
- Bottom Right: Cook's dist vs Leverage**
 - Y-axis:** Cook's distance (ranging from 0.00 to 0.30).
 - X-axis:** Leverage (ranging from 0.05 to 0.25).
 - Description:** A plot showing Cook's distance against leverage. Most points are near the origin, with one notable outlier at approximately (0.25, 0.30).


$$\text{stat} \sim \text{scale}(\text{tempanom} \mid \text{DW} \mid \text{tstat}) + \text{scale}(\text{prcp} \mid \text{DW} \mid \text{totstat}) \sim \text{scale}(\text{tempanom} \mid \text{DW} \mid \text{tstat}) + \text{scale}(\text{prcp} \mid \text{DW} \mid \text{totstat})$$

The figure displays four diagnostic plots for a linear regression model, arranged in a 2x2 grid. The model is defined as $\text{stat} \sim \text{scale}(\text{tempanom_DW_tstat}) + \text{scale}(\text{prcp_DW_totastat})$.

- Top Left: Standardized Residuals vs Fitted**
 - Y-axis:** Standardized residuals (ranging from -2 to 2).
 - X-axis:** Fitted values (ranging from -1.0 to 1.0).
 - Description:** A scatter plot showing the distribution of standardized residuals against fitted values. A horizontal dotted line is drawn at y=0. Most points are clustered between y=-1 and y=1, with a few outliers at higher values.
- Top Right: Normal Q-Q**
 - Y-axis:** Standardized residuals (ranging from -2 to 2).
 - X-axis:** Theoretical Quantiles (ranging from -2 to 2).
 - Description:** A Q-Q plot showing standardized residuals against theoretical quantiles. The points closely follow a solid diagonal line, indicating that the residuals are approximately normally distributed.
- Bottom Left: Scale-Location**
 - Y-axis:** $\sqrt{|\text{Standardized residuals}|}$ (ranging from 0.4 to 1.2).
 - X-axis:** Fitted values (ranging from -1.0 to 1.0).
 - Description:** A plot showing the square root of the absolute value of standardized residuals against fitted values. The points are scattered around a value of 0.8, with no clear trend, suggesting constant variance.
- Bottom Right: Cook's dist vs Leverage**
 - Y-axis:** Cook's distance (ranging from 0.00 to 0.30).
 - X-axis:** Leverage (ranging from 0.05 to 0.25).
 - Description:** A plot showing Cook's distance against leverage. Most points are near the origin, with one notable outlier at approximately (0.25, 0.30).



The figure displays four diagnostic plots for a linear regression model, arranged in a 2x2 grid. The model is defined as $\text{stat} \sim \text{scale}(\text{tempanom_DW_tstat}) + \text{scale}(\text{prcp_DW_totastat})$.

- Top Left: Standardized Residuals vs Fitted**
 - Y-axis:** Standardized residuals (ranging from -2 to 2).
 - X-axis:** Fitted values (ranging from -1.0 to 1.0).
 - Description:** A scatter plot showing the distribution of standardized residuals against fitted values. A horizontal dotted line is drawn at y=0. Most points are clustered between y=-1 and y=1, with a few outliers at higher values.
- Top Right: Normal Q-Q**
 - Y-axis:** Standardized residuals (ranging from -2 to 2).
 - X-axis:** Theoretical Quantiles (ranging from -2 to 2).
 - Description:** A Q-Q plot showing standardized residuals against theoretical quantiles. The points closely follow a solid diagonal line, indicating that the residuals are approximately normally distributed.
- Bottom Left: Scale-Location**
 - Y-axis:** $\sqrt{|\text{Standardized residuals}|}$ (ranging from 0.4 to 1.2).
 - X-axis:** Fitted values (ranging from -1.0 to 1.0).
 - Description:** A plot showing the square root of the absolute value of standardized residuals against fitted values. The points are scattered around a value of 0.8, with no clear trend, suggesting constant variance (homoscedasticity).
- Bottom Right: Cook's dist vs Leverage**
 - Y-axis:** Cook's distance (ranging from 0.00 to 0.30).
 - X-axis:** Leverage (ranging from 0.05 to 0.25).
 - Description:** A plot showing Cook's distance against leverage. Most points have low leverage and low Cook's distance. One point at approximately (0.25, 0.30) is a notable outlier with high leverage and high Cook's distance.

