



The figure displays four diagnostic plots for a linear regression model, arranged in a 2x2 grid. The model equation is $\text{stat} \sim \text{scale}(\text{tempanom_DW_tstat}) + \text{scale}(\text{prcp_DW_totastat})$.

- Standardized Residuals vs Fitted:** A scatter plot showing standardized residuals on the y-axis (ranging from -2 to 2) against fitted values on the x-axis (ranging from -0.5 to 0.5). The residuals are scattered around a horizontal dashed line at zero.
- Normal Q-Q:** A Q-Q plot showing standardized residuals on the y-axis (ranging from -2 to 2) against theoretical quantiles on the x-axis (ranging from -2 to 2). The points closely follow a solid diagonal line, indicating approximate normality of the residuals.
- Scale-Location:** A scatter plot showing the square root of the absolute value of standardized residuals on the y-axis (ranging from 0.0 to 1.5) against fitted values on the x-axis (ranging from -0.5 to 0.5). The points are scattered, suggesting constant variance.
- Cook's dist vs Leverage:** A scatter plot showing Cook's distance on the y-axis (ranging from 0.0 to 0.4) against leverage on the x-axis (ranging from 0.0 to 0.4). Most points are clustered near the origin, with a few points showing higher leverage and Cook's distance.




The figure displays four diagnostic plots for a linear regression model, arranged in a 2x2 grid. The model equation is $\text{stat} \sim \text{scale}(\text{tempanom_DW_tstat}) + \text{scale}(\text{prcp_DW_totastat})$.

- Standardized Residuals vs Fitted:** A scatter plot showing standardized residuals on the y-axis (ranging from -2 to 2) against fitted values on the x-axis (ranging from -0.5 to 0.5). The residuals are scattered around a horizontal dashed line at zero.
- Normal Q-Q:** A Q-Q plot showing standardized residuals on the y-axis (ranging from -2 to 2) against theoretical quantiles on the x-axis (ranging from -2 to 2). The points follow a solid diagonal line, indicating approximate normality.
- Scale-Location:** A scatter plot showing the square root of the absolute value of standardized residuals on the y-axis (ranging from 0.0 to 1.5) against fitted values on the x-axis (ranging from -0.5 to 0.5). The points are scattered, suggesting constant variance.
- Cook's dist vs Leverage:** A scatter plot showing Cook's distance on the y-axis (ranging from 0.0 to 0.4) against leverage on the x-axis (ranging from 0.0 to 0.4). Most points are clustered near the origin, with a few points showing higher leverage and Cook's distance.



The figure displays four diagnostic plots for a linear regression model, arranged in a 2x2 grid. The model equation is $\text{stat} \sim \text{scale}(\text{tempanom_DW_tstat}) + \text{scale}(\text{prcp_DW_totastat})$.

- Standardized Residuals vs Fitted:** A scatter plot showing standardized residuals on the y-axis (ranging from -2 to 2) against fitted values on the x-axis (ranging from -0.5 to 0.5). The residuals are scattered around a horizontal dashed line at zero.
- Normal Q-Q:** A Q-Q plot showing standardized residuals on the y-axis (ranging from -2 to 2) against theoretical quantiles on the x-axis (ranging from -2 to 2). The points follow a solid diagonal line, indicating approximate normality.
- Scale-Location:** A scatter plot showing the square root of the absolute value of standardized residuals on the y-axis (ranging from 0.0 to 1.5) against fitted values on the x-axis (ranging from -0.5 to 0.5). The points are scattered, suggesting constant variance.
- Cook's dist vs Leverage:** A scatter plot showing Cook's distance on the y-axis (ranging from 0.0 to 0.4) against leverage on the x-axis (ranging from 0.0 to 0.4). Most points are clustered near the origin, with a few points showing higher leverage and Cook's distance.



The figure displays four diagnostic plots for a linear regression model, arranged in a 2x2 grid. The model equation is $\text{stat} \sim \text{scale}(\text{tempanom_DW_tstat}) + \text{scale}(\text{prcp_DW_totastat})$.

- Standardized Residuals vs Fitted:** A scatter plot showing standardized residuals on the y-axis (ranging from -2 to 2) against fitted values on the x-axis (ranging from -0.5 to 0.5). The residuals are scattered around a horizontal dashed line at zero.
- Normal Q-Q:** A Q-Q plot showing standardized residuals on the y-axis (ranging from -2 to 2) against theoretical quantiles on the x-axis (ranging from -2 to 2). The points closely follow a solid diagonal line, indicating approximate normality.
- Scale-Location:** A scatter plot showing the square root of the absolute value of standardized residuals on the y-axis (ranging from 0.0 to 1.5) against fitted values on the x-axis (ranging from -0.5 to 0.5). The points are scattered, suggesting constant variance.
- Cook's dist vs Leverage:** A scatter plot showing Cook's distance on the y-axis (ranging from 0.0 to 0.4) against leverage on the x-axis (ranging from 0.0 to 0.4). Most points are clustered near the origin, with a few points showing higher leverage and Cook's distance.

