

Hereditary Structural Completeness of Weakly Transitive Logics

Simon Lemal

Work supervised by Nick Bezhanishvili and Tommaso Moraschini

Wednesday 11th October, 2023

Deductive systems

Let Fm be a set of propositional formulas.

Definition

A **deductive system** is a relation $\vdash \subseteq \mathcal{P}(Fm) \times Fm$ such that

- $\phi \in \Gamma$ implies $\Gamma \vdash \phi$,

Deductive systems

Let Fm be a set of propositional formulas.

Definition

A **deductive system** is a relation $\vdash \subseteq \mathcal{P}(Fm) \times Fm$ such that

- $\phi \in \Gamma$ implies $\Gamma \vdash \phi$,
- $\Gamma \vdash \phi$ and $\Delta \vdash \psi$ for all $\psi \in \Gamma$ imply $\Delta \vdash \phi$.

Deductive systems

Let Fm be a set of propositional formulas.

Definition

A **deductive system** is a relation $\vdash \subseteq \mathcal{P}(Fm) \times Fm$ such that

- $\phi \in \Gamma$ implies $\Gamma \vdash \phi$,
- $\Gamma \vdash \phi$ and $\Delta \vdash \psi$ for all $\psi \in \Gamma$ imply $\Delta \vdash \phi$.
- $\Gamma \vdash \phi$ implies that there is a finite set $\Delta \subseteq \Gamma$ such that $\Delta \vdash \phi$.

Deductive systems

Let Fm be a set of propositional formulas.

Definition

A **deductive system** is a relation $\vdash \subseteq \mathcal{P}(Fm) \times Fm$ such that

- $\phi \in \Gamma$ implies $\Gamma \vdash \phi$,
- $\Gamma \vdash \phi$ and $\Delta \vdash \psi$ for all $\psi \in \Gamma$ imply $\Delta \vdash \phi$.
- $\Gamma \vdash \phi$ implies that there is a finite set $\Delta \subseteq \Gamma$ such that $\Delta \vdash \phi$.
- for any substitution $\sigma: Fm \rightarrow Fm$, $\Gamma \vdash \phi$ implies $\sigma[\Gamma] \vdash \sigma(\phi)$.

Admissible and derivable rules

Let \vdash be a deductive system.

Definition

A **rule** is an expression of the form $\Gamma \triangleright \phi$, where Γ is finite.

Admissible and derivable rules

Let \vdash be a deductive system.

Definition

A **rule** is an expression of the form $\Gamma \triangleright \phi$, where Γ is finite.

Definition

A rule $\Gamma \triangleright \phi$ is **admissible** in \vdash if for any substitution σ , $\emptyset \vdash \sigma(\psi)$ for all $\psi \in \Gamma$ implies $\emptyset \vdash \sigma(\phi)$.

In other words, the set of theorems of \vdash is closed under $\Gamma \triangleright \phi$, and adding $\Gamma \triangleright \phi$ to \vdash does not add new theorems.

Admissible and derivable rules

Let \vdash be a deductive system.

Definition

A **rule** is an expression of the form $\Gamma \triangleright \phi$, where Γ is finite.

Definition

A rule $\Gamma \triangleright \phi$ is **admissible** in \vdash if for any substitution σ , $\emptyset \vdash \sigma(\psi)$ for all $\psi \in \Gamma$ implies $\emptyset \vdash \sigma(\phi)$.

In other words, the set of theorems of \vdash is closed under $\Gamma \triangleright \phi$, and adding $\Gamma \triangleright \phi$ to \vdash does not add new theorems.

Definition

A rule $\Gamma \triangleright \psi$ is **derivable** in \vdash if $\Gamma \vdash \phi$.

Structural completeness

Proposition

In a deductive system \vdash , every derivable rule is admissible.

Structural completeness

Proposition

In a deductive system \vdash , every derivable rule is admissible.

In CPC, the converse is true. In IPC, it is not.

Structural completeness

Proposition

In a deductive system \vdash , every derivable rule is admissible.

In CPC, the converse is true. In IPC, it is not.

Definition

A deductive system \vdash is **structurally complete (SC)** if every admissible rule in \vdash is derivable in \vdash .

Extensions of a deductive system

Definition

A deductive system \vdash' is an **extension** of a deduction system \vdash if $\vdash \subseteq \vdash'$, i.e. if $\Gamma \vdash \phi$ implies $\Gamma \vdash' \phi$.

Extensions of a deductive system

Definition

A deductive system \vdash' is an **extension** of a deduction system \vdash if $\vdash \subseteq \vdash'$, i.e. if $\Gamma \vdash \phi$ implies $\Gamma \vdash' \phi$.

Definition

An extension \vdash' of a deductive system \vdash is an **axiomatic** extension if there is a set of formulas Δ such that

$$\Gamma \vdash' \phi \quad \text{iff} \quad \Gamma, \Delta \vdash \phi.$$

Extensions of a deductive system

Definition

A deductive system \vdash' is an **extension** of a deduction system \vdash if $\vdash \subseteq \vdash'$, i.e. if $\Gamma \vdash \phi$ implies $\Gamma \vdash' \phi$.

Definition

An extension \vdash' of a deductive system \vdash is an **axiomatic** extension if there is a set of formulas Δ such that

$$\Gamma \vdash' \phi \quad \text{iff} \quad \Gamma, \Delta \vdash \phi.$$

Remark


If \vdash' is an axiomatic extension of \vdash , we can always take Δ to be the theorems of \vdash' .

Hereditary structural completeness

Theorem (Olson, Raftery, Van Alten)

Let \vdash be a deductive system, TFEA.

- Every extension of \vdash is SC.
- Every axiomatic extension of \vdash is SC.
- Every extension of \vdash is an axiomatic extension of \vdash .



Hereditary structural completeness

Theorem (Olson, Raftery, Van Alten)

Let \vdash be a deductive system, TFEA.

- Every extension of \vdash is SC.
- Every axiomatic extension of \vdash is SC.
- Every extension of \vdash is an axiomatic extension of \vdash .

Definition

A deductive system \vdash is **hereditarily structurally complete (HSC)** if it validates any of the above.

Deductive systems from modal logics

Definition

Given a NML Λ , we define a deductive system \vdash_{Λ} by $\Gamma \vdash_{\Lambda} \phi$ iff ϕ is derivable from Γ using

- the theorems of Λ ,
- Modus Ponens,
- Necessitation

Deductive systems from modal logics

Definition

Given a NML Λ , we define a deductive system \vdash_{Λ} by $\Gamma \vdash_{\Lambda} \phi$ iff ϕ is derivable from Γ using

- the theorems of Λ ,
- Modus Ponens,
- Necessitation

Remark

Given a NML Λ , the axiomatic extension of \vdash_{Λ} are those systems of the form $\vdash_{\Lambda'}$ where Λ' is a NML extending Λ .

1-transitive logics

Definition

wK4 is the modal logic

$$K + p \wedge \Box p \rightarrow \Box \Box p.$$

1-transitive logics

Definition

wK4 is the modal logic

$$K + p \wedge \Box p \rightarrow \Box \Box p.$$

Definition

A Kripke frame is a wK4-frame if its relation is 1-transitive, i.e.

$$x R y \quad \text{and} \quad y R z \quad \text{implies} \quad x R z \quad \text{or} \quad x = z$$

1-transitive logics

Definition

wK4 is the modal logic

$$K + p \wedge \Box p \rightarrow \Box\Box p.$$

Definition

A Kripke frame is a wK4-frame if its relation is 1-transitive, i.e.

$$x R y \quad \text{and} \quad y R z \quad \text{implies} \quad x R z \quad \text{or} \quad x = z$$

Definition

A modal algebra is a wK4-algebra if it validates $a \wedge \Box a \leq \Box\Box a$ for all a .

1-transitive logics

Definition

wK4 is the modal logic

$$K + p \wedge \Box p \rightarrow \Box\Box p.$$

Remark

The master modality is definable as $\Box^+ p := p \wedge \Box p$.

1-transitive logics

Definition

wK4 is the modal logic

$$K + p \wedge \Box p \rightarrow \Box \Box p.$$

Remark

The master modality is definable as $\Box^+ p := p \wedge \Box p$.

Proposition (Blok, Pigozzi)

If Λ is a 1-transitive logic, then \vdash_Λ has a *deduction detachment theorem (DDT)* witnessed by $\Box^+ p \rightarrow q$:

$$\Gamma, \phi \vdash_\Lambda \psi \quad \text{iff} \quad \Gamma \vdash_\Lambda \Box^+ \phi \rightarrow \psi.$$

Timeline

- Friedman, 1975: One hundred and two problems in mathematical logic.

Timeline

- Friedman, 1975: One hundred and two problems in mathematical logic.
- Rybakov, 1980's: Decidability of admissible rules in IPC.

Timeline

- Friedman, 1975: One hundred and two problems in mathematical logic.
- Rybakov, 1980's: Decidability of admissible rules in IPC.
- Ghilardi, Iemhoff, Jeřábek, Metcalfe, Rozière, 1990's—: In depth study of admissibility in various contexts.

Timeline

- Friedman, 1975: One hundred and two problems in mathematical logic.
- Rybakov, 1980's: Decidability of admissible rules in IPC.
- Ghilardi, Iemhoff, Jeřábek, Metcalfe, Rozière, 1990's—: In depth study of admissibility in various contexts.
- Citkin, 1978: HSC of intermediate logics (extensions of IPC).

Timeline

- Friedman, 1975: One hundred and two problems in mathematical logic.
- Rybakov, 1980's: Decidability of admissible rules in IPC.
- Ghilardi, Iemhoff, Jeřábek, Metcalfe, Rozière, 1990's—: In depth study of admissibility in various contexts.
- Citkin, 1978: HSC of intermediate logics (extensions of IPC).
- Rybakov, 1995: HSC of transitive modal logics (extensions of K4).

Timeline

- Friedman, 1975: One hundred and two problems in mathematical logic.
- Rybakov, 1980's: Decidability of admissible rules in IPC.
- Ghilardi, Iemhoff, Jeřábek, Metcalfe, Rozière, 1990's—: In depth study of admissibility in various contexts.
- Citkin, 1978: HSC of intermediate logics (extensions of IPC).
- Rybakov, 1995: HSC of transitive modal logics (extensions of K4).
- Bezhanishvili, Moraschini, 2020: new proof of Citkin's result.


Timeline

- Friedman, 1975: One hundred and two problems in mathematical logic.
- Rybakov, 1980's: Decidability of admissible rules in IPC.
- Ghilardi, Iemhoff, Jeřábek, Metcalfe, Rozière, 1990's—: In depth study of admissibility in various contexts.
- Citkin, 1978: HSC of intermediate logics (extensions of IPC).
- Rybakov, 1995: HSC of transitive modal logics (extensions of K4).
- Bezhanishvili, Moraschini, 2020: new proof of Citkin's result.
- Carr, 2022: new proof of Rybakov's (corrected) result.

Timeline

- Friedman, 1975: One hundred and two problems in mathematical logic.
- Rybakov, 1980's: Decidability of admissible rules in IPC.
- Ghilardi, Iemhoff, Jeřábek, Metcalfe, Rozière, 1990's—: In depth study of admissibility in various contexts.
- Citkin, 1978: HSC of intermediate logics (extensions of IPC).
- Rybakov, 1995: HSC of transitive modal logics (extensions of K4).
- Bezhanishvili, Moraschini, 2020: new proof of Citkin's result.
- Carr, 2022: new proof of Rybakov's (corrected) result.
- SL, 2023: HSC of weakly transitive modal logics (extension of wK4)

Method



Algebraisable logics

Definition

A deductive system \vdash is **algebraisable** if there exist

- a quasi-variety K ,
- a set of equations $\tau(x)$,
- a set of formulas $\Delta(x, y)$,

such that for all set of equation Θ , all equation $\varepsilon \approx \delta$, all set of formulas Γ and all formula ϕ , we have

- $\Gamma \vdash \phi$ iff $\tau[\Gamma] \models_K \tau(\phi)$,
- $\Theta \models_K \varepsilon \approx \delta$ iff $\Delta[\Theta] \vdash \Delta(\varepsilon, \delta)$,
- $\phi \vdash \Delta[\tau(\phi)]$ and $\Delta[\tau(\phi)] \vdash \phi$,
- $\varepsilon \approx \delta \models_K \tau[\Delta(\varepsilon, \delta)]$ and $\tau[\Delta(\varepsilon, \delta)] \models_K \varepsilon \approx \delta$.

The quasi-variety K is the **equivalent algebraic semantics (EAS)** of \vdash . It is unique when it exists.

Rules and quasi-equations

Let \vdash be a deductive system with quasi-variety K as its EAS.

Remark

A rule $\Gamma \triangleright \phi$ corresponds to a quasi-equation $\bigwedge \tau[\Gamma] \rightarrow \tau(\phi)$.

Conversely, a quasi-equation $\bigwedge \Theta \rightarrow \varepsilon \approx \delta$ corresponds to a rule $\Delta[\Theta] \triangleright \Delta(\varepsilon, \delta)$

Rules and quasi-equations

Let \vdash be a deductive system with quasi-variety K as its EAS.

Remark

A rule $\Gamma \triangleright \phi$ corresponds to a quasi-equation $\bigwedge \tau[\Gamma] \rightarrow \tau(\phi)$.

Conversely, a quasi-equation $\bigwedge \Theta \rightarrow \varepsilon \approx \delta$ corresponds to a rule $\Delta[\Theta] \triangleright \Delta(\varepsilon, \delta)$

Proposition

A rule is admissible in \vdash iff the corresponding quasi-equation is valid in the free algebra $F_K(\omega)$.

Rules and quasi-equations

Let \vdash be a deductive system with quasi-variety K as its EAS.

Remark

A rule $\Gamma \triangleright \phi$ corresponds to a quasi-equation $\bigwedge \tau[\Gamma] \rightarrow \tau(\phi)$.

Conversely, a quasi-equation $\bigwedge \Theta \rightarrow \varepsilon \approx \delta$ corresponds to a rule $\Delta[\Theta] \triangleright \Delta(\varepsilon, \delta)$

Proposition

A rule is admissible in \vdash iff the corresponding quasi-equation is valid in the free algebra $F_K(\omega)$.

Proposition

A rule is derivable in \vdash iff the corresponding quasi-equation is valid in K

Structural completeness

Let \vdash be a deductive system with quasi-variety K as its EAS.

Corollary

\vdash is SC iff every quasi-equation which is valid in $F_K(\omega)$ is valid in K .

Theorem (Prucnal, Wroński)

\vdash is SC iff $K = \mathbb{Q}(F_K(\omega))$.

Extensions and subquasi-varieties

Let \vdash be a deductive system with quasi-variety K as its EAS.

Theorem (Blok, Pigozzi)

The lattice of extensions of \vdash is dually isomorphic to the lattice of subquasi-varieties of K

Extensions and subquasi-varieties

Let \vdash be a deductive system with quasi-variety K as its EAS.

Theorem (Blok, Pigozzi)

The lattice of extensions of \vdash is dually isomorphic to the lattice of subquasi-varieties of K

Theorem (Blok, Pigozzi)

The lattice of axiomatic extensions of \vdash is dually isomorphic to the lattice of relative subvarieties of K .

Hereditary structural completeness

Let \vdash be a deductive system with **variety** K as its EAS.

Corollary

\vdash is HSC iff every subquasi-variety of K is a variety.

Hereditary structural completeness

Let \vdash be a deductive system with **variety** K as its EAS.

Corollary


\vdash is HSC iff every subquasi-variety of K is a variety.

Definition

A variety is **primitive** if all of its subquasi-varieties are varieties.

Dictionary

Logic	Algebra
Deductive system	EAS (variety)
Rules	Quasi-equations
Admissible	Valid in $F_K(\omega)$
Derivable	Valid in K
SC	$K = \mathbb{Q}(F_K(\omega))$
Extensions	Subquasi-variety
Axiomatic extensions	Subvarieties
HSC	Primitive



Wrapping up

The following problems are equivalent:

- Characterising the 1-transitive modal logics Λ such that \vdash_{Λ} is HSC.
- Characterising the axiomatic extensions of \vdash_{wK4} which are HSC.
- Characterising the primitive varieties of wK4-algebras.