


Flashback: Agents of Spatial Pattern

Considerations for marine systems


- Greater challenges than in terrestrial systems
- Temporal rates of change can be much more rapid (compare forest patch mosaics to eddies with patches of entrained phytoplankton)
- Importance of chemical seascapes and soundscapes that generate spatial and temporal patterns & are modified by human activities



Seascapes are...

1. Lagrangian, transient, dilute and patchy
2. Size-structured
3. Vertically heterogenous
4. Tractable with observing system capabilities
 1. Dynamic Ocean Management
 2. Offshore wind farm observatories

Lagrangian: Flying along, going with the flow



Dilute and Patchy:


How do aquatic food webs differ than terrestrial ones?

Table 1. Global comparison of marine and terrestrial food web attributes. Estimates are from Cohen (1994). Mean depth of life zone and related statistics are modified from Cohen's stipulated depth of 1.37 km.



Attribute	Marine	Terrestrial
Surface (% Earth)	71	29
Mean Depth of Life Zone (m)	500	50
Volume of Life Zone (10^8 km^3)	1.8	0.0075
Living Biomass (10^{15} g carbon)	2	560
Living Biomass Density ($10^7 \text{ g carbon km}^{-3}$)	1.1	7,500
Net Primary Production ($10^{15} \text{ g yr}^{-1}$)	20-44	48
Residence Time of Carbon in Biomass (yr)	0.08	11.2
Animal Biomass (% total)	10	0.01
Predator Animal Body Mass (g)	4700	140
Number of Trophic Links	High	Low

Size structured - scale-free attribute

How do aquatic food webs differ than terrestrial ones?




How do aquatic food webs differ than terrestrial ones?
Contrast a striped bass larva with an ant.
Teleost fish exhibit complex life cycles (life cycle omnivory)




Patchiness - ecotones (frontal systems)

Striped bass - salinity



Secor, D.H., E.D. Houde, and L.L. Kellogg. 2017. Estuarine retention and production of striped bass larvae: a mark recapture experiment. ICES Journal of Marine Science 74:1735-1748.



Eckman (spiral) ecocline: seasonal transience in vertical shear







Patuxent Estuary

Dissolved Oxygen



Temperature




Vertical ecotone

Bottom hypoxia in Chesapeake


Kraus, R.T., D.H. Secor, and R.L. Wingate. 2015. Testing the thermal-niche oxygen-squeeze hypothesis for estuarine striped bass. *Environmental Biology of Fishes* 98: 2083-2092

Chesapeake seascapes managed on basis of vertical structure

Designated Uses




Biological Criteria (EPA 2003)




Batiuk, R. D. Breitburg, G. Thursby, D. Secor, R. Diaz, and T. Cronin. 2009. Derivation of habitat-specific dissolved oxygen criteria for Chesapeake Bay and its tidal tributaries. J. Exp. Mar. Biol. Ecol. 381, Suppl. 1: 204-215.

Seascapes forced by wind regimes



- US NW Atlantic wind velocity regime: low in summer; high in fall
- Climate change will impact seasonal distribution of wind velocities
- Increased hurricane amplitude and frequency is predicted




Mid-Atlantic Bight (MAB)


Cold Pool: May-Sept

- Spring warming “traps” winter water
- Quiescent summer period intensifies stratification
- Summer storms disturb the stratification
- Fall storm(s) destroy the stratification


Most productive black sea bass habitats in MAB occur within the bottom “cold pool”




MARACOOS
H. Roarty, J. Keroot



Hurricane Hermine




Middle Reference Site






FV-COM Output

Study design




Telemetry Study


- Undersized fish implanted across two sites with pressure sensing tags
- Acoustic release VEMCO receivers at 800 m distances 0°, 120°, 240° N from release site




Mid-July week: hourly depths, 8 individuals




a) June




b) July




c) August




d) September




e) October




Wind Disturbance Regime




Heupel et
al. 2003



Fabrizio et
al. 2005




Bailey &
Secor 2016



Bacheler et
al. 2019



Biggs et al.
2018



Secor et al. 2019
Wiernicki et al. 2020
O'Brien and Secor
2021

Black sea bass

- In MAB storms regularly impact black sea bass habitat through recurring and permanent destratification (destruction of the cold pool)
- Incomplete evacuations
- Depressed vertical movement behaviors
- Telemetry can assay fish behaviors during, and in the wake of storms (with caveats)

Secor, D.H., F. Zhang, M. O'Brien, and M. Li. 2019. Ocean destratification and fish evacuation caused by a Mid-Atlantic Tropical Storm. *ICES Journal of Marine Science* 76:573-584.

Wiernicki, C. M.O. Brien, F. Zhang, V. Lyubchich, M. Li and D. Secor. 2020. The recurring role of storm disturbance on black sea bass (*Centropristis striata*) movement behaviors in the Mid-Atlantic Bight. *Plos One* 15(12):e0239919.


O'Brien, M.H.P. and D.H. Secor. 2021. Influence of thermal stratification and storms on acoustic telemetry detection efficiency: a year-long test in the US Southern Mid-Atlantic Bight. *Animal Biotelemetry*

Seascapes are...

1. Lagrangian, transient, dilute and patchy
2. Size-structured
3. Vertically heterogenous
4. **Tractable with observing system capabilities**
 1. **Dynamic Ocean Management**
 2. **Offshore wind farm observatories**


Seascape ecology entails a recursive approach, where observed oceanographic variables are made operational (accessible) and synoptic through wide bandwidth telecommunication.

Knowledge of ecological and oceanographic processes is incorporated into spatially explicit models that permit predictions of the dynamic distributions of marine populations and related ecosystem properties.



Manuel Hidalgo, D. Secor and H. Browman 2016. Observing and managing seascapes: linking synoptic oceanography, ecological processes, and geospatial modeling **ICES J. Mar. Sci.** 2016;icesjms.fsw079


Movement Ecology: Reduce, reconstitute, predict



<http://www.westcoast.fisheries.noaa.gov/whalewatch/>

Likelihood of Occurrence

Average



Dynamic Marine Management

Telemetry (state-space movement model)



Observed Ocean Variables



Species Distribution Model



Operational synoptic oceanography
(ocean observing x telecommunications)




Near real-time collision risk maps

Bailey, H., et al. 2009. Endangered Species Research 10:93-106.


Wind Harvest Regime

UMCES TAILWINDS Project






OWFs as flyway observatories

- **Site passage** - Eulerian
- **Flying along** - Lagrangian




Fall-Spring Migrations - Passing through (Eulerian). Flux of different striped bass populations



Movement and Habitat Selection by Migratory Fishes within the Maryland Wind Energy Area and Adjacent Reference Sites.

https://espis.boem.gov/final%20reports/BOEM_2020-030.pdf



Striped Bass Flyway

- Hudson River striped bass
- Receiver arrayed across latitudes: coming attractions for OWFs
- Flyway information on.
 - Migrations: spawning, spring, and fall
 - Destinations: summer feeding and overwintering habitats



Secor, et al. 2021. Plos One 15(11):e0242797


Key Challenge: How does one model species distributions on the basis of Lagrangian data? (Animals and encountered environments simultaneously changing in time).

Point Counterpoint: Landscape v. Seascape

Four primary drivers

1. **Abiotic conditions:** Climate, topography, landform, soils, flow, depth/light, salinity, temperature, nutrients, turbidity
2. **Biotic processes:** species interactions, demographics including dispersal
3. **Disturbance and succession**
4. **Land use and history**

Landscape Ecology: Emergent patterns from underlying ecological processes



Take home points: agents of spatial ecology (terrestrial)

1. Physical template (abiotic environment) is the arena in which biotic processes and disturbance regimes interact.
2. Landform (terrain) modifies climate to produce microclimate variations
3. Historical contingencies and legacies are critical to understanding current landscape patterns
4. Spatial hierarchy allows tractable analysis of spatial ecology: local, landscape, regional, and global scales.

Take home points: agents of spatial ecology (marine/aquatic)

1. Physical template **transience** (abiotic environment) is the arena in which biotic processes and disturbance regimes interact.
2. ~~Landform (terrain)~~ **Wind and ocean current regimes** modify climate to produce **mesoscale** ~~microclimate~~ variations
3. Historical contingencies and legacies are **less** critical to understanding current patterns
4. ~~Spatial hierarchy allows tractable analysis~~ **is confounded by transience and vertical heterogeneity** at local, ~~landscape~~ **seascape**, regional, and global scales.
5. **Distribution of biota is dilute, size-structured and characterized by high patchiness.**

Point Counterpoint

Contribution to the Themed Section: '*Seascape Ecology*'

Food for Thought

Seascapes are not landscapes: an analysis performed using Bernhard Riemann's rules

John Pilling Manderson*


Comment

Seascapes are landscapes after all; Comment on Manderson (2016): Seascapes are not landscapes: an analysis performed using Bernhard Riemann's rules. ICES Journal of Marine Science, 73:1831–1838

Susan S. Bell^{1,*} and Bradley T. Furman²

Reply

Response to Bell and Furman (2017): Seascapes are landscapes after all; Comment on Manderson (2016): Seascapes are not landscapes: an analysis performed using Bernhard Riemann's rules: ICES Journal of Marine Science, 73:1831–1838



Discussion Questions

The point-counterpoint on the overlap between landscape and seascape ecology seems to center on dimension and scale. Describe how types of dimension and types of scale define key points of difference between Manderson's and Bell and Furmans' views.

Would subdividing seascape ecology into “demersal (seabed) seascape ecology” and “pelagic seascape ecology” resolve issues raised in the point-counterpoint?

Landscape ecology emphasizes emergent properties and synthesis. Would the bioenergetics (cellular) approach adopted by Manderson to define seascapes offer improved perspectives in landscape ecology?

Are applications in movement ecology similar between landscapes and seascapes?