PRACTITIONER PROTOCOL · PARAMEDICAL

Areola Restoration

Your device and pigment maker's directions come first. Follow your state and local body art rules.

CLEARANCE

Written surgeon or physician clearance on file and dated. Typically 6+ months after final surgery or radiation, fully healed. Note implant, expander or flap type.

PRIVACY

Private room, door closed, sign on the door. Gown open at front. Drape so only the working side is exposed. Ask before you touch.

CONSULT

Intake, consent and photo choices signed. Torso-only photos, never the face. Patch test offered, or waiver signed.

SET UP FIRST ☐ Drape & gown ☐ Barrier film ☐ Single-use cartridges ☐ Pigment cups ☐ Sterile pads ☐ Surgical marker ☐ Ruler or calipers ☐ Sharps container

#	STEP	TIME	HOW
1	**Review & inspect**	5 min	Confirm clearance and health changes. Inspect under good light. Stop for any open, weeping, red or unhealed tissue, or a new lump or change; refer to the surgeon.
2	**Photograph**	2 min	Torso only, same light and angle every visit. Never include the face.
3	**Measure & map**	10–15 min	Measure from the midline and the other side if present. Draw the shape with the client standing and seated. Client checks in a mirror and approves.
4	**Color match**	5–10 min	Match the other areola, or choose a tone that suits the skin. Swatch on skin. Scar and grafted skin often heal lighter; record the mix.
5	**Prep & numb**	Per label	Cleanse and prep the skin. Topical numbing only as your state allows; sensation may already be reduced.
6	**Pigment**	45–90 min	Light hand, shallow passes. Go extra gently over scars and thin skin; reconstructed tissue can be fragile. Build color in layers, not depth.
7	**Check in**	Throughout	Ask for comfort 0–10 and offer breaks. Stop for heavy bleeding, blanching or skin that looks stressed.
8	**Clean & cover**	3 min	Wipe with sterile saline. Thin layer of balm and a sterile non-stick pad. Client puts on a soft, clean bra.
9	**Educate & rebook**	5 min	Walk through aftercare and daily checks. Book the next session 6–8 weeks out.

IF SOMETHING GOES WRONG

- Heavy bleeding: firm pressure with sterile gauze; stop the session.
- Hives or swelling: stop. Call 911 if breathing is affected.
- Implant area pain or swelling: stop and refer to the surgeon same day.
- Signs of infection later: refer to the client's physician promptly.
- Always write an incident report and check in with the client within 24–48 hours.

RECORD FOR THIS SESSION

Clearance from · date	Side(s) · size mm	Pigment brand · color	Lot # · expiry
Needle config · machine	**Numbing · time on**	**Tissue notes · scars**	**Comfort 0–10 · initials**