PRACTITIONER PROTOCOL

Lip Blush

Follow your state and local body-art rules and your pigment and numbing labels. Times are typical.

SCREEN

Photo ID (18+). No cold sore in 2 weeks. Lip filler at least 4 weeks ago.

COLD SORE HISTORY

Client asked their physician about preventive medication. Chart it.

PATCH TEST

Offered 24 to 48 hours ahead, 7-day final check. Waiver if declined.

SET UP FIRST ☐ BBP precautions, PPE on ☐ Barrier film on bed, lamp, machine ☐ Sharps container in reach ☐ Sealed single-use needle cartridge ☐ Pigment decanted to a single-use cup ☐ Brand, color, lot, expiry recorded ☐ Sterile gauze & cool compress ☐ Mirror, lip liner, timer

#	STEP	TIME	HOW
1	**Consult & screen**	10–15 min	Confirm health changes. Check for cracks or a cold sore. Defer if present.
2	**Consent & photos**	5 min	Consent signed. Photos front, sides and close-up, lips relaxed.
3	**Mapping & approval**	15–20 min	Outline lip line for symmetry. Client approves in a mirror and initials.
4	**Numbing**	20–30 min	Topical only as state rules and label allow. Record product, times.
5	**Pigment selection**	5 min	Lips heal cooler, lighter. Dark or cool lips: warm corrector first.
6	**Prep**	2 min	Remove numbing, antiseptic prep. Protect the mouth corners.
7	**Lip line & fill**	60–90 min	Define the border, then shade in soft, even, superficial layers. Avoid blowout past the border.
8	**Wipe & assess**	Ongoing	Sterile gauze. Check symmetry and even color, lips relaxed.
9	**Secondary numbing**	If used	Only on opened skin, only if state rules and the label permit.
10	**After photos**	5 min	Client mirror check. Same angles and light as the before photos.
11	**Aftercare & touch-up**	5 min	Thin balm, cool compress, review sheet and card. Touch-up 6 to 8 weeks.

ENDPOINT AND SAFETY

- Soft, even color; pinpoint bleeding at most
- Dark lips: corrector first, 2 to 3 sessions
- Lips swell fast: work efficiently
- Blister or tingling during visit: stop, refer

IF SOMETHING GOES WRONG

- Hives, wheezing or swelling: stop, remove product. Call 911 if breathing is affected.
- Fainting or dizziness: stop, lay the client flat, raise legs, monitor. 911 if slow to recover.
- Needlestick or blood exposure: follow your written exposure control plan.
- Heavy bleeding: stop, firm pressure with sterile gauze. End the session if it doesn't settle.
- Severe or one-sided swelling: cool compress; if it worsens, refer to a physician.
- Blisters later (possible cold sore): refer the client to their physician.

Always write an incident report and check in with the client within 24 to 48 hours.

RECORD FOR THIS SESSION

Pigment · corrector	Lot · expiry	Needle config · settings	Numbing · times applied
Passes · lip response	**Bleeding · swelling**	**Retention notes**	**Touch-up date · initials**