

Faddeev–Skyrme soliton energies from controlled lattice extrapolation

Paul Vodrazka

Independent researcher

(Dated: August 14, 2026)

Abstract

We recompute the low-charge Faddeev–Skyrme soliton spectrum with both lattice systematics under explicit control, and find a disagreement with the accepted charge-one energy that we are unable to remove. Two observations organise the computation. First, a discretisation whose derivative symbol vanishes at the Nyquist frequency gives the discrete energy exact flat directions: a checkerboard mode costs nothing, the Hopf charge unwinds under gradient flow, and the energy falls below the Vakulenko–Kapitanskii bound. Symmetrising the energy over one-sided differences removes the null mode while keeping the analytic variation the exact gradient of the discrete energy. Second, finite spacing and finite volume bias the energy in *opposite* directions, so neither can be assessed alone; fitting $E = E_\infty - ah^2 + cL^{-p}$ to nine runs separates them. The result is $E_1/c_0 = 1.231 \pm 0.003$ against the published 1.204, an excess of 2.2%. Three independent checks support it rather than the alternative that our lattice is at fault: at matched spacing $h = 0.100$ we obtain 1.2257, a 1.8% gap of which our finite box can account for at most 0.7%; a fourth-order accurate symmetrisation, whose error has the opposite sign, extrapolates from above to 1.2371 against the second-order scheme’s 1.2354 from below, agreeing to 0.14%; and the identical procedure applied to the Skyrme model, where the charge-one energy follows from an ordinary differential equation, recovers that exact value to 0.03% from runs whose individual errors reach 2%. In the same frame sector we obtain $B = 1$ –4 energies within 0.6–0.8% of the tabulated values and binding energies per baryon within 0.15 percentage points. We reproduce the published charge-one to charge-four configuration types, the sub-linear scaling exponent 0.765 against $3/4$, and binding against fission in every sector, so the discrepancy is confined to the absolute normalisation and not to the structure of the spectrum.

I. INTRODUCTION

The Faddeev–Skyrme model [1] is the simplest field theory in three dimensions whose solitons are classified by a linking number rather than a winding number. For a unit three-vector field $\mathbf{n} : \mathbb{R}^3 \rightarrow S^2$ with $\mathbf{n} \rightarrow \mathbf{n}_0$ at infinity, configurations fall into sectors labelled by the Hopf invariant $Q_H \in \pi_3(S^2) = \mathbb{Z}$, and the minimal energy in each sector is a well-posed numerical question that has been studied for two decades [2–7]. The charge-one energy is the reference against which all other sectors are quoted, and it is usually taken to be 1.204

in the normalisation fixed by Ward’s conjectured bound.

This paper recomputes that number with the two lattice systematics separated and extrapolated, and reports that we cannot reproduce it. Our value is 2.2% higher. Because a discrepancy of this size is more often a defect of the computation than of the literature, most of what follows is an attempt to break our own result.

Two things make the exercise worth reporting. The first is a discretisation pitfall that is easy to fall into and whose symptom is dramatic—the topological charge simply dissolves—but which we have not seen stated in these terms. The second is that the two lattice systematics push in opposite directions, so a study that varies only the spacing, or only the box, will appear better converged than it is. Section III treats the first, Section IV the second.

Section V is the control. Applied to the Skyrme model, where the charge-one energy is available from an ordinary differential equation, the same code and the same extrapolation recover the exact answer to three parts in ten thousand. Section VI reports the $Q_H \leq 4$ spectrum, which agrees with the published one in every structural respect. All source is available [8].

II. MODEL, BOUND AND NORMALISATION

The static energy is

$$E[\mathbf{n}] = \int d^3x \left[c_2 \partial_i \mathbf{n} \cdot \partial_i \mathbf{n} + \frac{c_4}{2} (\partial_i \mathbf{n} \times \partial_j \mathbf{n}) \cdot (\partial_i \mathbf{n} \times \partial_j \mathbf{n}) \right], \quad (1)$$

and we set $c_2 = c_4 = 1$ throughout, so that lengths and energies are in model units. Under $\mathbf{n}(x) \rightarrow \mathbf{n}(x/\mu)$ the two terms scale as $E(\mu) = \mu E_2 + \mu^{-1} E_4$, so every stationary point satisfies the virial identity $E_2 = E_4$; with either constant absent the soliton collapses or spreads, and Derrick’s theorem is satisfied rather than evaded.

Finite-energy configurations obey the bound of Vakulenko and Kapitanskii [9],

$$E \geq c_0 |Q_H|^{3/4}, \quad (2)$$

whose proof does not fix the constant. Ward [6] conjectured the sharp value, which in the normalisation of (1) reads

$$c_0 = 32\pi^2\sqrt{2} = 446.65. \quad (3)$$

Reference [7] divides its energies by exactly this factor and states the conjecture with unit coefficient; its integrand is identical to (1), so its tabulated energies are directly comparable with ours with no conversion. We have verified the identity $\frac{1}{2}(\partial_i \mathbf{n} \times \partial_j \mathbf{n})^2 = \frac{1}{2}[(\partial_i \mathbf{n} \cdot \partial_i \mathbf{n})^2 - (\partial_i \mathbf{n} \cdot \partial_j \mathbf{n})^2]$ symbolically, so the two normalisations agree term by term.

Because the exponent $3/4$ is sub-linear, $E_Q < |Q|E_1$ is permitted and multi-charge solitons may be bound against fission; whether they are is settled in Sec. VI. For the general theory we refer to Manton and Sutcliffe [10].

III. LATTICE CONSTRUCTION, AND A NULL MODE THAT DESTROYS TOPOLOGY

We minimise (1) on a periodic cubic lattice. One implementation choice carries more than numerical significance.

With central differences *of any order*, the Fourier symbol of the derivative operator vanishes at the Nyquist wavenumber. For the fourth-order antisymmetric stencil, setting $f_{i\pm 1} \rightarrow -f_i$ and $f_{i\pm 2} \rightarrow f_i$ gives identically zero. A checkerboard mode therefore costs *no energy*. The discrete energy acquires exact flat directions, and gradient flow carries any Hopfion monotonically to the vacuum: in our first runs Q_H fell from 1 to 0 while E fell far below the bound $c_0|Q_H|^{3/4}$, which is supposed to be inviolable. The topological protection is lost not because the physics permits it but because the discretisation has a zero-cost path out of the sector.

The cure is a scheme whose symbol vanishes only at $k = 0$. We use one-sided differences and symmetrise the *energy* rather than the derivative,

$$E = \frac{1}{2}[E(D^+ \mathbf{n}) + E(D^- \mathbf{n})], \quad (D^\pm f)_i = \pm \frac{f_{i\pm 1} - f_i}{h}, \quad (4)$$

which has no null mode and, because $(D^\pm)^\top = -D^\mp$ holds exactly on a periodic lattice, makes the analytic variation the *exact* gradient of the discrete energy. We verify this against finite differences of the energy density to a maximum relative error of 5×10^{-7} .

The symmetrisation also fixes the form of the extrapolation used below. Expanding $D_i^\pm = \partial_i \pm \frac{h}{2}\partial_i^2 + \frac{h^2}{6}\partial_i^3 + \dots$ and averaging removes the $O(h)$ term, leaving

$$\frac{1}{2} \int [|D^+ \mathbf{n}|^2 + |D^- \mathbf{n}|^2] = \int |\partial \mathbf{n}|^2 - \frac{h^2}{12} \int |\partial^2 \mathbf{n}|^2 + O(h^4), \quad (5)$$

after integrating $\int \partial \mathbf{n} \cdot \partial^3 \mathbf{n} = -\int |\partial^2 \mathbf{n}|^2$ by parts. The discrete energy is therefore below the continuum one, by $O(h^2)$ – both the sign and the order seen in the fits of Sec. IV.

A fourth-order variant. Nothing forces the one-sided differences in (4) to be first-order accurate. Built from third-order one-sided differences,

$$(D_3^+ f)_i = \frac{-11f_i + 18f_{i+1} - 9f_{i+2} + 2f_{i+3}}{6h}, \quad (6)$$

and its reflection, the leading errors are equal and opposite, $D_3^\pm = \partial \pm \frac{h^3}{4}\partial^4 + \dots$, so they cancel in the symmetrised energy and leave an $O(h^4)$ scheme. Its Nyquist symbol is $-20/3h$ rather than zero, so the null mode is still absent, and $(D_3^+)^\top = -D_3^-$ still holds, so the variation remains exact. We refer to the two schemes as $O(h^2)$ and $O(h^4)$; crucially their errors have opposite sign, so together they bracket the continuum limit.

The Hopf charge is evaluated by reconstructing $\mathbf{A}$ spectrally in Coulomb gauge from $\mathbf{B}$, with $\|\nabla \times \mathbf{A} - \mathbf{B}\|/\|\mathbf{B}\| < 10^{-4}$. Seeds of known charge come from composing a degree- d hedgehog with the Hopf projection (giving $Q_H = d$, since the Chern–Simons three-form pulls back with the degree) and from the toroidal $A_{n,m}$ ansatz [3, 5] (giving $Q_H = nm$).

IV. THE CHARGE-ONE ENERGY

Two systematics act in opposite directions, and both must be removed before any comparison is meaningful.

Finite spacing lowers the energy, by (5). Repeating the $Q_H = 1$ relaxation at fixed box $L = 12$, with every lattice initialised by resampling the converged solution so that all points share the same convergence quality, gives Table I. The measured charge approaches the integer and the virial ratio approaches unity, both as they must.

Finite volume raises it. The director is massless, so the soliton has power-law tails and interacts with its own periodic images. Holding $h = 1/6$ and enlarging the box gives $E = 540.50, 538.77, 538.40$ for $L = 12, 16, 20$, with the energy fraction within six cells of the boundary falling from 4.0×10^{-3} to 1.3×10^{-4} .

Neither series alone is a convergence test, because the two effects partially cancel. Fitting them together,

$$\frac{E}{c_0} = E_\infty - ah^2 + cL^{-p}, \quad (7)$$

TABLE I. Lattice-spacing dependence of the $Q_H = 1$ soliton at $L = 12$.

N	h	E	E/c_0	Q_H	E_2/E_4
56	0.2143	532.28	1.1917	0.9833	0.823
72	0.1667	540.42	1.2100	0.9951	0.883
88	0.1364	544.12	1.2182	0.9980	0.904
96	0.1250	545.29	1.2208	0.9986	0.908
104	0.1154	546.18	1.2228	0.9990	0.910
120	0.1000	547.45	1.2257	0.9994	0.911

to all nine runs gives an h^2 coefficient $a = 0.94$, a box correction of $+0.4$ to $+0.7\%$ at $L = 12$, and

$$E_1 = 550 \pm 2, \quad E_1/c_0 = 1.231 \pm 0.003, \quad (8)$$

with residuals of 4×10^{-4} throughout. The quoted spread is the variation of the intercept as the box exponent is fixed anywhere between $p = 2$ and $p = 6$, widened to cover a second fit to an independently relaxed set of runs. The extrapolation reaches only 0.4% beyond the finest lattice computed.

Against the published 1.204 this is an excess of 2.2% . Three checks bear on whether the fault is ours.

Matched resolution. Our finest lattice uses $h = 0.100$, exactly the spacing of Ref. [7], and returns 1.2257 with no extrapolation at all. At equal spacing the gap is 1.8% , of which our finite box can account for at most 0.7% .

An independent discretisation. The remaining worry is that our energy is only $O(h^2)$ accurate while Ref. [7] uses fourth-order derivatives, so that the extrapolation rather than the data is at fault. Repeating the whole calculation with the $O(h^4)$ scheme of Sec. III, whose error enters with the opposite sign, gives Table II: the two schemes approach from opposite sides and agree with each other to 0.14% . Two discretisations of different order converging from opposite directions do not conspire.

A control with a known answer. This is Sec. V.

TABLE II. The two symmetrised schemes at $L = 12$. The $O(h^2)$ scheme approaches from below and the $O(h^4)$ scheme from above; the extrapolations agree to 0.14%. Values are E/c_0 before the box correction of Eq. (7).

h	$O(h^2)$	E_2/E_4	$O(h^4)$	E_2/E_4
0.1875	1.2026	0.862	1.2646	1.071
0.1500	1.2147	0.902	1.2492	0.975
0.1250	1.2207	0.920	1.2420	0.943
$\rightarrow 0$	1.2354		1.2371	

V. A CONTROL: THE FRAME SECTOR

The Faddeev–Skyrme model has no closed-form solution, so within it a lattice can only be compared against other lattices. The Skyrme model does have one, and it is close enough to be a genuine control: the same energy functional, the same code, the same boundary conditions and the same extrapolation, with only the target changed from S^2 to S^3 .

For $\varphi : \mathbb{R}^3 \rightarrow S^3$ the energy (1) is the Skyrme model [11] in Manton’s strain form [12], configurations are classified by $\pi_3(S^3) = \mathbb{Z}$ with baryon number

$$B = \frac{1}{2\pi^2} \int \det[\varphi, \partial_1\varphi, \partial_2\varphi, \partial_3\varphi] d^3x, \quad (9)$$

and the topological bound follows from two applications of AM–GM to the principal stretches, $E \geq 12\pi^2\sqrt{c_2c_4}|B|$. For $B = 1$ the minimiser is spherically symmetric, so with $\varphi = (\cos f, \sin f \hat{r})$ the energy reduces to

$$E = 4\pi \int_0^\infty \left[f'^2 r^2 + 2 \sin^2 f (1 + f'^2) + \frac{\sin^4 f}{r^2} \right] dr, \quad (10)$$

whose Euler–Lagrange equation we solve as a two-point boundary value problem with $f(0) = \pi$, $f(\infty) = 0$, obtaining

$$E/12\pi^2 = 1.23145, \quad (11)$$

consistent with the value 1.232 standard since Ref. [13]. Because we compute (11) ourselves it tests the lattice rather than our reading of a table.

Relaxing on the lattice at two box sizes and three spacings each, and fitting Eq. (7) to all six runs, gives

$$E_\infty/12\pi^2 = 1.2311, \quad (12)$$

TABLE III. Frame-sector solitons at $L = 12$, $N = 96$, from rational-map seeds of the indicated symmetry. Literature values are Ref. [14]. Binding is $1 - E_B/(BE_1)$.

B	symmetry	$E/12\pi^2 B$	Ref. [14]	binding	lit. binding
1	spherical	1.2231	1.2322	—	—
2	axial	1.1712	1.1791	4.25%	4.31%
3	tetrahedral	1.1396	1.1462	6.83%	6.98%
4	cubic	1.1110	1.1201	9.17%	9.10%

with an rms residual of 6.5×10^{-5} and a leave-one-out spread of ± 0.0008 . This is 0.03% from the exact answer (11), obtained from runs whose individual errors reach 2%. The box exponent comes out $p \simeq 2.2$, and the $L = 8$ series overshoots the exact value while still rising with resolution—direct confirmation that a periodic box biases upward, an effect also noted in the Skyrme literature [14].

A procedure accurate to three parts in ten thousand on a problem with a known answer is unlikely to be wrong by 2% on a neighbouring one. This is our main reason for reporting the discrepancy of Sec. IV as a discrepancy.

The same machinery reproduces the multi-baryon sector. Table III gives $B = 1$ –4 from symmetric rational-map seeds [15]; the energies sit 0.6–0.8% below the tabulated values [14], uniformly, so the binding energies—being ratios, in which the systematic cancels—come out within 0.15 percentage points.

VI. THE CHARGE-ONE TO CHARGE-FOUR SPECTRUM

Table IV gives the Hopf spectrum on a common lattice, and Table V compares it with Ref. [7]; Fig. 1 shows the relaxed configurations and Fig. 2 the energies against the bound and the fission threshold.

Four structural results, none of them imposed.

Every multi-charge sector is bound against fission, $E_Q < QE_1$ by 18.5%, 23.4% and 28.0% for $Q = 2, 3, 4$.

The sub-linear exponent is recovered. A power-law fit gives $E_Q \propto |Q_H|^{0.765}$ against the Vakulenko–Kapitanskii 3/4; the same fit to the published energies gives 0.760.




FIG. 1. Mid-plane sections of the relaxed Hopfions. Top: energy density. Bottom: Hopf density $\mathbf{A} \cdot \mathbf{B}$ on a symmetric colour scale, so the vacuum is neutral in every panel; the $Q_H = 4$ solution was found with negative charge, which is equivalent to the positive one by a spatial reflection. The $Q_H = 2, 3$ ground states are toroidal; the $Q_H = 4$ ground state is the more compact linked $A_{2,2}$ configuration.

TABLE IV. Relaxed Hopfion spectrum ($N = 80$, $L = 12$, $h = 0.15$). Several independent seeds were relaxed in each sector and the lowest is quoted.

Q_H	seed	E	$E/c_0 Q ^{3/4}$	R_{rms}	E_2/E_4	$E_Q/(QE_1)$
1	hedgehog $d=1$	542.65	1.215	1.705	0.887	1.000
2	hedgehog $d=2$	884.15	1.177	2.089	0.950	0.815
3	hedgehog $d=3$	1247.30	1.225	2.489	0.955	0.766
4	$A_{2,2}$ (linked)	1563.24	1.237	2.331	0.956	0.720

The $Q_H = 4$ ground state is linked. The $A_{2,2}$ seed relaxes to 1563.2, below both the hedgehog $d=4$ (1621.6) and $A_{4,1}$ (1621.8), and it is more compact than the $Q_H = 3$ soliton ($R_{\text{rms}} = 2.33$ against 2.49). Linking, not size, lowers the energy.

Seeds that should agree do. $A_{2,1} \rightarrow 884.20$ against the hedgehog's 884.15, and $A_{3,1} \rightarrow 1248.34$ against 1247.30: agreement at the 10^{-3} level from entirely different initial data. The $A_{1,m}$ family stalls higher (1006.9, 1478.9, 1898.9), which we report as upper stationary




FIG. 2. Left: computed energies against the Ward bound $c_0 Q^{3/4}$ and the fission threshold QE_1 . Right: $E_Q/(QE_1)$; values below unity indicate binding against fission.

TABLE V. Comparison with Ref. [7] in the common normalisation E/c_0 , on the lattice of Table IV. The ground-state configuration type is reproduced in every sector.

Q_H	this work	Ref. [7]	difference	type (ours / published)
1	1.215	1.204	+0.9%	$A_{1,1}$ / $A_{1,1}$
2	1.980	1.967	+0.6%	$A_{2,1}$ / $A_{2,1}$
3	2.793	2.754	+1.4%	$A_{3,1}$ / $\tilde{A}_{3,1}$
4	3.500	3.445	+1.6%	$A_{2,2}$ / $A_{2,2}$

configurations without claiming they are local minima.

The excess over Ward's bound is 21.5, 17.7, 22.5 and 23.7% here against 20.4, 17.0, 20.8 and 21.8% in Ref. [7]: not only the overall $\sim 20\%$ level but the characteristic dip at $Q_H = 2$ is reproduced. The structure of the spectrum is therefore not in question; the disagreement is confined to the absolute normalisation.

VII. DISCUSSION

We are unable to reduce the charge-one discrepancy below 2%, and we have not found a defect in our own calculation that would produce it. The candidate explanations we can test have been tested: the spacing, by matching Ref. [7] exactly; the order of the scheme, by an

independent $O(h^4)$ discretisation approaching from the other side; the box, by an explicit two-parameter fit; and the whole procedure, by a control with an exact answer.

What we cannot test from here is the effect of fixed rather than periodic boundaries, which is the one methodological difference remaining. Reference [7] truncates the field to the vacuum at the boundary of a box of side 15.1; for a massless field with power-law tails this removes gradient energy that a periodic box retains, and the sign is right to explain part of a gap. Against that, Ref. [7] itself argues the opposite way, noting that its grid “appear[s] to underestimate the energy by an amount of the order of 1%” and quoting $E \simeq 1.21$ from higher-resolution studies. Our result would put the underestimate nearer 2%.

The cleanest resolution would be an independent computation with fixed boundaries and an $h \rightarrow 0$ extrapolation, which we encourage; the code needed to reproduce everything here, including both schemes, is available [8]. Until then we report $E_1/c_0 = 1.231 \pm 0.003$ as our result and the difference from 1.204 as open.

ACKNOWLEDGMENTS

Numerical work used NumPy, SciPy, SymPy, Matplotlib and CuPy. The author used an AI assistant (GitHub Copilot) for code development, numerical analysis and manuscript preparation; all claims, and responsibility for them, are the author’s.

Appendix A: Numerical methods

Discretisation. Periodic cubic lattice; the energy is the symmetrisation of Eq. (4) in either the $O(h^2)$ or the $O(h^4)$ form. Because $(D^\pm)^\top = -D^\mp$, the analytic variation is the exact gradient of the discrete energy, verified against finite differences of the energy density to 5×10^{-7} .

Minimisation. Adaptive damped flow: the step is capped so that no site rotates by more than δ radians per iteration; a heavy-ball term with energy arrest accelerates convergence; and the momentum is projected back onto the tangent space each step—omitting this projection allows a spurious radial component to accumulate and drives the field out of its sector. Every hundred steps the exact Derrick dilation $\mu = \sqrt{E_4/E_2}$ is applied and accepted only if it lowers the energy *and* does not push the configuration into the boundary collar; the

second condition matters, because unbounded dilation samples only the core and returns a uniform field, whose energy is zero and which the energy test alone would therefore accept.

Convergence. Stationarity is checked by continued relaxation: the stored $Q_H = 1$ field moves by 2×10^{-4} in relative energy over 1800 further steps, and the residual virial deficit is a property of the discrete functional rather than of incomplete relaxation, since it persists while the energy is stationary and falls with both h and L .

Topological charge. $\mathbf{B}$ from fourth-order central differences, $\mathbf{A}$ from $\hat{\mathbf{A}} = i\mathbf{k} \times \hat{\mathbf{B}}/k^2$ in Coulomb gauge; the residual $\|\nabla \times \mathbf{A} - \mathbf{B}\|/\|\mathbf{B}\|$ is below 10^{-4} at $N = 96$. The baryon number of the frame sector is the determinant of $[\varphi, \partial_1\varphi, \partial_2\varphi, \partial_3\varphi]$ evaluated at every site, and is integral to 5×10^{-4} at the finest lattice.

Appendix B: Reproducibility

The supplement [8] comprises a library and one driver per result, with `verify_claims.py` re-deriving every quoted quantity from the stored fields rather than from intermediate summaries, and `verify_manuscript.py` checking that the numbers printed in this paper are the numbers those drivers produced. The lattice kernels are written against a swappable array backend and run unchanged on a CPU or, with `--gpu`, on a CUDA device; double precision is used throughout, since the kernels are limited by memory bandwidth rather than by floating-point throughput. The GPU path agrees with the CPU path to rounding for every energy, charge and radius reported here, except in the Derrick rescaling step, where the host and device cubic-spline interpolators differ at the 10^{-7} level; because that step is accepted only when it lowers the energy, this perturbs the relaxation path but not the minimum reached. All numbers quoted in this paper were produced on the CPU path.

-
- [1] L. Faddeev and A. J. Niemi, [Phys. Rev. Lett. **82**, 1624 \(1999\)](#).
 - [2] L. Faddeev and A. J. Niemi, [Nature **387**, 58 \(1997\)](#).
 - [3] J. Gladikowski and M. Hellmund, [Phys. Rev. D **56**, 5194 \(1997\)](#).
 - [4] R. A. Battye and P. M. Sutcliffe, [Phys. Rev. Lett. **81**, 4798 \(1998\)](#).
 - [5] J. Hietarinta and P. Salo, [Phys. Lett. B **451**, 60 \(1999\)](#).
 - [6] R. S. Ward, [Nonlinearity **12**, 241 \(1999\)](#).

- [7] P. Sutcliffe, [Proc. R. Soc. A](#) **463**, 3001 (2007).
- [8] P. Vodrazka, Computational supplement: Topological solitons of a hyperelastic vacuum, <https://github.com/slickdextric/faddeev-skyrme-solitons> (2026), library, drivers and manuscript source.
- [9] A. F. Vakulenko and L. V. Kapitanskii, *Sov. Phys. Dokl.* **24**, 433 (1979).
- [10] N. Manton and P. Sutcliffe, *Topological Solitons* (Cambridge University Press, 2004).
- [11] T. H. R. Skyrme, [Proc. R. Soc. A](#) **260**, 127 (1961).
- [12] N. S. Manton, [Commun. Math. Phys.](#) **111**, 469 (1987).
- [13] G. S. Adkins, C. R. Nappi, and E. Witten, [Nucl. Phys. B](#) **228**, 552 (1983).
- [14] R. A. Battye and P. M. Sutcliffe, [Rev. Math. Phys.](#) **14**, 29 (2002).
- [15] C. J. Houghton, N. S. Manton, and P. M. Sutcliffe, [Nucl. Phys. B](#) **510**, 507 (1998).