Introduction to the Fractional Distribution Families

Stephen Horng-Twu Lihn
stevelihn@gmail.com
Version: Draft 2026-10

ORFE, Princeton University

Dedicated to Professor John M. Mulvey for his 80th birthday on Oct 27th, 2026.

(as of October 26, 2026)

Contents

Chapter 1. Introduction 1
1.1. Motivation 1
1.2. Elements of Probability Distributions 2
1.3. Organization of This Book 3
1.4. Unresolved Issues 4

Part 1. Mathematical Functions 5

Chapter 2. Mellin Transform 7
2.1. One-Sided Distribution 8
2.2. Two-Sided Distribution 9
2.3. Ramanujan's Master Theorem 10

Chapter 3. The Wright Function and The Inverse Stable Law 13
3.1. Definition 13
3.2. Classic Results 13
3.3. The M-Wright Functions 14
3.4. The Inverse Stable Law 15
3.5. The Fractional Gamma-Star Function 16
3.6. The Elasticity Operator 19
3.7. The Elasticity of the M-Wright Functions 19
3.8. Numerical Methods of the M-Wright Functions 21

Chapter 4. The Alpha-Stable Distribution - Review 25
4.1. Classic Result 25
4.2. Extremal Distributions 26
4.3. Ratio Distribution Approach 26
4.4. SaS 28

Chapter 5. Fractional Hypergeometric Functions 29
5.1. Fractional Confluent Hypergeometric Function 29
5.2. Fractional Gauss Hypergeometric Function 31

Part 2. One-Sided Distributions 35

Chapter 6. FG: Fractional Gamma Distribution and Tilted Stable Law 37
6.1. Definition 37
6.2. Determination of C 37
6.3. FG Subsumes Generalized Gamma Distribution 38
6.4. Mellin Transform 38
6.5. The Inverse Tilted Stable Law 38

6.6. The Lognormal Limit 40
6.7. CDF and Fractional Incomplete Gamma Function 41
6.8. Inverse Expression of Several Fractional Distributions 43
6.9. Alternative Definition 43

Chapter 7. Fractional Chi Distributions 45
7.1. Introduction to Fractional Chi Distribution 45
7.2. FCM: Fractional Chi-Mean Distribution 45
7.3. FCM Moments 46
7.4. FCM Reflection Formula and Negative k 47
7.5. FCM: Unified Algebra of Cumulants of the Log Variable 47
7.6. FCM: Lognormal Asymptotics and Density by Log Inversion 49
7.7. FCM Moment Coordinates 55
7.8. FCM2: Fractional Chi-Squared-Mean Distribution 61
7.9. FCM2 Mellin Transform 62
7.10. FCM2 Moments 62
7.11. FCM2 Increment of k 63
7.12. Sum of Two Chi-Squares with Correlation 63

Chapter 8. Fractional F Distribution 65
8.1. Definition 65
8.2. PDF at Zero 66
8.3. Mellin Transform 66
8.4. Moments 66

Chapter 9. The Framework of Continuous Gaussian Mixture 69
9.1. The Inverse Chi Distribution 69
9.2. The Characteristic Distributions 70
9.3. Summary of Gaussian Mixture 71
9.4. FCM Extensions 71
9.5. Application of FCM Variants 73

Part 3. Two-Sided Univariate Distributions 77

Chapter 10. SN: The Skew-Normal Distribution - Review 79
10.1. Definition 79
10.2. The Location-Scale Family 80
10.3. Invariant Quantities 80
10.4. Mellin Transform 81
10.5. Moments 82

Chapter 11. GAS: Generalized Alpha-Stable Distribution (Experimental) 85
11.1. Definition 85
11.2. Limitation 86
11.3. Workaround 87
11.4. Moments 89

Chapter 12. GAS-SN: Generalized Alpha-Stable Distribution with Skew-Normal 91
12.1. Definition 91
12.2. The Location-Scale Family 92
12.3. Mellin Transform 92

12.4. Moments 92
12.5. GSaS Version of the Central Limit Theorem 97
12.6. Tail Behavior 106
12.7. Maximum Skewness and Half GSaS 108
12.8. Fractional Skew Exponential Power Distribution 109
12.9. Quadratic Form 109
12.10. Univariate MLE 109
12.11. Examples of Univariate MLE Fits 110

Chapter 13. Fractional Feller Square-Root Process 113
13.1. Generation of Random Variables for FCM 114
13.2. Generation of Random Variables for FCM2 115

Part 4. Multivariate Distributions 117

Chapter 14. Multivariate SN Distribution - Review 119
14.1. Definition 119
14.2. The Location-Scale Family 120
14.3. Quadratic Form 120
14.4. Stochastic Representation 120
14.5. Moments 121
14.6. Canonical Form 122
14.7. 1D Marginal Distribution 122

Chapter 15. Multivariate GAS-SN Elliptical Distribution 125
15.1. Definition 125
15.2. Location-Scale Family 125
15.3. Moments 125
15.4. Canonical Form 126
15.5. Marginal 1D Distribution 126
15.6. Quadratic Form 126
15.7. Multivariate MLE 127

Chapter 16. Multivariate GAS-SN Adaptive Distribution (Experimental) 129
16.1. Definition 129
16.2. Location-Scale Family 129
16.3. Moments 129
16.4. Canonical Form 130
16.5. Marginal 1D Distribution 130
16.6. 2D Adaptive MLE 130

Chapter 17. Fitting SPX-VIX Daily Returns with Bivariate Distributions 133
17.1. Elliptical Fit 133
17.2. Adaptive Fit 138

Appendix A. List of Useful Formula 143
A.1. Gamma Function 143
A.2. Transformation 143
A.3. Half-Normal Distribution 144

Bibliography 145

CHAPTER 1

Introduction

1.1. Motivation

In quantitative finance, asset return data often exhibit substantial skewness and kurtosis. In portfolio optimization and market-regime modeling[18, 38, 29], a standard example is the joint behavior of the S&P 500 Index (SPX) and the CBOE Volatility Index (VIX), whose daily prices have been publicly available since 1990[1]. Although such data sets are readily available, fitting them with existing parametric distributions remains difficult. Even with the many probability distributions implemented in modern statistical software, such as **scipy.stats**, the fit is often inadequate.

This book develops a multivariate elliptical distribution system based on the Wright function[43, 44, 2]. As shown in the next section, this subordinator approach is identical to the tilted stable law[8]. The construction combines and extends the α-stable distribution[17] and the multivariate skew-t distribution[1]. The resulting family is intended to model real-world data sets with pronounced tail behavior with more accuracy.

For example, the daily return distribution of VIX has kurtosis about 16 and skewness about 2.0. Its standardized peak density is approximately 0.55 (see Figure 12.10). For comparison, the excess kurtosis of the t distribution[40] is $6/(k-4)$ for $k > 4$, so matching this level of kurtosis would require k to be very close to 4. However, the corresponding standardized peak density is only 0.53 at $k = 4$. Thus, the VIX data lie near the boundary of what the t distribution can represent.

The daily return distribution of SPX is even more challenging (see Figure 12.11). In addition to kurtosis of about 11, its standardized peak density is approximately 0.65. A reasonable fit requires a t distribution with about 3 degrees of freedom ($k \approx 3$). However, from a theoretical standpoint, finite kurtosis does not exist for the t distribution until $k > 4$.

These two examples illustrate a basic difficulty in fitting existing parametric distributions. In particular, it is difficult to match both the kurtosis and the peak density simultaneously.

Our new multivariate distribution is designed to fit both data sets with satisfactory accuracy while matching empirical skewness, kurtosis, and peak density. Goodness of fit is evaluated not only through the density function but also through the extent to which the distribution captures tail behavior via the quadratic form. These fits are presented in Chapter 17.

[1]SPX data: Courtesy of S&P Dow Jones Indices LLC, from https://fred.stlouisfed.org/series/SP500. VIX data: Courtesy of Chicago Board Options Exchange (CBOE), from https://fred.stlouisfed.org/series/VIXCLS). Retrieved from FRED, Federal Reserve Bank of St. Louis. Not for commercial use.

1.2. Elements of Probability Distributions

The word "fractional" can be understood as adding the Lévy stability index α to a known classic distribution. Depending on the context, α could be in $[0, 1]$ or $[0, 2]$.

For example, in the Mellin transform of the PDF of a distribution, $\Gamma(s + c)$ in the classic world becomes $\Gamma(\alpha s + c)$ or $\Gamma(s/\alpha + c)$ in the fractional world. When the coefficient of s is $\frac{1}{2}$, 1, or 2, the fractional distribution subsumes the classic distribution, since the Legendre duplication formula (A.2) becomes applicable. But where do these fractional $\Gamma(\alpha\,...)$ functions come from?

The most fundamental and elegant representation is based on the **Tilted Stable law** (TS). The tilted stable law emerged from the study of the Poisson-Dirichlet distribution by Pitman and Yor[34]. In the limit of infinite samplings, the normalized distribution of the size of the cluster approaches the tilted stable law: $T^{-\alpha}_{\alpha,\beta} = (T_{\alpha,\beta})^{-\alpha}$, where α is the stability index in $[0, 1]$ and $\beta \geq 0$ is the power of the polynomial tilt. T_α is the stable law without tilt $T_{\alpha,0}$.

Take the Chinese restaurant process as an example (Section 3 of [33]). Let K_n be the number of tables occupied after n customers, and n_j be customers already in table j. The Pitman-Yor probability for the next customer to join the table j or to open a new table is

$$\Pr(\text{join table } j) = \frac{n_j - \alpha}{\beta + n},$$
$$\Pr(\text{new table} \mid K_n) = \frac{\beta + \alpha K_n}{\beta + n}.$$

Its α-diversity limit is $\dfrac{K_n}{n^\alpha} \longrightarrow T^{-\alpha}_{\alpha,\beta}$. Hence, α plays a discount role as the larger α discounts the attractiveness of joining the existing tables. On the other hand, β is a baseline strength for creating a new table, independent of n and K_n.

In [8] and [9], the superscript is generalized to form a three-parameter distribution family $T^{-\gamma}_{\alpha,\beta}$ where $\gamma \in \mathbb{R}$. To avoid collisions of the greeks, we also use the text-based notation: $TS(\alpha, \beta, \gamma)$. When $\gamma > 0$, it represents a local time process. When $\gamma < 0$, it represents a volatility process.

This three-parameter distribution law is identical to our fractional gamma distribution with a different parameterization. In Table 1, we list all the one-sided distributions in this book that have direct mapping to $T^{-\gamma}_{\alpha,\beta}$.

Fractional Distribution	Section	Notation	$T^{-\gamma}_{\alpha,\beta}$ or $TS(\alpha,\beta,\gamma)$			
			α	β	γ	scale
One-sided stable	4.2	L_α (aka T_α)	α	0	-1	1
M-Wright (or Mittag-Leffler)	3.4	M_α (aka $T^{-\alpha}_\alpha$)	α	0	α	1
Fractional gamma	6.5	$\mathfrak{N}_\alpha(\sigma, d, p)$	α	$\alpha d/p$	α/p	σ
Fractional chi (FCM)	7.2.1	$\overline{\chi}_{\alpha,k}\,(k > 0)$	$\alpha/2$	(k-1)/2	1/2	$\sigma_{\alpha,k}$
Inverse FCM	9.4	$\overline{\chi}^{\dagger}_{\alpha,k}$	$\alpha/2$	(k-1)/2	-1/2	$\sigma^{-1}_{\alpha,k}$
Characteristic FCM	7.4.2	$\overline{\chi}_{\alpha,-k}$	$\alpha/2$	k/2	-1/2	$\sigma^{-1}_{\alpha,k}$
Fractional chi-squared (FCM2)	7.8.1	$\overline{\chi}^2_{\alpha,k}$	$\alpha/2$	(k-1)/2	1	$\sigma^2_{\alpha,k}$
Characteristic FCM2	7.8.1	$\overline{\chi}^2_{\alpha,-k}$	$\alpha/2$	$k/2$	-1	$\sigma^{-2}_{\alpha,k}$

TABLE 1. The tilted stable law for the fractional distributions in this book. k is the "degree of freedom" parameter as in Student's t and χ^2.

Therefore, $T_{\alpha,\beta}$ is seen as **the atom** of all one-sided distributions. The polynomial tilt is expressed in its density function: $\frac{\Gamma(1+\beta)}{\Gamma(1+\beta/\alpha)}\, x^{-\beta}\, L_\alpha(x)$ $(x > 0)$.[**33, 8**] And importantly, $\overline{\chi}^2_{\alpha,-k}$ is equal to $T_{\alpha/2,k/2}$ according to the last row of the table.

In addition to Table 1, two fractional distributions are formed by a ratio of two tilted stable laws.

(1) Fractional F distribution in Section 8.1.2: It is $F_{\alpha,d,k} \sim \overline{\chi}^2_{1,d}/\overline{\chi}^2_{\alpha,k}$
(2) The α-stable distribution in Section 4.3.2: We have

$$L^\theta_\alpha \sim M_\gamma / T^{-\gamma}_{\gamma\alpha,\,0} = (L_{\gamma\alpha}/L_\gamma)^\gamma, \text{ where } \quad \gamma = \frac{\alpha-\theta}{2\alpha}, \alpha \in (0,2].$$

This is for the γ portion of the population in $x > 0$. And (α, θ) is in Feller's parametrization (4.1)[**12, 13**]. (A similar formula was derived in [**9**].)

To construct the two-sided distributions, we need the **second atom**, **Selective Sampling**, to produce the skewness[**1**].

We take the well-known *multivariate normal distribution* $\mathcal{N}_d(0, \bar{\Omega})$, where d is the dimension, and $\bar{\Omega}$ is a $d \times d$ correlation matrix. The selective sampling works as follows (Section 14.4).

Assuming $X_0 \sim \mathcal{N}_d(0, \bar{\Omega})$, $X_1 \sim \mathcal{N}(0,1)$, and $\boldsymbol{\beta} \in \mathbb{R}^d$ is the skew parameter, they form the skew-normal distribution $Z \sim SN_d(0, \bar{\Omega}, \boldsymbol{\beta})$ such that (Section 2.1 of [**1**])

$$Z = \begin{cases} X_0 & \text{if } X_1 > \boldsymbol{\beta}^\intercal X_0, \\ -X_0 & \text{otherwise.} \end{cases}$$

A two-sided stable distribution is built by properly assembling the ratio $Z/TS(...)$ or the product $Z \times TS(...)$.

The crown jewel is the skew multivariate elliptical distribution, based on the ratio:

$$SN_d(0, \bar{\Omega}, \boldsymbol{\beta}) / \overline{\chi}_{\alpha,k} \qquad \text{(Definition 15.1)}$$

The most important chapters of the book are

- Chapter 12 on the univariate GAS-SN distribution; and
- Chapter 15 on the multivariate GAS-SN elliptical distribution.

The reader can think of the entire book as aiming at developing tools (atoms) to create these two distributions.

Our univariate GAS-SN distribution is supposed to be the most flexible two-sided distribution up to date for statisticians to fit a univariate data set, such as return distributions in finance.

Our multivariate GAS-SN elliptical distribution is intended to be the most flexible multivariate distribution to date that extends the multivariate skew-t and skew-normal distributions[**1**].

A reference implementation in **python** can be found on Github at: https://github.com/slihn/gas-impl. This provides a baseline if these distributions make their way to standard libraries, such as **scipy.stats** in the future.

1.3. Organization of This Book

This book is divided into three parts.

Part I describes the mathematical foundation needed for the construction of fractional distributions. It contains several higher transcendental functions. Several classic special functions are extended with a fractional parameter.

Each distribution has its density function (PDF) and distribution function (CDF). Its Mellin transform. The squared variable or quadratic forms. Therefore, new mathematical tools are needed to address them.

Part II contains the univariate one-sided fractional distributions that are invented. All of them have their classic counterparts. For example, the generalized gamma distribution (GG) is upgraded. All the χ and F related distributions are also upgraded.

Part III contains the two-sided univariate fractional distributions. The Azzalini (2013) book is used as the blueprint[1]. It is integrated with the symmetric distributions developed in my 2024 work[20].

This book can be viewed as an integration between the two works, literally going chapter-by-chapter. The consistency of such integration and harmony speaks volumes.

The fourth part introduces the multivariate fractional distributions. These families extend the univariate constructions in Part III and subsume the skew-normal and skew-t families discussed in Azzalini's book.

Their main advantage is the ability to represent a wide range of skewness, kurtosis, and peak density within a unified framework. This flexibility makes them suitable for data sets whose distributional features cannot be adequately captured by more classical parametric models.

Modern data sets are often large in both sample size and dimension, making tail behavior increasingly important to model accurately. In finance, this is especially true for the left tail, where risk is concentrated.

An adaptive version of the multivariate distribution is therefore developed so that each dimension may carry its own shape parameters. This construction is used to fit one of the most challenging financial data sets considered in this book: daily SPX and VIX returns since 1990. The corresponding methodology is presented in Part IV.

1.4. Unresolved Issues

Although the two multivariate distributions present new opportunities to fit the data sets that were thought impossible previously, the outcomes post new challenges.

On the one hand, maximum-likelihood estimation (MLE) for the elliptical distribution is relatively straightforward to implement. The resulting fit (Figures 17.1, 17.2, 17.3) appears satisfactory at the visual level. However, the fitted values of (α, k) lie near a region of effectively infinite kurtosis when the bivariate distribution is projected onto its two one-dimensional marginals. This tension between visual fit and implied marginal behavior remains an important unresolved issue.

On the other hand, the adaptive distribution suffers from the curse of dimensionality, which makes direct MLE computationally prohibitive. Therefore, a modified fitting algorithm is used. The resulting fit (Figures 17.4, 17.5, 17.6) is qualitatively reasonable, but several limitations remain. In particular, the theoretical correlation coefficient remains smaller in absolute value than the empirical value of about -0.7, and the quadratic form does not yet have a matching F distribution.

The broader mathematical and statistical questions raised by these constructions remain open.

Part 1

Mathematical Functions

CHAPTER 2

Mellin Transform

This chapter begins with the Mellin-transform tools used throughout the book. Readers who are mainly interested in the statistical models may treat this and the following chapter as technical background, but the notation introduced here is used repeatedly in the later constructions.

The Mellin transform, introduced by Hjalmar Mellin in 1897 [27], is a central tool for analyzing one-sided probability distributions. Once the Mellin transform of a density is known, the moments and several derivative relations can often be obtained directly. This makes the transform especially useful for the Wright function, the α-stable distribution, and the fractional χ distribution, whose connections are most naturally expressed in Mellin-transform form.

DEFINITION 2.1. This chapter provides an overview of the Mellin transform. Following the notation of [24], the Mellin transform of a function $f(x)$ properly defined for $x \geq 0$ is

$$f^*(s) := \mathcal{M}\{f(x); s\} = \int_0^\infty f(x)\, x^{s-1}\, dx, \qquad c_1 < \mathbb{R}(s) < c_2. \tag{2.1}$$

The role of c_1, c_2 will be explained in the following.

If $f^*(s)$ has analytic continuation on the complex plane, the inverse Mellin transform is

$$f(x) = \mathcal{M}^{-1}\{f^*(s); x\} = \frac{1}{2\pi i} \int_{C-i\infty}^{C+i\infty} f^*(s)\, x^{-s}\, ds, \qquad c_1 < C < c_2. \tag{2.2}$$

Equation (2.1) shows the direct connection between the Mellin transform and distributional moments. If $f(x)$ is the PDF of a one-sided distribution, then its n-th moment is $\mathbb{E}(X^n|f) = f^*(n+1)$. Thus, modifying the Mellin transform $f^*(s)$ provides a systematic way to construct new distributions from existing ones.

For compactness, we use the juxtaposition notation $\overset{\mathcal{M}}{\longleftrightarrow}$ and write $f(x) \overset{\mathcal{M}}{\longleftrightarrow} f^*(s)$, with the validity strip inherited from (2.1) and (2.2). The precise choice of the contour parameter C is needed for numerical inversion in (2.2), but otherwise the notation mainly serves to keep the transform identities concise.

LEMMA 2.2. The main rules of Mellin transform used in this paper are:

$$f(ax) \overset{\mathcal{M}}{\longleftrightarrow} a^{-s}\, f^*(s), \qquad a > 0 \tag{2.3}$$

$$x^k f(x) \overset{\mathcal{M}}{\longleftrightarrow} f^*(s+k), \tag{2.4}$$

$$f(x^p) \overset{\mathcal{M}}{\longleftrightarrow} \frac{1}{p}\, f^*(s/p), \qquad p \neq 0 \tag{2.5}$$

and the following ones involving an integral,

$$h(x) = \int_0^\infty f(xs)g(s)\, s\, ds \overset{\mathcal{M}}{\longleftrightarrow} h^*(s) = f^*(s)g^*(2-s), \qquad \text{(ratio distribution)} \tag{2.6}$$

$$\gamma_f(x) = \int_0^x f(x)\, dx \overset{\mathcal{M}}{\longleftrightarrow} -s^{-1}f^*(s+1), \qquad \text{(lower incomplete function)} \tag{2.7}$$

$$\Gamma_f(x) = \int_x^\infty f(x)\, dx \overset{\mathcal{M}}{\longleftrightarrow} s^{-1}f^*(s+1). \qquad \text{(upper incomplete function)} \tag{2.8}$$

The ratio distribution rule (2.6) is widely used in our fractional distribution system. Notice that the argument of $g^*(s)$ is transformed via $s \to 2 - s$.

For (2.7) and (2.8), the valid range of C is decremented by one: $c_1 - 1 < C < c_2 - 1$.

△

Example 2.3. A simple exercise is the Mellin transform of the standard normal distribution. It starts with

$$e^{-x} \overset{\mathcal{M}}{\longleftrightarrow} \Gamma(s)$$

via the definition of the gamma function itself.

By applying (2.5) then (2.3), we get

$$\mathcal{N}(x) \overset{\mathcal{M}}{\longleftrightarrow} \mathcal{N}^*(s) = \frac{2^{(s-3)/2}}{\sqrt{\pi}}\, \Gamma\left(\frac{s}{2}\right) \tag{2.9}$$

where $\mathcal{N}(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$ is our notation for the PDF of a standard normal distribution.

Example 2.4. A slightly more complicated exercise is the Mellin transform of the PDF of the fractional gamma distribution (FG) in Chapter 6. But we only work out its skeleton here.

Assume we have a function $F_\alpha(x)$ whose Mellin transform is

$$F_\alpha(x) \overset{\mathcal{M}}{\longleftrightarrow} \frac{\Gamma(s)}{\Gamma(\alpha s)}.$$

It undergoes the following transforms:

$$F_\alpha(x^p) \overset{\mathcal{M}}{\longleftrightarrow} \frac{1}{|p|}\frac{\Gamma(s/p)}{\Gamma(\alpha s/p)},$$

$$x^{d-1}\, F_\alpha(x^p) \overset{\mathcal{M}}{\longleftrightarrow} \frac{1}{|p|}\frac{\Gamma((s+d-1)/p)}{\Gamma(\alpha(s+d-1)/p)},$$

which is the prototype of FG before further normalization.

2.1. One-Sided Distribution

When the subject matter is a probability distribution, the two rules of incomplete functions, (2.7) and (2.8), provide a clear path to obtain its distribution function (CDF) while $f(x)$ is associated with its density function (PDF).

For a one-sided distribution denoted f, it is straightforward to define it, since the domain of the distribution $x \geq 0$ is exactly the same as the domain of the Mellin transform. Hence, $f(x)$ is exactly its density function and $\gamma_f(x)$ is its CDF.

By assigning $s = n + 1$, its n-th moment is

$$\mathbb{E}(X^n|f) = f^*(n+1) \tag{2.10}$$

The zeroth moment is the total density that should be normalized to one. That is $f^*(1) = 1$.

The mean of the distribution is $\mathbb{E}(X|f) = f^*(2)$. For a distribution whose mean needs to be determined, for instance, the fractional χ distribution in Chapter 7, this formula is useful.

2.2. Two-Sided Distribution

When the distribution is two-sided, one more rule is needed.

DEFINITION 2.5 (The Reflection Rule). Assume the distribution is two sided, its domain of support is $x \in \mathbb{R}$. The reflection rule requires that its density function $f(x)$ is based on a skew parameter $\beta \in \mathbb{R}$ that satisfies

$$f(-x;\beta) := f(x;-\beta) \quad \text{for } x > 0. \tag{2.11}$$

Let $c_\beta < 1$ be the one-sided integral of

$$c_\beta = \int_0^\infty f(x;\beta)\,dx.$$

(2.11) leads to $c_{-\beta} + c_\beta = 1$ since $\int_{-\infty}^{\infty} f(x;\beta)\,dx = 1$.

2.2.1. Mellin Transform of a Two-sided CDF.

LEMMA 2.6. The Mellin transform of the CDF $\Phi(x)$ of a two-sided distribution has two parts. Both can be derived from its density function transform, $f(x;\beta) \overset{\mathcal{M}}{\longleftrightarrow} f^*(s;\beta)$, in the positive domain.

From (2.7), let $\gamma_f(x;\beta) \overset{\mathcal{M}}{\longleftrightarrow} \Phi^*(s;\beta) := -s^{-1} f^*(s+1;\beta)$. Then for $x > 0$, the Mellin transform of the CDF can be expressed as

$$\Phi(x) - \Phi(0) \overset{\mathcal{M}}{\longleftrightarrow} \Phi^*(s;\beta),$$
$$1 - \Phi(0) - \Phi(-x) \overset{\mathcal{M}}{\longleftrightarrow} \Phi^*(s;-\beta).$$

△

PROOF. Note that $\Phi(0) = c_{-\beta} = 1 - c_\beta$. When $x \geq 0$, its CDF is

$$\Phi(x) = \int_{-\infty}^{x} f(x;\beta)\,dx \;= c_{-\beta} + \int_0^x f(x;\beta)\,dx \;= \Phi(0) + \gamma_f(x;\beta).$$

In the negative domain, its CDF is

$$\begin{aligned}\Phi(-x) = \int_{-\infty}^{-x} f(x;\beta)\,dx \;&= \int_x^\infty f(x;-\beta)\,dx \\ &= 1 - \Phi(0) - \int_0^x f(x;-\beta)\,dx \;= 1 - \Phi(0) - \gamma_f(x;-\beta).\end{aligned}$$

□

The point is that, once the Mellin transform of either the PDF or CDF is known, the other one can be derived by simple algebraic rules.

2.2.2. From Mellin Transform to Moments. By assigning $s = n+1$, it is easy to show that its n-th moment is

$$\mathbb{E}(X^n|f) = f^*(n+1;\beta) + (-1)^n f^*(n+1;-\beta) \tag{2.12}$$
$$= -n\big[\Phi^*(n;\beta) + (-1)^n \Phi^*(n;-\beta)\big] \tag{2.13}$$

The moment formula is tightly linked to $\Phi^*(n;\beta)$.

The Zeroth Moment. The total density can be regarded as the zeroth moment. Hence, c_β can be determined by

$$c_\beta = \int_0^\infty f(x;\beta)dx = f^*(1;\beta). \tag{2.14}$$

Its application is in (10.9).

2.3. Ramanujan's Master Theorem

In order to keep things simple, we describe all the distributions via the Mellin transform of their PDFs. Due to Ramanujan's master theorem[3], not only can the moments be obtained from the Mellin transform but also all the derivatives of the PDF at $x = 0$. We get its series representation "for free", so to speak.

LEMMA 2.7 (Ramanujan's master theorem). If $f(x)$ has an expansion of the form

$$f(x) = \sum_{n=0}^{\infty} \frac{\varphi(n)}{n!}(-x)^n \tag{2.15}$$

then its Mellin transform is given by

$$f(x) \overset{\mathcal{M}}{\longleftrightarrow} f^*(s) = \Gamma(s)\,\varphi(-s) \tag{2.16}$$

△

Assume that $g^*(s) := f^*(s)/\Gamma(s)$ exists on the complex plane, $s \in \mathbb{C}$. Its connection to the derivatives of the PDF at $x = 0$ is as follow.

LEMMA 2.8. The Taylor series of $f(x)$ is

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!}x^n$$

where $f^{(n)}(0)$ is the n-th derivative of $f(x)$ at $x = 0$.

Then $f^{(n)}(0)$ can be obtained from $g^*(s)$ by

$$f^{(n)}(0) = (-1)^n g^*(-n) \tag{2.17}$$

At $x = 0$, we have $f(0) = g^*(0)$.

△

The power of the master theorem is that, once the Mellin transform is known, the Taylor series is also known immediately. We provide a contrived example from next chapter as a showcase.

EXAMPLE 2.9. The Mellin transform of the Wright function from (3.5) is $f(-x) \overset{\mathcal{M}}{\longleftrightarrow} f^*(s) = \Gamma(s)/\Gamma(\delta - \lambda s)$. Then its $g^*(s) = 1/\Gamma(\delta - \lambda s)$.

According to Lemma 2.8, its Taylor series should be

$$f(-x) := \sum_{n=0}^{\infty} \frac{(-1)^n g^*(-n)}{n!}x^n = \sum_{n=0}^{\infty} \frac{g^*(-n)}{n!}(-x)^n$$

Replace $-x$ with z, and plug in $g^*(-n)$, we have

$$f(z) := \sum_{n=0}^{\infty} \frac{z^n}{n!\,\Gamma(\lambda n + \delta)}$$

This is the series representation (3.1) where we essentially "derived" it from the master theorem.

The major application in this book is in Chapter 11. In the experimental construction of the generalized α-stable distribution, the theorem is used to remedy the discontinuity of the PDF in $x = 0$.

2.3.1. Distribution Function. The form of the Mellin transform in (2.16) has an important implication when $f(x)$ is a density function.

LEMMA 2.10. Assume $x > 0$, its complimentary distribution function $\Gamma_f(x) := \int_x^\infty f(x)\,dx$ has the series representation of

$$\Gamma_f(x) = \sum_{n=0}^{\infty} \frac{\varphi(n-1)}{n!}(-x)^n \tag{2.18}$$

△

PROOF. From (2.8), the Mellin transform of $\Gamma_f(x)$ is

$$\Gamma_f(x) = \int_x^\infty f(x)\,dx \overset{\mathcal{M}}{\longleftrightarrow} s^{-1} f^*(s+1)$$

which can be simplified to

$$\begin{aligned} s^{-1} f^*(s+1) &= s^{-1}\Gamma(s+1)\,\varphi(-s-1) \\ &= \Gamma(s)\,\varphi(-s-1). \end{aligned}$$

This is still in the form of (2.16), with a transformation rule of $s \to s+1$ in the function $\varphi(-s)$.
Applying the master theorem of (2.15), we get (2.18).

□

We use the CDF of the M-Wright function from (3.17) as an example.

LEMMA 2.11. The goal is to show

$$\int_x^\infty M_\alpha(t)dt = W_{-\alpha,1}(-x). \tag{2.19}$$

△

PROOF. We start with the Mellin transform of $M_\alpha(x)$ from (3.13),

$$M_\alpha(x) \overset{\mathcal{M}}{\longleftrightarrow} M_\alpha^*(s) = \frac{\Gamma(s)}{\Gamma((1-\alpha)+\alpha s)}$$

which yields $\varphi(-s) = 1/\Gamma((1-\alpha)+\alpha s)$.
Therefore, its $\Gamma_f(x)$ should be

$$\Gamma_f(x) = \sum_{n=0}^{\infty} \frac{1}{n!\,\Gamma((1-\alpha)-\alpha(n-1))}(-x)^n = \sum_{n=0}^{\infty} \frac{1}{n!\,\Gamma(-\alpha n+1)}(-x)^n$$

which is $W_{-\alpha,1}(-x)$ according to (3.1).

□

CHAPTER 3

The Wright Function and The Inverse Stable Law

3.1. Definition

The Wright function is a basic building block in the fractional-distribution system developed in this book. It was introduced by E. M. Wright in the 1930s[43, 44], and Bateman later catalogued it together with the Mittag-Leffler function[2].

Its importance became much clearer in the later literature, especially through the work of F. Mainardi, who introduced the M-Wright function $M_\alpha(x)$. In this framework, $M_\alpha(x)$ plays the role of a fractional analogue of the exponential function e^{-x}. This chapter collects the main facts about the Wright function needed in the remainder of the book.

DEFINITION 3.1. The series representation of the Wright function is

$$W_{\lambda,\delta}(z) := \sum_{n=0}^{\infty} \frac{z^n}{n!\,\Gamma(\lambda n + \delta)} \qquad (\lambda \geq -1, z \in \mathbb{C}) \tag{3.1}$$

Its shape parameters are pairs (λ, δ). The apparent limit is $W_{0,1}(z) = e^z$.

Four variants of the Wright function are used repeatedly in this book. The first pair is

- $M_\alpha(z) := W_{-\alpha,1-\alpha}(-z)$
- $F_\alpha(z) := W_{-\alpha,0}(-z)$

with $\alpha \in [0, 1]$. They satisfy $M_\alpha(z) = F_\alpha(z)/(\alpha z)$.

The function $M_\alpha(z)$ is known as *the M-Wright function*, or equivalently *the Mainardi function*[21, 26, 22]. See Section 3.3 for further details. Its role as a fractional extension of classical exponential behavior is reflected in the two reference cases $M_0(z) = \exp(-z)$ and $M_{\frac{1}{2}}(z) = \frac{1}{\sqrt{\pi}} \exp(-z^2/4)$.

The second pair is

- $W_{-\alpha,-1}(-z)$
- $-W_{-\alpha,1-2\alpha}(-z)$

These functions arise naturally from derivatives of $F_\alpha(z)$ and $M_\alpha(z)$. They are useful in random-variable generation, as in (3.19) and Section 11 of [20], and in some cases they reduce to polynomial expressions.

3.2. Classic Results

The recurrence relations of the Wright function are (Chapter 18, Vol 3 of [2])

$$\lambda z\, W_{\lambda,\lambda+\mu}(z) = W_{\lambda,\mu-1}(z) + (1-\mu) W_{\lambda,\mu}(z) \tag{3.2}$$

$$\frac{d}{dz} W_{\lambda,\mu}(z) = W_{\lambda,\lambda+\mu}(z) \tag{3.3}$$

The moments of the Wright function are (See (1.4.28) of [26])

$$\mathbb{E}(X^{d-1}) = \int_0^\infty x^{d-1} W_{-\lambda,\delta}(-x)dx = \frac{\Gamma(d)}{\Gamma(d\lambda + \delta)} \tag{3.4}$$

The way it is written is in fact its Mellin transform:

$$W_{\lambda,\delta}(-x) \overset{\mathcal{M}}{\longleftrightarrow} W^*_{\lambda,\delta}(s) = \frac{\Gamma(s)}{\Gamma(\delta - \lambda s)} \tag{3.5}$$

$W_{\lambda,\delta}(z)$ has the following Hankel integral representation:

$$W_{\lambda,\delta}(z) = \frac{1}{2\pi i} \int_H dt\, \frac{\exp\left(t + z\, t^{-\lambda}\right)}{t^{\delta}} \tag{3.6}$$

Prodanov[35] derived an integral representation of the Wright function. We use the branch with $\lambda < 0$ and $\delta \leq 1$ from Theorem 1, namely

$$W_{\lambda,\delta}(z) = \frac{1}{\pi} \int_0^{\infty} \frac{dr}{r^{\delta}}\, \sin(\sin(\lambda\pi)w + \delta\pi)\, e^{\cos(\lambda\pi)w - r}, \quad \text{where } w = z\, r^{-\lambda}. \tag{3.7}$$

This representation is well suited to numerical evaluation by tanh-sinh quadrature.

The four-parameter Wright function is defined as

$$W\begin{bmatrix} a, & b \\ \lambda, & \mu \end{bmatrix}(z) := \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(n+1)} \frac{\Gamma(an+b)}{\Gamma(\lambda n + \mu)} \tag{3.8}$$

This function is a higher-order Wright function. It was used seriously for the first time by the author[20].

3.3. The M-Wright Functions

Mainardi has introduced two auxiliary functions of Wright type (see F.2 of [21]). Assume $\alpha \in [0, 1]$,

$$F_\alpha(z) := W_{-\alpha,0}(-z) \qquad (z > 0) \tag{3.9}$$

$$M_\alpha(z) := W_{-\alpha,1-\alpha}(-z) = \frac{1}{\alpha z} F_\alpha(z) \qquad (z > 0) \tag{3.10}$$

The relation between $M_\alpha(z)$ and $F_\alpha(z)$ in (3.10) is an application of (3.2) by setting $\lambda = -\alpha, \mu = 1$.

$F_\alpha(z)$ has the following Hankel integral representation:

$$F_\alpha(z) = \frac{1}{2\pi i} \int_H dt\, \exp\left(t - z t^{\alpha}\right) \tag{3.11}$$

Both functions have simple Mellin transforms from (3.5):

$$F_\alpha(x) \overset{\mathcal{M}}{\longleftrightarrow} F^*_\alpha(s) = \frac{\Gamma(s)}{\Gamma(\alpha s)} \tag{3.12}$$

$$M_\alpha(x) \overset{\mathcal{M}}{\longleftrightarrow} M^*_\alpha(s) = \frac{\Gamma(s)}{\Gamma((1-\alpha) + \alpha s)} \tag{3.13}$$

The function $F_\alpha(z)$ is used to define fractional one-sided distributions. However, its series representation is not especially convenient for numerical computation, because many terms may be required to reach a given precision.

By contrast, $M_\alpha(z)$ admits a series representation that is more computationally practical, especially for small values of α:

$$M_\alpha(z) = \sum_{n=0}^{\infty} \frac{(-z)^n}{n!\,\Gamma(-\alpha n + (1-\alpha))} = \frac{1}{\pi} \sum_{n=1}^{\infty} \frac{(-z)^{n-1}}{(n-1)!} \Gamma(\alpha n) \sin(\alpha n \pi) \qquad (0 < \alpha < 1) \tag{3.14}$$

$M_\alpha(0) = 1/\Gamma(1-\alpha)$ monotonically decreases from 1 to 0 as α increases from 0 to 1.

The function $M_\alpha(z)$ also has simple closed-form expressions in the special cases $\alpha = 0$ and $\alpha = \frac{1}{2}$:

$$M_0(z) = \exp(-z) \text{ and } M_{\frac{1}{2}}(z) = \frac{1}{\sqrt{\pi}} \exp(-z^2/4). \tag{3.15}$$

$M_\alpha(z)$ can be computed to high accuracy when properly implemented with arbitrary-precision floating point library, such as the `mpmath` package[28]. In this regard, it is much more "useful" than $F_\alpha(z)$.

This is particularly important in working with large degrees of freedom and extreme values of α, mainly close to 0 and 1. The typical 64-bit floating-point algorithm suffers from overflow and/or underflow. See Section 3.8 for more details.

$M_\alpha(z)$ has the asymptotic representation in the *generalized gamma* (GG) style: (see F.20 of [21])

$$M_\alpha\left(\frac{x}{\alpha}\right) = A\, x^{d-1}\, e^{-B\, x^p} \tag{3.16}$$
$$\text{where } p = 1/(1-\alpha), d = p/2,\ A = \sqrt{p/(2\pi)},\ B = 1/(\alpha p).$$

Additional correction terms in the asymptotic expansion have been derived up to the order $x^{-6/(1-\alpha)}$[32]. This formula is important in guiding (3.14) to high precision for large x, where the series representation often fails to converge.

$M_\alpha(x)$ can be used as the density function of a one-sided distribution M_α[22], because $\int_0^\infty M_\alpha(x)dx = 1$ and $M_\alpha(x)$ for $x \geq 0$. Its CDF is another Wright function:

$$\int_0^x M_\alpha(t)dt = 1 - W_{-\alpha,1}(-x). \tag{3.17}$$

This is proved in Lemma 2.11.

The absolute moments of $M_\alpha(x)$ in $\mathbb{R}^+$ are

$$\mathbb{E}(M_\alpha^n) = \int_0^\infty t^n\, M_\alpha(t)dt = \frac{\Gamma(n+1)}{\Gamma(n\alpha+1)}, \quad n > -1. \tag{3.18}$$

Hence, its mean is located at $1/\Gamma(\alpha+1)$, which is equal to 1 when $\alpha = 0, 1$. Its variance is $2/\Gamma(2\alpha+1) - 1/\Gamma(\alpha+1)^2$. The variance becomes zero when $\alpha = 1$, consistent with $M_1(x) = \delta(x-1)$.

Differentiating $M_\alpha(z)$, and from (3.14), we get

$$\frac{d}{dz}M_\alpha(z) = -W_{-\alpha,1-2\alpha}(-z) = \frac{-1}{\pi}\sum_{n=2}^{\infty}\frac{(-z)^{n-2}}{(n-2)!}\Gamma(\alpha n)\sin(\alpha n\pi) \tag{3.19}$$

Note that $\frac{d}{dz}M_\alpha(0) = -\frac{1}{\pi}\Gamma(2\alpha)\sin(2\alpha\pi)$. This also indicates that

$$\frac{d}{dz}F_\alpha(z) = \alpha\left(1 + z\frac{d}{dz}\right)M_\alpha(z) \tag{3.20}$$

which can be implemented from $M_\alpha(z)$ through (3.14) and (3.19). These differential forms lead to the concept of elasticity in Section 3.6 and below.

3.4. The Inverse Stable Law

M_α is said to follow the *inverse stable law*. In Section 4.2, the extremal stable distribution L_α will be introduced [1]. The law is

$$M_\alpha \sim (L_\alpha)^{-\alpha}. \tag{3.21}$$

This can be proved by the moment formula. Let $X = M_\alpha, Y = L_\alpha$. Then $\mathbb{E}(Y^{-q}) = \dfrac{\Gamma(q/\alpha)}{\alpha\Gamma(q)}$. Set $q = \alpha n$. We get $Y^{-q} = (Y^{-\alpha})^n = X^n$. It leads to $\mathbb{E}(X^n) = \dfrac{\Gamma(n)}{\alpha\Gamma(\alpha n)} = \dfrac{\Gamma(n+1)}{\Gamma(\alpha n+1)}$. That is (3.18).

Since $M_{1/2}/\sqrt{2}$ has the half-normal density on $(0,\infty)$, it provides a simple reference case for the inverse stable law.

[1] Historically, people also used the notation S_α because S is the first letter of "stable".

The calculation of M_α follows the framework in Theorem 1 of [8]. This sets the foundation for the tilted stable law in Section 6.5.

DEFINITION 3.2. The factor $A_\alpha(q)$ is often called *the Zolotarev function*. It is defined as

$$A_\alpha(q) = \frac{\sin(\pi\alpha q)}{\sin(\pi q)^{1/\alpha}} \sin(\pi(1-\alpha)q)^c, \quad c = \frac{1-\alpha}{\alpha}. \tag{3.22}$$

LEMMA 3.3. For the inverse stable law M_α, the sampling is performed via Kanter's method:

$$M_\alpha = A_\alpha(Q)^{-\alpha}\, G^{1-\alpha}.$$

(1) Q is from a uniform random variable in $(0,1)$.
(2) G is a gamma variable: $G \sim \text{Gamma}(1, \text{scale} = 1)$.

△

3.5. The Fractional Gamma-Star Function

The so-called γ^* function is documented in 8.2.6 and 8.2.7 of DLMF[10]. It is defined as follows:

$$\gamma^*(s,x) := \frac{1}{\Gamma(s)} \int_0^1 t^{s-1}\, e^{-xt}\, dt = \frac{x^{-s}}{\Gamma(s)}\gamma(s,x)$$

The finite integral in $t \in [0,1]$ is transformed from the incomplete gamma function, which takes the form of $\gamma(s,x) = \int_0^x t^{s-1}\, e^{-t}\, dt$.

$\gamma^*(s,x)$ can be extended fractionally in a straightforward manner. It is used to calculate the CDF of the FG in Chapter 6. See (6.17) for details.

DEFINITION 3.4 (The fractional γ^* function). It is defined by replacing e^{-xt} with $M_\alpha(xt)$ such that

$$\gamma^*_\alpha(s,x) := \frac{\Gamma((1-\alpha)+\alpha s)}{\Gamma(s)} \int_0^1 dt\, t^{s-1}\, M_\alpha(xt) \tag{3.23}$$

The $\alpha \to 0$ limit of $\gamma^*_\alpha(s,x)$ subsumes the classic γ^* function, that is, $\gamma^*_0(s,x) = \gamma^*(s,x)$. This is reflected in the simple fact that $M_0(xt) = \exp(-xt)$.

The γ^* function is a subset of the fractional confluent hypergeometric function in Lemma 5.4.

The supplementary γ^*_α function is defined as

$$\Gamma^*_\alpha(s,x) := \frac{\Gamma((1-\alpha)+\alpha s)}{\Gamma(s)} \int_1^\infty dt\, t^{s-1}\, M_\alpha(xt) \tag{3.24}$$

LEMMA 3.5. It has an analytic relation such as

$$\Gamma^*_\alpha(s,x) = x^{-s} - \gamma^*_\alpha(s,x) \tag{3.25}$$

△

PROOF. The total integral of $t \in \mathbb{R}$ has an analytic solution due to (3.18):

$$\begin{aligned}
\Gamma^*_\alpha(s) &:= \frac{\Gamma((1-\alpha)+\alpha s)}{\Gamma(s)} \int_0^\infty dt\, t^{s-1}\, M_\alpha(xt) \\
&= \frac{\Gamma((1-\alpha)+\alpha s)}{\Gamma(s)\, x^s} \int_0^\infty dz\, z^{s-1}\, M_\alpha(z) \\
&= x^{-s}.
\end{aligned}$$

Split the integral into two parts: $\int_0^1$ and $\int_1^\infty$. We obtain (3.25). □

The identity formula is

$$x^s\Gamma^*_\alpha(s,x) + x^s\gamma^*_\alpha(s,x) = 1 \tag{3.26}$$

This identity is critical in understanding how to use γ^*_α versus Γ^*_α when calculating the CDF of a FG distribution.

Numerically, it is better to use γ^*_α when $x \le 1$ and to use Γ^*_α when $x > 1$.

3.5.1. The lognormal limit. The fractional gamma-star functions represent the CDF of FG through (6.17), including its FCM specializations in Section 7.1. The error-function limit below therefore describes the lognormal CDF limit for these families after the appropriate rescaling, while its proof remains purely analytic.

Although $M_\alpha(x) \to M_0(x) = e^{-x}$ at fixed x, the limit below also lets the gamma-star parameter and its argument vary. It follows directly from the M-Wright Mellin integral.

Define the "log-gamma difference" helper function

$$\Delta_\alpha(x) := \log\Gamma(x/\alpha) - \log\Gamma(x). \tag{3.27}$$

For $0 < \alpha \le 1$ and $x > 0$, its first three derivatives are

$$\Delta^{(j)}_\alpha(x) = \alpha^{-j}\psi^{(j-1)}(x/\alpha) - \psi^{(j-1)}(x), \qquad j = 1,2,3, \tag{3.28}$$

where $\psi = (\log\Gamma)'$ is the digamma function and $\psi^{(m)}(x) := \frac{d^m}{dx^m}\psi(x)$, for $m = 0,1,2,\ldots$, denotes its polygamma derivatives; in particular, $\psi^{(0)} = \psi$.

LEMMA 3.6 (Polygamma expansions). The polygamma expansions (DLMF 5.11) give, as $x \to \infty$,

$$\begin{aligned}
\Delta'_\alpha(x) &= \left(\frac{1}{\alpha} - 1\right)\log x - \frac{\log\alpha}{\alpha} + \frac{1-\alpha}{12x^2} + O(x^{-4}),\\
\Delta''_\alpha(x) &= \frac{1-\alpha}{\alpha x} - \frac{1-\alpha}{6x^3} + O(x^{-5}),\\
\Delta'''_\alpha(x) &= -\frac{1-\alpha}{\alpha x^2} + \frac{1-\alpha}{2x^4} + O(x^{-6}).
\end{aligned} \tag{3.29}$$

The remainders are uniform for $0 < \alpha \le 1$, so these expansions also apply when α tends to zero. They remain valid for $x = \beta + w$ as $\beta \to \infty$ with bounded complex w. △

LEMMA 3.7 (Error-function limit of the fractional gamma-star functions). Fix $\gamma \in \mathbb{R} \setminus \{0\}$ and $\tau_{LN} > 0$.[2] Let $0 < \alpha_n < 1$ and $s_n > 1$ satisfy

$$\alpha_n \longrightarrow 0, \qquad \alpha_n^2(s_n - 1) \longrightarrow \frac{\gamma^2}{\tau_{LN}^2}. \tag{3.30}$$

Put $\beta_n = \alpha_n(s_n - 1)$, which tends to infinity, and define

$$\begin{aligned}
\ell_n &:= \Delta_{\alpha_n}(\beta_n + \gamma) - \Delta_{\alpha_n}(\beta_n),\\
x_n(t) &:= \exp\left\{\frac{\alpha_n}{\gamma}\left(\ell_n + \tau_{LN}t - \frac{\tau_{LN}^2}{2}\right)\right\}, \qquad t \in \mathbb{R}.
\end{aligned}$$

The shift ℓ_n is defined for all sufficiently large β_n. It absorbs the diverging logarithmic drift into $x_n(t)$. Along the stated path, Taylor expansion gives

$$\ell_n = \gamma\Delta'_{\alpha_n}(\beta_n) + \frac{\tau_{LN}^2}{2} + o(1).$$

[2]For FG, $\gamma = \alpha/p$; ordinary and characteristic FCM use $\gamma = 1/2$ and $\gamma = -1/2$, respectively. Writing the power explicitly shows how γ enters the logarithmic coordinate and how the diverging shift is subtracted:

$$\frac{1}{\alpha_n}\log\big(x_n(t)^\gamma\big) - \ell_n = \tau_{LN}t - \frac{\tau_{LN}^2}{2}.$$

The two diverging terms on the left cancel exactly, leaving a finite centered coordinate for each fixed t.

Thus $\ell_n - \tau_{LN}^2/2$ in $x_n(t)$ is asymptotically the logarithmic center $\gamma\Delta'_{\alpha_n}(\beta_n)$.

For each fixed $t \in \mathbb{R}$, the regularized products satisfy

$$\begin{cases} x_n(t)^{s_n}\, \gamma^*_{\alpha_n}(s_n, x_n(t))\,, & \gamma > 0, \\ x_n(t)^{s_n}\, \Gamma^*_{\alpha_n}(s_n, x_n(t))\,, & \gamma < 0 \end{cases} \longrightarrow \frac{1}{2}\left[1 + \operatorname{erf}\left(\frac{t}{\sqrt{2}}\right)\right]. \tag{3.31}$$

The prefactors and moving arguments are essential; this is not a fixed-argument limit of either gamma-star function alone. △

Proof. Write $u_n(y) = \exp[\alpha_n(y + \ell_n)/\gamma]$ and introduce the nonnegative kernel

$$h_n(y) := \frac{\alpha_n}{|\gamma|}\frac{\Gamma(1+\beta_n)}{\Gamma(s_n)} u_n(y)^{s_n} M_{\alpha_n}(u_n(y))\,.$$

Changing variables in (3.18) gives $\int_{\mathbb{R}} h_n(y)\,dy = 1$ and the exact transform identity

$$\int_{\mathbb{R}} e^{qy} h_n(y)\,dy = \exp\big[\Delta_{\alpha_n}(\beta_n + \gamma q) - \Delta_{\alpha_n}(\beta_n) - q\ell_n\big].$$

It holds for complex q with $\Re(\beta_n + \gamma q) > 0$, by analytic continuation within the convergence domain of the Mellin integral.

Suppressing the sequence index, (3.29) and the path (3.30) give $\gamma^2\Delta''_\alpha(\beta) \to \tau_{LN}^2$ and $\Delta'''_\alpha(\beta + w) = O(\beta^{-1})$ uniformly for bounded complex w. Taylor expansion cancels the linear term to give, uniformly for bounded complex q,

$$\Delta_\alpha(\beta + \gamma q) - \Delta_\alpha(\beta) - q\ell = \frac{\gamma^2}{2}\Delta''_\alpha(\beta)(q^2 - q) + O(\beta^{-1}) \longrightarrow \frac{\tau_{LN}^2}{2}(q^2 - q).$$

At $q = i\xi$, the transforms therefore converge to $\exp[-i\xi\tau_{LN}^2/2 - \tau_{LN}^2\xi^2/2]$. The Fourier continuity theorem for nonnegative unit-mass kernels gives convergence of their cumulative integrals to the continuous Gaussian integral:

$$\int_{-\infty}^{\tau_{LN}t - \tau_{LN}^2/2} h_n(y)\,dy \longrightarrow \frac{1}{\sqrt{2\pi}}\int_{-\infty}^{t} e^{-r^2/2}\,dr = \frac{1}{2}\left[1 + \operatorname{erf}\left(\frac{t}{\sqrt{2}}\right)\right].$$

Changing variables back to $u = u_n(y)$ identifies the left-hand integral with the lower regularized expression for $\gamma > 0$ and the supplementary expression for $\gamma < 0$, proving (3.31). □

Why the limit is lognormal. The error function in (3.31) is written in the standardized logarithmic coordinate t. To display the corresponding positive variable, set

$$v = e^{\tau_{LN}t - \tau_{LN}^2/2}, \qquad t = \frac{\log v + \tau_{LN}^2/2}{\tau_{LN}}, \qquad v > 0.$$

In this coordinate, the limiting function is

$$\Phi_{\tau_{LN}}(v) := \frac{1}{2}\left[1 + \operatorname{erf}\left(\frac{\log v + \tau_{LN}^2/2}{\sqrt{2}\,\tau_{LN}}\right)\right]. \tag{3.32}$$

Its derivative is explicitly the lognormal density:

$$\Phi'_{\tau_{LN}}(v) = \frac{1}{v\tau_{LN}\sqrt{2\pi}} \exp\left[-\frac{(\log v + \tau_{LN}^2/2)^2}{2\tau_{LN}^2}\right].$$

Thus $\Phi_{\tau_{LN}}$ is the CDF of Lognormal$(-\tau_{LN}^2/2, \tau_{LN}^2)$. The lognormal form appears in the rescaled positive coordinate $v = e^{-\ell_n} x^{\gamma/\alpha_n}$; it is the same limit as the error-function form in t, expressed in a different coordinate.

3.6. The Elasticity Operator

In (3.19) and (3.20), we encountered an important mathematical structure called "elasticity" which will be used in Chapter 13. It provides an elegant view of the inner structure of the FG density functions.

DEFINITION 3.8 (The elasticity operator). Assume $f(x)$ is differentiable for $x \in \mathbb{R}$. The elasticity of $f(x)$ is defined as

$$\mathcal{L}\, f(x) := \frac{x}{f(x)} \frac{d}{dx} f(x) \tag{3.33}$$

$$= \frac{d \log f(x)}{d \log x}, \quad \text{when } x > 0 \text{ and } f(x) > 0. \tag{3.34}$$

The second line can be interpreted as the percentage change of $f(x)$ over a percentage change of x. This is often used in statistics and economics. (It is an extension of the Euler dilation operator, $x \frac{d}{dx}$.)

To illustrate its property, if $f(x) \sim x^k$ locally, then $\mathcal{L}\, f(x) \approx k$. It informs *local degree of homogeneity* in the scaling analysis.

More generally, some algebraic rules of $\mathcal{L}$ are

- $\mathcal{L}\,[f(x)\, g(x)] = \mathcal{L}\, f(x) + \mathcal{L}\, g(x)$; multiplication becomes addition.
- $\mathcal{L}\,[f(g(x))] = \mathcal{L}\, g(x) \times [\mathcal{L}\, f](g(x))$; composition becomes multiplication.
- $\mathcal{L}\,(x^k) = k$; the trivial case is $\mathcal{L}\,(x) = 1$.
- $\mathcal{L}\,(e^{-x}) = -x$;
- $\mathcal{L}\,(\text{constant})$ is zero;

As an application, it is a good exercise to derive $\mathcal{L}\,[f((x/\sigma)^p)] = p\,[\mathcal{L}\, f]((x/\sigma)^p)$.

The recurrence relations of the Wright function, (3.2) and (3.3), can be rewritten using the $\mathcal{L}$ operator. They become two expressions of the elasticity of the Wright function.

Define the ratio of two Wright functions as

$$Q_{\lambda,\mu,\delta}(z) = \frac{W_{\lambda,\mu+\delta}(z)}{W_{\lambda,\mu}(z)}. \tag{3.35}$$

It follows immediately that (3.2) becomes

$$\lambda z\, Q_{\lambda,\mu,\lambda}(z) = Q_{\lambda,\mu,-1}(z) + 1 - \mu. \tag{3.36}$$

LEMMA 3.9. The elasticity of the Wright function is expressed by the following ratios:

$$\mathcal{L}\, W_{\lambda,\mu}(z) = \frac{1}{\lambda} Q_{\lambda,\mu,-1}(z) + \frac{1-\mu}{\lambda}, \tag{3.37}$$

$$\mathcal{L}\, W_{\lambda,\mu}(z) = z\, Q_{\lambda,\mu,\lambda}(z). \tag{3.38}$$

△

PROOF. The second line is straightforward from (3.3). The first line is derived from the second line by replacing the $z\, Q_{\lambda,\mu,\lambda}(z)$ term on the RHS with (3.36).

□

3.7. The Elasticity of the M-Wright Functions

What we are most interested in is the elasticity of $M_\alpha(x)$:

$$\mathcal{L}\, M_\alpha(x) = [\mathcal{L}\, W_{-\alpha,1-\alpha}](-x) \tag{3.39}$$

which is from (3.10). Note that $\mathcal{L}\,F_\alpha(x)$ is trivial if $\mathcal{L}\,M_\alpha(x)$ is known. This is due to (3.20), we have

$$\mathcal{L}\,F_\alpha(x) = \mathcal{L}\,M_\alpha(x) + 1. \tag{3.40}$$

However, $\mathcal{L}\,F_\alpha(x)$ has a representation that is more friendly to FCM. From (3.37),

$$\mathcal{L}\,F_\alpha(x) = [\mathcal{L}\,W_{-\alpha,0}](-x) = \frac{1}{\alpha}Q_\alpha(x) - \frac{1}{\alpha}, \tag{3.41}$$

$$\text{where} \quad Q_\alpha(x) := -Q_{-\alpha,0,-1}(-x) = -\frac{W_{-\alpha,-1}(-x)}{W_{-\alpha,0}(-x)}. \tag{3.42}$$

It follows that $Q_\alpha(x) = \alpha\,\mathcal{L}\,M_\alpha(x) + (1+\alpha)$.

The following lemma converts the elasticity of the FG PDF to either $\mathcal{L}\,M_\alpha(x)$ or $Q_\alpha(x)$.

LEMMA 3.10. Let $\mathfrak{N}(x)$ represent the functional form of FG PDF (6.1) where $\mathfrak{N}(x) = x^{d-1}F_\alpha\left(\left(\frac{x}{\sigma}\right)^p\right)$ (apart from a constant multiplier). The elasticity of $\mathfrak{N}(x)$ is

$$\mathcal{L}\,\mathfrak{N}(x) = p\,[\mathcal{L}\,M_\alpha]\big((x/\sigma)^p\big) + (d+p-1). \tag{3.43}$$

Alternatively, a useful ratio form for the FCM where p/α is a constant is

$$\mathcal{L}\,\mathfrak{N}(x) = \frac{p}{\alpha}\,Q_\alpha\left(\left(\frac{x}{\sigma}\right)^p\right) - \frac{p}{\alpha} + (d-1). \tag{3.44}$$

We observe that the role of the degrees of freedom d is very simple in $\mathcal{L}\,\mathfrak{N}(x)$. It shifts the constant level, but it does not affect the shape of $\mathcal{L}\,\mathfrak{N}(x)$.

△

$\mathcal{L}\,M_\alpha(x)$ has simple behaviors in a few cases. For example,

$$\mathcal{L}\,M_0(x) = -x;$$
$$\mathcal{L}\,M_{1/2}(x) = -x^2/2.$$

When $x \to 0$,

$$\mathcal{L}\,M_\alpha(x) \sim -b_1 x, \qquad \text{where } b_1 := \frac{\Gamma(1-\alpha)}{\Gamma(1-2\alpha)}. \tag{3.45}$$

When $\alpha \in [0, 1/2]$, $\mathcal{L}\,M_\alpha(x) < 0$ for all $x > 0$. It is a monotonically decreasing function for $x \in [0, \infty)$.

When $x \to \infty$, the GG-style asymptotic form in (3.16) leads to

$$\mathcal{L}\,M_\alpha(x) \sim -\alpha^{\alpha/(1-\alpha)}\,x^{1/(1-\alpha)} + \frac{\alpha - 1/2}{1-\alpha}, \tag{3.46}$$

in which the first term is dominant. It leads to the asymptotic limit of second-order elasticity:

$$\lim_{x\to\infty} \mathcal{L}[-\mathcal{L}\,M_\alpha](x) \to \frac{1}{1-\alpha}. \tag{3.47}$$

It follows immediately from (3.38) that (with $z \to -x$)

$$\mathcal{L}\,M_\alpha(x) = -x\,\frac{W_{-\alpha,1-2\alpha}(-x)}{W_{-\alpha,1-\alpha}(-x)} = -x\,Q_{-\alpha,1-\alpha,-\alpha}(-x) \tag{3.48}$$

where the series form of the numerator is in (3.19). We can compute the numerator and denominator individually, then take the ratio. Or we can derive its series representation as follows.

LEMMA 3.11. The series representation of $\mathcal{L}\,M_\alpha(x) = -x\,Q_{-\alpha,1-\alpha,-\alpha}(-x)$ is

$$\mathcal{L}\,M_\alpha(x) = \sum_{k=1}^{\infty} c_k x^k$$

where

$$c_k = \frac{(-1)^k}{(k-1)!}\, b_k + \sum_{j=1}^{k-1} \frac{(-1)^{(j+1)}}{j!}\, b_j\, c_{k-j}, \qquad k \geq 1; \tag{3.49}$$

$$b_n = \frac{\Gamma\big(1-\alpha\big)}{\Gamma\big(1-\alpha(n+1)\big)}, \qquad n \geq 1. \tag{3.50}$$

△

PROOF. From (3.14), we have

$$M_\alpha(x) = \sum_{n=0}^{\infty} a_n x^n, \qquad a_n = \frac{(-1)^n}{n!\,\Gamma\big(1-\alpha(n+1)\big)}.$$

Then (3.19) can be written as

$$\frac{d}{dx} M_\alpha(x) = -W_{-\alpha,1-2\alpha}(-x) = \sum_{n=1}^{\infty} n\, a_n x^{n-1}.$$

And

$$x\frac{M'_\alpha(x)}{M_\alpha(x)} = \frac{\sum\limits_{n\geq 1} n\, a_n\, x^n}{\sum\limits_{n\geq 0} a_n\, x^n}.$$

The coefficients satisfy the standard recurrence of series divisions, which becomes

$$c_k = \frac{1}{a_0}\left(k\, a_k - \sum_{j=1}^{k-1} a_j\, c_{k-j}\right), \qquad k \geq 1.$$

With $a_0 = \dfrac{1}{\Gamma(1-\alpha)}$, and $\dfrac{a_n}{a_0} = \dfrac{(-1)^n}{n!} b_n$, it leads to (3.49) and (3.50).

□

REMARK 3.12. The first three coefficients are explicitly derived as follows.

$$c_1 = -b_1 = -\frac{\Gamma(1-\alpha)}{\Gamma(1-2\alpha)},$$

$$c_2 = b_2 - b_1^2 = \frac{\Gamma(1-\alpha)}{\Gamma(1-3\alpha)} - \left(\frac{\Gamma(1-\alpha)}{\Gamma(1-2\alpha)}\right)^2,$$

$$c_3 = -\frac{1}{2}b_3 + \frac{3}{2}b_2 b_1 - b_1^3 = -\frac{\Gamma(1-\alpha)}{2\,\Gamma(1-4\alpha)} + \frac{3}{2}\,\frac{\Gamma(1-\alpha)^2}{\Gamma(1-2\alpha)\,\Gamma(1-3\alpha)} - \left(\frac{\Gamma(1-\alpha)}{\Gamma(1-2\alpha)}\right)^3.$$

The small-x expansion up to the x^3 term is

$$\mathcal{L}\, M_\alpha(x) = \left[-\frac{\Gamma(1-\alpha)}{\Gamma(1-2\alpha)}\right] x + \, c_2 x^2 + \, c_3 x^3 + O(x^4).$$

3.8. Numerical Methods of the M-Wright Functions

To properly compute the subsequent special functions and distributions in this book, we need a very robust numerical implementation of $F_\alpha(x)$ and $M_\alpha(x)$ for *the entire range* of $\alpha \in [0,1]$ and $x \geq 0$. Since $F_\alpha(x) = \alpha x\, M_\alpha(x)$, we can easily compute one from the other in most cases. It is a matter of which approach is faster, more convenient, and precise.

3.8.1. Handling alpha for zero and one. When $\alpha = 0$, we should use $M_0(x) = e^{-x}$. $\lim_{\alpha\to 0} F_\alpha(x)$ should be handled carefully in the fractional gamma distribution.

When $\alpha = 1$, we could use a normal distribution to simulate the delta function: $M_1(x) = \mathcal{N}(x; 1, \sigma^2)$ where $\sigma = 0.001$. This is to ensure that $\int_0^\infty M_1(x)\,dx = 1$.

3.8.2. Using scipy.stats.levy-stable. Both functions can be derived from the one-sided α-stable distribution $L_\alpha(x)$ of Section 4.2, which is implemented in `scipy.stats.levy_stable` package[42].

For example, due to (3.21), $M_\alpha(x)$ can be computed using $L_\alpha(x) = \alpha x^{-\alpha-1} M_\alpha(x^{-\alpha})$ where $x > 0$. On the other hand, for $\beta > 1/2$, we can also use $M_\beta(x) = \alpha L_\alpha^{\alpha-2}(x)$ where $\beta = 1/\alpha$.

These two numerical methods are good for the bulk of α and x. However, they begin to lose precision for small $\alpha < 0.08$ and large $\alpha > 0.99$. They are also not good enough for small $x < 0.01$.

3.8.3. Using the series sum in numpy. Based on (3.14), we define the sum of the series of finite terms as

$$M_\alpha^{(m)}(x) = \frac{1}{\pi}\sum_{n=1}^{m} \frac{(-x)^{n-1}}{(n-1)!}\Gamma(\alpha n)\sin(\alpha n\pi). \qquad (0 < \alpha < 1) \tag{3.51}$$

This method implemented in `numpy` and `scipy` is good for several scenarios. First, to cover the small x area ($x < 0.01$), use $M_\alpha^{(7)}(x)$ if $\alpha < 0.9$.

Otherwise, we could use $M_\alpha^{(80)}(x)$ for $\alpha \le 0.998$ and $x < 0.85$. The sum of 80 terms takes more time to compute. But it is a necessary path when `scipy.stats.levy_stable` approach loses precision.

3.8.4. Using the series sum in mpmath. $M_\alpha^{(m)}(x)$ implemented in `mpmath` is our de facto implementation to calibrate the precision of other approaches. In order to make it a good baseline implementation, we must carefully choose `mp.prec` and m to use.

After rigorous testing, it was found that `mp.prec` >= 64 provides sufficient precision. Therefore, `mp.prec` = 128 is more than abundant up to three decimal points in α. `mpmath` is smart about handling summing many small terms, especially with large amount of cancellation due to the $\sin(\alpha n\pi)$ factor in (3.51). This feature can be used during the calibration phase.

The more important numerical choice is the truncation level m. In practice, m = 40,000 is sufficient for $\alpha < 0.9$, whereas a much larger value, m = 80,000, is needed when α is very close to 1, such as $\alpha = 0.998$. Larger values of m substantially increase the computational cost of the series summation. This approach is therefore appropriate during calibration, but it is too expensive for the final `NumPy`-style implementation.

This will be elaborated on in the next section.

3.8.5. Using the asymptotic approximation. Paris et al. [32] derived a more refined asymptotic formula, where (3.16) is simply its first term. Theorem 2.2 of that paper is recaptured in the following.

LEMMA 3.13.

$$M_\alpha(x) \sim \frac{A(\alpha)}{2\pi} X^{\alpha-1/2} e^{-X} \sum_{n=0}^{\infty} c_j(\alpha)(-X)^{-j}. \qquad (0 < \alpha < 1) \tag{3.52}$$

where $c_j(\alpha)$ is in its (2.4) up to $j = 6$. Other parameters are $A(\alpha) = \sqrt{\frac{2\pi}{\alpha}}\left(\frac{\alpha}{\kappa}\right)^\alpha$ and $X = \kappa(hx)^{1/\kappa}$ with $\kappa = 1 - \alpha$ and $h = \alpha^\alpha$.

△

When $M_\alpha(x)$ is small, (3.52) could be very precise with an error as small as 10^{-5}. Our strategy is to use other implementations to get $M_\alpha(x)$ to a small number, e.g. 10^{-6} in most cases, and at least 10^{-3} in some difficult cases. Then use (3.52) for larger x up to infinity (the maximum 64-bit float).

This right-tail strategy works for the bulk of α from 0.1 to 0.9. The transition interval (defined as $M_\alpha(x) \in [10^{-5}, 10^{-6}]$) could be precomputed by the faster (3.16).

For α from 0.9 to 0.99, the `mpmath` version of $M_\alpha^{(m)}(x)$ is more precise to determine the transition interval.

For α from 0.001 to 0.1, the asymptotic form is adjusted to

$$M_\alpha(x) \sim A'(\alpha)\, e^{-B'(\alpha)X'}, \quad \text{where } X' = x^{1/\kappa}. \tag{3.53}$$

$A'(\alpha)$ and $B'(\alpha)$ are obtained from a linear regression in the transition interval: $\log M_\alpha(x) \sim \log A'(\alpha) - B'(\alpha)X'$.

3.8.6. Using the integral form. For α in the range from 0.99 to 0.998, the `scipy` implementation of $M_\alpha(x)$ loses precision rapidly in the right tail. In this regime, we use (3.7) to supplement the computation as long as $M_\alpha(x) > 10^{-3}$.

The main numerical difficulty arises when the target value of $M_\alpha(x)$ is very small. In that case, the integrand in (3.7) becomes highly oscillatory and is effectively nonzero only over a very narrow range of r^δ. Standard integration algorithms have difficulty identifying this small interval, resolving the oscillations, and performing the required cancellation accurately. A more sophisticated quadrature method would be needed, and this is left for future research.

For $M_\alpha(x) < 10^{-3}$ at large x, we still use (3.52) asymptotically. This equation is fine for large α, as long as the numeric overflow is handled properly.

CHAPTER 4

The Alpha-Stable Distribution - Review

The two-sided distributions developed in this book are built on the α-stable distribution, introduced in Paul Lévy's 1925 monograph[17]. Among its parameters, the central one is the *stability index* $\alpha \in (0, 2]$, which we interpret throughout the book as the principal *fractional* parameter.

This chapter reviews the α-stable distribution from the Mellin transform perspective. That viewpoint is used repeatedly in the later construction of fractional extensions. The ratio-distribution representation of the density in Section 4.3 is due to the author.

4.1. Classic Result

The α-stable distribution has two shape parameters. There are many parameterizations that have been studied (see p.5 of [30]). We are primarily concerned with Feller's (α, θ) parameterization [12, 13], where α is called the stability index with a range of $0 < \alpha \leq 2$, and θ is an angle that injects skewness into the distribution when it is not zero [1]

An innovative approach is to study its Mellin transform. This presentation is used because it is *simpler* and provides great insight into its structure.

LEMMA 4.1. The Mellin transform of its PDF is

$$L_\alpha^\theta(x) \overset{\mathcal{M}}{\longleftrightarrow} \epsilon \frac{\Gamma(s)\Gamma(\epsilon(1-s))}{\Gamma(\gamma(1-s))\Gamma(1-\gamma+\gamma s)} \tag{4.2}$$

$$\text{where } \epsilon = \frac{1}{\alpha}, \gamma = \frac{\alpha - \theta}{2\alpha}.$$

where $0 < C < 1$ implicitly. This is defined for $x \geq 0$. The reflection rule is used for $x < 0$ such that $L_\alpha^\theta(x) := L_\alpha^{-\theta}(-x)$.

△

This formula was first derived by Schneider in 1986[37], rediscovered by Mainardi et al. in 2001[23], and later summarized by Mainardi and Pagnini in (2.8) of [24].

Equation (4.2) is written not directly in (α, θ), but in the auxiliary variables (ϵ, γ). This reparameterization is especially convenient in Mellin space, where the transform takes a more symmetric form.

In the Feller parameterization, the admissible range $|\theta| \leq \min\{\alpha, 2 - \alpha\}$ is known as the Feller-Takayasu diamond. In the (ϵ, γ) variables, the same constraint becomes (a) $0 \leq \gamma \leq 1$ when $\epsilon > 1$, and (b) $1 - \epsilon \leq \gamma \leq \epsilon$ when $\epsilon \leq 1$.[2]

---

[1] The conversion from/to the S1 parameterization is summarized as

$$\beta \tan\left(\frac{\alpha\pi}{2}\right) = -\tan\left(\frac{\theta\pi}{2}\right), \qquad \text{scale} = \cos^{1/\alpha}\left(\frac{\theta\pi}{2}\right). \tag{4.1}$$

where $\beta \in [-1, 1]$ is the standard skew parameter implemented in `scipy.stats.levy_stable`.

[2] Conversely, if γ is fixed, condition (b) implies $\alpha \leq \min\{1/\gamma, 1/(1-\gamma)\}$.

4.1.1. The Reflection Rule. Note that the reflection of $\theta \to -\theta$ in the (α, θ) parametrization is equivalent to the reflection of $\gamma \to 1-\gamma$ in the (ϵ, γ) parametrization.

Since we often mingle the two parameterizations, this alternative view can be very helpful in certain scenarios. For example, the total density in the positive domain is $\int_0^\infty L_\alpha^\theta(x) = \gamma$. By the reflection rule, $\int_0^\infty L_\alpha^{-\theta}(x) = 1-\gamma$. Hence, the total density $\int_{-\infty}^\infty L_\alpha^\theta(x) = \gamma + (1-\gamma) = 1$.

4.2. Extremal Distributions

There are two extremal regimes in which the skew parameter θ reaches the boundary of its admissible range. These cases are important because the M-Wright functions $F_\alpha(x)$ and $M_\alpha(x)$ from Section 3.3 arise naturally from them.

The first extremal regime corresponds to $\gamma = 0$ or $\gamma = 1$, or equivalently $\theta = \pm\alpha \le 1$. By the reflection rule, it is sufficient to study the case $\theta = -\alpha$, or $\gamma = 1$.

This yields the one-sided α-stable distribution:

$$L_\alpha(x) := L_\alpha^{-\alpha}(x) \overset{\mathcal{M}}{\longleftrightarrow} \epsilon \, \frac{\Gamma(\epsilon(1-s))}{\Gamma(1-s)}$$

In the literature [8, 9], this distribution is often characterized by the Laplace transform

$$\mathbb{E}\left[e^{-tL_\alpha}\right] = e^{-t^\alpha}, \quad t \ge 0.$$

Remark 4.2. Starting from (3.12) where $F_\alpha(x) \overset{\mathcal{M}}{\longleftrightarrow} F_\alpha^*(s) = \Gamma(s)/\Gamma(\alpha s)$, and applying three Mellin-transform operations, namely $x \to x^\alpha$, multiplication by x, and then $x \to x^{-1}$, we recover the classical identity

$$L_\alpha(x) = x^{-1} F_\alpha(x^{-\alpha}) \quad (x \ge 0 \text{ and } 0 < \alpha \le 1) \tag{4.3}$$

with the endpoint case $L_1(x) = \delta(x-1)$. This leads to the important random variable relation of $M_\alpha \sim (L_\alpha)^{-\alpha}$ in (3.21).

Remark 4.3. $L_\alpha(x)$ can be computed with `scipy.stats.levy_stable`[42] in the S1 parameterization by setting `beta`=1 and `scale`=$\cos(\alpha\pi/2)^{1/\alpha}$ for $0 < \alpha < 1$.[3] This provides a practical numerical starting point for developing the new fractional distributions and testing the constructions in proof-of-concept form.

The second extremal regime occurs when $\theta = \alpha - 2$. In this case $\epsilon = \gamma = 1/\alpha$, and

$$L_\alpha^{\alpha-2}(x) \overset{\mathcal{M}}{\longleftrightarrow} \epsilon \, \frac{\Gamma(s)}{\Gamma(1-\epsilon+\epsilon s)}$$

Comparing this expression with (3.13) gives the classical identity: (See, for example, formula (F.49) in [21])

$$L_\alpha^{\alpha-2}(x) = \frac{1}{\alpha} M_{1/\alpha}(x) \quad (x \in \mathbb{R} \text{ and } 1 < \alpha \le 2) \tag{4.4}$$

This identity extends the M-Wright function to $x < 0$, because $L_\alpha^{\alpha-2}(x)$ is two-sided.

4.3. Ratio Distribution Approach

Important insight can be obtained by interpreting (4.2) as a ratio distribution (2.6). We split (4.2) into two components:

$$L_\alpha^\theta(x) \overset{\mathcal{M}}{\longleftrightarrow} \epsilon \left[\frac{\Gamma(s)}{\Gamma(1-\gamma+\gamma s)}\right] \left[\frac{\Gamma(\epsilon(1-s))}{\Gamma(\gamma(1-s))}\right] \tag{4.5}$$

The first bracket is the Mellin transform of the M-Wright function (3.13).

[3] See Chapter 1 of [30] for further discussion of stable-law parameterizations. We do not pursue that issue here.

The second bracket is related to the Mellin transform of the fractional χ-mean distribution (FCM) at $k = 1$:

$$\overline{\chi}^{\theta}_{\alpha,1}(x) \overset{\mathcal{M}}{\longleftrightarrow} \overline{\chi}^{\theta}_{\alpha,1}{}^{*}(s) = \epsilon\gamma^{\gamma(s-1)-1}\,\frac{\Gamma(\epsilon(s-1))}{\Gamma(\gamma(s-1))} \tag{4.6}$$

For a ratio distribution, the Mellin rule requires replacing s with $2 - s$ in $\overline{\chi}^{\theta}_{\alpha,1}{}^{*}(s)$. Consequently, the factor $s - 1$ in the second line of (4.6) becomes $1 - s$ in the second bracket of (4.5).

4.3.1. Rescaled M-Wright Function. Additionally, a small nuance here is to deal with scaling factors. Define the rescaled M-Wright function

$$\tilde{M}_\gamma(x) := \gamma^{1-\gamma} M_\gamma(x/\gamma^\gamma), \quad x \geq 0 \tag{4.7}$$

$$\tilde{M}_\gamma(-x) := \tilde{M}_{1-\gamma}(x) \qquad \text{(reflection rule)}. \tag{4.8}$$

Since $\int_0^\infty \tilde{M}_\gamma(x)\,dx = \gamma$ and $\int_{-\infty}^0 \tilde{M}_\gamma(x)\,dx = \int_0^\infty \tilde{M}_{1-\gamma}(x)\,dx = 1 - \gamma$, we have $\int_0^\infty M_\gamma(x) = 1$. Hence, $\tilde{M}_\gamma(x)$ is a valid two-sided density function.

The scale is adjusted so that it matches the standard normal distribution: $\tilde{M}_{1/2}(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$ of $\mathcal{N}(0, 1)$.

According to (2.3), the rescaling of PDF modifies the Mellin transform from (3.13) to

$$\tilde{M}_\gamma(x) \overset{\mathcal{M}}{\longleftrightarrow} \tilde{M}^*_\gamma(s) = \gamma^{1-\gamma+\gamma s}\,\frac{\Gamma(s)}{\Gamma(1-\gamma+\gamma s)} \tag{4.9}$$

from which the $\gamma^{1-\gamma+\gamma s}$ term cancels out its counterpart in $\overline{\chi}^{\theta}_{\alpha,1}{}^{*}(2 - s)$ nicely.

Therefore, we find a new method to construct the α-stable distribution using the following integral.

LEMMA 4.4 (The ratio-distribution representation of the α-stable distribution). The Mellin transform of the PDF (4.2) becomes

$$L^\theta_\alpha(x) \overset{\mathcal{M}}{\longleftrightarrow} \tilde{M}^*_\gamma(s)\,\overline{\chi}^{\theta}_{\alpha,1}{}^{*}(2 - s) \tag{4.10}$$

from which the PDF can be written in a ratio distribution form of

$$L^\theta_\alpha(x) := \int_0^\infty \tilde{M}_\gamma(xs)\,\overline{\chi}^{\theta}_{\alpha,1}(s)\,s\,ds \quad (x \geq 0) \tag{4.11}$$

Since the Mellin integral is only valid for $x > 0$, it is supplemented with *the reflection rule*:

$$L^\theta_\alpha(-x) := L^{-\theta}_\alpha(x) \tag{4.12}$$

△

4.3.2. Ratio of Stable Law Approach. It can be expressed elegantly in terms of the stable law. The normalization term γ^γ cancels out between $\tilde{M}_\gamma$ and $\overline{\chi}^{\theta}_{\gamma\alpha,1}$ (See Section 7.2.1). For $x \geq 0$, we obtain the following law of ratio:

$$L^\theta_\alpha \sim M_\gamma / T^{-\gamma}_{\gamma\alpha,\,0} = (L_{\gamma\alpha}/L_\gamma)^\gamma \tag{4.13}$$

for the γ portion of the population. For $x < 0$, the law is modified by replacing γ with $1 - \gamma$ for the $(1 - \gamma)$ portion of the population due to the reflection rule.

This construction places $\overline{\chi}^{\theta}_{\alpha,1}$ in the central role. We define it at one degree of freedom $k = 1$. In Chapter 7, we will add *degrees of freedom* k to it and make it $\overline{\chi}^{\theta}_{\alpha,k}$, which is the fractional extension of the classic χ distribution.

Subsequently, in Chapter 11, we will add *degrees of freedom* k to the α-stable distribution and merge it with Student's t distribution.

4.4. SaS

Note that $\theta = 0$ is equivalent to $\gamma = 1/2$. The distribution is symmetric, with the nickname of "SaS", which stands for "Symmetric α-Stable".

Its Mellin transform is simplified to

$$\begin{aligned} L^0_\alpha(x) &\overset{\mathcal{M}}{\longleftrightarrow} \epsilon \left[\frac{\Gamma(s)}{\Gamma((1+s)/2)}\right] \left[\frac{\Gamma(\epsilon(1-s))}{\Gamma((1-s)/2)}\right] \\ &= \epsilon \left[\frac{2^{s-1}}{\sqrt{\pi}}\Gamma\left(\frac{s}{2}\right)\right] \left[\frac{\Gamma(\epsilon(1-s))}{\Gamma((1-s)/2)}\right]. \end{aligned} \tag{4.14}$$

The first bracket is the Mellin transform of a normal distribution (2.9) with a scale. The second bracket is ${\overline{\chi}^{0}_{\alpha,1}}^{*}(2-s)$ from above.

Hence, the PDF of SaS is

$$L^0_\alpha(x) = \int_0^\infty \mathcal{N}(xs)\, \overline{\chi}^0_{\alpha,1}(s)\, s\, ds. \tag{4.15}$$

This is one of the foundations of GAS-SN in (12.1).

4.4.1. Method of Normal Mixture. SaS in (4.15) will be generalized to GSaS in (12.3) in Chapter 12. Both integrals are in the normal mixture structure (9.1) that enjoys several nice properties described in Chapter 9.

The classic exponential power distribution (Section 3.11.1 of [**30**]) is the characteristic function transform in Lemma 9.2.

CHAPTER 5

Fractional Hypergeometric Functions

This chapter develops fractional versions of two classical hypergeometric functions: the confluent hypergeometric function ${}_1F_1(a,b;x)$, also denoted by $M(a,b;x)$ (Chapter 13 of DLMF[10]), and the Gauss hypergeometric function ${}_2F_1(a,b,c;x)$ (Chapter 15 of DLMF).

The confluent case appears in the CDFs of the FG and FCM distributions, whereas the Gauss case appears in the CDFs of the GSaS and F distributions. We therefore first record the relevant DLMF formulas and then convert them to the Mellin-transform convention used in (2.2).

From DLMF 13.2.4 and 13.4.16, the Mellin transform of the Kummer function is

$$M(a,b;-x) = \frac{\Gamma(b)}{2\pi i\,\Gamma(a)} \int_{C-i\infty}^{C+i\infty} \frac{\Gamma(a-s)\Gamma(s)}{\Gamma(b-s)}\, x^{-s}\, ds,$$

where $a \neq 0, -1, -2, \ldots$.

From DLMF 15.1.2 and 15.6.6, the Mellin transform of the Gauss hypergeometric function is

$${}_2F_1(a,b,c;-x) = \frac{\Gamma(c)}{2\pi i\,\Gamma(a)\Gamma(b)} \int_{C-i\infty}^{C+i\infty} \frac{\Gamma(a-s)\Gamma(b-s)\Gamma(s)}{\Gamma(c-s)}\, x^{-s}\, ds,$$

where $a, b \neq 0, -1, -2, \ldots$.

In the Mellin-transform notation used in this book, these formulas become

$$M(a,b;-x) \overset{\mathcal{M}}{\longleftrightarrow} M^*(a,b;s) = \frac{\Gamma(b)}{\Gamma(a)} \frac{\Gamma(a-s)\Gamma(s)}{\Gamma(b-s)}, \tag{5.1}$$

$${}_2F_1(a,b,c;-x) \overset{\mathcal{M}}{\longleftrightarrow} {}_2F_1^*(a,b,c;s) = \frac{\Gamma(c)}{\Gamma(a)\Gamma(b)} \frac{\Gamma(a-s)\Gamma(b-s)\Gamma(s)}{\Gamma(c-s)}. \tag{5.2}$$

We now introduce the corresponding fractional extensions.

5.1. Fractional Confluent Hypergeometric Function

The fractional confluent hypergeometric function (FCHF) is the union of the Kummer function and the Wright function. It allows us to extend many classic functions to their fractional forms.

We start with its Mellin transform. And we follow with the integral and series representations.

Definition 5.1. The Mellin transform of the FCHF is

$$M_{\lambda,\delta}(a,b;-x) \overset{\mathcal{M}}{\longleftrightarrow} M^*_{\lambda,\delta}(a,b;s) = \frac{\Gamma(b)}{\Gamma(a)} \frac{\Gamma(a-s)\Gamma(s)}{\Gamma(\delta-\lambda s)\Gamma(b-s)} \tag{5.3}$$

where the $\Gamma(\delta-\lambda s)$ term is from the Wright function (3.5).

Lemma 5.2. The integral representation from DLMF 13.4.1 is extended to

$$M_{\lambda,\delta}(a,b;z) = \frac{\Gamma(b)}{\Gamma(a)\Gamma(b-a)} \int_0^1 W_{\lambda,\delta}(zt)\, t^{a-1}(1-t)^{b-a-1}\, dt \tag{5.4}$$

The obvious limit $W_{0,1}(zt) = e^{zt}$ restores it to the classic DLMF formula.

△

PROOF. Replace the Wright function in (5.4) with its Hankel integral (3.6),

$$M_{\lambda,\delta}(a,b;z) = \frac{\Gamma(b)}{2\pi i\,\Gamma(a)} \int_0^1 \int_{Ha} \left(\frac{e^{s+zt\,s^{-\lambda}}}{s^{\delta}} ds \right) t^{a-1}(1-t)^{b-a-1}\,dt$$

which can be simplified to

$$M_{\lambda,\delta}(a,b;z) = \frac{1}{2\pi i} \int_{Ha} \left(s^{-\delta}\, e^{s}\, ds \right)\; M(a,b;-z\,s^{-\lambda})$$

Substitute the Mellin integral from (5.1) to it,

$$\begin{aligned} M_{\lambda,\delta}(b,c;-z) &= \frac{1}{2\pi i} \int_{Ha} \left(s^{-\delta}\, e^{s}\, ds \right)\; \frac{\Gamma(b)}{2\pi i\Gamma(a)} \int_{C-i\infty}^{C+i\infty} \frac{\Gamma(b-t)\Gamma(t)}{\Gamma(c-t)}\,(z\,s^{-\lambda})^{-t}\,dt \\ &= \frac{\Gamma(b)}{2\pi i\Gamma(a)} \int_{C-i\infty}^{C+i\infty} \left[\frac{1}{2\pi i} \int_{Ha} s^{\lambda t-\delta}\, e^{s}\, ds \right] \frac{\Gamma(b-t)\Gamma(t)}{\Gamma(c-t)}\, z^{-t}\,dt \\ &= \frac{\Gamma(b)}{2\pi i\Gamma(a)} \int_{C-i\infty}^{C+i\infty} \left[\frac{1}{\Gamma(\delta-\lambda t)} \right] \frac{\Gamma(b-t)\Gamma(t)}{\Gamma(c-t)}\, z^{-t}\,dt \end{aligned}$$

which is the Mellin transform in (5.3).

From the second line to the third line, we use the well-known Hankel integral of the reciprocal gamma function:

$$\frac{1}{2\pi i} \int_{Ha} s^{-z}\, e^{s}\, ds = \frac{1}{\Gamma(z)}$$

□

LEMMA 5.3. The series representation is

$$M_{\lambda,\delta}(a,b;z) := \sum_{n=0}^{\infty} \left[\frac{(a)_n}{(b)_n\,\Gamma(\lambda n+\delta)} \right] \frac{z^n}{n!} \tag{5.5}$$

where $(a)_n, (b)_n$ are Pochhammer symbols.

△

PROOF. Take (5.3) and apply Ramanujan's master theorem from Section 2.3. This produces $(M^{*}_{\lambda,\delta}(a,b;s)/\Gamma(s))|_{s=-n}$, which is equal to the bracket term, since $(x)_n = \Gamma(x+n)/\Gamma(x)$. □

5.1.1. FCHF Subsumes the Kummer Function. It is obvious that $M_{0,1}(a,b;x) = M(a,b;x)$.

5.1.2. FCHF Subsumes the M-Wright Function. By using the same setting from (3.10), we get

$$M_{\alpha}(z) = M_{-\alpha,1-\alpha}(c,c;-z) \qquad (c \neq 0)$$

5.1.3. FCHF Subsumes Fractional Gamma-Star Function. An important variant of FCHF is the fractionalization of the incomplete gamma function. The reader is referred to Sections 8 and 13 of DLMF[**10**] and Wikipedia for background information.

We are mainly concerned with the following setup:

$$M_{-\alpha,1-\alpha}(c,c+1;-x) = c \int_0^1 M_{\alpha}(xt)\,t^{c-1}\,dt$$

This integral is found in (3.23). Hence, we obtain -

LEMMA 5.4. The fractional γ^* function (3.23) has the following FCHF representation:

$$\gamma^*_\alpha(s,x) = \frac{\Gamma(\alpha s-\alpha+1)}{\Gamma(s+1)}\, M_{-\alpha,1-\alpha}(s,s+1;-x) \tag{5.6}$$

△

The fractional γ^* function is the basis for expressing the CDF of the fractional gamma distribution in Section 6.7. In fact, this was the main motivation to enrich the classic confluent hypergeometric function.

5.2. Fractional Gauss Hypergeometric Function

The fractional Gauss hypergeometric function (FGHF) arises from the ratio distribution between an elementary function and FCM2 ($\widehat{\chi}^2_{\alpha,k}$) in Section 7.8.

When $\alpha = 1$, the Mellin transform of FCM2 is reduced from a fractional form to a classic form in (7.72). The ratio distribution is reduced to a Gauss hypergeometric function ${}_2F_1$. Hence, we consider the general form of such a ratio distribution as fractional ${}_2F_1$.

We start by modifying the Mellin transform from (5.2) (DLMF 15.6.6). Then we derive the integral and series representations from it.

DEFINITION 5.5. The Mellin transform of the fractional Gauss hypergeometric function is

$$\begin{aligned} {}_2F_1(a,b,c,\epsilon;-x) &\overset{\mathcal{M}}{\longleftrightarrow} {}_2F_1^*(a,b,c,\epsilon;s) \\ &= \Big[M^*\left(a,c;s\right)\Big]\,\left[\frac{B(k/2,1/2)}{\Gamma(1/2)}\,\widehat{\chi}^{2*}_{\alpha,k}(3/2-s)\right] \end{aligned} \tag{5.7}$$

where $\epsilon = 1/\alpha$ is the convention from (4.2), and $b = (k+1)/2$. $M^*\left(a,c;s\right)$ is from (5.1), and $\widehat{\chi}^{2*}_{\alpha,k}(s)$ is from (7.71) (we jump ahead). And $B(x,y)$ is the beta function.

This structure is a fractional form of the generalized hypergeometric function ${}_3F_2$ (DLMF 16.5.1, replace s with $-s$). To see this, expand (5.7) and we get

$${}_2F_1^*(a,b,c,\epsilon;s) = \left[\frac{\Gamma(c)}{\Gamma(a)}\frac{\Gamma(a-s)\Gamma(s)}{\Gamma(c-s)}\right]\left[2^{2s-1}\,\frac{B(k/2,1/2)}{\Gamma(1/2)}\,\frac{\Gamma((k-1)/2)}{\Gamma(\epsilon(k-1))}\frac{\Gamma(2\epsilon\,(k/2-s))}{\Gamma(k/2-s)}\right]. \tag{5.8}$$

There are five gamma functions that contain s: three in the numerator, two in the denominator. And the $\Gamma(2\epsilon\,(k/2-s))$ term is fractional.

5.2.1. FGHF Subsumes the Gauss Hypergeometric Function.

LEMMA 5.6. When $\epsilon = 1$,

$${}_2F_1^*(a,b,c,\epsilon=1;s) = {}_2F_1^*(a,b,c;s)$$

△

PROOF. Let $\epsilon = 1$, the second bracket becomes

$$\frac{B(k/2,1/2)}{\Gamma(1/2)}\frac{\Gamma(k/2+1/2-s)}{\Gamma(k/2)} = \frac{\Gamma(k/2+1/2-s)}{\Gamma(k/2+1/2)} = \frac{\Gamma(b-s)}{\Gamma(b)}. \tag{5.9}$$

Hence, (5.7) is reduced to the classic limit of ${}_2F_1^*(a,b,c;s)$ in (5.2). □

5.2.2. The Integral Form.

LEMMA 5.7. The integral form of FGHF is

$$ {}_2F_1(a,b,c,\epsilon;-x) := \frac{B(k/2,1/2)}{\Gamma(1/2)} \int_0^\infty M(a,c;-x\nu)\, \widehat{\chi}^2_{\alpha,k}(\nu)\, \sqrt{\nu} d\nu \tag{5.10} $$

where $\epsilon = 1/\alpha$ and $b = (k+1)/2$. $M\,(a,c;x)$ is the Kummer function (Chapter 13, DLMF). $\widehat{\chi}^2_{\alpha,k}(x)$ is from (7.63).

△

PROOF. We use the generalized convolution formula:

$$ h(x) = \int_0^\infty f(xs) g(s)\, s^p\, ds \overset{\mathcal{M}}{\longleftrightarrow} h^*(s) = f^*(s) g^*(1+p-s), $$

Clearly f is M, and g is $\widehat{\chi}^2_{\alpha,k}$. Substitute $p = 1/2$ due to the $\sqrt{\nu}$ term. The Mellin transform of (5.10) is

$$ {}_2F_1(a,b,c,\epsilon;-x) \overset{\mathcal{M}}{\longleftrightarrow} \frac{B(k/2,1/2)}{\Gamma(1/2)} M^*(a,c;s)\, \widehat{\chi}^{2*}_{\alpha,k}(3/2-s) $$

This is exactly (5.7).

□

5.2.3. Relation between FGHF and Real-World Usage. This section explains how FGHF connects to the later FCM, GAS, and GAS-SN constructions. The connection is most transparent at the level of ratio-distribution integrals: many of the univariate PDFs used later can be viewed as specializations of the FGHF integral representation.

To isolate this structure, suppress the parameters that are not needed for the present comparison. Define $F(-x) := {}_2F_1(a,b,c,\epsilon;-x)$ and $M(-x) := M^*(a,c;-x)$ for $x \geq 0$. Then (5.10) becomes

$$ F(-x) := B \int_0^\infty M(-x\nu)\, \overline{\chi}^2_{\alpha,k}(\nu;\sigma=1/4)\, \sqrt{\nu} d\nu \tag{5.11} $$

where we employ the notation $\widehat{\chi}^2_{\alpha,k}(x) = \overline{\chi}^2_{\alpha,k}(x;\sigma=\frac{1}{4})$ from (7.63), and $B := B(\frac{k}{2},\frac{1}{2})/\Gamma(\frac{1}{2})$.

LEMMA 5.8. Let $F'(-x)$ denote the scaled FGHF used in the later ratio-distribution applications. The following ratio-distribution integrals can be expressed in terms of F':

$$ \left.\begin{array}{l} \int_0^\infty M(-xs)\, \overline{\chi}^2_{\alpha,k}(s)\, \sqrt{s} ds \\ \int_0^\infty M(-xs^2)\, \overline{\chi}_{\alpha,k}(s)\, s\, ds \end{array}\right\} = F'(-x) := \frac{2\sqrt{\pi}\,\sigma_{\alpha,k}}{B(\frac{k}{2},\frac{1}{2})}\, F\left(-4\sigma^2_{\alpha,k}\, x\right) \tag{5.12} $$

Equivalently, in the full FGHF notation,

$$ \left.\begin{array}{l} \int_0^\infty M(a,c;-xs)\, \overline{\chi}^2_{\alpha,k}(s)\, \sqrt{s} ds \\ \int_0^\infty M(a,c;-xs^2)\, \overline{\chi}_{\alpha,k}(s)\, s\, ds \end{array}\right\} = F'_{\alpha,k}(a,c;-x) := \frac{2\sqrt{\pi}\,\sigma_{\alpha,k}}{B(\frac{k}{2},\frac{1}{2})}\, {}_2F_1\left(a,b,c,\epsilon;-4\sigma^2_{\alpha,k}\, x\right) \tag{5.13} $$

where $\epsilon = 1/\alpha$ and $b = (k+1)/2$ on the RHS.

△

PROOF. Let Q be the scale that we want to solve. (5.11) is rewritten to $F'(-x)$ such that

$$ F'(-x) := \frac{\sqrt{Q}}{B} F\left(-Qx\right) = \sqrt{Q} \int_0^\infty M(-Qx\nu)\, \overline{\chi}^2_{\alpha,k}(\nu;\sigma=1/4)\, \sqrt{\nu} d\nu. $$

Let $s = Q\nu$,

$$F'(-x) = \int_0^\infty M(-xs)\,\overline{\chi}^2_{\alpha,k}(s/Q;\sigma = 1/4)/Q\,\sqrt{s}ds$$
$$= \int_0^\infty M(-xs)\,\overline{\chi}^2_{\alpha,k}(s;\sigma = Q/4)\,\sqrt{s}ds$$

Let $Q = 4\sigma^2_{\alpha,k}$, we obtain the integral form in terms of FCM2,

$$F'(-x) = \int_0^\infty M(-xs)\,\overline{\chi}^2_{\alpha,k}(s)\,\sqrt{s}ds$$

This is the first line of (5.12). Then apply (7.65) and (7.66) to get the second line. And on the FGHF side, we have

$$F'(-x) = \frac{\sqrt{Q}}{B}F\left(-Qx\right) = \frac{2\sqrt{\pi}\,\sigma_{\alpha,k}}{B(\frac{k}{2},\frac{1}{2})}\,F(-4\sigma^2_{\alpha,k}\,x)$$

□

5.2.4. Example 1: GSaS. In Lemma 8.3 of [20], a fractional extension was explored for the CDF of GSaS. We formalized it further here. However, we note that the $M(-x)$ function needed to describe GAS-SN is more complicated than a Kummer function. See (10.2) and (10.3).

LEMMA 5.9. Assume $\Phi[L_{\alpha,k}](x)$ is the CDF of a GSaS, which is (12.2) with $\beta = 0$. It can be expressed by the scaled FGHF via

$$\Phi[L_{\alpha,k}](x) = \frac{1}{2} + \frac{x}{\sqrt{2\pi}}\,F'_{\alpha,k}\left(\frac{1}{2},\frac{3}{2};-\frac{x^2}{2}\right). \tag{5.14}$$

△

PROOF. From Lemma 8.3 of [20],

$$\Phi[L_{\alpha,k}](x) = \frac{1}{2} + \frac{x}{\sqrt{k}}M_{\alpha,k}\left(a,c;-\frac{x^2}{k}\right),$$

where $a = \frac{1}{2}, c = \frac{3}{2}$ and

$$M_{\alpha,k}(a,c;x) := \sqrt{\frac{k}{2\pi}}\int_0^\infty s\,ds\,M\left(a,c;\frac{xks^2}{2}\right)\,\overline{\chi}_{\alpha,k}(s).$$

This pattern fits right in with the second line of (5.13). It is immediately clear that its $M_{\alpha,k}(a,c;x)$ is our $\sqrt{k/2\pi}F'_{\alpha,k}\,(a,c;kx/2)$. Therefore,

$$\Phi[L_{\alpha,k}](x) = \frac{1}{2} + \frac{x}{\sqrt{2\pi}}\,F'_{\alpha,k}\left(a,c;-\frac{x^2}{2}\right),$$

where $a = \frac{1}{2}, c = \frac{3}{2}$.

□

Notice that this formula is much cleaner, without the cluttering of k in the previous attempt in [20].

5.2.5. Example 2: Fractional F.

LEMMA 5.10. From (8.2), the standard CDF of a fractional F distribution $F_{\alpha,d,k}$ is

$$\Phi[F_{\alpha,d,k}](x) = \frac{1}{\Gamma\left(\frac{d}{2}\right)} \int_0^\infty ds\, \gamma\left(\frac{d}{2}, \frac{dxs}{2}\right) \overline{\chi}^2_{\alpha,k}(s).$$

It can be expressed by the scaled FGHF via

$$\Phi[F_{\alpha,d,k}](x) = \left[C_{\alpha,d,k}\, \frac{(dx/2)^{d/2}}{\Gamma\left(\frac{d}{2}+1\right)}\right] F'_{\alpha,k+d-1}\left(\frac{d}{2}, \frac{d}{2}+1; -\frac{dx}{2\Sigma}\right). \tag{5.15}$$

where $C_{\alpha,d,k}$ is defined in (5.16) and $\Sigma := \sigma^2_{\alpha,k+d-1}/\sigma^2_{\alpha,k}$. △

PROOF. Note that

$$\frac{1}{\Gamma\left(\frac{d}{2}\right)}\gamma\left(\frac{d}{2}, \frac{x}{2}\right) = \frac{(x/2)^{d/2}}{\Gamma\left(\frac{d}{2}+1\right)} M\left(\frac{d}{2}, \frac{d}{2}+1; -\frac{x}{2}\right).$$

Then

$$\begin{aligned}\Phi[F_{\alpha,d,k}](x) &= \int_0^\infty \left[\frac{(dxs/2)^{d/2}}{\Gamma\left(\frac{d}{2}+1\right)} M\left(\frac{d}{2}, \frac{d}{2}+1; -\frac{dxs}{2}\right)\right] \overline{\chi}^2_{\alpha,k}(s)\, ds \\ &= \frac{(dx/2)^{d/2}}{\Gamma\left(\frac{d}{2}+1\right)} \int_0^\infty M\left(\frac{d}{2}, \frac{d}{2}+1; -\frac{dxs}{2}\right) s^{(d-1)/2}\, \overline{\chi}^2_{\alpha,k}(s)\, \sqrt{s}\, ds.\end{aligned}$$

When $d = 1$, it fits right in with FGHF. When $d > 1$, it needs more work.

From (7.17), let $m = (d-1)/2$, then $k + 2m = k + d - 1$ and

$$\Phi[F_{\alpha,d,k}](x) = C_{\alpha,d,k}\, \frac{(dx/2)^{d/2}}{\Gamma\left(\frac{d}{2}+1\right)} \int_0^\infty M\left(\frac{d}{2}, \frac{d}{2}+1; -\frac{dxy}{2\Sigma}\right) \overline{\chi}^2_{\alpha,k+d-1}(y)\, \sqrt{y}\, dy,$$

where $\Sigma := \sigma^2_{\alpha,k+d-1}/\sigma^2_{\alpha,k}$ and $y = \Sigma\, s$, and

$$C_{\alpha,d,k} := \frac{\sigma^{d-1}_{\alpha,k}}{\sqrt{\Sigma}}\, \frac{C_{\alpha,k}}{C_{\alpha,k+d-1}} = \frac{\sigma^d_{\alpha,k}}{\sigma_{\alpha,k+d-1}}\, \frac{C_{\alpha,k}}{C_{\alpha,k+d-1}}. \tag{5.16}$$

The integral matches the FGHF pattern in Lemma 5.12, and we get (5.15).

□

REMARK 5.11. One final note. There is a connection between (5.14) and (5.15). When $d = 1$, $\Sigma = 1$ and $C_{\alpha,d,k} = 1$. Then

$$\Phi[F_{\alpha,1,k}](x^2) = \frac{2x}{\sqrt{2\pi}}\, F'_{\alpha,k}\left(\frac{1}{2}, \frac{3}{2}; -\frac{x^2}{2}\right) \tag{5.17}$$

which is $2\,\Phi[L_{\alpha,k}](x) - 1$ in (5.14).

This is a reflection of Lemma 8.3. If the variable X distributes as a GSaS $L_{\alpha,k}$, then X^2 distributes as a one-dimensional F, aka $F_{\alpha,1,k}$. It is particularly easy to see this relation in the FGHF form above.

Part 2

One-Sided Distributions

CHAPTER 6

FG: Fractional Gamma Distribution and Tilted Stable Law

The fractional gamma distribution (FG) is one of the central one-sided families used throughout this book. It contains the fractional χ-mean distribution (FCM) as a special case and provides a fractional extension of the generalized gamma distribution.

The same family was called *the generalized stable count distribution* in [20], extending the "stable count distribution" terminology introduced in [19]. In the present formulation, the name *fractional gamma distribution* is more natural because the construction is organized around a fractional analogue of the gamma family.

A key result of this chapter is that FG is exactly a reparameterized inverse tilted stable law. This identification provides the probabilistic foundation for FG and for the distributional constructions built from it in later chapters.

6.1. Definition

DEFINITION 6.1 (Fractional Gamma distribution (FG)). FG is a four-parameter one-sided distribution family, whose PDF is defined as

$$\mathfrak{N}_\alpha(x;\sigma,d,p) := C\left(\frac{x}{\sigma}\right)^{d-1} F_\alpha\left(\left(\frac{x}{\sigma}\right)^p\right) \qquad (x \geq 0) \tag{6.1}$$

where $F_\alpha(x) = W_{-\alpha,0}(-x) = \alpha x M_\alpha(x)$ from (3.9) and $\alpha \in [0,1]$ controls the shape of the Wright function; σ is the scale parameter; p is also the shape parameter controlling the tail behavior ($p \neq 0, dp \geq 0$); d is the *degree of freedom* parameter. When $\alpha \to 1$, the PDF becomes a Dirac delta function: $\delta(x-\sigma)$ assuming σ is finite. When $d \geq 1$, all the moments of the FG exist and have closed forms.

6.2. Determination of C

The normalization constant C is:

$$C = \begin{cases} \frac{|p|}{\sigma}\frac{\Gamma(\alpha d/p)}{\Gamma(d/p)} & , \text{ for } \alpha \neq 0, d \neq 0. \\ \frac{|p|}{\sigma\alpha} & , \text{ for } \alpha \neq 0, d = 0. \end{cases} \tag{6.2}$$

It is important to note that d and p are allowed to be negative, as long as $dp \geq 0$.

PROOF. The normalization constant C in (6.1) is obtained from the requirement that the integral of the PDF must be 1:

$$\int_0^\infty \mathfrak{N}_\alpha(x;\sigma,d,p)\,dx = \frac{C\sigma}{|p|}\frac{\Gamma(\frac{d}{p})}{\Gamma(\frac{d}{p}\alpha)} = 1$$

where the integral is carried out by the moment formula of the Wright function.

We typically constrain $dp \geq 0$ and p is typically positive. However, it becomes negative in the inverse distribution and/or characteristic distribution types. So we need $|p|$ to ensure that C is positive.

For the case of $\alpha \neq 0$ and $d \to 0$, due to (A.3), we have

$$C = \frac{|p|}{\sigma\alpha} \qquad (\alpha \neq 0, d = 0)$$

These two cases are combined to form (6.2). □

6.3. FG Subsumes Generalized Gamma Distribution

Since the Wright function extends an exponential function to the fractional space, FG is the fractional extension of the generalized gamma (GG) distribution[39], whose PDF is defined as:

$$f_{\text{GG}}(x; a, d, p) = \frac{|p|}{a\Gamma(d/p)} \left(\frac{x}{a}\right)^{d-1} e^{-(x/a)^p}. \tag{6.3}$$

The parallel use of parameters is obvious, except that a in GG is replaced by σ in FG to avoid confusion with α.

GG is subsumed to FG in two ways:

$$f_{\text{GG}}(x; \sigma, d, p) := \begin{cases} \mathfrak{N}_0(x; \sigma, d = d - p, p) & , \text{ at } \alpha = 0. \\ \mathfrak{N}_{\frac{1}{2}}\left(x; \sigma = \frac{\sigma}{2^{2/p}}, d = d - \frac{p}{2}, p = \frac{p}{2}\right) & , \text{ at } \alpha = \frac{1}{2}. \end{cases} \tag{6.4}$$

The first line is treated as the definition of FG at $\alpha = 0$. The proof is given in [20].

Although the first line is more obvious, it is the second line that leads to the fractional extension of the χ distribution.

6.4. Mellin Transform

From Example 2.4, we add σ and C. The Mellin transform of the PDF of the fractional gamma distribution is

$$\begin{aligned} \mathfrak{N}_\alpha(x; \sigma, d, p) &\overset{\mathcal{M}}{\longleftrightarrow} \frac{C\,\sigma^s}{|p|} \frac{\Gamma((s + d - 1)/p)}{\Gamma(\alpha(s + d - 1)/p)} \\ &= \sigma^{s-1} \frac{\Gamma(\alpha d/p)}{\Gamma(d/p)} \frac{\Gamma((s + d - 1)/p)}{\Gamma(\alpha(s + d - 1)/p)}, \end{aligned} \tag{6.5}$$

where C is from Section 6.2. The typical limiting case for the gamma functions shall be taken care in each scenario.

Substituting $s = n + 1$ into (6.5), we obtain the moment formula of the FG distribution,

$$\mathbb{E}[X^n] = \sigma^n \frac{\Gamma(\alpha d/p)}{\Gamma(d/p)} \frac{\Gamma((n + d)/p)}{\Gamma(\alpha(n + d)/p)}. \tag{6.6}$$

FG is often used in a ratio distribution, such as the role of $g^*(s)$ in (2.6), where $s \to 2 - s$. The term $s + d - 1$ becomes $d + 1 - s$. Furthermore, in the FCM case, since $d = k - 1$, it becomes the elegant $k - s$ term.

6.5. The Inverse Tilted Stable Law

FG is the foundation for all the distributions that follow in this book. It is beautifully rooted in the polynomially tilted inverse stable distribution[8, 15]. We will show that the FG random variable is exactly the re-parametrized *inverse tilted stable variable*.

We use the T notation from [8] where $T_{\alpha,\beta}$ is the polynomially tilted stable variable. Its density function is $\propto x^{-\beta} L_\alpha(x), x > 0$. β is the *tilted angle* due to polynomial bias.[1]

Polynomial tilting corresponds to a shift in Mellin space. By (2.4), $x^k f(x) \overset{\mathcal{M}}{\longleftrightarrow} f^*(s+k)$. This shift mechanism underlies the introduction of the degrees-of-freedom parameter in (11.1), which motivated the development of the fractional distribution families studied in this book.

[1] It will be shown in Section 7.8 that $\chi^2_{\alpha,-k}$ is equal to $T_{\alpha/2,k/2}$.

It is generalized by adding the third parameter γ so that $T_{\alpha,\beta}^{-\gamma} = (T_{\alpha,\beta})^{-\gamma}$. The inverse power of γ transforms the density function to a polynomially tilted M-Wright function, which will be proved in the Lemma 6.3. We might use the notion $TS(\alpha,\beta,\gamma)$ to avoid collisions between greeks since β,γ might have different meanings in other contexts later in the book.

Let $X \sim \mathfrak{N}_\alpha(\sigma,d,p)$ be the FG random variable. The conclusion is $\beta = \alpha d/p, \gamma = \alpha/p$. That is,

$$X = \sigma\, T_{\alpha,\,\alpha d/p}^{-\alpha/p}. \tag{6.7}$$

The calculation of $T_{\alpha,\beta}$ is carried out in Theorem 1 of [8]. We reproduce the formula for the more familiar $U = T_{\alpha,\beta}^{-\alpha}$ in the following way.

LEMMA 6.2. For the tilted stable law $T_{\alpha,\beta}$, the sampling is performed via the modified Kanter's method,

$$T_{\alpha,\beta} = A_\alpha(Q_\beta)\, G_\beta^{-c}, \quad c = \frac{1-\alpha}{\alpha}.$$

(1) The factor $A_\alpha(q)$ is *the Zolotarev function* from (3.22).
(2) Q_β is the sampling of A_α with angle β, whose density is

$$f_{Q_\beta}(q) = I_\beta\, A_\alpha(q)^{-\beta}, \quad \text{where } I_\beta = \frac{\Gamma(1+\beta)\Gamma(1+c\beta)}{\Gamma(1+\beta/\alpha)}, \qquad 0<q<1.$$

When $\beta = 0$, Q_0 is uniformly distributed on $(0,1)$.
(3) G_β is a gamma variable

$$G_\beta \sim \text{Gamma}(1+c\beta, \text{scale} = 1)\,.$$

Then

$$U = T_{\alpha,\beta}^{-\alpha} = A_\alpha(Q_\beta)^{-\alpha}\, G_\beta^{1-\alpha}. \tag{6.8}$$

△

The case of $\beta = 0$ is added because it is an interesting special case: $T_{\alpha,0}^{-\alpha} = M_\alpha$ where M_α represents the M-Wright function treated as a one-sided distribution.

Moving from U to X is straightforward,

$$X = \sigma\, U^{1/p} \tag{6.9}$$

LEMMA 6.3. For FG, the corresponding tilted stable parameters are $\beta = \alpha d/p$ and $\gamma = \alpha/p$.

△

PROOF. It follows immediately from (6.7) that $\gamma = \alpha/p$. From [8, 15], $U = T_{\alpha,\beta}^{-\alpha}$ has density

$$f_U(u) = J_\beta\, u^{\beta/\alpha} M_\alpha(u) = \frac{J_\beta}{\alpha}\, u^{\beta/\alpha-1} F_\alpha(u), \quad \text{where } J_\beta = \frac{\Gamma(\beta+1)}{\Gamma(\beta/\alpha+1)}.$$

Let

$$u = \left(\frac{x}{\sigma}\right)^p.$$

Then

$$\left|\frac{du}{dx}\right| = \frac{|p|}{\sigma}\left(\frac{x}{\sigma}\right)^{p-1}.$$

Hence the density of $X = \sigma U^{1/p}$ is

$$\begin{aligned} f_X(x) &= f_U(u)\left|\frac{du}{dx}\right| \\ &= \frac{|p| J_\beta}{\alpha\sigma}\left(\frac{x}{\sigma}\right)^{p\beta/\alpha-1} F_\alpha\left(\left(\frac{x}{\sigma}\right)^p\right). \end{aligned}$$

Comparing this with (6.1), we obtain

$$\frac{p\beta}{\alpha} - 1 = d - 1,$$

hence

$$\beta = \frac{\alpha d}{p}.$$

The normalization constant is also consistent, because

$$\begin{aligned}\frac{|p|J_\beta}{\alpha\sigma} &= \frac{|p|}{\sigma}\frac{\Gamma(\beta)}{\Gamma(\beta/\alpha)} \\ &= \frac{|p|}{\sigma}\frac{\Gamma(\alpha d/p)}{\Gamma(d/p)},\end{aligned}$$

which agrees with (6.2). The case $d = 0$ follows by the same limiting argument used in Section 6.2. □

The numeric sampler implements the logarithmic form:

$$\log U = -\alpha \log T_{\alpha,\beta} = (1-\alpha)\log G_\beta - \alpha \log A_\alpha(Q_\beta). \tag{6.10}$$

Then

$$\begin{aligned}\log X = -\gamma \log T_{\alpha,\beta} &= \log\sigma + c\gamma \log G_\beta - \gamma \log A_\alpha(Q_\beta) \\ \text{where } \beta = \frac{\alpha d}{p}, \gamma = \frac{\alpha}{p}, c &= \frac{1-\alpha}{\alpha}.\end{aligned} \tag{6.11}$$

6.6. The Lognormal Limit

Under a suitable joint parameter limit, the tilted-stable representation (6.7) yields a lognormal (LN) limit for FG after normalization by its ordinary mean. We establish this general result here; it underlies the lognormal limit of FCM and the lognormal-normal (LNN) limit of GSaS developed in subsequent chapters.

Throughout this section, use the FG parameters (α, σ, d, p) and their tilted-stable counterparts from Lemma 6.3:

$$\beta = \frac{\alpha d}{p}, \qquad \gamma = \frac{\alpha}{p}, \qquad X \overset{d}{=} \sigma T_{\alpha,\beta}^{-\gamma}.$$

LEMMA 6.4 (The mean-normalized FG variable). Let $X \sim \mathfrak{N}_\alpha(\sigma, d, p)$ with $0 < \alpha < 1$, $\sigma > 0$, $dp > 0$, and $\beta + \gamma > 0$, so that its ordinary mean is finite. Define $V = X/\mathbb{E}[X]$ and let f_V denote its density. In the notation of (2.1),

$$f_V(v) \overset{\mathcal{M}}{\longleftrightarrow} f_V^*(s) = \mathbb{E}[V^{s-1}].$$

For real s, put $q = s - 1$ and assume $\beta + \gamma q > 0$. Then (6.5) gives the logarithm of the Mellin transform as

$$\begin{aligned}\log f_V^*(s) = \log \mathbb{E}[V^q] &= \Delta_\alpha(\beta + \gamma q) - \Delta_\alpha(\beta) \\ &\quad - q\big[\Delta_\alpha(\beta + \gamma) - \Delta_\alpha(\beta)\big].\end{aligned} \tag{6.12}$$

where Δ_α is the log-gamma difference defined in (3.27).

Equivalently, $\log \mathbb{E}[V^q]$ is the cumulant-generating function of $\log V$ evaluated at q. △

LEMMA 6.5 (The mean-one lognormal limit of FG). Consider a sequence of FG variables X_n with $0 < \alpha_n < 1$ and arbitrary positive scales σ_n. Fix $\gamma = \alpha_n/p_n \neq 0$ and suppose

$$\beta_n = \frac{\alpha_n d_n}{p_n} \longrightarrow \infty, \qquad \frac{1-\alpha_n}{p_n d_n} \longrightarrow \tau_{LN}^2 \in (0, \infty). \tag{6.13}$$

Then $V_n = X_n/\mathbb{E}[X_n]$ is well defined for all sufficiently large n, and

$$\begin{aligned} \log V_n &\xrightarrow{d} \mathcal{N}(-\tau_{LN}^2/2, \tau_{LN}^2), \\ V_n &\xrightarrow{d} \text{Lognormal}(-\tau_{LN}^2/2, \tau_{LN}^2). \end{aligned} \tag{6.14}$$

For every fixed real s, the Mellin transforms also converge:

$$\begin{aligned} &f_{V_n}(v) \xleftrightarrow{\mathcal{M}} f^*_{V_n}(s) = \mathbb{E}[V_n^{s-1}], \\ &f^*_{V_n}(s) \longrightarrow \exp\left[\frac{\tau_{LN}^2}{2}(s-1)(s-2)\right]. \end{aligned} \tag{6.15}$$

△

Proof. Suppress the sequence index. Since $\beta \to \infty$ and γ is fixed, the moment condition $\beta+\gamma q > 0$ eventually holds uniformly on every fixed bounded real q-interval, including $q = 1$ for the mean.

By the shared expansion (3.29), the path (6.13) implies

$$\gamma^2 \Delta''_\alpha(\beta) = \frac{1-\alpha}{pd} + O(\beta^{-3}) \longrightarrow \tau_{LN}^2.$$

The third-derivative expansion in (3.29) also gives $\Delta'''_\alpha(\beta+u) = O(\beta^{-1})$ uniformly for bounded u.

Taylor expansion of the normalized log-moment formula (6.12) yields, uniformly for q in a fixed bounded real interval,

$$\log \mathbb{E}[V^q] = \frac{\gamma^2}{2}\Delta''_\alpha(\beta)(q^2-q) + O(\beta^{-1}) \longrightarrow \frac{\tau_{LN}^2}{2}(q^2-q).$$

The linear term $\gamma q \Delta'_\alpha(\beta)$ cancels because of the ordinary-mean normalization. The limit is the log-MGF of $\mathcal{N}(-\tau_{LN}^2/2, \tau_{LN}^2)$, proving the normal limit by MGF convergence and the lognormal limit by exponentiation. Setting $q = s-1$ gives (6.15). □

The limiting lognormal has ordinary mean one, as also seen by setting $s = 2$ in (6.15). This result covers either sign of p with d of the same sign. It is a joint limit with $p_n = \alpha_n/\gamma$, not a limit with p held fixed.

6.7. CDF and Fractional Incomplete Gamma Function

The CDF of FG is

$$\Phi(x) := \int_0^x \mathfrak{N}_\alpha(s; \sigma, d, p)\, ds \quad (x \geq 0). \tag{6.16}$$

This integral leads to fractionalization of the incomplete gamma function in Section 3.5.

Lemma 6.6. The CDF of FG can be represented by γ^*_α in (3.23) or Γ^*_α in (3.24) as

$$\Phi(x) = \begin{cases} z^{d+p}\, \gamma^*_\alpha\left(d/p+1, z^p\right) & , \text{ when } p > 0. \\ \\ z^{d+p}\, \Gamma^*_\alpha\left(d/p+1, z^p\right) & , \text{ when } p < 0. \end{cases} \tag{6.17}$$

where $z = x/\sigma$ is the standardized variable.

This could be viewed as one form of fractional extension to the regularized lower incomplete gamma function, $\gamma(s,z)/\Gamma(s)$, which is the CDF of GG mentioned above.

Due to this result, this distribution is called the *fractional gamma distribution* (FG).

△

PROOF. The CDF of FG is

$$\begin{aligned}\Phi(x) &= \int_0^x \mathfrak{N}_\alpha(s;\sigma,d,p)\,ds \\ &= C\int_0^x ds\left(\frac{s}{\sigma}\right)^{d-1} W_{-\alpha,0}\left(-\left(\frac{s}{\sigma}\right)^p\right). \\ &= C\int_0^x ds\left(\frac{s}{\sigma}\right)^{d-1} F_\alpha\left(\left(\frac{s}{\sigma}\right)^p\right).\end{aligned}$$

Since $F_\alpha(x) = \alpha x\, M_\alpha(x)$ from (3.9), and let $u = s/x$, then $u \in [0,1]$ and

$$\begin{aligned}\Phi(x) &= \alpha C\int_0^x ds\left(\frac{s}{\sigma}\right)^{d+p-1} M_\alpha\left(\left(\frac{s}{\sigma}\right)^p\right) \\ &= \alpha C\, x\int_0^1 du\left(\frac{xu}{\sigma}\right)^{d+p-1} M_\alpha\left(\left(\frac{xu}{\sigma}\right)^p\right)\end{aligned}$$

Recognize that $u^p \in [0,1]$ when $p > 0$. Let $t = u^p$, and $dt/t = p\,du/u$,

$$\Phi(x) = \frac{\alpha\sigma C}{p}\, z^{d+p}\int_0^1 dt\, t^{d/p}\, M_\alpha\left(z^p t\right)$$

Compare the last line with γ_α^* in (3.23), and we get

$$\Phi(x) = \frac{\alpha\sigma C}{p}\,\frac{\Gamma(\frac{d}{p}+1)}{\Gamma((1-\alpha)+\alpha(\frac{d}{p}+1))}\, z^{d+p}\,\gamma_\alpha^*\left(d/p+1, z^p\right)$$

Using the case of $\alpha \neq 0, d \neq 0$ for C, it can be shown that the constant part is just 1. Hence,

$$\Phi(x) = z^{d+p}\,\gamma_\alpha^*\left(d/p+1, z^p\right), \quad \text{for } p > 0.$$

On the other hand, $u^p \in [1,\infty]$ when $p < 0$. The range of the integral changes to $\int_1^\infty$. It leads to the use of the supplementary function (3.24):

$$\Phi(x) = z^{d+p}\,\Gamma_\alpha^*\left(d/p+1, z^p\right).$$

□

Numerically, it is better to use γ_α^* when $z^p \le 1$ and to use Γ_α^* when $z^p > 1$, as long as the identity relation (3.25) is preserved.

6.7.1. The lognormal limit in CDF. The error-function limit is established directly for the special functions in Lemma 3.7. In FG parameters, substitute

$$s_n = \frac{d_n}{p_n} + 1, \qquad \beta_n = \frac{\alpha_n d_n}{p_n}, \qquad \gamma = \frac{\alpha_n}{p_n}.$$

Then $\alpha_n^2(s_n - 1) = \gamma\alpha_n d_n$, so the small-$\alpha$ path corresponds to $\alpha_n d_n \to \gamma/\tau_{LN}^2$ with γ fixed. By (6.6), the gamma-ratio shift in that lemma satisfies $e^{\ell_n} = \mathbb{E}[X_n]/\sigma_n$. Its moving argument is therefore

$$x_n(t) = \left[\frac{\mathbb{E}[X_n]}{\sigma_n} e^{\tau_{LN} t - \tau_{LN}^2/2}\right]^{p_n}.$$

Since $p_n s_n = d_n + p_n$, the regularized expressions in (3.31) coincide with the FG CDF (6.17) evaluated at $\mathbb{E}[X_n]e^{\tau_{LN} t - \tau_{LN}^2/2}$.

6.8. Inverse Expression of Several Fractional Distributions

Several known fractional distributions could be expressed in the FG in Table 1. This shows that the FG is the super set of the one-sided fractional distribution system. Its parametrization provides immense flexibility to express other formerly known one-sided distributions.

Distribution (PDF)	Wright Equiv.	FG: $\mathfrak{N}_\alpha(x;\sigma,d,p)$			
		α	σ	d	p
One-sided stable: $L_\alpha(x)$	$x^{-1}W_{-\alpha,0}(-x^{-\alpha})$	α	1	0	$-\alpha$
Stable Count: $\mathfrak{N}_\alpha(x)$		α	1	1	α
Stable Vol: $V_\alpha(x)$		$\frac{\alpha}{2}$	$\frac{1}{\sqrt{2}}$	1	α
M-Wright: $M_\alpha(x)$	$\frac{1}{\alpha x}W_{-\alpha,0}(-x)$	α	1	0	1
M-Wright II: $\Gamma(\alpha)F_\alpha(x)$	$\Gamma(\alpha)W_{-\alpha,0}(-x)$	α	1	1	1

TABLE 1. FG mapping of several known fractional distributions in the literature. $\mathfrak{N}_\alpha(x)$ and $V_\alpha(x)$ first appeared in [19], which led to this work.

6.9. Alternative Definition

DEFINITION 6.7. It is reasonable to argue that the PDF of FG can be defined via the M-Wright function directly, such that

$$\mathfrak{N}'_\alpha(x;\sigma,d',p) := C' \left(\frac{x}{\sigma}\right)^{d'-1} M_\alpha\left(\left(\frac{x}{\sigma}\right)^p\right). \qquad (x \geq 0) \tag{6.18}$$

However, since $F_\alpha(z) = \alpha z\, M_\alpha(z)$, it is easy to see that

$$\mathfrak{N}'_\alpha(x;\sigma,d',p) = \alpha\, C' \left(\frac{x}{\sigma}\right)^{d'+p-1} F_\alpha\left(\left(\frac{x}{\sigma}\right)^p\right).$$

Therefore, this is merely a reparameterization of $d = d' + p$. This definition will encounter some issues in FCM later due to the assignment of $d \to k-1$, $\alpha \to \alpha/2$ and $p \to \alpha$ (see (7.4)). We learn from Figures 12.1 and 12.2 that there is a natural linear relation between k and $\epsilon = 1/\alpha$. Mixing the role of d with α from p is not a good idea.

CHAPTER 7

Fractional Chi Distributions

7.1. Introduction to Fractional Chi Distribution

Chapter 4 showed that the Mellin transform of the α-stable density can be interpreted as the transform of a ratio distribution. This observation motivates the fractional χ construction. The two factors are

$$L_\alpha^\theta(x) \overset{\mathcal{M}}{\longleftrightarrow} \epsilon \left[\frac{\Gamma(s)}{\Gamma(1-\gamma+\gamma s)}\right] \left[\frac{\Gamma(\epsilon(1-s))}{\Gamma(\gamma(1-s))}\right]$$
$$\text{where } \epsilon = \frac{1}{\alpha}, \gamma = \frac{\alpha-\theta}{2\alpha}.$$

The first bracket is the Mellin transform of the M-Wright function. The second bracket is interpreted as the Mellin transform of the fractional χ-mean distribution (FCM) at $k = 1$:

$$\overline{\chi}_{\alpha,1}^\theta(x) \overset{\mathcal{M}}{\longleftrightarrow} \overline{\chi}_{\alpha,1}^\theta{}^*(s) \propto \frac{\Gamma(\epsilon(s-1))}{\Gamma(\gamma(s-1))},$$

up to the normalization constant and scale in the PDF.

The connection follows from the Mellin-transform rule for ratio distributions: the transform of the denominator is evaluated at $2-s$. Thus the term $s-1$ in $\overline{\chi}_{\alpha,1}^\theta{}^*(s)$ becomes $1-s$ in the stable-density factor above. In this chapter, the degrees-of-freedom parameter k is introduced by replacing $s-1$ with $s+k-2$, giving

$$\overline{\chi}_{\alpha,k}^\theta(x) \overset{\mathcal{M}}{\longleftrightarrow} \overline{\chi}_{\alpha,k}^\theta{}^*(s) \propto \frac{\Gamma(\epsilon(s+k-2))}{\Gamma(\gamma(s+k-2))}. \tag{7.1}$$

This prototype is the starting point for the full definition of FCM.

7.2. FCM: Fractional Chi-Mean Distribution

There are two ways to define FCM. The first approach is to define it via Mellin transform. The second approach is to define the shape of its PDF.

Definition 7.1 (Fractional χ-mean distribution (FCM) via Mellin Transform). The Mellin transform of FCM's PDF is enriched from (7.1) to

$$\begin{aligned}\overline{\chi}_{\alpha,k}^\theta(x) &\overset{\mathcal{M}}{\longleftrightarrow} \overline{\chi}_{\alpha,k}^\theta{}^*(s) \\ &= (\sigma_{\alpha,k}^\theta)^{s-1} \frac{\Gamma(\gamma(k-1))}{\Gamma(\epsilon(k-1))} \frac{\Gamma(\epsilon(s+k-2))}{\Gamma(\gamma(s+k-2))}, \\ &\text{where } \sigma_{\alpha,k}^\theta := \gamma^\gamma k^{\gamma-\epsilon}.\end{aligned} \tag{7.2}$$

The main differences are (1) to address the normalization of the total density, and (2) to have a proper scale $\sigma_{\alpha,k}^\theta$ such that it is consistent with the classic χ distribution ans α-stable distribution.

For positive k, the PDF of an FCM is

$$\overline{\chi}^{\theta}_{\alpha,k}(x) := \mathfrak{N}_{\gamma\alpha}(x;\sigma=\sigma^{\theta}_{\alpha,k},d=k-1,p=\alpha) \qquad (x\geq 0) \tag{7.3}$$
$$= \frac{\Gamma(\gamma(k-1))}{\epsilon\Gamma(\epsilon(k-1))}\left(\sigma^{\theta}_{\alpha,k}\right)^{1-k} x^{k-2}\, F_{\gamma\alpha}\left(\left(\frac{x}{\sigma^{\theta}_{\alpha,k}}\right)^{\alpha}\right),$$

where $\mathfrak{N}_\lambda(x;\sigma,d,p)$ is FG (6.1), and $F_\lambda(x) := W_{-\lambda,0}(-x) = \lambda x M_\lambda(x)$ is the Wright function of the second kind (3.9).

Notice the appearances of γ that replaces all the $1/2$ in Section 7.6 of [**20**]. That is is how θ comes into play in the upgraded FCM. This full representation is used in Chapter 11.

However, for GAS-SN in Chapter 12 and beyond, such θ upgrade is unnecessary. The skew-normal framework is based on modulation of normal distributions. It is required to have $\theta = 0$, aka $\gamma = 1/2$.

Hence, we recite the original definition of FCM PDF ($k > 0$):

$$\overline{\chi}_{\alpha,k}(x) = \overline{\chi}^{0}_{\alpha,k}(x) := \mathfrak{N}_{\alpha/2}(x;\sigma=\sigma_{\alpha,k},d=k-1,p=\alpha) \qquad (x\geq 0) \tag{7.4}$$
$$= (C_{\alpha,k})\left(\sigma_{\alpha,k}\right)^{1-k} x^{k-2}\, F_{\frac{\alpha}{2}}\left(\left(\frac{x}{\sigma_{\alpha,k}}\right)^{\alpha}\right),$$

where

$$C_{\alpha,k} := \frac{\alpha\Gamma((k-1)/2)}{\Gamma((k-1)/\alpha)} \tag{7.5}$$

$$\sigma_{\alpha,k} := \frac{|k|^{1/2-1/\alpha}}{\sqrt{2}}. \tag{7.6}$$

Note that the difference between (7.3) and (7.4) is very small: Just replace $\mathfrak{N}_{\gamma\alpha}(...)$ to $\mathfrak{N}_{\alpha/2}(...)$.

7.2.1. Inverse Tilted Stable Law. According to Section 6.5, (7.4) implies $\overline{\chi}_{\alpha,k} = \sigma_{\alpha,k}\, T^{-1/2}_{\alpha/2,\,(k-1)/2}$. The special case for stable distribution decomposition (4.11) is $\overline{\chi}^{\theta}_{\alpha,1} = \gamma^{\gamma}\, T^{-\gamma}_{\gamma\alpha,\,0}$ and $\overline{\chi}_{\alpha,1} = T^{-1/2}_{\alpha/2,\,0}/\sqrt{2}$.

7.2.2. FCM CDF. Extending directly from Lemma 6.6, we have

LEMMA 7.2. The CDF of FCM can be represented by γ^*_α in (3.23) as

$$\Phi[\overline{\chi}_{\alpha,k}](x) = z^{k-1+\alpha}\,\gamma^*_{\alpha/2}\left(\frac{k-1+\alpha}{\alpha}, z^{\alpha}\right), \qquad (k>0,\alpha\in[0,2]) \tag{7.7}$$

where $z = x/\sigma_{\alpha,k}$.

△

7.2.3. FCM Subsumes Chi. It is easy to show that $\overline{\chi}_{1,k}$ is χ_k with a scale of $1/\sqrt{k}$, due to (3.15).

7.3. FCM Moments

By letting $s = n+1$ and $\theta = 0$ in (7.2), its n-th moment is

$$\mathbb{E}(X^n|\overline{\chi}_{\alpha,k}) = (\sigma_{\alpha,k})^n\,\frac{\Gamma((k-1)/2)}{\Gamma((k-1)/\alpha)}\,\frac{\Gamma((n+k-1)/\alpha)}{\Gamma((n+k-1)/2)}, \qquad (k>0,\alpha>0) \tag{7.8}$$

which requires $k > 1$ and $n + k > 1$ to avoid the singularity of the gamma functions (See Section 7.6 of [**20**]).

The explicit form of the first moment is

$$\mathbb{E}(X|\overline{\chi}_{\alpha,k}) = \sigma_{\alpha,k}\,\frac{\Gamma((k-1)/2)}{\Gamma((k-1)/\alpha)}\,\frac{\Gamma(k/\alpha)}{\Gamma(k/2)} = \sigma_{\alpha,k}\,\frac{C_{\alpha,k}}{C_{\alpha,k+1}}. \qquad (k>0,\alpha>0) \tag{7.9}$$

Notice that it can be used as a bridge connecting the coefficient C between k and $k+1$.

The moment formula of FCM is fundamental to all the fractional distributions built on top of it. However, ironically, due to the nature of a ratio distribution, it is often evaluated as negative moments $n<0$. Hence, n is restricted in the range of $1-k<n<0$.

This results in non-existing moments when k is not "large enough", which happens to be a core feature of the α-stable distribution and Student's t distribution. Our two-dimensional parameter space (α,k) adds more complexity to it.

7.3.1. FCM at Infinite Degrees of Freedom. The choice of $\sigma_{\alpha,k}$ is intentional, such that

$$\lim_{k\to\infty}\mathbb{E}(X^n|\overline{\chi}_{\alpha,k}) = \alpha^{-n/\alpha}. \qquad (k>0,\alpha>0) \tag{7.10}$$

Under such conditions, its variance is zero. That is, FCM becomes a delta function, $\delta(x-\alpha^{-1/\alpha})$, as $k\to\infty$.

7.4. FCM Reflection Formula and Negative k

7.4.1. FCM for Negative k. We quote Definition 3.2 of [20] for FCM in the negative k space. Its PDF defined by FG is

$$\overline{\chi}_{\alpha,-k}(x) := \mathfrak{N}_{\alpha/2}(x;\sigma=(\sigma_{\alpha,k})^{-1},d=-k,p=-\alpha). \qquad (x\geq 0,k>0) \tag{7.11}$$

This is defined as the *characteristic FCM* discussed in Chapter 9. That is, $\left[\overline{\chi}_{\phi}\right]_{\alpha,k} := \overline{\chi}_{\alpha,-k}$ in Lemma 9.6. It is derived from the properties of the α-stable characteristic function.

7.4.2. Tilted Stable Law. According to Section 6.5, (7.11) implies $\overline{\chi}_{\alpha,-k} = (\sigma_{\alpha,k})^{-1}\,T^{1/2}_{\alpha/2,\,k/2}$. Notice that the superscript $(-\gamma=1/2)$ is positive. The distribution is based on L_α, instead of M_α.

The explicit PDF derivation is documented in Section 9.4. We quote the inverse relation:

$$\overline{\chi}_{\alpha,-k}(x) = \frac{x^{-3}}{\mathbb{E}(X|\overline{\chi}_{\alpha,k})}\,\overline{\chi}_{\alpha,k}\left(\frac{1}{x}\right). \qquad (x\geq 0,k>0) \tag{7.12}$$

which is proved in (9.18) in Section 9.4.

$\overline{\chi}_{\alpha,-k}$ is used to define the fractional exponential power distribution within the GSaS (and GAS-SN) nomenclature, which subsumes the exponential power distribution (Section 3.11.1 of [30]). See Section 12.8. Readers interested in more details are referred to the FCM sections in [20].

We quote the FCM reflection formula from Section 7 of [20] to summarize the relation:

$$\mathbb{E}(X^n|\overline{\chi}_{\alpha,-k}) = \frac{\mathbb{E}(X^{-n+1}|\overline{\chi}_{\alpha,k})}{\mathbb{E}(X|\overline{\chi}_{\alpha,k})},\ k>0. \tag{7.13}$$

This indicates an elegant relation for the first moment that $\mathbb{E}(X|\overline{\chi}_{\alpha,-k}) = 1/\mathbb{E}(X|\overline{\chi}_{\alpha,k})$.

7.5. FCM: Unified Algebra of Cumulants of the Log Variable

The logarithm of an FCM variable admits a common cumulant algebra for both the ordinary and characteristic branches. This formulation serves two purposes. First, it establishes the Gaussian limit of the logarithmic variable, leading to the lognormal limit of FCM and the lognormal-normal (LNN) limit of GSaS. Second, it provides the log-moment and characteristic functions needed for log-domain inversion in the large-$|k|$, small-α region.

7.5.1. The Cumulant Generating Function. Let $V_k \sim \overline{\chi}_{\alpha,k}$, with $k > 1$ for ordinary FCM or $k < 0$ for characteristic FCM.

For the two branches of FCM, we define $\zeta = \operatorname{sign}(k) \in \{+1, -1\}$ and let

$$z = \begin{cases} k - 1, & k \geq 1, \\ |k|, & k < 0. \end{cases} \tag{7.14}$$

V_k is assigned to the tilted stable law as

$$V_k \overset{d}{=} \sigma^{\zeta} T^{-\zeta/2}_{\alpha/2,z/2}. \tag{7.15}$$

This is the elegant one-liner for the FCM lines in Table 1. Leave $\sigma > 0$ as the branch scale; its value depends on the FCM scaling convention and special situations.

We still retain $\epsilon = 1/\alpha$, but we use $g = 1/2$ for $\theta = 0$. Define the *Mellin moment function* as

$$H(s; \epsilon, z, \zeta, g, \sigma) := \sigma^{\zeta s} \frac{\Gamma(\epsilon(z + \zeta s))}{\Gamma(\epsilon z)} \frac{\Gamma(gz)}{\Gamma(g(z + \zeta s))}. \tag{7.16}$$

This holds for real s in the moment domain $z + \zeta s > -\alpha$, with the value at the removable point $z + \zeta s = 0$ defined by continuity.

Then, according to the moment formula of the tilted stable law[8], we have

$$\mathbb{E}[V_k^s] = \sigma^{\zeta s} \mathbb{E}[T^{-\zeta s/2}_{\alpha/2,z/2}] = H(s; \epsilon, z, \zeta, g, \sigma). \tag{7.17}$$

Here, H treats z, ζ, σ as independent arguments for which we can substitute different values. For example, we can use $\zeta = -1$ for the inverse of a positive-k FCM.

From this point on, we assume that $\epsilon, z, \zeta, g, \sigma$ are constants. They are therefore treated as fixed parameters in the functions below, such as $A(x)$ and $K(s)$.

Define

$$A(x) := \log \Gamma(\epsilon x) - \log \Gamma(gx), \tag{7.18}$$

$$K(s) := A(z + \zeta s) - A(z). \tag{7.19}$$

The log-moment, $\log \mathbb{E}[V_k^s]$, then becomes

$$\log H(s; \epsilon, z, \zeta, g, \sigma) = \zeta s \log \sigma + K(s). \tag{7.20}$$

The term $\zeta \log \sigma$ is the shift of the logarithmic FCM variable, and $K(s)$ is its cumulant generating function (CGF) without the shift. The algebra of $K(s)$ is our main concern in this section.

7.5.2. The cumulants. Calculating the cumulants from the CGF involves differentiating $A(x)$ and $K(s)$.

First,

$$A^{(n)}(x) = \epsilon^n \psi^{(n-1)}(\epsilon x) - g^n \psi^{(n-1)}(gx), \qquad (n \geq 1, x > 0) \tag{7.21}$$

where $\psi^{(n)}(x) := \frac{d^n}{dx^n}\psi(x)$ is the polygamma function of order n. $\psi^{(1)}$ is the trigamma function, and $\psi^{(0)} = \psi$ is the digamma function, where $\psi(x) := \frac{d}{dx} \log \Gamma(x) = \Gamma'(x)/\Gamma(x)$.

Differentiating $A(x)$ twice and applying the chain rule gives the exact trigamma expression below, with factors ϵ^2 and g^2.

LEMMA 7.3 ($A^{(n)}(x)$ in polygamma integrals). For fixed $\epsilon, g > 0$ and $r > 0$, define the *Bernoulli difference kernel* (our terminology) by

$$\begin{aligned} B(r) &:= \frac{1}{1 - e^{-r/\epsilon}} - \frac{1}{1 - e^{-r/g}} \\ &= \frac{1}{2} \left[\coth\left(\frac{r}{2\epsilon}\right) - \coth\left(\frac{r}{2g}\right) \right]. \end{aligned} \tag{7.22}$$

The integral representations in DLMF §5.9(ii), equations (5.9.13) and (5.9.16)[10], yield

$$\begin{aligned} A'(x) &= (\epsilon - g)\log x + \epsilon\log\epsilon - g\log g \\ &\quad - \int_0^\infty e^{-xr}\left[B(r) - \frac{\epsilon - g}{r}\right] dr, \\ A''(x) &= \int_0^\infty re^{-xr}B(r)\,dr. \end{aligned} \tag{7.23}$$

Both integrals converge absolutely. The subtraction in the first integrand is essential when $\epsilon \neq g$, since $B(r) = (\epsilon - g)/r + O(r)$ as $r \to 0$.

△

Proof. DLMF (5.9.13) gives

$$\psi(x) = \log x + \int_0^\infty e^{-xr}\left[\frac{1}{r} - \frac{1}{1 - e^{-r}}\right] dr.$$

Substitute this into $A'(x) = \epsilon\psi(\epsilon x) - g\psi(gx)$. Rescale $r = u/\epsilon$ and $r = u/g$ in the respective integrals to obtain the first identity.

Differentiating DLMF (5.9.16) under the integral sign gives

$$\psi'(x) = \int_0^\infty \frac{re^{-xr}}{1 - e^{-r}}\,dr.$$

The same rescalings in $A''(x) = \epsilon^2\psi'(\epsilon x) - g^2\psi'(gx)$ give the second identity.

For fixed ϵ, g, $B(r) = (\epsilon - g)/r + O(r)$ as $r \to 0$, so both integrands are integrable at zero. Exponential decay ensures convergence at infinity and justifies differentiation under the integral sign.

□

For large positive u, the expansion $\psi'(u) = 1/u + 1/(2u^2) + 1/(6u^3) + O(u^{-5})$ shows that the x^{-2} terms cancel. Thus, for fixed $\epsilon, g > 0$, as $x \to \infty$,

$$A''(x) = \frac{\epsilon - g}{x} + \frac{\epsilon^{-1} - g^{-1}}{6x^3} + O(x^{-5}). \tag{7.24}$$

Second,

$$K^{(n)}(s) = \zeta^n A^{(n)}(z + \zeta s) = \zeta^n\left[\epsilon^n\psi^{(n-1)}(\epsilon(z + \zeta s)) - g^n\psi^{(n-1)}(g(z + \zeta s))\right], \qquad n \geq 1. \tag{7.25}$$

At removable zeros, including $z = s = 0$ for $k = 1$, take the limit of the complete bracket. However, this is not an issue here, since we focus on large z.

The first cumulant is $\kappa_1 = \zeta\log\sigma + K'(0)$, and $K'(0) = \zeta[\epsilon\psi(\epsilon z) - g\psi(gz)]$ for $z > 0$. This is the mean of the log variable.

For higher cumulants κ_n with $n \geq 2$, further differentiation removes the linear scale term and gives

$$\kappa_n = K^{(n)}(0) = \zeta^n A^{(n)}(z) = \zeta^n\left[\epsilon^n\psi^{(n-1)}(\epsilon z) - g^n\psi^{(n-1)}(gz)\right]. \tag{7.26}$$

7.6. FCM: Lognormal Asymptotics and Density by Log Inversion

Definition 7.4 (The Lognormal Distribution). Let $Z \sim \mathcal{N}(\mu, \tau^2)$. Then $X = e^Z$ has distribution Lognormal(μ, τ^2), where τ is the shape parameter and μ is the log-location parameter, which affects only the scale of the distribution. The case $\mu = 0$ is geometrically centered.

The original FCM scaling does not yield a nondegenerate lognormal limit along the path considered below, because its log-location diverges. We introduce the following rescaling to remove this drift, which also facilitates stable numerical implementation of the log-inversion algorithm.

DEFINITION 7.5 (Lognormal-compatible scaling of FCM). For $0 < \alpha \le 2$ and $|k| \ge 1$, let $V \sim \overline{\chi}_{\alpha,k}$ and define

$$D_{\alpha,k} := \left[\alpha^{1/\alpha}\, 2^{(1-\alpha)/2}\right]^{1-1/k^2}, \qquad \hat{V} := D_{\alpha,k}^{\zeta} V. \tag{7.27}$$

The correction is even in k and equals 1 at $\alpha = 1$, $\alpha = 2$, and $|k| = 1$, preserving these specializations. The original FCM definition is unchanged. Substituting (7.6) and (7.27), its effective FG scale is explicitly

$$\begin{aligned} \sigma &= (\sigma_{\alpha,k} D_{\alpha,k})^{\zeta} \\ &= \left\{ \frac{\sqrt{|k|}}{2^{\alpha/2}} \left(\frac{\alpha}{|k|}\right)^{1/\alpha} \left[\alpha^{1/\alpha}\, 2^{(1-\alpha)/2}\right]^{-1/k^2} \right\}^{\zeta}. \end{aligned} \tag{7.28}$$

This technical rescaling preserves the original scale at the special cases listed above, while canceling the log-location drift to obtain a nondegenerate lognormal limit along the path specified below.

The two branches use reciprocal scale due to the ζ exponent. Applying the correction reciprocally on the two branches also preserves (7.13).

LEMMA 7.6. Let $\hat{V}$ be defined by (7.27), and $\delta = \epsilon - \frac{1}{2}$. On either fixed-sign branch, let $z \to \infty$ with $\delta/z \to \tau_{LN}^2 \in (0, \infty)$; equivalently, $\alpha|k| \to 1/\tau_{LN}^2$. Then

$$\log \hat{V} \xrightarrow{d} \mathcal{N}(\mu_{LN}^{\zeta}, \tau_{LN}^2), \qquad \hat{V} \xrightarrow{d} \text{Lognormal}(\mu_{LN}^{\zeta}, \tau_{LN}^2), \tag{7.29}$$

$$\mu_{LN}^{\zeta} := \begin{cases} \frac{1}{2}\log 2 - \tau_{LN}^2, & \zeta = 1, \\ -\frac{1}{2}\log 2, & \zeta = -1. \end{cases}$$

Thus both rescaled branches have nondegenerate lognormal limits without parameter-dependent expectation centering. Equivalently, $e^{-\mu_{LN}^{\zeta}} \hat{V} \xrightarrow{d} \text{Lognormal}(0, \tau_{LN}^2)$. The lognormal belongs to the closure of the rescaled family, not to its finite-parameter members. △

PROOF. By (7.20), with $\sigma = \sigma_{\alpha,k} D_{\alpha,k}$, the log-MGF of $\log \hat{V}$ is

$$\log H(s) = \zeta s \log(\sigma_{\alpha,k} D_{\alpha,k}) + K(s).$$

Writing $r = |k|$, the digamma expansion of κ_1 gives

$$\kappa_1 = \zeta\left[\epsilon \log \epsilon + \delta \log(z/r) + \log D_{\alpha,k}\right] + O(z^{-2}) \longrightarrow \mu_{LN}^{\zeta}.$$

Indeed, $\log D_{\alpha,k} = -\epsilon \log \epsilon + \frac{1}{2}\log 2 + o(1)$, while $\delta \log(z/r) \to -\tau_{LN}^2$ on the positive branch and is zero on the negative branch. Also $A''(z) \to \tau_{LN}^2$ and $A'''(z+u) = O(z^{-1})$ uniformly for bounded u. Taylor expansion therefore yields

$$\log H(s) = s\kappa_1 + \frac{s^2}{2} A''(z) + O(z^{-1}) \longrightarrow s\mu_{LN}^{\zeta} + \frac{s^2}{2}\tau_{LN}^2$$

uniformly on every bounded real s-interval. MGF convergence gives the normal limit, and exponentiation gives the lognormal limit. □

Relation to the original scaling. The original log-mean is

$$\mathbb{E}[\log V] = \zeta\left[\log \sigma_{\alpha,k} + \epsilon\psi(\epsilon z) - \tfrac{1}{2}\psi(z/2)\right] = \zeta\, \epsilon \log \epsilon + O(1),$$

where $\psi(x) = \frac{d}{dx}\log\Gamma(x)$ is the digamma function. The deterministic correction cancels this drift: $\mathbb{E}[\log \hat{V}] = \mathbb{E}[\log V] + \zeta \log D_{\alpha,k} \to \mu_{LN}^{\zeta}$.

7.6.1. The Centered Logarithmic Variable. For $0 < \alpha < 2$ and $z > 0$, let $U = \log V - \kappa_1$, where $\kappa_1 = \mathbb{E}[\log V]$ is evaluated at the chosen scale. The cumulant generating function (CGF) of this centered logarithmic variable is

$$C(s) := \log \mathbb{E}[e^{sU}] = \log H(s) - s\kappa_1 = K(s) - sK'(0). \tag{7.30}$$

LEMMA 7.7 (Integral representation of the centered log-CGF). For $0 < \alpha < 2$, $z > 0$, and complex s with $z + \zeta \operatorname{Re} s > 0$, the centered log-CGF has the absolutely convergent integral representation

$$C(s) = \int_0^\infty \left(e^{-\zeta sr} - 1 + \zeta sr\right) \frac{e^{-zr} B(r)}{r}\, dr. \tag{7.31}$$

△

PROOF. Indeed, integrating the representation of A'' in Lemma 7.3 twice gives $A(z+\zeta s) - A(z) - \zeta s A'(z) = C(s)$. The compensation $-1 + \zeta sr$ makes the integrand bounded at $r = 0$ and ensures $C(0) = C'(0) = 0$; exponential decay gives convergence at infinity. □

REMARK 7.8 (On the Lévy measure). The Lévy measure of jump magnitudes in this log-space representation of the tilted stable law is

$$\rho(dr) = \frac{e^{-zr} B(r)}{r}\, dr, \qquad r > 0. \tag{7.32}$$

Here $\epsilon = 1/\alpha > g = 1/2$, so $B(r) > 0$; also $\int_0^\infty (1 \wedge r^2)\rho(dr) < \infty$. The Lévy measure ν_U on the logarithmic axis is the image of ρ under $r \mapsto -\zeta r$: ordinary FCM has negative jumps and characteristic FCM has positive jumps.

Thus (7.31) is a compensated Lévy–Khintchine representation[4], with no Gaussian component. In particular, U is infinitely divisible and can be realized as the time-one marginal $U_1 \overset{d}{=} U$ of a centered additive Lévy process $(U_t)_{t \geq 0}$ satisfying $\log \mathbb{E}[e^{sU_t}] = tC(s)$. The process has stationary independent increments; U itself denotes its time-one random variable, not the entire process.

Differentiating (7.31) at $s = 0$ gives the unchanged higher cumulants directly:

$$\begin{aligned} \kappa_n = C^{(n)}(0) = K^{(n)}(0) &= \zeta^n A^{(n)}(z) \\ &= (-1)^n \zeta^n \int_0^\infty r^{n-1} e^{-zr} B(r)\, dr, \qquad n \geq 2. \end{aligned} \tag{7.33}$$

LEMMA 7.9 (Large-z expansion of κ_n). For each fixed integer $n \geq 2$ and fixed $\tau_{LN}^2 > 0$, consider either fixed-sign branch $\zeta \in \{-1, 1\}$ on the exact path

$$h := \frac{1}{z} \longrightarrow 0, \qquad g = \frac{1}{2}, \qquad \delta = \epsilon - g = \frac{\tau_{LN}^2}{h}. \tag{7.34}$$

Then

$$\begin{aligned} \kappa_n &= (n-2)!\, \tau_{LN}^2 (-\zeta h)^{n-2} (1 - R_n), \\ 0 \leq R_n &\leq \frac{n(n-1)h^2}{6\epsilon} \leq \frac{n(n-1)}{6\tau_{LN}^2} h^3. \end{aligned} \tag{7.35}$$

That is, $R_n \approx O_{n,\tau_{LN}}(h^3)$. △

PROOF. To separate its leading term, put

$$b(u) := \frac{1}{1 - e^{-u}} - \frac{1}{u} - \frac{1}{2}, \qquad u > 0.$$

Then the kernel admits the exact decomposition and global bound

$$B(r) = \frac{\tau_{LN}^2}{hr} - q_\epsilon(r), \qquad q_\epsilon(r) := \int_{r/\epsilon}^{2r} b'(u)\,du, \tag{7.36}$$
$$0 \le q_\epsilon(r) \le \frac{(2-1/\epsilon)r}{12} = \frac{\tau_{LN}^2 r}{6\epsilon h}, \qquad r > 0.$$

In particular, $q_\epsilon(r) \le \tau_{LN}^2 r/(6h)$ when $\epsilon \ge 1$, as holds for all sufficiently small h on this path.

In (7.33), rescale $r = ht$ to obtain

$$\kappa_n = (-\zeta)^n h^n \int_0^\infty t^{n-1} e^{-t} B(ht)\,dt.$$

By (7.36),

$$B(ht) = \frac{\tau_{LN}^2}{h^2 t} - q_\epsilon(ht), \qquad 0 \le q_\epsilon(ht) \le \frac{\tau_{LN}^2 t}{6\epsilon}.$$

Inserting this decomposition and using $\int_0^\infty t^{n-2}e^{-t}\,dt = \Gamma(n-1) = (n-2)!$ gives

$$\begin{aligned}\kappa_n &= (-\zeta)^n \tau_{LN}^2 h^{n-2}\left[(n-2)! - \frac{h^2}{\tau_{LN}^2}\int_0^\infty t^{n-1}e^{-t} q_\epsilon(ht)\,dt\right] \\ &= (n-2)!\,\tau_{LN}^2(-\zeta h)^{n-2}(1-R_n),\end{aligned}$$

where

$$R_n := \frac{h^2}{\tau_{LN}^2 (n-2)!}\int_0^\infty t^{n-1}e^{-t} q_\epsilon(ht)\,dt. \tag{7.37}$$

Here $(-\zeta)^n = (-\zeta)^{n-2}$ on both branches. The global kernel bound and $\epsilon = 1/2 + \tau_{LN}^2/h \ge \tau_{LN}^2/h$ give

$$0 \le R_n \le \frac{h^2}{6\epsilon(n-2)!}\int_0^\infty t^n e^{-t}\,dt = \frac{n(n-1)h^2}{6\epsilon} \le \frac{n(n-1)}{6\tau_{LN}^2}h^3,$$

□

The exact-path assumption is needed for this relative error rate: $\delta h \to \tau_{LN}^2$ alone does not control the rate at which the leading cumulant magnitude approaches its limit.

7.6.2. Density Function by Log Inversion. Direct evaluation of the FCM density can lose numerical accuracy at large $|k|$, because the density formula involves products of very large and very small quantities. Fourier inversion in centered log space avoids evaluating these quantities separately.

The characteristic function of U is therefore

$$\varphi_U(t) = e^{C(it)} = \exp(K(it) - itK'(0))\,. \tag{7.38}$$

The deterministic scale term cancels exactly, so the same inversion applies to either branch and to the rescaled variable $\hat{V}$; only κ_1 changes. By conjugate symmetry and the change-of-variable formula,

$$f_U(u) = \frac{1}{\pi}\int_0^\infty \mathrm{Re}\big[\varphi_U(t)e^{-itu}\big]\,dt, \tag{7.39}$$
$$\log f_V(v) = \log f_U(\log v - \kappa_1) - \log v, \qquad v > 0. \tag{7.40}$$

Thus the density can be computed from $\log v$ even when v lies outside the floating-point range.

For stable evaluation, group the log-gamma terms by nearby arguments:

$$\begin{aligned}K(it) = {} & [\log\Gamma(\epsilon z + i\epsilon\zeta t) - \log\Gamma(\epsilon z)] \\ & - [\log\Gamma(gz + ig\zeta t) - \log\Gamma(gz)]\,.\end{aligned}$$

Cancellation-free log-gamma differences. For large positive x and a purely imaginary increment h, direct subtraction of $\log\Gamma(x+h)$ and $\log\Gamma(x)$ loses precision, especially when $|h| \ll x$. Instead, subtract their Stirling expansions algebraically before numerical evaluation. With $r = \log(1+h/x)$ and $c_n = B_{2n}/[2n(2n-1)]$, where B_{2n} are Bernoulli numbers, this gives

$$\log\Gamma(x+h) - \log\Gamma(x) \approx \left(x - \tfrac{1}{2}\right) r + h(\log x + r) - h + \sum_{n=1}^{N} c_n x^{1-2n} \operatorname{expm1}((1-2n)r)\,.$$

Here r is evaluated with `log1p`, while $\operatorname{expm1}(w) = e^w - 1$ avoids subtracting nearly equal numbers in the correction terms. The common terms of order $x\log x$ cancel symbolically, rather than after rounding. Apply this formula to the two brackets above with $(x,h) = (\epsilon z, i\epsilon\zeta t)$ and $(gz, ig\zeta t)$, respectively. The series is asymptotic and must be truncated; for small arguments, use the direct log-gamma difference instead.

A reusable frequency grid gives the trapezoidal approximation

$$f_U(u) \approx \frac{h}{\pi}\left[\frac{1}{2} + \sum_{j=1}^{J} \operatorname{Re}\big(\varphi_U(jh)e^{-ijhu}\big)\right]. \tag{7.41}$$

The frequency step h produces spatial aliases separated by $2\pi/h$; the cutoff Jh controls the omitted Fourier tail. Both must be checked independently, especially near $\alpha = 2$, where decay can be slow. The implementation uses the defaults

$$h = \frac{2\pi}{60\sqrt{\kappa_2}}, \qquad t_{\max} = \frac{40}{\sqrt{\kappa_2}} \text{ initially}, \qquad J = \left\lceil \frac{t_{\max}}{h} \right\rceil.$$

These choices give $J = 382$ when the initial cutoff suffices, as is typical along the lognormal path. The cutoff is increased until $|\varphi_U(t_{\max})| < 10^{-17}$, increasing J accordingly when decay is slow. The spatial period is $2\pi/h = 60\sqrt{\kappa_2}$. A small integrand at the cutoff alone does not bound the integrated tail, and absolute accuracy does not guarantee relative accuracy in the density tails. At $\alpha = 2$, V is a point mass, so ordinary density inversion does not apply.

7.6.3. Relation to the Tempered Stable Law. Near the lognormal limit, along the joint parameter path below, the leading kernel in (7.32) gives a classical tempered stable law with index one. This approximates the centered logarithmic variable associated with the tilted stable representation, not the tilted stable variable itself.

LEMMA 7.10 (Tempered-stable approximation of the centered log-law). This is a continuation of Lemma 7.9. Given $\delta = \epsilon - g$ and $g = \frac{1}{2}$, consider the exact lognormal-compatible path

$$h := \frac{1}{z} > 0, \qquad \delta\, h = \tau_{LN}^2. \tag{7.42}$$

with a fixed $\tau_{LN}^2 > 0$. Thus $h \to 0$ is the large-z limit.

We retain only the leading term of (7.35) as

$$\kappa_n^{TS} = (n-2)!\,\tau_{LN}^2(-\zeta h)^{n-2}, \quad n \geq 2, \tag{7.43}$$

which defines a new log variable denoted as U_{TS}. It is also centered, $\mathbb{E}[U_{TS}] = \kappa_1^{TS} = 0$.

U_{TS} is the time-one law of a Lévy process with the measure of jump magnitudes

$$\rho_{TS}(dr) = \frac{\tau_{LN}^2}{h}\frac{e^{-r/h}}{r^2}\,dr, \qquad r > 0, \tag{7.44}$$

with jumps of sign $-\zeta$.

It satisfies $\int_0^\infty (1 \wedge r^2)\rho_{TS}(dr) < \infty$ and, under full compensation with no Gaussian component, defines a centered, spectrally one-sided, exponentially tempered stable law of stability index one[25, Section 2.1, equation (2.3)].

△

PROOF. U_{TS} has the centered log-CGF

$$\begin{aligned} C_{TS}(s) &:= \log \mathbb{E}[e^{sU_{TS}}] \\ &= \frac{\tau_{LN}^2}{h} \int_0^\infty \left(e^{-\zeta sr} - 1 + \zeta sr\right) \frac{e^{-r/h}}{r^2}\, dr, \qquad h^{-1} + \zeta \operatorname{Re} s > 0. \end{aligned} \tag{7.45}$$

This is the fully compensated Lévy–Khintchine integral with (7.44) and jumps of sign $-\zeta$. Differentiating twice under the integral sign gives

$$C''_{TS}(s) = \frac{\tau_{LN}^2}{h} \int_0^\infty e^{-(h^{-1}+\zeta s)r}\, dr = \frac{\tau_{LN}^2}{1+\zeta hs}.$$

Since $C_{TS}(0) = C'_{TS}(0) = 0$, integration twice yields

$$\begin{aligned} C_{TS}(s) &= \frac{\tau_{LN}^2}{h^2} \left[(1+\zeta hs)\log(1+\zeta hs) - \zeta hs\right] \\ &= \frac{\tau_{LN}^2 s^2}{2}\, {}_2F_1(1,1;3;-\zeta hs). \end{aligned} \tag{7.46}$$

The logarithm is on its principal branch; the stated domain gives $\operatorname{Re}(1+\zeta hs) > 0$. This agrees, after scaling and reflection, with the index-one characteristic exponent in Rosiński[36, Theorem 2.9, equation (2.12)].

For the hypergeometric form, Euler's integral (DLMF 15.6.1[10]), with $q = \zeta hs$, gives

$$\frac{1}{2}\, {}_2F_1(1,1;3;-q) = \int_0^1 \frac{1-t}{1+qt}\, dt = \frac{(1+q)\log(1+q) - q}{q^2},$$

where the value at $q = 0$ is $1/2$ by continuity. Finally, differentiating C''_{TS} another $n-2$ times at zero gives

$$C_{TS}^{(n)}(0) = (n-2)!\, \tau_{LN}^2 (-\zeta h)^{n-2}, \qquad n \geq 2,$$

which proves (7.43); the first cumulant is zero by compensation.

Comparing (7.44) with the Lévy density in Massing[25]

$$\delta_{TS}\, \frac{e^{-\lambda_{TS}\, r}}{r^{1+\alpha_{TS}}}$$

gives the parameter correspondence

$$\alpha_{TS} = 1, \qquad \delta_{TS} = \frac{\tau_{LN}^2}{h} = \delta, \qquad \lambda_{TS} = \frac{1}{h} = z. \tag{7.47}$$

The resulting positive measure is Lévy-integrable at zero because its density is of order r^{-2}, and at infinity because of exponential decay. □

Its variance is $\operatorname{Var}(U_{TS}) = \kappa_2^{TS} = \tau_{LN}^2$. This variance identity is exact for U_{TS}. The original centered FCM log-variable has $\operatorname{Var}(U) = A''(z)$, which only tends to τ_{LN}^2, according to (7.24), along the lognormal path.

7.7. FCM Moment Coordinates

Continuing with the centered logarithmic variable U and its CGF $C(s)$ from Section 7.6.2, we define scale-free moment coordinates for comparing FCM shapes with the lognormal limit.

Definition 7.11 (Moment coordinates of order n).

$$v_{\pm}^{(n)} := \frac{2C(\pm n)}{n^2} = \frac{2}{n^2}\left(\log \mathbb{E}[V^{\pm n}] \mp n\mathbb{E}[\log V]\right), \qquad n > 0. \tag{7.48}$$

These coordinates are nonnegative by Jensen's inequality and invariant under $V \mapsto cV$, $c > 0$.

Lemma 7.12. Equation (7.30) gives

$$v_{\pm}^{(n)} = \frac{2}{n^2}\left[A(z \pm \zeta n) - A(z) \mp \zeta n A'(z)\right]. \tag{7.49}$$

For $0 < \alpha < 2$ and $z > 0$, both moments exist when $n < z + \alpha$. At zero, use $A(0) = -\log(2\epsilon)$; for $-\alpha < x < 0$, use the real logarithm of the positive gamma ratio defining $A(x)$.

Expanding in the logarithmic cumulants from (7.26) gives

$$v_{\pm}^{(n)} = \kappa_2 \pm n\kappa_3/3 + n^2\kappa_4/12 + \cdots$$

This expansion shows how increasing n gives greater weight to log-asymmetry (κ_3) and higher-order log cumulants, including κ_4, making the coordinates more sensitive to the tails. △

Every lognormal law lies on the common diagonal $v_{-}^{(n)} = v_{+}^{(n)} = \tau_{LN}^2$, as shown in Figure 7.2. To describe a point's position relative to this diagonal, we introduce the polar-type coordinates (R_n, θ_n) below.

Lemma 7.13 (Polar-type coordinates).

$$R_n := \frac{1}{2}\left(v_{+}^{(n)} + v_{-}^{(n)}\right), \tag{7.50}$$

$$\theta_n := \operatorname{atan2}\left(v_{+}^{(n)}, v_{-}^{(n)}\right). \tag{7.51}$$

On the lognormal diagonal, R_n equals the log-variance τ_{LN}^2. Away from this diagonal, it need not equal the log-variance, and it is not the Euclidean radius.

Away from the point-mass origin, the polar angle satisfies

$$\tan\theta_n = \frac{C(n)}{C(-n)} = \frac{A(z+\zeta n) - A(z) - \zeta n A'(z)}{A(z-\zeta n) - A(z) + \zeta n A'(z)}. \tag{7.52}$$

The exact angle depends on both α and k. △

Lemma 7.14 (Reflection of moment coordinates). For fixed $0 < \alpha < 2$, $z > 0$, and $0 < n < z + \alpha$, the paired branches $(k_+, k_-) = (z+1, -z)$ satisfy

$$v_{\pm}^{(n)}(\alpha, -z) = v_{\mp}^{(n)}(\alpha, z+1).$$

The corresponding points are mirror images across the lognormal diagonal: they have the same R_n, and their angles sum to $\pi/2$. △

Proof. For $k = -z$, substitute $\zeta = -1$ into (7.49) to obtain

$$v_{\pm}^{(n)}(\alpha, -z) = \frac{2}{n^2}\left[A(z \mp n) - A(z) \pm nA'(z)\right] = v_{\mp}^{(n)}(\alpha, z+1),$$

where the last equality uses $\zeta = +1$ for $k = z + 1$.

Swapping the two coordinates preserves their average and changes the angle to $\pi/2$ minus its original value, proving the remaining claims. □

In these moment planes, ordinary FCM lies below the diagonal, while characteristic FCM lies above it. For example, $k = 6$ is paired with $k = -5$, not $k = -6$.

For fixed n and $\tau_{LN}^2 > 0$, expanding (7.49) along the exact lognormal path $\delta = \tau_{LN}^2 z$ as $z \to \infty$ gives

$$R_n = \tau_{LN}^2 + \frac{n^2 \tau_{LN}^2}{6z^2} + O(z^{-3}),$$
$$\theta_n = \pi/4 - \frac{n\zeta}{3z} + O(z^{-3}).$$

Averaging the two moment coordinates cancels the odd cumulants. Consequently, R_n approaches its lognormal value with a correction of order z^{-2}, while the angular departure from the diagonal is of order z^{-1}. Near the limit, increasing n magnifies this leading angular departure.

Figure 7.2 illustrates the effect for $n = 1, 2, 3, 4$: curves for different values of k are more tightly clustered at lower moment orders and more widely separated at higher orders.

TODO There is a new version of the mixed moment coordinates that is better. It uses CV of V and 1/V, is simpler, match GSaS angle at k=8 out of box. the estimate preserves the arithmetic mean naturally, better MLLK than n=2+4 and current equal blend. It doesn't need complicated explanation.

For presentation, we choose the equal-blend coordinates (b_-, b_+), shown in Figure 7.1, and define

$$b_\pm := \tfrac{1}{2}\left(v_\pm^{(2)} + v_\pm^{(4)}\right), \qquad R_b := \tfrac{1}{2}(b_- + b_+), \qquad \theta_b := \operatorname{atan2}(b_+, b_-). \tag{7.53}$$

This choice is motivated by comparison with the GSaS angle at $k = 8$ in Figure 12.4. We seek similar angles in the two charts while keeping the blending weights simple. Although the $n = 2$ plane is useful on its own, a half-half blend of the $n = 2$ and $n = 4$ coordinates gives a closer angular match at $k = 8$.

The blend is a plotting convention; it does not reproduce the GSaS coordinates exactly for every k. Figure 7.1 shows curves for selected values of k on both sides of the lognormal diagonal.

As an empirical example, the star marks the moment coordinates of the historical VIX sample. The corresponding (α, k) estimates are obtained using the FCM method of moments described in the next subsection.

7.7.1. Method of Moments and the VIX Fit. TODO the CV uses a different estimate. It uses arithmetic mean.

The method of moments estimates distribution parameters by equating selected theoretical moments to their sample counterparts, rather than maximizing the likelihood.

Here the selected statistics are the scale-free coordinates $v_-^{(n)}$ and $v_+^{(n)}$. They are constructed from the ordinary and inverse nth moments of the geometrically normalized variable e^U.

For observations $y_i > 0$, let $\ell_i = \log y_i$ and $\bar{\ell} = N^{-1} \sum_i \ell_i$. The empirical counterparts are

$$\widehat{v}_\pm^{(n)} = \frac{2}{n^2} \log \left[\frac{1}{N} \sum_{i=1}^{N} \exp\{\pm n(\ell_i - \bar{\ell})\} \right]. \tag{7.54}$$

Centering the logarithms is equivalent to dividing the observations by their geometric mean, which removes the multiplicative scale. For a chosen order n, we solve

$$v_-^{(n)}(\widehat{\alpha}_n, \widehat{k}_n) = \widehat{v}_-^{(n)}, \qquad v_+^{(n)}(\widehat{\alpha}_n, \widehat{k}_n) = \widehat{v}_+^{(n)}. \tag{7.55}$$

These two equations are used to estimate the two shape parameters on the chosen branch, within the domain where the required moments exist. The blended estimate instead matches the empirical counterparts of b_- and b_+.




FIGURE 7.1. The equal-blend FCM moment plane with the empirical VIX coordinates marked by a star. The sample contains 8,945 lagged VIX levels dated January 3, 1990–June 30, 2025. Solid and dashed curves denote positive and negative k, respectively; the diagonal is the lognormal limit. Matching the star's coordinates gives the moment-based estimate $(\alpha, k) \approx (0.94543, -4.85905)$.

Once $(\widehat{\alpha}_n, \widehat{k}_n)$ fixes the shape, a separate variance match determines the multiplicative scale, as described below. The procedure is therefore moment-based, even though it uses logarithms of moments; it does not necessarily match the arithmetic mean or moments other than those selected. Likelihood scores are used afterward to assess the fitted densities, not to determine these estimates.

The historical sample consists of 8,945 positive VIX levels. The sample dates range from January 3, 1990, to June 30, 2025. Its blended point, $(\widehat{b}_-, \widehat{b}_+) = (0.09593, 0.15041)$, has $R_b = 0.12317$ and angle $57.47°$ and is marked by the star in Figure 7.1.

Matching these two coordinates on the characteristic branch gives $(\widehat{\alpha}_b, \widehat{k}_b) = (0.94543, -4.85905)$. This estimate satisfies the fourth-moment condition $4 < |k| + \alpha$. The plotted curves are truncated wherever this condition fails.

Next, we examine how the shape estimates depend on the moment order. Here n selects the moments being matched, not a power applied to the FCM variable. We use the FCM family without a power transformation or an additive shift; only a multiplicative scale is fitted after the shape parameters. Figure 7.2 shows the four separate matches summarized below:

n	$\widehat{v}_-^{(n)}$	$\widehat{v}_+^{(n)}$	$\widehat{\alpha}_n$	$\widehat{k}_n$
1	0.11008	0.12799	0.98200	-4.31900
2	0.10249	0.13875	0.97770	-4.41282
3	0.09561	0.15054	0.96689	-4.61368
4	0.08938	0.16207	0.94684	-4.96903

The estimates remain in the same neighborhood across $n = 1, 2, 3, 4$: $\widehat{\alpha}_n$ ranges from about 0.95 to 0.98, and $\widehat{k}_n$ from about -5.0 to -4.3. The empirical coordinates also remain close, with R_n ranging only from 0.119 to 0.126. Thus the inferred FCM shape is relatively insensitive to the choice of moment order over this range. Figure 7.2 shows that the stars occupy a similar region of the characteristic branch in all four planes. The similar SD-matched density profiles in Figure 7.3 provide further support for this stability.

To complete each FCM fit and express its density in VIX units, we estimate a multiplicative scale. For candidate $c \in \{1, 2, 3, 4, \text{half-half}\}$, let $V_c \sim \overline{\chi}_{\widehat{\alpha}_c, \widehat{k}_c}$ have the canonical FCM scale, and write $m_{j,c} = \mathbb{E}[V_c^j]$, where $j = 1, 2$ denotes its first and second moments. The matching of the sample standard deviation $s_y = 7.82840$ gives

$$a_c = \frac{s_y}{\sqrt{m_{2,c} - m_{1,c}^2}}, \qquad Y_c = a_c V_c. \tag{7.56}$$

Figure 7.3 compares all five scaled densities with the same VIX histogram. Matching the standard deviation alone does not ensure that the model mean equals the sample mean, 19.49451; the additive location is fixed at zero. We evaluate each fit using the mean negative log-likelihood (MLLK) on every observation, without excluding the tails:

$$\mathrm{MLLK}_{\mathrm{raw},c} = -\frac{1}{N} \sum_i \log f_{Y_c}(y_i), \tag{7.57}$$

$$\mathrm{MLLK}_{\mathrm{std},c} = \mathrm{MLLK}_{\mathrm{raw},c} - \log s_y. \tag{7.58}$$

The standardized score corresponds to measuring VIX in units of its sample standard deviation; this change of units does not affect the ranking. For $n = 1, 2, 3, 4$ and the blend, these scores are $1.16320, 1.16194, 1.16355, 1.17585, 1.16704$, respectively.

The similar density profiles and likelihood scores indicate that the overall fit is relatively insensitive to the chosen moment order. Lower MLLK is better: the $n = 2$ candidate scores best, with the first three scores especially close. The half-half blend scores between these three candidates and the somewhat higher-scoring $n = 4$ fit.

All five densities capture the broad empirical profile but do not fully reproduce the sharp concentration near VIX 12–13. These moment-derived fits provide stable starting points for likelihood refinement. Serial dependence and changing regimes should also be kept in mind when interpreting this long-run marginal comparison.




FIGURE 7.2. The same VIX sample in the $n = 1, 2, 3, 4$ moment planes. Stars denote empirical coordinates; panel titles give the corresponding characteristic FCM estimates. The model curves use the same color scheme as in Figure 7.1, therefore their labels are ignored to simplify the graphs. The curves spread farther apart at higher orders, while the empirical points remain in the same characteristic-branch neighborhood with similar R_n values.




FIGURE 7.3. The VIX histogram compared with five characteristic FCM densities, each scaled to match the sample standard deviation, with additive location fixed at zero. The four single-order fits and the equal-blend fit have similar density profiles. The final panel compares MLLK scores for VIX divided by its sample standard deviation; lower values are better.

7.8. FCM2: Fractional Chi-Squared-Mean Distribution

If $Z \sim \overline{\chi}_{\alpha,k}$, then $X \sim Z^2$ is FCM2, denoted as $X \sim \overline{\chi}^2_{\alpha,k}$. This is the fractional extension of the classic χ^2_k/k, which is subsumed by it at $\alpha = 1$.

$\overline{\chi}^2_{\alpha,k}$ is used in the fractional F distribution in the area of the squared variable and the quadratic form in the multivariate elliptical distribution.

DEFINITION 7.15. The PDF of FCM2 is

$$\overline{\chi}^2_{\alpha,k}(x) = \frac{1}{2\sqrt{x}}\,\overline{\chi}_{\alpha,k}(\sqrt{x}) \qquad (x \ge 0, \alpha \in [0,2]) \tag{7.59}$$

Expressed in FG and (7.4), it is

$$\begin{aligned}\overline{\chi}^2_{\alpha,k}(x) &:= \mathfrak{N}_{\alpha/2}(x; \sigma = \sigma^2_{\alpha,k}, d = (k-1)/2, p = \alpha/2) \qquad (k > 0) \\ &= \frac{C_{\alpha,k}}{2\,\sigma^2_{\alpha,k}} \left(\frac{x}{\sigma^2_{\alpha,k}}\right)^{k/2-3/2} F_{\frac{\alpha}{2}}\left(\left(\frac{x}{\sigma^2_{\alpha,k}}\right)^{\alpha/2}\right).\end{aligned} \tag{7.60}$$

Or for the negative degree of freedom,

$$\overline{\chi}^2_{\alpha,-k}(x) := \mathfrak{N}_{\alpha/2}(x; \sigma = \sigma^{-2}_{\alpha,k}, d = -k/2, p = -\alpha/2) \tag{7.61}$$

7.8.1. Inverse Tilted Stable Law. According to Section 6.5, (7.60) implies $\overline{\chi}^2_{\alpha,k} = \sigma^2_{\alpha,k}\, T^{-1}_{\alpha/2,\,(k-1)/2}$ $(k > 0)$. The inverse power is one.

For the negative degree of freedom, we have $\overline{\chi}^2_{\alpha,-k} = \sigma^{-2}_{\alpha,k}\, T_{\alpha/2,\,k/2}$. That is, without interference of the scale, the characteristic FCM2 is equivalent to the tilted stable variable: $\chi^2_{\alpha,-k} = T_{\alpha/2,k/2}$.

When dealing with the fractional Gauss hypergeometric function (FGHF) in Section 5.2, we need two more variations from FCM2. The first allows an FCM2 to take a different scale:

$$\overline{\chi}^2_{\alpha,k}(x;\sigma) := \mathfrak{N}_{\alpha/2}(x; \sigma = \sigma, d = (k-1)/2, p = \alpha/2) \qquad (k > 0) \tag{7.62}$$

from which the constant-scale variant is defined by replacing $\sigma_{\alpha,k}$ with $1/2$,

$$\widehat{\chi}^2_{\alpha,k}(x) := \overline{\chi}^2_{\alpha,k}(x; \sigma = 1/4) = \mathfrak{N}_{\alpha/2}(x; \sigma = 1/4, d = (k-1)/2, p = \alpha/2) \qquad (k > 0) \tag{7.63}$$

Notice the hat symbol replaces the bar symbol.

7.8.2. FCM2 CDF. Extending directly from Lemma 6.6, we have:

LEMMA 7.16. The CDF of FCM2 can be represented by γ^*_α as

$$\Phi[\overline{\chi}^2_{\alpha,k}](x) = z^{(k-1+\alpha)/2}\,\gamma^*_{\alpha/2}\left(\frac{k-1+\alpha}{\alpha}, z^{\alpha/2}\right) \qquad (k > 0, \alpha \in [0,2]) \tag{7.64}$$

where $z = x/\sigma^2_{\alpha,k}$. △

7.8.3. FCM2 Subsumes Chi-squared. It is easy to show that $\overline{\chi}^2_{1,k}$ is χ^2_k with a scale of $1/k$, due to (3.15).

7.8.4. Representing FCM by FCM2. In (7.59), let $s = \sqrt{x}$, we get the inverse relation:

$$\overline{\chi}_{\alpha,k}(s) = 2s\,\overline{\chi}^2_{\alpha,k}(s^2) \qquad (s \ge 0) \tag{7.65}$$

Many ratio distribution integrals involving FCM can be rewritten in terms of FCM2, such that

$$\begin{aligned} f(x) &:= \int_0^\infty g(xs)\,\overline{\chi}_{\alpha,k}(s)\,s\,ds \\ &= \int_0^\infty g(x\sqrt{\nu})\,\overline{\chi}^2_{\alpha,k}(\nu)\,\sqrt{\nu}\,d\nu \end{aligned} \tag{7.66}$$

For the CDF case, the incomplete integral can be transformed as

$$\begin{aligned} F(x) := \int_0^x f(x)\,dx &= \int_0^\infty G(xs)\,\overline{\chi}_{\alpha,k}(s)\,ds \\ &= \int_0^\infty G(x\sqrt{\nu})\,\overline{\chi}^2_{\alpha,k}(\nu)\,d\nu \end{aligned} \tag{7.67}$$

where $G(x) := \int_0^x g(x)\,dx$. The lower bound of the incomplete integrals can be $-\infty$ such as $\int_{-\infty}^x dx$ too.

7.8.5. Universal Expression. Assume $x \geq 0$, let $M(x^2) := G(x)/x$ in (7.67) or $g(x)$ in (7.66), we get the universal expression of

$$F(x) = x\int_0^\infty M(x^2\nu)\,\overline{\chi}^2_{\alpha,k}(\nu)\,\sqrt{\nu}\,d\nu \tag{7.68}$$

$$f(x) = \int_0^\infty M(x^2\nu)\,\overline{\chi}^2_{\alpha,k}(\nu)\,\sqrt{\nu}\,d\nu \tag{7.69}$$

Most of the univariate PDFs and CDFs in subsequent chapters can be understood in such framework. It is just a matter of what $M(x)$ is.

When $M(x)$ can be expressed by a Kummer function (apart from a negative sign), these integrals are members of the FGHF in Section 5.2.

7.9. FCM2 Mellin Transform

From (6.5), the Mellin transform of FCM2's PDF is

$$\begin{aligned} &\overline{\chi}^2_{\alpha,k}(x) \overset{\mathcal{M}}{\longleftrightarrow} \overline{\chi}^{2*}_{\alpha,k}(s) \\ &\qquad = (\sigma_{\alpha,k})^{2s-2}\,\frac{\Gamma((k-1)/2)}{\Gamma((k-1)/\alpha)}\frac{\Gamma((s+k/2-3/2)\ \times 2/\alpha)}{\Gamma(s+k/2-3/2)}. \end{aligned} \qquad (k>0) \tag{7.70}$$

Likewise, for the constant-scale variant, it becomes

$$\begin{aligned} &\widehat{\chi}^2_{\alpha,k}(x) \overset{\mathcal{M}}{\longleftrightarrow} \widehat{\chi}^{2*}_{\alpha,k}(s) \\ &\qquad = 2^{2-2s}\,\frac{\Gamma((k-1)/2)}{\Gamma((k-1)/\alpha)}\frac{\Gamma((s+k/2-3/2)\ \times 2/\alpha)}{\Gamma(s+k/2-3/2)}, \end{aligned} \qquad (k>0) \tag{7.71}$$

whose most important special case is $\alpha = 1$,

$$\widehat{\chi}^2_{1,k}(x) \overset{\mathcal{M}}{\longleftrightarrow} \widehat{\chi}^{2*}_{1,k}(s) = \frac{\Gamma(s+k/2-1)}{\Gamma(k/2)} \tag{7.72}$$

$\Gamma(s+k/2-1)$ in $\widehat{\chi}^{2*}_{1,k}(s)$ is just an ordinary gamma function without a fractional coefficient in front of s. This property is the basis that connects the fractional Gauss hypergeometric function to its classic form in Section 5.2.

7.10. FCM2 Moments

From the Mellin transform by $s = n+1$, its n-th moment is

$$\begin{aligned} \mathbb{E}(X^n|\overline{\chi}^2_{\alpha,k}) &= \mathbb{E}(X^{2n}|\overline{\chi}_{\alpha,k}) \\ &= (\sigma_{\alpha,k})^{2n}\,\frac{\Gamma((k-1)/2)}{\Gamma((k-1)/\alpha)}\frac{\Gamma((n+k/2-1/2)\ \times 2/\alpha)}{\Gamma(n+k/2-1/2)}. \end{aligned} \qquad (k>0) \tag{7.73}$$

As mentioned in Section 7.3, due to the nature of a ratio distribution, it is often evaluated as negative moments, $n < 0$. Hence, n is confined in the range of $1/2 - k/2 < n < 0$.

This puts stricter constraint on non-existing moments than FCM when k is not "large enough". For instance, in the case of fractional F distribution in Section 8.4, $k \approx 3$ is in the neighborhood where it second moment barely exists. This makes it rather hard for the statistics of the SPX daily return data set, since its k is just slightly larger than 3 while α is slightly below 1.

7.11. FCM2 Increment of k

LEMMA 7.17. When x^m is multiplied to $\overline{\chi}^2_{\alpha,k}(x)$, it follows a scaling rule where k is incremented to $k+2m$ in the parametrization.

$$x^m\, \overline{\chi}^2_{\alpha,k}(x) = \sigma^{2m}_{\alpha,k}\, Q\, \frac{C_{\alpha,k}}{C_{\alpha,k+2m}}\, \overline{\chi}^2_{\alpha,k+2m}(y). \tag{7.74}$$

where $Q := \sigma^2_{\alpha,k+2m}/\sigma^2_{\alpha,k}$ and $y = Q\,x$. △

PROOF. From (7.60),

$$x^m\, \overline{\chi}^2_{\alpha,k}(x) = \sigma^{2m}_{\alpha,k}\, \frac{C_{\alpha,k}}{2\,\sigma^2_{\alpha,k}} \left(\frac{x}{\sigma^2_{\alpha,k}}\right)^{(k+2m)/2-3/2} F_{\frac{\alpha}{2}}\left(\left(\frac{x}{\sigma^2_{\alpha,k}}\right)^{\alpha/2}\right).$$

We see that $\overline{\chi}^2_{\alpha,k}$ should become $\overline{\chi}^2_{\alpha,k+2m}$ according to the power in the $x^{(k+2m)/2-3/2}$ term, but other parts of the formula need to be adjusted too.

Since

$$\overline{\chi}^2_{\alpha,k+2m}(y) = \frac{C_{\alpha,k+2m}}{2\,\sigma^2_{\alpha,k+2m}} \left(\frac{y}{\sigma^2_{\alpha,k+2m}}\right)^{(k+2m)/2-3/2} F_{\frac{\alpha}{2}}\left(\left(\frac{y}{\sigma^2_{\alpha,k+2m}}\right)^{\alpha/2}\right),$$

we obtain $y = x\,\sigma^2_{\alpha,k+2m}/\sigma^2_{\alpha,k}$ in order to match the two structurally.

Then take the ratio of $x^m\, \overline{\chi}^2_{\alpha,k}(x)/\overline{\chi}^2_{\alpha,k+2m}(y)$ to determine the needed constant, we arrive at (7.74).

□

7.12. Sum of Two Chi-Squares with Correlation

The sum of bivariate variables is studied here.

LEMMA 7.18. Let $Z = Z_1/s_1 + Z_2/s_2$ where Z_1, Z_2 are two independent χ^2_1 variables. The PDF of Z is

$$\begin{aligned}\chi^2_{11}(z, s_1, s_2) &= \frac{\sqrt{s_1 s_2}}{2} e^{-s_2 z/2}\, {}_1F_1\left(\frac{1}{2}, 1; \frac{(s_2 - s_1)z}{2}\right) \\ &= \frac{\sqrt{s_1 s_2}}{2} e^{-(s_1+s_2)z/4} I_0(|s_2 - s_1| z/4)\end{aligned}$$

We apply DLMF 12.6.9 to get the second line, where the symmetry of a, b is explicit since $I_0(x)$ is symmetric. For $x \gg 1$, $I_0(x) \approx e^x/\sqrt{2\pi x}$ (DLMF 10.40.5). △

When $Z_1 = U_1^2$, $Z_2 = U_2^2$, and U_1, U_2 has correlation ρ, then s_1, s_2 must be modified by the eigenvalue solution of $\bar{\Omega}^{-1}\mathrm{diag}(\boldsymbol{s})$ such that

$$\begin{aligned}&\chi^2_{11}(z, s_1, s_2, \rho) = \chi^2_{11}(z, s_1', s_2') \\ &\qquad \text{where} \quad (s_1', s_2') = \frac{(s_1 + s_2) \pm \sqrt{(s_1 - s_2)^2 - 4\rho^2 s_1 s_2}}{2(1-\rho^2)}\end{aligned}$$

CHAPTER 8

Fractional F Distribution

The classic F distribution comes from the ratio of two χ^2 distributions. Assume $U_1 \sim \chi^2_d/d$ and $U_2 \sim \chi^2_k/k$, then $F \sim U_1/U_2$ is an F distribution, $F_{d,k}$.

Two use cases were mentioned in Azzalini (2013)[1]. In Section 4.3 there, the squared variable of a univariate skew-t with k degrees of freedom is distributed as $F_{1,k}$.

In Section 6.2 there, the quadratic from a $d \times d$ multivariate skew-t with k degrees of freedom is distributed as $F_{d,k}$.

Thus, the meaning of d and k is quite clear in such a context: d is the dimension of the multivariate skew-normal process; k is the degree of freedom in the denominator of the ratio distribution. This chapter extends it fractionally.

8.1. Definition

DEFINITION 8.1. Assume $U_1 \sim \chi^2_d/d$ and $U_2 \sim \overline{\chi}^2_{\alpha,k}$, then $F \sim U_1/U_2$ is a fractional F distribution. We use the notation $F \sim F_{\alpha,d,k}$.

The standard PDF of $F_{\alpha,d,k}$ is

$$F_{\alpha,d,k}(x) = \int_0^\infty s\,ds\left[d\,\chi^2_d(dxs)\right]\overline{\chi}^2_{\alpha,k}(s) \tag{8.1}$$

and note that the classic term in the integrand, $d\,\chi^2_d(d\,z)$, is equivalent to our $\bar{\chi}^2_{1,d}(z)$.

The reader should be aware of the subtlety that "ds" in "$s\,ds$" is the calculus notation, while d in $\left[d\,\chi^2_d(dxs)\right]$ is the constant from $F_{\alpha,d,k}$.

The standard CDF of $F_{\alpha,d,k}$ is

$$\Phi[F_{\alpha,d,k}](x) = \int_0^x F_{\alpha,d,k}(x)\,dx \tag{8.2}$$

$$= \int_0^\infty \left[\frac{1}{\Gamma\left(\frac{d}{2}\right)}\gamma\left(\frac{d}{2},\frac{dxs}{2}\right)\right]\overline{\chi}^2_{\alpha,k}(s)\,ds \tag{8.3}$$

since the CDF of a χ^2_d is the regularized lower incomplete gamma function of $\gamma\left(\frac{d}{2},\frac{x}{2}\right)/\Gamma\left(\frac{d}{2}\right)$.

It can also be represented by a fractional Gauss hypergeometric function. See Section 5.2.5.

8.1.1. The Origin of Fractional F. $F_{\alpha,d,k}$ is connected to the quadratic form of a d-dimensional multivariate GAS-SN distribution, $L_{\alpha,k}(0,\bar{\Omega},\boldsymbol{\beta})$. Indeed, its three parameters, α, d, k, are designated such that the symbols convey the same meanings. However, $\bar{\Omega}$ and β doesn't affect the outcome of $F_{\alpha,d,k}$.

To elaborate from Section 15.6, assume Z is a $d \times d$ multivariate skew-normal (SN) distribution $SN(0,\bar{\Omega},\beta)$, and $\overline{\chi}_{\alpha,k}$ is a standard FCM. Then $X = Z/\overline{\chi}_{\alpha,k}$ is an $L_{\alpha,k}(0,\bar{\Omega},\boldsymbol{\beta})$.

The quadratic form of X is $Q = \frac{1}{d}\,X^\intercal\bar{\Omega}^{-1}X$. And $Q \sim F_{\alpha,d,k}$ is a fractional F distribution.

8.1.2. Ratio of Titled Stable Law. It is obvious that Fractional F is a ratio of tilted stable law.

$$\begin{aligned} F_{\alpha,d,k} &\sim \overline{\chi}^2_{1,d} / \overline{\chi}^2_{\alpha,k} \\ &= \frac{\sigma^2_{\alpha,k}}{d} \frac{T_{\alpha/2,(k-1)/2}}{T_{1/2,(d-1)/2}} \end{aligned}$$

8.1.3. Fractional F Subsumes F.

LEMMA 8.2. When $\alpha = 1$, it becomes a classic F. That is, $F_{1,d,k} = F_{d,k}$. △

8.1.4. Fractional F Subsumes GSaS-Squared and GAS-SN-Squared. The following cases are for $d = 1$:

LEMMA 8.3. If $X_1 \sim L_{\alpha,k}$, then $X_1^2 \sim F_{\alpha,1,k}$. △

LEMMA 8.4. If $X_2 \sim L_{\alpha,k}(\beta)$, then $X_2^2 \sim F_{\alpha,1,k}$, independent of β. △

They will be discussed in Chapter 12.

8.2. PDF at Zero

The PDF of an F distribution is singular as $x \to 0$ when $d < 2$. We can see that from

$$\begin{aligned} F_{\alpha,d,k}(x) &\approx \frac{(d/2)^{d/2}}{\Gamma(d/2)} x^{d/2-1} \int_0^\infty s^{d/2}\, ds\, \overline{\chi}^2_{\alpha,k}(s) \\ &= \frac{(d/2)^{d/2}}{\Gamma(d/2)} \mathbb{E}(X^d | \overline{\chi}_{\alpha,k})\, x^{d/2-1} \end{aligned} \tag{8.4}$$

for very small x.

When $d = 1$, the peak is divergent as $F_{\alpha,1,k}(x) \approx \frac{1}{\sqrt{2\pi}} \mathbb{E}(X|\overline{\chi}_{\alpha,k}) \sqrt{x}^{-1}$. But its CDF $\propto \sqrt{x}$.

When $d = 2$, this peak is finite. $F_{\alpha,2,k}(0) = \mathbb{E}(X^2|\overline{\chi}_{\alpha,k})$.

When $d > 2$, $F_{\alpha,d,k}(x)$ drops to zero at $x = 0$. This strange phenomenon seems to indicate that the bivariate system is the lowest dimension to have stable quadratic statistics. And a three dimension system is likely more stable. But we only analyze the bivariate case in this book.

8.3. Mellin Transform

From (7.70), and note that $\bar{\chi}^2_d = \bar{\chi}^2_{1,d}$, the Mellin transform of Fractional F's PDF is

$$F_{\alpha,d,k}(x) \overset{\mathcal{M}}{\longleftrightarrow} (\bar{\chi}^2_{1,d})^*(s)\, (\bar{\chi}^2_{\alpha,k})^*(2-s) \qquad (d > 0, k > 0) \tag{8.5}$$

$$= \left(\sqrt{2d}\, \sigma_{\alpha,k}\right)^{2-2s} \left[\frac{\Gamma((d-1)/2)}{\Gamma(d-1)} \frac{\Gamma((k-1)/2)}{\Gamma((k-1)/\alpha)}\right] \left[\frac{\Gamma(2p(s))}{\Gamma(p(s))} \frac{\Gamma(2q(s)/\alpha)}{\Gamma(q(s))}\right], \tag{8.6}$$

$$\text{where } \; p(s) := s + d/2 - 3/2, \; q(s) := 1/2 + k/2 - s.$$

The number of gamma functions can be reduced via the Legendre duplication formula (A.2).

8.4. Moments

Its n-th moment is

$$\begin{aligned} \mathbb{E}(X^n | F_{\alpha,d,k}) &= d^{-n}\, \mathbb{E}(X^n|\chi^2_d)\, \mathbb{E}(X^{-n}|\overline{\chi}^2_{\alpha,k}) \\ &= \left(\frac{2}{d}\right)^n (d/2)_n\, \mathbb{E}(X^{-n}|\overline{\chi}^2_{\alpha,k}). \end{aligned} \tag{8.7}$$

where $(d/2)_n$ is the Pochhammer symbol, $(a)_n := \Gamma(a+n)/\Gamma(a)$.

Its first moment is $\mathbb{E}(X^{-1}|\overline{\chi}^2_{\alpha,k}) = \mathbb{E}(X^{-2}|\overline{\chi}_{\alpha,k})$, independent of d. This is due to $\mathbb{E}(X|\chi^2_d) = d$.

Note that this first moment is also the second moment of an univariate GAS-SN in (12.9), or simply the variance of the corresponding GSaS.

Its second moment is $(1+2/d)\,\mathbb{E}(X^{-2}|\overline{\chi}^2_{\alpha,k})$. Hence, its variance is

$$\begin{aligned}\operatorname{var}\{F_{\alpha,d,k}\} &= (1+2/d)\,\mathbb{E}(X^{-2}|\overline{\chi}^2_{\alpha,k}) - \mathbb{E}(X^{-1}|\overline{\chi}^2_{\alpha,k})^2 \\ &= (1+2/d)\,\mathbb{E}(X^{-4}|\overline{\chi}_{\alpha,k}) - \mathbb{E}(X^{-2}|\overline{\chi}_{\alpha,k})^2.\end{aligned} \tag{8.8}$$

8.4.1. Stability Issue of the Second Moment. The moment formula appears to be straightforward. But the devil is in the detail.

The stability of moments symbolizes the challenge of stability in the α-stable distribution. Even the second moment has dramatic behavior when k is smaller than 4.

First, we recognize that the first moment of F is actually the second moment of the underlying two-sided distribution because the variable of F is squared. Having a finite and stable first moment in F is quite meaningful. But it is much harder to make sense of the variance when k is too small.

Notice that, when $d \to \infty$, the variance is independent of d,

$$\operatorname{var}\{F_{\alpha,\infty,k}\} = \mathbb{E}(X^{-2}|\overline{\chi}^2_{\alpha,k}) - \mathbb{E}(X^{-1}|\overline{\chi}^2_{\alpha,k})^2$$

This is the most relevant quantity if it exists. Other variance of finite d could be expressed as the following:

$$\operatorname{var}\{F_{\alpha,d,k}\} - \operatorname{var}\{F_{\alpha,\infty,k}\} = \frac{2}{d}\,\mathbb{E}(X^{-2}|\overline{\chi}^2_{\alpha,k}).$$

CHAPTER 9

The Framework of Continuous Gaussian Mixture

The construction of a symmetric two-sided distribution is in the form of a continuous Gaussian mixture. Both the ratio and product distribution methods could be used.

In the case of the symmetric α-stable distribution (SaS)[7], the exponential power distribution comes from its characteristic function (CF)[30]. In our framework, both distributions will be expanded by adding the parameter for the degrees of freedom (k).

In addition, we use positive k for the former and negative k for the latter. Hence, the "$k \in \mathbb{R}$" domain will consolidate both distributions into one.

We would like to present a unified framework and familiarize the reader with the notations, which would otherwise be subtle and confusing.

The results of this chapter are the following. First, the symmetric two-sided distribution can be enriched using the characteristic function transform.

Second, the features of the two-sided distribution are transferred to the one-sided distribution because of the Gaussian mixture.

Third, that one-sided distribution could be transformed to the inverse distribution that represents the marginal distribution of a volatility process. This is more meaningful in applications, such as quantitative finance.

9.1. The Inverse Chi Distribution

Assume the PDF of a two-sided symmetric distribution is $L(x)$ where $x \in \mathbb{R}$. It has zero mean, $\mathbb{E}(X|L) = 0$. Assume the PDF of a one-sided distribution is $\chi(x)\,(x > 0)$ such that

$$L(x) := \int_0^\infty s\,ds\,\mathcal{N}(xs)\,\chi(s) \tag{9.1}$$

This is not new. It is the definition of a ratio distribution with a standard normal variable $\mathcal{N}$. This is the first form of the Gaussian mixture: $L \sim \mathcal{N}/\chi$. A typical example is that L is a Student's t distribution when χ is $\sqrt{\chi^2_k}$.

The skewness is added by replacing the normal distribution $\mathcal{N}$ with its skew-normal counterpart $\mathcal{N}(\beta)$. See next chapter for more details.

It has the equivalent expression in terms of a product distribution using *the inverse distribution* $\chi^\dagger$ such that $L \sim \mathcal{N}\chi^\dagger$[14]. This is the second form of the Gaussian mixture.

The advantage of this expression is that $\chi^\dagger$ is closer to our typical understanding of the marginal distribution of a volatility process. For example, when the Brownian motion process $dX_t = \sigma_t\,dW_t$ is measured in a particular time interval Δt, we have $\Delta X_t \sim L$ and $\sigma_t \sim \chi^\dagger$.

Since χ in the ratio form is more natural in the expression of the α-stable distribution, we are more inclined to use the ratio distribution. The reader should keep this subtlety in mind.

LEMMA 9.1. (Inverse distribution) Let the inverse distribution of χ be $\chi^\dagger$, the relation between their density functions is

$$\chi^\dagger(s) := s^{-2}\,\chi\left(\frac{1}{s}\right) \tag{9.2}$$

such that

$$\int_0^\infty s\,ds\,\mathcal{N}(xs)\,\chi(s) = \int_0^\infty \frac{ds}{s}\,\mathcal{N}\left(\frac{x}{s}\right)\,\chi^\dagger(s) \qquad (x \in \mathbb{R}) \tag{9.3}$$

$$\int_0^\infty s\,ds\,\mathcal{N}(xs)\,\chi^\dagger(s) = \int_0^\infty \frac{ds}{s}\,\mathcal{N}\left(\frac{x}{s}\right)\,\chi(s) \tag{9.4}$$

The proof is straightforward by a change of variable $t = 1/s$. You can move between LHS and RHS easily.

△

9.2. The Characteristic Distributions

The characteristic transform of a symmetric α-stable distribution (SaS) leads to an exponential power distribution. We would like to generalize this concept in this section.

When the PDF of a distribution is represented by a Gaussian mixture, the features of a characteristic transform are transferred to its χ distribution counterpart.

We use the notation $\text{CF}\{g\}(t) = \mathbb{E}(e^{itX}|g)$ to represent the characteristic function transform of the PDF $g(x)$. Note that $\mathcal{N}$ has a special property that its CF is still itself: $\text{CF}\{\mathcal{N}\}(t) = \sqrt{2\pi}\,\mathcal{N}(t)$.

LEMMA 9.2. (Characteristic function transform of L) Let $\phi(t)$ be the CF of L such that $\phi(t) := \text{CF}\{L\}(t) = \int_{-\infty}^{\infty} dx\,\exp(itx)\,L(x)$. (9.1) is transformed to

$$\phi(t) = \sqrt{2\pi}\int_0^\infty ds\,\mathcal{N}\left(\frac{t}{s}\right)\,\chi(s) \qquad (t \in \mathbb{R}) \tag{9.5}$$

The proof is straightforward from the fact that $\text{CF}\{\mathcal{N}(0,\sigma^2)\}(t) = \sqrt{2\pi}\,\mathcal{N}(\sigma\,t)$.

△

(9.5) is quite similar to the RHS of (9.4). It inspires us to define a new distribution pair, L_ϕ and $\chi_\phi^\dagger$, in terms of a product distribution, such that

$$L_\phi(x) := \int_0^\infty \frac{ds}{s}\,\mathcal{N}\left(\frac{x}{s}\right)\,\chi_\phi^\dagger(s) \qquad (x \in \mathbb{R}) \tag{9.6}$$

$$\chi_\phi^\dagger(s) := \frac{s\,\chi(s)}{\mathbb{E}(X|\chi)} \tag{9.7}$$

where $\mathbb{E}(X|\chi)$ is the first moment of χ. Here $\chi_\phi^\dagger$ is the inverse distribution of χ_ϕ, which can be reverse-engineered according to (9.2),

$$\chi_\phi(s) := \frac{s^{-3}}{\mathbb{E}(X|\chi)}\,\chi\left(\frac{1}{s}\right) \tag{9.8}$$

We are in an interesting place: We start with a one-sided distribution χ, we derive two variants from it: χ_ϕ and $\chi_\phi^\dagger$. We also obtain two two-sided distributions: L and L_ϕ.

We shall call χ_ϕ *the characteristic distribution* of χ as it facilitates $L_\phi \sim \mathcal{N}/\chi_\phi$.

9.3. Summary of Gaussian Mixture

In summary, considering the inverse distribution and characteristic transform, we obtain four Gaussian mixture relations. The first two parallel relations in terms of ratio distribution:

$$L \sim \mathcal{N}/\chi;$$

$$L_\phi \sim \mathcal{N}/\chi_\phi, \qquad \chi_\phi(s) := \frac{s^{-3}}{\mathbb{E}(X|\chi)}\,\chi\left(\frac{1}{s}\right).$$

In terms of production distribution, they become

$$L \sim \mathcal{N}\,\chi^\dagger, \qquad \chi^\dagger(s) := s^{-2}\,\chi\left(\frac{1}{s}\right) = \frac{s}{\mathbb{E}(X|\chi_\phi)}\,\chi_\phi(s);$$

$$L_\phi \sim \mathcal{N}\,\chi_\phi^\dagger, \qquad \chi_\phi^\dagger(s) := \frac{s}{\mathbb{E}(X|\chi)}\,\chi(s).$$

Both $\chi^\dagger$ and $\chi_\phi^\dagger$ are the volatility processes that generate L and L_ϕ.

Next, the ϕ suffix in $\chi_\phi(s)$ will be replaced by the *negation* (sign change) of the degree of freedom $(k \to -k)$.

The multiplication of $\dfrac{s}{\mathbb{E}(X|f)}$ is closely related to an increase in the degree of freedom $(k \to k+1)$ of the underlying FG, which is equivalent to the scaling of χ_ϕ and χ.

$$\begin{aligned} g(s) &= \frac{s}{\mathbb{E}(X|\chi)}\,f_k(s;\sigma_k) \\ &= \frac{1}{\sigma}\,f_{k+1}\left(\frac{s}{\sigma};\sigma_{k+1}\right) \end{aligned}$$

9.4. FCM Extensions

In this section, we explicitly define the three extensions of $\overline{\chi}_{\alpha,k}$ and prove their connections. They are sub-families of FG. Hence, their density functions should be expressed by $\mathfrak{N}_{\alpha/2}(x;...)$ in (6.1).

First, we quote the PDF of $\overline{\chi}_{\alpha,k}$ from (7.4) below. For $x \geq 0$ and $k > 0$,

$$\begin{aligned} \overline{\chi}_{\alpha,k}(x) &:= \mathfrak{N}_{\alpha/2}(x;\sigma=\sigma_{\alpha,k}, d=k-1, p=\alpha) \qquad (x\geq 0, k>0) \\ &= (C_{\alpha,k})\,(\sigma_{\alpha,k})^{1-k}x^{k-2}\,F_{\frac{\alpha}{2}}\left(\left(\frac{x}{\sigma_{\alpha,k}}\right)^\alpha\right). \end{aligned} \tag{9.9}$$

The characteristic FCM $\left[\overline{\chi}_\phi\right]_{\alpha,k}$ uses the $-k$ index of $\overline{\chi}_{\alpha,k}$. Its PDF is

$$\begin{aligned} \left[\overline{\chi}_\phi\right]_{\alpha,k}(x) = \overline{\chi}_{\alpha,-k}(x) &:= \mathfrak{N}_{\alpha/2}(x;\sigma=(\sigma_{\alpha,k})^{-1}, d=-k, p=-\alpha). \\ &= (C_{\alpha,k+1})\,(\sigma_{\alpha,k})^{-k}x^{-k-1}\,F_{\frac{\alpha}{2}}\left((\sigma_{\alpha,k}\,x)^{-\alpha}\right). \end{aligned} \tag{9.10}$$

Definition 9.3. The density functions of the two inverse distributions are as follows.

$$\begin{aligned} \overline{\chi}^\dagger_{\alpha,k}(x) &:= \mathfrak{N}_{\alpha/2}(x;\sigma=(\sigma_{\alpha,k})^{-1}, d=-k+1, p=-\alpha). \\ &= (C_{\alpha,k})\,(\sigma_{\alpha,k})^{-k+1}x^{-k}\,F_{\frac{\alpha}{2}}\left((\sigma_{\alpha,k}\,x)^{-\alpha}\right). \end{aligned} \tag{9.11}$$

And

$$\begin{aligned} \left[\overline{\chi}^\dagger_\phi\right]_{\alpha,k}(x) = \overline{\chi}^\dagger_{\alpha,-k}(x) &:= \mathfrak{N}_{\alpha/2}(x;\sigma=\sigma_{\alpha,k}, d=k, p=\alpha). \\ &= (C_{\alpha,k+1})\,(\sigma_{\alpha,k})^{-k}x^{k-1}\,F_{\frac{\alpha}{2}}\left(\left(\frac{x}{\sigma_{\alpha,k}}\right)^\alpha\right). \end{aligned} \tag{9.12}$$

REMARK 9.4. In the following subsections, we will show that only the characteristic FCM is a new distribution family, which we use $-k$ to align its parameter such that $\left[\overline{\chi}_{\phi}\right]_{\alpha,k}(x) := \overline{\chi}_{\alpha,-k}(x)$ in (9.14).

The other two variations are just rescaled $k \pm 1$ of FCM. Since we will add location and scale to a distribution family, they don't add new explanatory power to the data in real-world applications.

9.4.1. Expressed via Alpha-Stable Extremal Distribution. From (4.3), we have $L_{\alpha}(x) = x^{-1}F_{\alpha}(x^{-\alpha})$. The two density functions, $\overline{\chi}^{\dagger}_{\alpha,k}(x)$ and $\left[\overline{\chi}_{\phi}\right]_{\alpha,k}(x)$, can be expressed by $L_{\alpha}(x)$ more elegantly since

$$F_{\frac{\alpha}{2}}(x^{-\alpha}) = x^2\, L_{\frac{\alpha}{2}}(x^2). \tag{9.13}$$

Therefore, the PDF of the characteristic distribution becomes

$$\left[\overline{\chi}_{\phi}\right]_{\alpha,k}(x) := \overline{\chi}_{\alpha,-k}(x) = (C_{\alpha,k+1})\,(\sigma_{\alpha,k})^{-k+2} x^{-k+1}\, L_{\frac{\alpha}{2}}\left((\sigma_{\alpha,k}\,x)^2\right). \tag{9.14}$$

which is a FG-type manipulation in the α-stable extremal distribution.

Likewise, the PDF of the inverse distribution becomes

$$\overline{\chi}^{\dagger}_{\alpha,k}(x) = (C_{\alpha,k})\,(\sigma_{\alpha,k})^{-k+3} x^{-k+2}\, L_{\frac{\alpha}{2}}\left((\sigma_{\alpha,k}\,x)^2\right). \tag{9.15}$$

LEMMA 9.5. Show that

$$\overline{\chi}^{\dagger}_{\alpha,k}(x) = \frac{x}{\mathbb{E}(X|\left[\overline{\chi}_{\phi}\right]_{\alpha,k})}\ \left[\overline{\chi}_{\phi}\right]_{\alpha,k}(x) \tag{9.16}$$

△

PROOF. Divide (9.15) by (9.14). The result is $\overline{\chi}^{\dagger}_{\alpha,k}(x)/\left[\overline{\chi}_{\phi}\right]_{\alpha,k}(x) = x\,\sigma_{\alpha,k}\,C_{\alpha,k}/C_{\alpha,k+1}$, which is $x\,\mathbb{E}(X|\overline{\chi}_{\alpha,k})$ according to (7.9). And $\mathbb{E}(X|\overline{\chi}_{\alpha,k}) = 1/\mathbb{E}(X|\left[\overline{\chi}_{\phi}\right]_{\alpha,k})$. We arrive at the desired result. □

Furthermore, this indicates the following.

LEMMA 9.6. $\overline{\chi}^{\dagger}_{\alpha,k}(x)$ is a rescaled $\left[\overline{\chi}_{\phi}\right]_{\alpha,k-1}(x)$, which is $\overline{\chi}_{\alpha,-k+1}(x)$. That is,

$$\overline{\chi}^{\dagger}_{\alpha,k}(x) = \frac{1}{\sigma}\ \left[\overline{\chi}_{\phi}\right]_{\alpha,k-1}\left(\frac{x}{\sigma}\right), \quad \text{where } \sigma = \frac{\sigma_{\alpha,k-1}}{\sigma_{\alpha,k}}. \tag{9.17}$$

△

PROOF. From (9.14), we have

$$\left[\overline{\chi}_{\phi}\right]_{\alpha,k-1}(x) = (C_{\alpha,k})\,(\sigma_{\alpha,k-1})^{-k+3} x^{-k+2}\, L_{\frac{\alpha}{2}}\left((\sigma_{\alpha,k-1}\,x)^2\right).$$

Compare the argument in $L_{\frac{\alpha}{2}}()$ to (9.15) after the substitution x/σ. We obtain $\sigma^2_{\alpha,k-1}/\sigma^2 = \sigma^2_{\alpha,k}$ and arrive at the desired result for σ. □

9.4.2. Proofs of Other Relations.

LEMMA 9.7. Show that (9.8) is true in the FCM implementation.

$$\left[\overline{\chi}_{\phi}\right]_{\alpha,k}(x) = \frac{x^{-3}}{\mathbb{E}(X|\overline{\chi}_{\alpha,k})}\,\overline{\chi}_{\alpha,k}\left(\frac{1}{x}\right) \tag{9.18}$$

△

PROOF. The proof is straightforward. First, replace x with $1/x$ in (9.9). Second, divide it by (9.10). The result is $\left[\overline{\chi}_{\phi}\right]_{\alpha,k}(x)\,/\left[\overline{\chi}_{\phi}\right]_{\alpha,k}(x) = \sigma_{\alpha,k}\,x^3\,C_{\alpha,k}/C_{\alpha,k+1}$, which is $x^3\,\mathbb{E}(X|\overline{\chi}_{\alpha,k})$ according to (7.9). We arrive at the desired result. □

LEMMA 9.8. Show that

$$\left[\overline{\chi}_{\phi}^{\dagger}\right]_{\alpha,k}(x)=\frac{x}{\mathbb{E}(X|\overline{\chi}_{\alpha,k})}\,\overline{\chi}_{\alpha,k}(x) \tag{9.19}$$

This indicates that $\left[\overline{\chi}_{\phi}^{\dagger}\right]_{\alpha,k}(x)$ is a rescaled $\overline{\chi}_{\alpha,k+1}(x)$. This makes it a trivial case. △

PROOF. Divide (9.9) by (9.12). The result is $\overline{\chi}_{\alpha,k}(x)\,/\left[\overline{\chi}_{\phi}^{\dagger}\right]_{\alpha,k}(x)=\sigma_{\alpha,k}\,x^{-1}\,C_{\alpha,k}/C_{\alpha,k+1}$, which is $x^{-1}\,\mathbb{E}(X|\overline{\chi}_{\alpha,k})$ according to (7.9). We arrive at the desired result. □

These relations from different angles ensure that all the formulas are consistent with each other.

9.5. Application of FCM Variants

I have hypothesized that the distribution of the daily VIX indexes can be fitted by the FCM-type distributions[19]. Now we have a more sophisticated parameterization system. We can attempt to fit the data and study how the FCM variants behave.

The data set is the daily VIX data from 1/1990 to 4/2026. The data is converted to a histogram with about 230 bins. Each VIX cycle could stretch over more than a decade. And we only have less than three decades of data. The shape of the data could be somewhat twisted, depending on where the cutoff is in the current cycle.

9.5.1. The FCM Fit. We fit the data by two methods. The first fit uses the FCM itself with the location parameter, whose PDF is

$$f_1(x)=\frac{1}{\omega}\overline{\chi}_{\alpha,k}\left(\frac{x-\varepsilon}{\omega}\right) \tag{9.20}$$

as shown in Figure 9.1.

The distribution fits the peak nicely. But the peak of the data before the lower bound is sharper than the distribution can explain.

The distribution follows the shape of the right tail in the data until $x=7$. In the far right tail of $x>7$, the data have a fatter tail than the distribution.

9.5.2. The Characteristic FCM Fit. The second fit uses the characteristic FCM without the location parameter, whose PDF is

$$f_2(x)=\frac{1}{\omega}\left[\overline{\chi}_{\phi}\right]_{\alpha,k}\left(\frac{x}{\omega}\right) \tag{9.21}$$

as shown in Figure 9.2.

It is intentional to force the location parameter to zero. We want to examine the behavior of the form $x^{-k}\,L_{\frac{\alpha}{2}}\left(x^2\right)$, which suppresses the distribution to zero when x is small. This provides an equivalent of a location parameter that shifts the density to the right.

The distribution fits the peak nicely. However, the data reach their peak at a location lower than that of the distribution. We also observe that the cutoff point of the data at the lower bound is sharper than that of the distribution.

An interesting result of the fit is $\alpha\sim 1$ which is an exponential tail like the classic χ distribution.

The distribution follows the shape of the right tail in the data until $x=9$. In the far right tail of $x>9$, the distribution has a fatter tail than the data.

REMARK 9.9. The fatter tail of the $\left[\overline{\chi}_{\phi}\right]_{\alpha,k}$ distribution is interesting. In the real world, an extreme VIX reading indicates a collapse of the equity market, which is catastrophic. This can not happen often, or even once if it is extremely large. Indeed, after every large market crash, the U.S. regulator implemented mechanisms to slow the market panic.




FIGURE 9.1. FCM fit of VIX daily data from 1990 to 4/2026. Data is standardized to one standard deviation. The optimal parameters are: $\{\alpha = 0.645, k = 2.39, \varepsilon = 1.19, \omega = 0.86\}$. The top left chart is the PDF vs x. The top right chart is the PDF in the log scale vs x. The bottom left chart is the complimentary CDF vs x. The bottom right chart is the complimentary CDF in the log scale vs x.

It appears that the FCM distribution is too conservative in the right tail, while the characteristic FCM is too aggressive. The right tail behavior of real-world data lies between the two, which is likely determined empirically and could evolve over time.




FIGURE 9.2. Characteristic FCM fit of VIX daily data from 1990 to 4/2026. Data is standardized to one standard deviation. This fit forces the location parameter to zero. The optimal parameters are: $\{\alpha = 1.035, k = 4.45, \omega = 2.33\}$. The top left chart is the PDF vs x. The top right chart is the PDF in the log scale vs x. The bottom left chart is the complimentary CDF vs x. The bottom right chart is the complimentary CDF in the log scale vs x.

Part 3

Two-Sided Univariate Distributions

CHAPTER 10

SN: The Skew-Normal Distribution - Review

10.1. Definition

The skew-normal distribution family is well documented in A. Azzalini's 2013 monograph[1]. We recap the results and clarify the symbology. My contribution is to incorporate the skew-normal methodology into the fractional distributions wherever suitable. The enhanced distributions are flexible and can adapt to many different shapes and tails with high skewness and kurtosis.

10.1.1. The Selective Sampling. The *selective sampling* method is used to inject skewness into the stochastic system, which is otherwise symmetric. This mechanism is fairly common in an applied context, for example, in social sciences, where a variable X_0 is observed only when a correlated variable X_1, which is usually unobserved, satisfies a certain condition (p.128 of [1]).

In quantitative finance, the condition could be market regimes. In a two-regime model, a market index such as the S&P 500 index (SPX) is classified into the growth regime or the crash regime at a given time. It is well known that the volatility of the market behaves differently in each regime. In the growth regime, volatility tends to be low, and the market is trending upward. In the crash regime, volatility tends to be high, and the market is trending downward.

A univariate random variable $Z \sim SN(0, 1, \beta)$ is a standard skew-normal variable with skew parameter $\beta \in \mathbb{R}$ (Section 2.1 of [1]). The sign of β determines the sign of its skewness (10.14).

One of its stochastic representations is

$$Z = \begin{cases} X_0 & \text{if } X_1 < \beta X_0, \\ -X_0 & \text{otherwise.} \end{cases} \tag{10.1}$$

where X_0, X_1 are independent $\mathcal{N}(0, 1)$ variables.

An alternative representation uses filtering, or rejection, such that $Z = (X_0 | X_1 < \beta X_0)$. That is, X_0 is accepted as Z only when the condition $X_1 < \beta X_0$ is satisfied. Otherwise, it is discarded.

10.1.2. The PDF and CDF. The standard PDF is

$$\mathcal{N}(x; \beta) := 2\,\mathcal{N}(x)\Phi_{\mathcal{N}}(\beta x), \quad (x \in \mathbb{R}) \tag{10.2}$$

where $\mathcal{N}(x)$ and $\Phi_{\mathcal{N}}(x)$ are the PDF and CDF of $\mathcal{N}(0, 1)$.

Its extremal distribution occurs at $\beta \to \infty$, where $\Phi_{\mathcal{N}}(\beta x)$ becomes a step function. The PDF becomes that of a half-normal distribution.

The standard CDF is

$$\Phi_{SN}(x; \beta) := \Phi_{\mathcal{N}}(x) - 2\,T(x, \beta) \tag{10.3}$$

where $T(h, a)$ is called the Owen's T function[31]. Its numerical methods are widely implemented in modern software packages.

Several important properties are quoted from Proposition 2.1 of [1]:

- $\mathcal{N}(0; 0) = 1/\sqrt{2\pi}$. Universal anchor at $x = 0, \beta = 0$.
- $\mathcal{N}(x; 0) = \mathcal{N}(x)$. Continuity at $\beta = 0$.
- $\mathcal{N}(-x; \beta) = \mathcal{N}(x; -\beta)$. This is the reflection rule.

- $Z^2 \sim \chi^2_1$, irrespective of β.

Notice that Z^2 is independent of β. This is an important property, but may not be intuitive for new students. This is due to the fact that the squares of X_0 and $-X_0$ are the same in (10.1). This property is carried into the quadratic form of the multivariate elliptical distribution.

10.2. The Location-Scale Family

Its location-scale family is $Y = \xi + \omega Z \sim SN(\xi, \omega^2, \beta)$, where $\xi \in \mathbb{R}$ and $\omega > 0$. Its PDF becomes

$$\frac{1}{\omega}\mathcal{N}\left(\frac{x-\xi}{\omega};\beta\right). \tag{10.4}$$

10.3. Invariant Quantities

The following quantity plays an important role in the selective sampling concept of SN:

$$\delta = \frac{\beta}{\sqrt{1+\beta^2}}, \quad \delta \in (-1,1). \tag{10.5}$$

It can be thought of as some kind of correlation in the following. Inversely, β can be calculated from

$$\beta = \frac{\delta}{\sqrt{1-\delta^2}}. \tag{10.6}$$

These two quantities will appear in many places in the ensuing chapters. They are invariants in the context of the multivariate elliptical distribution, called the Canonical Form.

In a trigonometry representation, one can think of δ as $\sin(\theta)$ of a right triangle, where one leg is 1, the other leg is β, and θ is the angle facing β.

Three representations use δ as the correlation coefficient to generate SN. (Section 2.1.3 of [1]) First, designate the correlation matrix as

$$\bar{\Omega} = \begin{pmatrix} 1 & \delta \\ \delta & 1 \end{pmatrix}.$$

The Cholesky factor of $\bar{\Omega}$ is

$$L \;=\; \begin{pmatrix} 1 & 0 \\ \delta & \sqrt{1-\delta^2} \end{pmatrix},$$

so that $L\,L^T = \bar{\Omega}$.

Assume U_0 and U_1 are two independent $\mathcal{N}(0,1)$ variates. The first representation of $Z \sim SN(0,1,\beta)$ is

$$Z = \begin{cases} X_0 & \text{if } X_1 > 0, \\ -X_0 & \text{otherwise.} \end{cases} \tag{10.7}$$

where X_0, X_1 are marginals of a standard correlated normal bivariate with $\text{cor}\{X_0, X_1\} = \delta$ such that

$$\begin{pmatrix} X_0 \\ X_1 \end{pmatrix} \;=\; L \begin{pmatrix} U_0 \\ U_1 \end{pmatrix}.$$

The second representation is from

$$\begin{pmatrix} - \\ Z \end{pmatrix} \;=\; L \begin{pmatrix} U_0 \\ |U_1| \end{pmatrix}$$

such that $Z = \sqrt{1-\rho^2}\,U_0 + \delta\,|U_1| \sim SN(0,1,\beta)$.

The third representation is $Z = \max\{X_0, X_1\} \sim SN(0, 1, \beta)$, where X_0, X_1 are marginals of a standard correlated bivariate such that

$$\begin{pmatrix} X_0 \\ X_1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ \rho & \sqrt{1-\rho^2} \end{pmatrix} \begin{pmatrix} U_0 \\ U_1 \end{pmatrix}.$$

and $\text{cor}\{X_0, X_1\} = \rho = 1 - 2\delta^2$.

10.4. Mellin Transform

The following result is elegant, but also peculiar. It is discovered by the author.

LEMMA 10.1. The Mellin transform of the SN PDF is

$$\mathcal{N}(x;\beta) \overset{\mathcal{M}}{\longleftrightarrow} \mathcal{N}^*(s;\beta) := 2\,\mathcal{N}^*(s)\,\Phi[t_s](\beta\sqrt{s}), \tag{10.8}$$
$$\text{where} \quad \mathcal{N}^*(s) = \frac{1}{2}\frac{2^{s/2}}{\sqrt{2\pi}}\Gamma\left(\frac{s}{2}\right)$$

is the Mellin transform of the PDF of $\mathcal{N}(0,1)$ in (2.9). And $\Phi[t_k](x)$ is the CDF of a Student's t distribution with k degrees of freedom. But it is used in a strange way, where s substitutes k and goes into x at the same time.

△

PROOF. We prove (10.8) via the CDF of GSaS with $\alpha = 1$. By definition,

$$\mathcal{N}^*(s;\beta) = \int_0^\infty x^{s-1}\ [2\mathcal{N}(x)\Phi_{\mathcal{N}}(\beta x)]\,dx.$$

We use the known result from $\bar{\chi}_{1,k}$ where

$$x^{k-1}\mathcal{N}(x) = \frac{2^{k/2-1}\Gamma(k/2)}{\sqrt{2\pi k}}\,\bar{\chi}_{1,k}(x/\sqrt{k}) = \frac{1}{\sqrt{k}}\,\mathcal{N}^*(k)\,\bar{\chi}_{1,k}(x/\sqrt{k}).$$

Then

$$\begin{aligned}\mathcal{N}^*(s;\beta) &= \frac{2\,\mathcal{N}^*(s)}{\sqrt{s}}\int_0^\infty \Phi_{\mathcal{N}}(\beta x)\,\bar{\chi}_{1,s}(x/\sqrt{s})\,dx \\ &= 2\,\mathcal{N}^*(s)\int_0^\infty \Phi_{\mathcal{N}}(\beta\sqrt{s}t)\,\bar{\chi}_{1,s}(t)\,dt \quad \text{via} \quad t = x/\sqrt{s}.\end{aligned}$$

The integral is exactly the CDF of a GSaS, $L_{1,s}$, with the argument $\beta\sqrt{s}$. That is, $\mathcal{N}^*(s;\beta) = 2\,\mathcal{N}^*(s)\,\Phi[L_{1,s}](\beta\sqrt{s})$.

When $\alpha = 1$, $L_{1,s}$ becomes t_s. Therefore, $\mathcal{N}^*(s;\beta) = 2\,\mathcal{N}^*(s)\,\Phi[t_s](\beta\sqrt{s})$.

□

The beauty of this lemma is that $\mathcal{N}^*(s;\beta)$ is the multiplication of a symmetric component and a skew component, just like its PDF counterpart.

From (2.14), we also obtain that

$$\Phi_{SN}(0;\beta) = 1 - \mathcal{N}^*(1;\beta) = \frac{1}{2} - \frac{1}{\pi}\arctan(\beta). \tag{10.9}$$

This is due to $\mathcal{N}^*(1) = \frac{1}{2}$ and $\Phi[t_1](\beta) = \frac{1}{2} + \frac{1}{\pi}\arctan(\beta)$. This result is stated in Proposition 2.7 of [1], and is proved here via the Mellin transform.

10.4.1. Mellin Transform of Owen's T Function. Another peculiar result from the Mellin transform is

LEMMA 10.2.

$$T(x,\beta) \overset{\mathcal{M}}{\longleftrightarrow} s^{-1}\,\mathcal{N}^*(s+1)\,\left[\Phi[t_{s+1}](\beta\sqrt{s+1})-\frac{1}{2}\right] \tag{10.10}$$

△

PROOF. Define the upper incomplete integral as

$$\begin{aligned}\Gamma_f(x) := \int_x^\infty \mathcal{N}(x;\beta)\,dx &= 1-\Phi_{SN}(x;\beta)\\ &= 1-\Phi_{\mathcal{N}}(x)+2\,T(x,\beta)\end{aligned}$$

According to Lemma 2.6, its Mellin transform is

$$\begin{aligned}\Gamma_f(x) &\overset{\mathcal{M}}{\longleftrightarrow} s^{-1}\,\mathcal{N}^*(s+1;\beta)\\ &= 2s^{-1}\,\mathcal{N}^*(s+1)\,\Phi[t_{s+1}](\beta\sqrt{s+1})\end{aligned}$$

Combining the two results above, we obtain

$$T(x,\beta)=\frac{\Gamma_f(x)-(1-\Phi_{\mathcal{N}}(x))}{2} \overset{\mathcal{M}}{\longleftrightarrow} s^{-1}\,\mathcal{N}^*(s+1)\,\left[\Phi[t_{s+1}](\beta\sqrt{s+1})-\frac{1}{2}\right]$$

where $1-\Phi_{\mathcal{N}}(x) \overset{\mathcal{M}}{\longleftrightarrow} s^{-1}\,\mathcal{N}^*(s+1)$.

□

10.5. Moments

LEMMA 10.3. According to Section 2.2.2, by assigning $s=n+1$, the Mellin transform is converted to the moment formula. It is easy to show that the n-th moment of Z is

$$\begin{aligned}\mathbb{E}(Z^n)=\mathbb{E}(X^n|\mathcal{N}(\beta)) &= \mathcal{N}^*(n+1;\beta)+(-1)^n\mathcal{N}^*(n+1;-\beta)\\ &= 2\mathcal{N}^*(n+1)\times\begin{cases}1, & \text{when } n \text{ is even},\\ 1-2\Phi[t_{n+1}](-\beta\sqrt{n+1}), & \text{when } n \text{ is odd}.\end{cases}\end{aligned} \tag{10.11}$$

The even moments are identical to those of $\mathcal{N}(0,1)$. It is the odd moments that make the difference when $\beta\neq 0$.

△

The first four moments of Z' have simple analytic forms. Its first moment is

$$\mu_z = b\,\delta, \quad \text{where } b=\sqrt{2/\pi}. \tag{10.12}$$

The second moment is simply 1. Its variance is

$$\sigma_z^2 = 1-(b\,\delta)^2. \tag{10.13}$$

The third moment is $b\,\delta(3-\delta^2)$. Its skewness is

$$\gamma_1\{Z\}=\frac{4-\pi}{2}\,\frac{\mu_z^3}{\sigma_z^3}. \tag{10.14}$$

The fourth moment is 3. Its kurtosis is

$$\gamma_2\{Z\}=2(\pi-3)\,\frac{\mu_z^4}{\sigma_z^4}. \tag{10.15}$$

The maximum skewness of SN is approximately 0.9953 and the maximum kurtosis is 0.8692. They are not very interesting, since the extremal distribution is just a half-normal distribution.

However, these analytical forms are useful when SN is extended to GAS-SN. Both skewness and kurtosis are extended to much wider ranges, or even infinity!

CHAPTER 11

GAS: Generalized Alpha-Stable Distribution (Experimental)

This chapter studies how a degrees-of-freedom parameter k can be added to the α-stable distribution L_α^θ through the Mellin transform framework. The construction leads to the generalized α-stable distribution (GAS), denoted by $L_{\alpha,k}^\theta$.

This experiment is an early attempt and one of the cleanest approaches to understanding how k interacts with skewness. It is a valuable lesson on the mathematical structure of the α-stable distribution. Therefore, it is documented in this chapter.

The GAS construction is useful because it exposes how the new parameter k interacts with skewness in the stable-law setting. At the same time, the resulting distribution has a discontinuity associated with the reflection rule, which limits its direct use as a final statistical model.

Readers who are mainly interested in the final skew-normal-based construction may treat this chapter as optional background.

Thus, the main role of this chapter is structural. It explains how the fractional χ distribution and GSaS arise from the stable-law framework, and it motivates the later shift to the skew-normal approach, where skewness can be incorporated while preserving continuity of the PDF.

11.1. Definition

First, we recap the Mellin transform (4.5) of the PDF of the α-stable distribution from Section 4.3,

$$L_\alpha^\theta(x) \overset{\mathcal{M}}{\longleftrightarrow} \epsilon \left[\frac{\Gamma(s)}{\Gamma(1-\gamma+\gamma s)}\right] \left[\frac{\Gamma(\epsilon(1-s))}{\Gamma(\gamma(1-s))}\right].$$

It is interpreted in Lemma 4.4 as a multiplication of two components,

$$L_\alpha^\theta(x) \overset{\mathcal{M}}{\longleftrightarrow} \tilde{M}_\gamma^*(s)\, {\overline{\chi}_{\alpha,1}^\theta}^*(2-s).$$

The PDF of the second term $\overline{\chi}_{\alpha,1}$ is defined as

$$\overline{\chi}_{\alpha,1}^\theta(x) \overset{\mathcal{M}}{\longleftrightarrow} {\overline{\chi}_{\alpha,1}^\theta}^*(s) \;\propto \frac{\Gamma(\epsilon(s-1))}{\Gamma(\gamma(s-1))},$$

apart from the normalization constant and scale in the PDF. It is interpreted as the FCM of "one degree of freedom" in Section 7.1.

In (7.1) it is shown that the "degrees of freedom" parameter k is added to the FCM by replacing $s-1$ with $s+k-2$ such that

$$\overline{\chi}_{\alpha,k}^\theta(x) \overset{\mathcal{M}}{\longleftrightarrow} {\overline{\chi}_{\alpha,k}^\theta}^*(s) \;\propto \frac{\Gamma(\epsilon(s+k-1))}{\Gamma(\gamma(s+k-1))}. \tag{11.1}$$

Next, it is natural to use ${\overline{\chi}_{\alpha,k}^\theta}^*(s)$ in the Mellin space to extend L_α^θ as follows.

Definition 11.1 (The ratio-distribution representation of (unadjusted) GAS). The Mellin transform of the PDF of (unadjusted) GAS is defined as

$$\tilde{L}_{\alpha,k}^\theta(x) \overset{\mathcal{M}}{\longleftrightarrow} \tilde{M}_\gamma^*(s)\, {\overline{\chi}_{\alpha,k}^\theta}^*(2-s) \tag{11.2}$$

Based on the Mellin transform, its PDF can be written in a ratio distribution form,

$$\tilde{L}^{\theta}_{\alpha,k}(x) := \int_0^{\infty} \tilde{M}_{\gamma}(xs)\, \bar{\chi}^{\theta}_{\alpha,k}(s)\, s\, ds \quad (x \geq 0) \tag{11.3}$$

Since the Mellin integral is only valid for $x \geq 0$, it is supplemented with *the reflection rule*:

$$\tilde{L}^{\theta}_{\alpha}(-x) := \tilde{L}^{-\theta}_{\alpha}(x) \tag{11.4}$$

Thus, we have constructed a version of GAS for $x \in \mathbb{R}$, which produces fat tails and skewness -

(1) $\tilde{L}^{\theta}_{\alpha,k}$ subsumes the α-stable distribution L^{θ}_{α}.
(2) $\tilde{L}^{\theta}_{\alpha,k}$ subsumes Student's t distribution t_k.
(3) $\tilde{L}^{\theta}_{\alpha,k}$ subsumes the power-exponential distribution, with the proper definition of negative k in FCM.

What is wrong with it? The problem is that the PDF and its derivatives are discontinuous at $x = \pm 0$ when $k \neq 1$ and $\theta \neq 0$.

The remaining sections of this chapter will explain this problem and provide a remediation. The reader who just wants to explore the skew-normal implementation can safely skip the rest of this chapter. The conclusion is that such discontinuity makes the PDF far from mathematical elegance, which motivates the author to explore other alternatives. The answer is to abandon the M-Wright kernel for skewness ($\tilde{M}_{\gamma}(xs)$ in (11.3)), and integrate with the skew-normal distribution, outlined in the next chapter.

11.2. Limitation

The issue of discontinuity of the PDF $\tilde{L}^{\theta}_{\alpha,k}(x)$ at $x = 0$ is encountered when $k \neq 1$. We lay out a generic framework to understand and address it.

Assume that the unadjusted two-sided density function is $\tilde{f}(x) := \tilde{L}^{\theta}_{\alpha,k}(x)$, which is discontinuous at $x = 0$. It also must satisfy the reflection rule, where, for $x > 0$, $\tilde{f}(x) := \tilde{f}^{+}(x)$ and $\tilde{f}(-x) := \tilde{f}^{-}(-x)$.

$\tilde{f}(x)$ can be expanded at $x = 0$ in terms of x by

$$\tilde{f}^{\pm}(x) := \tilde{L}^{\pm\theta}_{\alpha,k}(x) = \tilde{f}^{\pm}_0 + \tilde{f}^{\pm}_1\, x + ... \tag{11.5}$$

where $\tilde{f}^{\pm}_0$ are the densities at $x = 0$, and $\tilde{f}^{\pm}_1$ are the respective slopes (aka the first derivatives).

The series expansion can be achieved via either (11.3), or (11.2) in conjuction with Ramanujan's master theorem in Section 2.3, such that

$$\tilde{f}^{+}_0 = \frac{\gamma^{1-\gamma}}{\Gamma(1-\gamma)}\, E(X|\bar{\chi}^{\theta}_{\alpha,k}), \tag{11.6}$$

$$\tilde{f}^{+}_1 = \frac{-\gamma^{1-2\gamma}}{\Gamma(1-2\gamma)}\, E(X^2|\bar{\chi}^{\theta}_{\alpha,k}). \tag{11.7}$$

Notice that they are based on the first and second moments of $\bar{\chi}^{\theta}_{\alpha,k}$. $(\tilde{f}^{-}_0, \tilde{f}^{-}_1)$ are obtained by applying the reflection rule from $(\tilde{f}^{+}_0, \tilde{f}^{+}_1)$. That is, θ is replaced with $-\theta$, and γ with $1-\gamma$ in every occurrence of the formula.

Furthermore, it is known that

$$\int_0^{\infty} \tilde{f}^{+}(x)\, dx = \gamma, \qquad \int_0^{\infty} \tilde{f}^{-}(x)\, dx = 1-\gamma. \tag{11.8}$$

These two are the only conditions required for $\tilde{f}^{\pm}(x)$.

The discontinuity occurs because $\tilde{f}^{+}_0 \neq \tilde{f}^{-}_0$ and $\tilde{f}^{+}_1 \neq \tilde{f}^{-}_1$ when $k \neq 1$ and $\theta \neq 0$. In fact, this is true for all orders of derivatives $\tilde{f}^{+}_n \neq \tilde{f}^{-}_n$ in the n-th term, $\tilde{f}^{\pm}_n\, x^n$.

Obviously, when $\theta = 0$, the density function is symmetric by definition: $\tilde{f}^+(x) = \tilde{f}^-(x)$. There is no issue here. So the issue is specific to the injection of skewness from $\theta \neq 0$.

On the other hand, when $k = 1$, the density function is continuous under the reflection rule, regardless the value of θ. This is the original α-stable distribution. It is perfectly fine. So the issue is specific to our attempt of adding degrees of freedom $k \neq 1$.

Either one of θ or k are fine, but when we try to do both, the distribution is broken, so to speak. That is the limitation. The dilemma is that adding θ and k is exactly what we try to achieve.

11.3. Workaround

An adjustment algorithm is proposed such that the PDF and its first derivative are continuous.

DEFINITION 11.2 (The adjusted GAS). The PDF of the adjusted GAS is defined as

$$L^{\pm\theta}_{\alpha,k}(x) := \frac{1}{A^{\pm}\sigma^{\pm}}\,\tilde{f}^{\pm}(x)\left(\frac{x}{\sigma^{\pm}}\right) \quad (x \geq 0) \tag{11.9}$$

It is required that (a) the new density function satisfies the reflection rule of $L^{\theta}_{\alpha,k}(-x) := L^{-\theta}_{\alpha,k}(x)$; (b) $A^{\pm}, \sigma^{\pm}$ are constrained by the continuity conditions that, at $x = 0$, both its density is continuous: $L^{\theta}_{\alpha,k}(0) = L^{-\theta}_{\alpha,k}(0)$; and its slope is continuous: $\frac{d}{dx} L^{\theta}_{\alpha,k}(0) = -\frac{d}{dx} L^{-\theta}_{\alpha,k}(0)$.

With such definition, we proceed to find the solutions of $A^{\pm}, \sigma^{\pm}$. The solutions form a distribution family. There is a canonical solution, simple and elegant, from which all other solutions are derived as a member of the location-scale family.

A member in the location-scale family shares the same "shapes" such as the skewness and kurtosis. Apart from the location and scale, it brings nothing new to the table. Hence, we can focus on analyzing the canonical distribution.

DEFINITION 11.3 (Two essential quantities for the canonical distribution). We define two essential quantities:

$$\Sigma := -\frac{\tilde{f}^+_0}{\tilde{f}^-_0}\frac{\tilde{f}^-_1}{\tilde{f}^+_1} \tag{11.10}$$

$$\Psi := \Sigma\,\frac{\tilde{f}^+_0}{\tilde{f}^-_0} \quad = -\left(\frac{\tilde{f}^+_0}{\tilde{f}^-_0}\right)^2 \frac{\tilde{f}^-_1}{\tilde{f}^+_1} \tag{11.11}$$

Notice that $\tilde{f}^+_0/\tilde{f}^-_0$ is the ratio of the original densities from two sides of $x = 0$. And $\tilde{f}^-_1/\tilde{f}^+_1$ is the ratio of the slopes of the two sides. Since $\tilde{f}^-_1, \tilde{f}^+_1$ always have the opposite signs, Σ is a positive quantity.

Note that Σ is singular when $\gamma = 1/2$. Both $\tilde{f}^-_1, \tilde{f}^+_1$ approach zero at the same speed. Hence, $\Sigma \to 1$ and $\Psi \to 1$.

The most important contribution is the discovery of the canonical distribution.

DEFINITION 11.4 (The canonical GAS). The canonical GAS distribution is defined according to $\sigma^+ = 1$ and $\sigma^- = \Sigma$. Hence, its PDF for $x \geq 0$ is (with the hat symbol)

$$\widehat{L}^{\theta}_{\alpha,k}(x) := \frac{1}{A^+}\,\tilde{f}^+(x) \tag{11.12}$$

$$\widehat{L}^{-\theta}_{\alpha,k}(x) := \frac{1}{A^-\Sigma}\,\tilde{f}^-\left(\frac{x}{\Sigma}\right) \tag{11.13}$$

where $A^+ = \gamma + \Psi(1-\gamma)$ and $A^- = A^+/\Psi$ from Lemma 11.7.

The reflection rule applies: $\widehat{L}^{\theta}_{\alpha,k}(-x) := \widehat{L}^{-\theta}_{\alpha,k}(x)$.

11.3.1. The Location-scale Family. The following lemmas show that all other solutions must obey $\sigma^-/\sigma^+ = \Sigma$. They are just the location-scale family of the canonical distribution.

Briefly, all other solutions are defined by a choice of scale $\sigma^+ > 0$, such that

$$L^{\theta}_{\alpha,k}(x) := \frac{1}{\sigma^+}\,\widehat{L}^{\theta}_{\alpha,k}\left(\frac{x}{\sigma^+}\right) \tag{11.14}$$

For instance, we found that $\sigma^+ = \Sigma^\gamma$ to be a very good alternative. In the remark of Definition 11.9, we show that the n-th moment of $L^{\theta}_{\alpha,k}$ is just that of $\widehat{L}_{\alpha,k}$ multiplied by its scale $(\sigma^+)^n$.

LEMMA 11.5. The requirement that the density and slope of the *adjusted* density function should be smooth at $x = 0$ leads to

$$\frac{1}{A^+\sigma^+}\,\tilde{f}^+_0 = \frac{1}{A^-\sigma^-}\,\tilde{f}^-_0 \tag{11.15}$$

$$\frac{1}{A^+(\sigma^+)^2}\,\tilde{f}^+_1 = -\frac{1}{A^-(\sigma^-)^2}\,\tilde{f}^-_1 \tag{11.16}$$

△

PROOF. To solve $A^\pm$ and $\sigma^\pm$, take (11.9) and carry out the series expansions from (11.5):

$$L^{\pm\theta}_{\alpha,k}(x) = \frac{\tilde{f}^\pm_0}{A^\pm\sigma^\pm} + \frac{\tilde{f}^\pm_1}{A^\pm(\sigma^\pm)^2}\,x + ... \tag{11.17}$$

(11.15) is straightforward from requiring $L^{\theta}_{\alpha,k}(0) = L^{-\theta}_{\alpha,k}(0)$ in (11.17). Likewise, (11.16) is the result of $\frac{d}{dx}L^{\theta}_{\alpha,k}(0) = \frac{d}{dx}L^{-\theta}_{\alpha,k}(0)$ from (11.17). □

LEMMA 11.6. The equations in Lemma 11.5 lead to the following invariant:

$$\frac{\sigma^-}{\sigma^+} = \Sigma \tag{11.18}$$

△

PROOF. Divide the LHS and RHS of (11.15) by those of (11.16) respectively,

$$\sigma^+\frac{\tilde{f}^+_0}{\tilde{f}^+_1} = -\sigma^-\frac{\tilde{f}^-_0}{\tilde{f}^-_1}$$

Rearrange the items and we obtain (11.18). □

LEMMA 11.7. The solution for $A^\pm$ are

$$A^+ = \gamma + \Psi(1-\gamma) \tag{11.19}$$

$$A^+/A^- = \Psi \tag{11.20}$$

△

PROOF. (11.20) is derived by rearranging the items in (11.15) and following the definition of Ψ.

(11.19) is derived from the fact that the total density of the adjusted distribution should be equal to 1, that is, $\int_{-\infty}^{\infty} f(x)dx = 1$. Hence,

$$\int_0^\infty f^+(x)dx + \int_0^\infty f^-(x)dx = \frac{1}{A^+}\int_0^\infty \tilde{f}^+(x)dx + \frac{1}{A^-}\int_0^\infty \tilde{f}^-(x)dx = 1$$

Apply (11.8), we get $\frac{\gamma}{A^+} + \frac{1-\gamma}{A^-} = 1$. Multiply it by A^+ on both sides, we obtain (11.19). □

We've shown that $A^\pm$ are well-defined constants based on (α, k, θ), while $\sigma^\pm$ is a choice of parametrization, constrained by (11.18).

11.4. Moments

The structure of the *moments* reveals critical information about the adjusted distribution. We show the moment formula of the canonical distribution, and how the location-scale family relates to it.

To simplify the notations below, let

- $f^{\pm} = L^{\pm\theta}_{\alpha,k}$ be the adjusted distribution family,
- $\widehat{f}^{\pm} = \widehat{L}^{\pm\theta}_{\alpha,k}$ be the canonical distribution,
- $\tilde{f}^{\pm} = \tilde{L}^{\pm\theta}_{\alpha,k}$ be the original (unadjusted) distribution.

First, the n-th one-sided moments of the adjusted distribution are ($x > 0$)

$$E(X^n|f^{\pm}) = \frac{1}{A^{\pm}\sigma^{\pm}} \int_0^{\infty} x^n \, \tilde{f}^{\pm}(x/\sigma^{\pm}))dx = \frac{(\sigma^{\pm})^n}{A^{\pm}} E(X^n|\tilde{f}^{\pm}) \tag{11.21}$$

where $E(X^n|\tilde{f}^{\pm})$ are the original n-th one-sided moments. They can be obtained from the Mellin transform (11.2).

The n-th total moment, given the notation of m_n, is the sum of $E(X^n|f^+)$ and $(-1)^n E(X^n|f^-)$. We show the following.

LEMMA 11.8. The n-th total moment of the adjusted distribution is based on the original one-sided moments such as

$$m_n := E(X^n|f) = \frac{(\sigma^+)^n}{A^+} \left[E(X^n|\tilde{f}^+) + \Psi(-\Sigma)^n \, E(X^n|\tilde{f}^-)\right] \tag{11.22}$$

△

PROOF. By definition, we have

$$\begin{aligned} m_n := E(X^n|f) &= \int_{-\infty}^{\infty} x^n \, f(x)dx = \int_0^{\infty} x^n \, f^+(x)dx + (-1)^n \int_0^{\infty} x^n \, f^-(x)dx \\ &= E(X^n|f^+) + (-1)^n E(X^n|f^-) \end{aligned}$$

Apply (11.21), we get

$$m_n = \frac{(\sigma^+)^n}{A^+} E(X^n|\tilde{f}^+) + \frac{(-\sigma^-)^n}{A^-} E(X^n|\tilde{f}^-)$$

Factor out $\frac{(\sigma^+)^n}{A^+}$, apply $\sigma^-/\sigma^+ = \Sigma$ from Lemma 11.6, and $A^+/A^- = \Psi$ from 11.7, we obtain (11.22). □

LEMMA 11.9 (The moments of the canonical distribution). The n-th moment of the canonical distribution is

$$\widehat{m}_n := E(X^n|\widehat{f}) = \frac{1}{A^+} \left[E(X^n|\tilde{f}^+) + \Psi(-\Sigma)^n \, E(X^n|\tilde{f}^-)\right] \tag{11.23}$$

△

PROOF. Lemma 11.8 shows that the canonical distribution $\widehat{f}$ is obtained by letting $\sigma^+ = 1$ and $\sigma^- = \Sigma$. Put them to (11.22), we obtain (11.23). □

Lastly, compare (11.22) with (11.23). We reach $m_n = (\sigma^+)^n \, \widehat{m}_n$. That is, all other members in the adjusted distribution family are rescaled canonical distributions.

CHAPTER 12

GAS-SN: Generalized Alpha-Stable Distribution with Skew-Normal

This fractional univariate distribution combines the features from a classic skew-normal distribution that provides skewness and a fractional distribution that provides fatter tails. The resulting distribution is analytically tractable. The PDF and all of its derivatives are continuous everywhere in $\mathbb{R}$.

12.1. Definition

DEFINITION 12.1. Assume $Z_0 \sim SN(0,1,\beta)$ is a skew-normal variable and $V \sim \overline{\chi}_{\alpha,k}$ is an FCM variable.

Then $Z \sim Z_0/V$ is a variable with a GAS-SN distribution. We use the notation $Z \sim L_{\alpha,k}(\beta)$ for this standard distribution.

Assume $\mathcal{N}(x)$ and $\Phi_{\mathcal{N}}(x)$ are the PDF and CDF of $N(0,1)$. The PDF of Z is

$$L_{\alpha,k}(x;\beta) = 2\int_0^\infty \mathcal{N}(xs)\,\Phi_{\mathcal{N}}(\beta xs)\,\overline{\chi}_{\alpha,k}(s)\,s\,ds. \tag{12.1}$$

This is the fractional extension of (10.2).

Its CDF is

$$\begin{aligned}\Phi[L_{\alpha,k}(\beta)](x) &:= \int_0^\infty \Phi_{SN}(xs;\beta)\,\overline{\chi}_{\alpha,k}(s)\,ds. \\ &= \int_0^\infty \left[\Phi_{\mathcal{N}}(xs) - 2\,T(xs,\beta)\right]\overline{\chi}_{\alpha,k}(s)\,ds.\end{aligned} \tag{12.2}$$

where $\Phi_{SN}(xs;\beta)$ is the CDF of $SN(0,1,\beta)$ in (10.3), and $T(h,a)$ is the Owen's T function.

We can clearly see that the CDF has two components: One from the symmetric part, and the other skew. The second component vanishes due to $T(h,0) = 0$.

12.1.1. GAS-SN Subsumes GSaS.

LEMMA 12.2. When $\beta = 0$, it becomes a symmetric distribution, previously called GSaS. The notation of $L_{\alpha,k}$ is given in [20]. "GSaS" stands for the *generalized symmetric α-stable* distribution.

A GSaS variable $X \sim L_{\alpha,k}$ is simply $X \sim \mathcal{N}(0,1)/V$. Its elegance leads to many important analytic properties that will be explored later in the chapter.

The PDF of a GSaS is

$$L_{\alpha,k}(x) = \int_0^\infty \mathcal{N}(xs)\,\overline{\chi}_{\alpha,k}(s)\,s\,ds. \tag{12.3}$$

When $\alpha \to 2$ or $k \to \infty$, the symmetric distribution approaches a normal distribution $N(0,\alpha^{2/\alpha})$ (Section 8.2 of [20]).

△

This integral is a normal mixture (9.1) that enjoys several nice properties outlined in Chapter 9.

In particular, the generalized exponential power distribution can be obtained via the characteristic function transform in Lemma 9.2 (Section 9 of [20]). We point out that the skew extension is straightforward, but leave the detailed description to future research.

12.1.2. GAS-SN Subsumes Skew-t Distribution. An important bridge between SN and GAS-SN is the skew-t (ST) distribution. It is documented in Section 4.3 of [1].

ST is fully consistent with GAS-SN by setting $\alpha = 1$. That is, in his notation, $T(\beta, k) = L_{1,k}(\beta)$.

12.2. The Location-Scale Family

Its location scale family is $Y = \xi + \omega Z \sim L_{\alpha,k}(\xi, \omega^2, \beta)$. Its PDF becomes

$$\phi(x) = \frac{1}{\omega} L_{\alpha,k}\left(\frac{x-\xi}{\omega}; \beta\right). \quad (x \in \mathbb{R}) \tag{12.4}$$

In real-world applications, this PDF is used for optimization, e.g. in the maximum likelihood estimation (MLE). See Section 12.10.

12.3. Mellin Transform

The Mellin transform of the PDF follows the rule of the ratio distribution. From (10.8) and (7.2), we have

$$\begin{aligned} L_{\alpha,k}(x;\beta) &\overset{\mathcal{M}}{\longleftrightarrow} L^*_{\alpha,k}(s;\beta) \\ &= \mathcal{N}^*(s;\beta)\, \overline{\chi}^*_{\alpha,k}(2-s) \qquad (12.5) \\ &= \big[2\,\Phi[t_s](\beta\sqrt{s})\big] \times \big[\mathcal{N}^*(s)\, \overline{\chi}^*_{\alpha,k}(2-s)\big] \qquad (12.6) \end{aligned}$$

Notice that the contribution for the skewness is $2\,\Phi[t_s](\beta\sqrt{s})$ in the first bracket, which becomes one if $\beta = 0$.

The second bracket is the Mellin transform of the GSaS PDF. From (2.9) and (7.2), it is

$$\begin{aligned} L_{\alpha,k}(x) &\overset{\mathcal{M}}{\longleftrightarrow} L^*_{\alpha,k}(s) = \mathcal{N}^*(s)\, \overline{\chi}^*_{\alpha,k}(2-s) \qquad (12.7) \\ &= \frac{1}{2\sqrt{\pi}} \left(\frac{2}{\sigma}\right)^{s-1} \Gamma\left(\frac{s}{2}\right) \frac{\Gamma((k-1)/2)}{\Gamma((k-1)/\alpha)} \frac{\Gamma((k-s)/\alpha)}{\Gamma((k-s)/2)}, \end{aligned}$$

where $\sigma := k^{1/2-1/\alpha}$ and $k > 0$ is assumed.

12.4. Moments

Based on $\mathbb{E}(X^n|\mathcal{N}(\beta))$ from (10.11), the n-th moment of Z is

$$\begin{aligned} \mathbb{E}(X^n|L_{\alpha,k}(\beta)) &:= \mathbb{E}(X^n|\mathcal{N}(\beta))\, \mathbb{E}(X^{-n}|\overline{\chi}_{\alpha,k}) \qquad (12.8) \\ &= 2\,\mathcal{N}^*(n+1)\, \mathbb{E}(X^{-n}|\overline{\chi}_{\alpha,k}) \\ &\quad \times \begin{cases} 1, & \text{when } n \text{ is even,} \\ 1 - 2\Phi[t_{n+1}](-\beta\sqrt{n+1}), & \text{when } n \text{ is odd.} \end{cases} \end{aligned}$$

Its first moment is $\mu_z = b\,\delta$, where $b = \sqrt{2/\pi}\, \mathbb{E}(X^{-1}|\overline{\chi}_{\alpha,k})$.

The second moment is $\mathbb{E}(X^{-2}|\overline{\chi}_{\alpha,k})$. Its variance is

$$\sigma_z^2 = \mathbb{E}(X^{-2}|\overline{\chi}_{\alpha,k}) - (b\,\delta)^2. \tag{12.9}$$

To simplify the symbology, let $q_n := \mathbb{E}(X^{-n}|\overline{\chi}_{\alpha,k})$. The third moment is $\delta_3\, q_3$, where $\delta_3 = \sqrt{\frac{2}{\pi}}\delta(3-\delta^2)$. The fourth moment is $3\, q_4$. To carry out the skewness γ_1 and excess kurtosis γ_2,

$$\gamma_1 \times \sigma_z^{3/2} = \delta_3\, q_3 - 3\mu_z q_2 + 2\mu_z^3,$$
$$\gamma_2 \times \sigma_z^4 = 3(q_4 - q_2^2) - 4\mu_z(\gamma_1 \times \sigma_z^{3/2}) + 2\mu_z^4.$$

The maximum skewness and kurtosis can be infinite. Since $\delta = \sin\theta$, where $\beta = \tan\theta$, we have $\delta \in [-1, 1]$. Infinity has to come from q_3 and q_4.

A typical example is the skew-t distribution at $\alpha = 1$. It is well known that kurtosis approaches infinity when k approaches 4 from above, and the skewness approaches infinity when k approaches 3 from above.

12.4.1. Excess Kurtosis of GSaS. It is important to understand the behavior of excess kurtosis γ_2. However, the presence of skewness adds more complexity to γ_2. Consider the symmetric case where $\beta = 0$, and we quote the result from [20] below.

The excess kurtosis of GSaS is plotted in Figure 12.1 in the (k, α) coordinate. Notice that a major division occurs along the line of $k = 5 - \alpha$. In the region where $0 < k \le 5 - \alpha$, there are complicated patterns caused by the infinities of the gamma function. Only small pockets of valid kurtosis exist.

LEMMA 12.3. In the region where $k > 5 - \alpha$, the excess kurtosis of GSaS is a continuous function with positive values. At large k's, the closed form of the moments can be expanded by Sterling's formula. The excess kurtosis γ_2 becomes part of a linear equation:

$$\left(\epsilon - \frac{1}{2}\right) = \left(\frac{k-3}{4}\right) \log\left(1 + \frac{\gamma_2}{3}\right), \quad \text{where } \epsilon = 1/\alpha \tag{12.10}$$

This equation shows how GSaS works under the **Central Limit Theorem**. GSaS approaches a normal distribution when γ_2 becomes zero. This can happen from two directions: when $\alpha \to 2$ or when $k \to \infty$.

△

The contour plot of excess kurtosis is shown in the (k, ϵ) coordinate in Figure 12.2. It is visually amusing. Notice the singular point at $\epsilon = 1/2, k = 3$.

12.4.2. Standardized Peak Density of GSaS. For the symmetric case $\beta = 0$, the GAS-SN density reduces to the GSaS density in (12.3). Its mode is located at the center. Under the location-scale transformation

$$Y = \xi + \omega Z, \qquad \phi_Y(x) = \frac{1}{\omega}\, L_{\alpha,k}\left(\frac{x-\xi}{\omega}\right),$$

the peak height changes by the factor $1/\omega$. Hence the location-scale invariant peak height is obtained by multiplying the peak density by the standard deviation.

DEFINITION 12.4. Assume $Z \sim L_{\alpha,k}$ and that $\sigma_Z^2 < \infty$. The standardized peak density of GSaS is defined as

$$\mathrm{SPD}_{\alpha,k} := \sigma_Z\, L_{\alpha,k}(0) = \frac{1}{\sqrt{2\pi}} \sqrt{\mathbb{E}(X^{-2}|\overline{\chi}_{\alpha,k})}\, \mathbb{E}(X|\overline{\chi}_{\alpha,k}). \tag{12.11}$$

Equivalently, for any location-scale version $Y = \xi + \omega Z$, it is

$$\mathrm{SPD}_{\alpha,k} = \sigma_Y\, \phi_Y(\xi), \tag{12.12}$$

so the quantity is independent of ξ and ω.




FIGURE 12.1. The contour plot of excess kurtosis in GSaS by (k, α).

PROOF. From (12.3), since $\mathcal{N}(0) = 1/\sqrt{2\pi}$,

$$L_{\alpha,k}(0) = \frac{1}{\sqrt{2\pi}}\,\mathbb{E}(X|\overline{\chi}_{\alpha,k}).$$

In the symmetric case $\beta = 0$, (12.9) gives

$$\sigma_Z^2 = \mathbb{E}(X^{-2}|\overline{\chi}_{\alpha,k}).$$

Therefore,

$$\mathrm{SPD}_{\alpha,k} = \frac{1}{\sqrt{2\pi}}\sqrt{\mathbb{E}(X^{-2}|\overline{\chi}_{\alpha,k})}\,\mathbb{E}(X|\overline{\chi}_{\alpha,k}).$$

□

Using (7.8), the same expression can be written as

$$\mathrm{SPD}_{\alpha,k} = \frac{1}{\sqrt{2\pi}}\left[\frac{\Gamma((k-1)/2)}{\Gamma((k-1)/\alpha)}\right]^{3/2}\frac{\Gamma(k/\alpha)}{\Gamma(k/2)}\left[\frac{\Gamma((k-3)/\alpha)}{\Gamma((k-3)/2)}\right]^{1/2}. \tag{12.13}$$

As a probabilistic standardized peak density, this formula requires the underlying gamma functions to render a positive number. The analytic continuation of (12.13) has a more complicated finite-real region in the (k, α) plane, as shown below.

The contour plot of standardized peak density is shown in the (k, ϵ) coordinate in Figure 12.3. This view highlights the approximate linearity in (12.14) for large k's.




FIGURE 12.2. The contour plot of excess kurtosis in GSaS by (k, ϵ) where $\epsilon = 1/\alpha$. This best describes the linearity in (12.10) for large k's.

LEMMA 12.5. At large k's, the closed form (12.13) can be expanded by Stirling's formula. The standardized peak density becomes part of a linear equation:

$$\left(\epsilon - \frac{1}{2}\right) = \frac{2}{3}\left(k - \frac{4}{3}\right)\log\left(\sqrt{2\pi}\,\mathrm{SPD}_{\alpha,k}\right), \qquad \text{where } \epsilon = 1/\alpha. \tag{12.14}$$

This equation shows how the standardized peak density approaches the normal value under the **Central Limit Theorem**. The normal reference is $\mathrm{SPD}_{2,\infty} = 1/\sqrt{2\pi} \approx 0.398942$, for which the logarithmic term vanishes.

△

12.4.3. Lognormal-Normal Asymptotics of GSaS. Combining (12.10) and (12.14) under the assumption $\epsilon \neq 1/2$ eliminates the common factor $\epsilon - 1/2$. This gives

$$1 + \frac{\gamma_2}{3} = \left(\sqrt{2\pi}\,\mathrm{SPD}_{\alpha,k}\right)^H, \qquad \text{where } H = \frac{8}{3}\left(\frac{1 - 4/(3k)}{1 - 3/k}\right). \tag{12.15}$$

As $k \to \infty$, $H \to 8/3$. The same exponent appears in the lognormal-normal limit in (12.16) below. This limit has profound implications for GSaS.

DEFINITION 12.6 (The Lognormal-Normal Distribution (LNN)). Let $Z \sim \mathcal{N}(0,1)$ and $Y \sim \mathcal{N}(0,\tau^2)$ be independent. The random variable $X = Ze^Y$ belongs to the lognormal-normal distribution family, denoted by $\mathrm{LNN}(\tau^2)$, where τ is the shape parameter[5].[1]

[1]The author studied LNN in a 2008 working paper. See https://ssrn.com/abstract=1537686. However, the role of LNN was rediscovered in 2026, when I attempted to fit the bear state of the global regime in the financial market. See https://github.com/slihn/Mixture-VAE/blob/main/slides/frac_slides.md.




FIGURE 12.3. The contour plot of standardized peak density in GSaS by (k, ϵ), where $\epsilon = 1/\alpha$.

LEMMA 12.7 (The LNN properties). The LNN has exact even moments

$$\mathbb{E}(X^{2m}) = (2m-1)!!\, e^{2m^2\tau^2}.$$

Therefore, the variance is $e^{2\tau^2}$ and

$$1 + \frac{\gamma_2}{3} = e^{4\tau^2},$$
$$\sqrt{2\pi}\,\mathrm{SPD}_{\mathrm{LNN}} = e^{3\tau^2/2}.$$

Eliminating τ^2 gives

$$1 + \frac{\gamma_2}{3} = \left(\sqrt{2\pi}\,\mathrm{SPD}_{\mathrm{LNN}}\right)^{8/3}. \tag{12.16}$$

△

LEMMA 12.8 (The LNN limit of GSaS). Let $X = Z/V$ be a GSaS variable, where $Z \sim \mathcal{N}(0,1)$ and $V \sim \overline{\chi}_{\alpha,k}$ are independent. Set $\zeta = \mathrm{sign}(k)$, $\delta = 1/\alpha - \frac{1}{2}$, and

$$z = \begin{cases} k-1, & k > 0, \\ |k|, & k < 0. \end{cases}$$

In either fixed-sign branch, let $z \to \infty$ with $\delta/z \to \tau_{LN}^2 \in (0,\infty)$; or equivalently, $\alpha|k| \to 1/\tau_{LN}^2$. With $D_{\alpha,k}$ and μ_{LN}^{ζ} as in Lemma 7.6,

$$\frac{e^{\mu_{LN}^{\zeta}}}{D_{\alpha,k}^{\zeta}} X \xrightarrow{d} Z e^{Y}, \qquad Y \sim \mathcal{N}(0, \tau_{LN}^2), \tag{12.17}$$

where Z and Y are based on Definition 12.6. Thus, both branches converge to the same $\mathrm{LNN}(\tau_{LN}^2)$ distribution after a deterministic adjustment of the scale. △

Proof. By Lemma 7.6, $\hat{V} = D_{\alpha,k}^{\zeta} V$ satisfies $e^{-\mu_{LN}^{\zeta}} \hat{V} \xrightarrow{d} e^{Y}$. Independence and the continuous mapping theorem give

$$\frac{e^{\mu_{LN}^{\zeta}}}{D_{\alpha,k}^{\zeta}} X = \frac{Z}{e^{-\mu_{LN}^{\zeta}} \hat{V}} \xrightarrow{d} Z e^{-Y} \overset{d}{=} Z e^{Y},$$

since $-Y \overset{d}{=} Y$ for centered normal Y. □

12.5. GSaS Version of the Central Limit Theorem

This section explores the implications of Lemma 12.8. We propose a pair of canonical coordinates and a version of the Central Limit Theorem (CLT) tailored to GSaS in Lemma 12.11. This approach leads to a fitting method that determines the GSaS parameters for a given time-series data set. We tested this method on several stock return data sets. The resulting GSaS distribution provides a very good initial estimate that can be refined using a more sophisticated MLE algorithm.

12.5.1. Canonical Coordinates and CLT. For $0 < \alpha < 2$, the excess kurtosis γ_2 and standardized peak density (SPD) are both finite on the joint domain $k > 5 - \alpha$ on the $k > 0$ branch and $|k| > \max(0, 1-\alpha)$ on the $k < 0$ branch. On this domain, the mapping from (α, k) to (γ_2, SPD) is one-to-one. The normal boundary $\alpha = 2$ is excluded from this uniqueness statement: this is the point of standard normal. All values of k give the same kurtosis and SPD there.

Definition 12.9 (GSaS Canonical Coordinates). The canonical coordinates of GSaS are defined by

$$v_1 := \frac{1}{4} \log\left(1 + \frac{\gamma_2}{3}\right), \tag{12.18}$$

$$v_2 := \frac{2}{3} \log\left(\sqrt{2\pi}\, \mathrm{SPD}_{\alpha,k}\right). \tag{12.19}$$

The constants $\frac{1}{4}, \frac{2}{3}$ are chosen so that $v_1 = v_2 = \tau_{\mathrm{LN}}^2$ on the terminal LNN line, which is exactly diagonal; see Figure 12.4.

Here v stands for variance. At a finite GSaS point, v_1 and v_2 are implied LNN log-variances inferred from kurtosis and SPD, respectively, not generally the actual variance of $\log V$. On the terminal line, both coincide at τ_{LN}^2.

Recall the FCM moment function $H(s) = \mathbb{E}[V^s]$ and its logarithm in (7.20). Conditioning on V gives

$$\mathbb{E}[X^2] = H(-2), \qquad \mathbb{E}[X^4] = 3H(-4), \qquad f_X(0) = \frac{H(1)}{\sqrt{2\pi}}.$$

Consequently, the definitions of excess kurtosis and standardized peak density imply

$$1 + \frac{\gamma_2}{3} = \frac{H(-4)}{H(-2)^2}, \qquad \sqrt{2\pi}\, \mathrm{SPD}_{\alpha,k} = H(1) H(-2)^{1/2}.$$

Taking logarithms and applying the normalization constants in v_1, v_2 gives

$$\begin{aligned} v_1 &= \frac{1}{4}\left[\log H(-4) - 2\log H(-2)\right], \\ v_2 &= \frac{2}{3}\left[\log H(1) + \frac{1}{2}\log H(-2)\right]. \end{aligned} \tag{12.20}$$

The canonical coordinates use $H(1)$, $H(-2)$, and $H(-4)$, but do not match $H(-3)$. The latter determines the standardized absolute third moment of the centered mixture:

$$\frac{\mathbb{E}[|X|^3]}{\mathbb{E}[X^2]^{3/2}} = \frac{2\sqrt{2}}{\sqrt{\pi}}\,\frac{H(-3)}{H(-2)^{3/2}}.$$

Unlike the signed third moment, which vanishes by symmetry, this quantity is not fixed by symmetry. After fitting (α, k), comparing this model prediction with the standardized absolute third central moment of the data provides an additional goodness-of-fit check, since that statistic was not used to determine the canonical coordinates.

Figure 12.4 shows the GSaS implied log-variance plane (v_1, v_2). The thick black diagonal is the terminal LNN line. The positive-k branch lies below it and represents the finite-kurtosis region of the GSaS extensions of the α-stable and Student's t families.

The negative-k branch lies above the diagonal and includes generalizations of the exponential power distribution.

On the joint finite domain stated above, the mapping from (α, k) to (v_1, v_2) is one-to-one. A point in its image therefore determines a unique parameter pair (α, k) and the corresponding GSaS shape $L_{\alpha,k}$.

Data Perspective. This coordinate system represents the admissible GSaS shapes throughout the joint finite domain. For an approximately symmetric return series, we estimate excess kurtosis and standardized peak density and locate the resulting point in the (v_1, v_2) plane. If the point lies in the image of the finite-parameter GSaS family, inversion determines a unique (α, k). Estimating location and scale then completes an initial fit to the data.

The curves of constant k approach the normal reference point at the origin as $\delta \to 0$. Their approximately ray-like appearance, especially for large $|k|$, motivates the polar-type coordinates (R, θ).

Definition 12.10 (GSaS Polar-type Coordinates). Define

$$R := \tfrac{1}{2}(v_1 + v_2), \tag{12.21}$$

$$\theta := \operatorname{atan2}(v_2, v_1). \tag{12.22}$$

We also define $D := \frac{1}{2}(v_2 - v_1)$ as a signed diagonal-offset coordinate. The perpendicular distance to the terminal line is $\sqrt{2}|D|$. The angle θ is defined away from the normal origin.

Here R is a diagonal coordinate, not the Euclidean radius: (R, R) is the orthogonal projection of (v_1, v_2) onto the terminal line. On that line, $R = \tau_{\mathrm{LN}}^2$. Away from the origin,

$$\frac{D}{R} = \frac{v_2 - v_1}{v_1 + v_2} = \frac{\tan\theta - 1}{\tan\theta + 1} = \tan\Big(\theta - \frac{\pi}{4}\Big).$$

Thus θ measures direction, whereas D measures an absolute signed offset. Multiplying both coordinates by the same positive factor leaves θ unchanged but multiplies D by that factor. At fixed R, either D or θ determines the other; D is not a third independent coordinate. Both indicate the side of the terminal line: $D > 0$ and $\theta > \pi/4$ above it, where $k < 0$; and $D < 0$ and $\theta < \pi/4$ below it, where $k > 0$.

The following shape-limit result summarizes the two limits represented in Figure 12.4.

LEMMA 12.11 (GSaS Central Limit Theorem). Let $X = Z/V$ be a GSaS variable in the joint finite domain stated above.

- **LNN limit.** In either fixed-sign branch, let $z \to \infty$ with $\delta/z \to \tau_{LN}^2 \in (0, \infty)$. After the deterministic scale adjustment in (12.17), X converges in distribution to $\mathrm{LNN}(\tau_{LN}^2)$, and $(R, \theta) \to (\tau_{LN}^2, \pi/4)$.
- **Normal limit.** Along any parameter sequence for which $R \to 0$, the variance-normalized GSaS variable satisfies

$$\frac{X}{\sqrt{\mathbb{E}[X^2]}} \xrightarrow{d} \mathcal{N}(0, 1).$$

△

PROOF. The distributional LNN convergence follows from Lemma 12.8. For the coordinate limit, use the centered log-CGF in (7.30). The log-MGF expansion in the proof of the FCM Lemma 7.6 gives $C(s) \to s^2\tau_{LN}^2/2$ for every fixed real s. The mixture moment and SPD identities therefore yield

$$v_1 = \tfrac{1}{4}\left[C(-4) - 2C(-2)\right] \to \tau_{LN}^2,$$
$$v_2 = \tfrac{2}{3}\left[C(1) + \tfrac{1}{2}C(-2)\right] \to \tau_{LN}^2.$$

The stated limits of R and θ follow. For the normal limit, nonnegativity of the coordinates implies $0 \le v_1 \le 2R \to 0$. Independence of Z and V gives

$$\mathbb{E}\left[\left(\frac{V^{-2}}{\mathbb{E}[V^{-2}]} - 1\right)^2\right] = \frac{\mathbb{E}[V^{-4}]}{\mathbb{E}[V^{-2}]^2} - 1 = \frac{\gamma_2}{3} = e^{4v_1} - 1 \to 0.$$

Thus $V^{-1}/\sqrt{\mathbb{E}[V^{-2}]} \to 1$ in probability. Since $\mathbb{E}[X^2] = \mathbb{E}[V^{-2}]$, the variance-normalized mixture is $ZV^{-1}/\sqrt{\mathbb{E}[V^{-2}]}$; Slutsky's theorem gives the normal limit. □

The first limit approaches the diagonal at a positive limiting R; the second approaches the origin. Increasing $|k|$ alone does not ensure a nondegenerate LNN limit: at fixed α, it instead gives the normal limit after variance normalization. The familiar convergence of Student's t distributions as their degrees of freedom increase is one example. The equality $R = \tau_{LN}^2$ applies on the terminal LNN line, not at a general finite-parameter point.

12.5.2. Analysis of the Polar-type Coordinates. In Figure 12.4, the constant-k curves are approximately ray-like, while the diagonal represents the LNN limit. The following large-$|k|$ analysis explains how (R, θ) approximately separates the shape parameters: the angle primarily identifies k, and R then determines α along the corresponding curve.

LEMMA 12.12 (Large $|k|$ expressions). For large $|k|$, the canonical coordinates satisfy

$$v_1 \approx \frac{\delta}{z - 2\zeta}, \qquad v_2 \approx \frac{\delta}{z - \zeta/3} \tag{12.23}$$

The shifted denominators account for the leading difference between the two coordinates at finite k. For $k > 0$, they are $k - 3$ and $k - \frac{4}{3}$; for $k < 0$, they are $|k| + 2$ and $|k| + \frac{1}{3}$. Both coordinates have the same numerator, so their ratio is approximately independent of α. This separates the angular dependence on k from the magnitude controlled by α, motivating the polar description below. △

PROOF. In (12.20), substitute $\log H(s) = \zeta s \log \sigma + K(s)$ from (7.20). The linear scale terms cancel because $-4 - 2(-2) = 0$ and $1 + \frac{1}{2}(-2) = 0$. Using the FCM definition $K(s) = A(z + \zeta s) - A(z)$




FIGURE 12.4. The GSaS implied log-variance plane. The thick black diagonal is the terminal LNN line; positive-k curves lie below it and negative-k curves above it. As $|k|$ increases while R tends to a positive value, the curves approach the LNN line. Lines of constant R are perpendicular to the diagonal and approach the normal origin as R decreases to zero. On the diagonal only, $R = \tau^2_{\mathrm{LN}}$. Data points represent SPX global regimes and selected financial assets, discussed in the main text.

then gives the exact finite differences

$$v_1 = \frac{1}{4}\left[A(z-4\zeta) - 2A(z-2\zeta) + A(z)\right],$$
$$v_2 = \frac{2}{3}\left[A(z+\zeta) + \frac{1}{2}A(z-2\zeta) - \frac{3}{2}A(z)\right].$$

Taylor-expand the first expression about $z - 2\zeta$ and the second about $z - \zeta/3$. In each case, the constant, linear, and cubic terms cancel, and the coefficient of the second derivative is 1, giving

$$v_1 \approx A''(z - 2\zeta), \qquad v_2 \approx A''(z - \zeta/3).$$

Applying $A''(x) \approx \delta/x$ from (7.24) gives the stated approximations on both branches. □

LEMMA 12.13 (The separate roles of the polar coordinates). For large $|k|$, substituting (12.23) into the definitions of the polar-type coordinates gives

$$R = \tfrac{1}{2}(v_1 + v_2) \approx \frac{\delta}{2}\left(\frac{1}{z - 2\zeta} + \frac{1}{z - \zeta/3}\right) \approx \frac{\delta}{z - 7\zeta/6}. \tag{12.24}$$

Similarly,

$$\tan\theta \approx \frac{z - 2\zeta}{z - \zeta/3} = \begin{cases} \dfrac{k - 3}{k - 4/3}, & k > 0, \\ \dfrac{|k| + 2}{|k| + 1/3}, & k < 0. \end{cases} \tag{12.25}$$

The coordinates (R, θ) approximately separate the parameters: θ primarily identifies k, after which R determines α.

△

Diagonal coordinate. R retains the dependence on α through δ. To leading order, $R \approx \delta/z$, so it increases as α decreases along a constant-k curve. On the joint terminal path $z \to \infty$ with $\delta/z \to \tau_{\mathrm{LN}}^2 > 0$, we have $R \to \tau_{\mathrm{LN}}^2$.

Angle. The approximation for $\tan\theta$ depends on k but not on α; to this order, each constant-k curve is a ray. The exact angle can still vary with α at finite k. At positive limiting R, $\theta \to \pi/4$ describes the terminal LNN limit. The positive-k branch has $\theta < \pi/4$, while the negative-k branch has $\theta > \pi/4$.

Near the diagonal, the angular departure is proportional to $1/|k|$ to leading order, since

$$\tan\theta - 1 = \frac{2D}{R - D} \approx \frac{2D}{R} \approx -\frac{5\zeta}{3z}.$$

12.5.3. Application to Real-World Data. Figure 12.4 shows seven empirical points calculated from daily financial returns. Three come from the SPX global regime analysis[**29**]. The point labeled "SPX" gives the empirical (v_1, v_2) coordinates for the full SPX return sample from February 1, 1991, through September 4, 2026. The regime analysis separates these returns into bull and bear states; "SPX bull" and "SPX bear" give the coordinates estimated from the corresponding subsets.

The remaining four points represent GLD, TLT, BTC, and DXY, providing comparisons across asset classes beyond equities. GLD and TLT are US-listed ETFs providing exposure to gold and Treasury bonds, respectively. The return samples begin on November 19, 2004, for GLD and July 31, 2002, for TLT. BTC represents Bitcoin returns beginning on January 4, 2017, and DXY represents Dollar Index returns beginning on February 4, 1991. All four samples end on September 4, 2026. BTC and DXY are sampled on NYSE trading days to align their observations with the SPX calendar.

The pink ellipses summarize uncertainty in the four asset-coordinate estimates. We use 1,000 moving-block bootstrap resamples with circular blocks of 20 consecutive observations to retain short-range dependence and volatility clustering. Recomputing (v_1, v_2) on each resample gives a covariance matrix, from which we construct a nominal joint 50% uncertainty ellipse. The BTC sample begins in 2017 to focus on the period when we regard Bitcoin as an established financial asset. Its relatively large ellipse reflects the sampling variability of the estimates over this period, not the start-date choice alone.

12.5.4. Method of Moments and Density. The canonical-coordinate map gives a practical fitting procedure for an approximately symmetric return sample $x_1, \dots, x_N$ with nonzero variance and finite fourth moment. We match two shape features: excess kurtosis and standardized peak density. Since the latter is estimated from a density ordinate rather than an ordinary sample moment, this is a mixed moment–density matching method, not a likelihood fit.

Write

$$\bar{x} = \frac{1}{N}\sum_{i=1}^{N} x_i, \quad \widehat{m}_j = \frac{1}{N}\sum_{i=1}^{N}(x_i - \bar{x})^j, \tag{12.26}$$
$$\widehat{\sigma} = \sqrt{\widehat{m}_2}, \qquad \widehat{\gamma}_2 = \frac{\widehat{m}_4}{\widehat{m}_2^2} - 3.$$

These are the empirical-moment conventions used for the figures below; no finite-sample kurtosis correction is applied.

Bin-free peak density. Let $Q(p)$ be the population quantile function and $m = Q(1/2)$ its median. For a continuous density f that is positive at m, $Q'(1/2) = 1/f(m)$. Thus a symmetric quantile spacing gives the estimator

$$\widehat{f}_h := \frac{2h}{\widehat{Q}(1/2+h) - \widehat{Q}(1/2-h)}, \qquad 0 < h < \frac{1}{2}, \tag{12.27}$$

where $\widehat{Q}$ is the empirical quantile function, evaluated by linear interpolation between ordered observations. The numerator is the probability mass between the two quantiles, and the denominator is the corresponding return interval. Their ratio estimates the average density over that interval. For the symmetric, unimodal GSaS family, the median is also the mode, so $f(m)$ is the peak density. Without this shape assumption, the same procedure estimates central density, not necessarily the maximum of the density.

To reduce the smoothing error, we use several probability widths rather than a single h. If Q is sufficiently smooth near $1/2$, its population spacing satisfies

$$\frac{2h}{Q(1/2+h) - Q(1/2-h)} = f(m) - \frac{Q'''(1/2)}{6[Q'(1/2)]^2}h^2 + O(h^4). \tag{12.28}$$

The leading population smoothing error is therefore quadratic in h. Following the notebook implementation, we take $\mathcal{H} = \{0.04, 0.06, 0.08, 0.10, 0.12, 0.15, 0.20\}$ and define

$$(\widehat{f}_0, \widehat{c}) := \arg\min_{a,c} \sum_{h\in\mathcal{H}} \left(\widehat{f}_h - a - ch^2\right)^2, \qquad \widehat{\mathrm{SPD}} := \widehat{\sigma}\,\widehat{f}_0. \tag{12.29}$$

The fitted intercept $\widehat{f}_0$ extrapolates the central density to $h = 0$ and supplies the red peak markers at the sample medians in Figures 12.5 and 12.6. This procedure is independent of histogram bin edges, but it is not free of smoothing choices: the set $\mathcal{H}$ controls the intervals used. The extrapolation reduces the leading population smoothing error; it does not eliminate sampling error or guarantee an unbiased estimate. It also requires nonzero quantile spacings and a positive fitted intercept.

Shape, location, and scale. The empirical coordinates are

$$\widehat{v}_1 = \frac{1}{4}\log\left(1 + \frac{\widehat{\gamma}_2}{3}\right), \qquad \widehat{v}_2 = \frac{2}{3}\log\left(\sqrt{2\pi}\,\widehat{\mathrm{SPD}}\right). \tag{12.30}$$

If this point lies in the admissible finite-parameter GSaS image, we solve $v_1(\widehat{\alpha}, \widehat{k}) = \widehat{v}_1$ and $v_2(\widehat{\alpha}, \widehat{k}) = \widehat{v}_2$ for the unique shape pair. Figure 12.4 provides a graphical starting value for this numerical inversion. Points near the terminal diagonal may require very small α and large $|k|$, as in the SPX bear-state example; an empirical point outside the admissible image has no exact match in this family.

Let Y have the centered, unit-scale GSaS distribution with shape $(\widehat{\alpha}, \widehat{k})$ and density f_Y. Matching the sample mean and standard deviation then gives

$$\widehat{\mu} = \bar{x}, \qquad \widehat{a} = \frac{\widehat{\sigma}}{\sqrt{\mathbb{E}[Y^2]}}, \qquad \widehat{f}_X(x) = \frac{1}{\widehat{a}} f_Y\left(\frac{x - \widehat{\mu}}{\widehat{a}}\right). \tag{12.31}$$

Under the symmetric model, this also matches the estimated central peak height because $\widehat{a}^{-1} f_Y(0) = \widehat{f}_0$; the sample mean and median need not coincide exactly. The resulting density fit uses no histogram bins. Histograms serve only as visual comparisons, and the fit can subsequently be refined by maximum likelihood estimation.




FIGURE 12.5. The theoretical GSaS density vs the histogram of daily returns for GLD, TLT, BTC, DXY. The theoretical α, k is taken from Figure 12.4. The red dots are the bin-free peaks at the median. GLD and TLT are the ETFs traded in the US stock market. BTC represents the bitcoin prices since 1/4/2017. DXY is the dollar index closing prices since 2/4/1991. All four time series end on September 4, 2026. BTC and DXY are trimmed to NYSE trading days.




FIGURE 12.6. GSaS density fits to daily simple returns for the full SPX sample (upper left), the bull state (upper right), and the bear state (lower left). The full sample runs from February 1, 1991 to September 4, 2026; the state samples contain the observations assigned to each regime. The shape parameters (α, k) are obtained by matching empirical excess kurtosis and standardized peak density in the canonical-coordinate plane of Figure 12.4. Red dots mark bin-free density estimates at the sample medians. The full sample combines a sharp central peak with high kurtosis, while the bull-state fit lies on the negative-k branch. The bear-state fit has small α and large positive k, consistent with its coordinates lying close to the terminal LNN line. The lower-right panel compares the bull- and bear-state histograms and fitted densities on a logarithmic density scale, highlighting their tail behavior.




FIGURE 12.7. The theoretical LNN density vs the histogram of daily returns for the SPX bear state. Three panels show the density, log density, and Q-Q plot. This fit comes from a single shape parameter: $\tau^2_{LN} = 0.2574$.

12.6. Tail Behavior

The tail behavior of GAS-SN is a "modified GSaS" type. Hence, it is well within what was known. Without losing generality, assume $\beta > 0$, that the decay of the left tail is more pronounced than that of the right tail. But it still follows the same power law of x^{-k} as in a $L_{\alpha,k}$.

It takes a small tweak to GSaS to capture that behavior.

DEFINITION 12.14. The shifted GSaS is defined as

$$L_{\alpha,k}(x||\mu) = \int_0^\infty \mathcal{N}(xs-\mu)\,\overline{\chi}_{\alpha,k}(s)\,s\,ds \tag{12.32}$$

Note that the shift μ is not a location parameter that shifts x. It is a shift inside the argument of $\mathcal{N}(\,)$. When $\mu = 0$, it is restored to the PDF of GSaS, $L_{\alpha,k}(x)$.

We use the following approximation of the erf function in (12.1)[**16**]

$$1 - \mathrm{erf}(x) \approx \frac{1}{B\sqrt{\pi}\,x}(1 - e^{-Ax})\,e^{-x^2} \quad (x \geq 0) \tag{12.33}$$

where $A = 1.98$ and $B = 1.135$. It is much better than the first-order expansion of $e^{-x^2}/(\sqrt{\pi}x)$ for the entire range of $x \in [0, \infty)$.

LEMMA 12.15. For $\beta > 0$, the left tail ($x < 0$) of the PDF in (12.1) can be approximated by

$$\widehat{L}_{\alpha,k}(x;\beta) = \frac{G}{\beta x}\left[e^{\mu^2/2}\,L_{\alpha,k-1}\,(qx||\mu) - L_{\alpha,k-1}\,(qx)\right] \tag{12.34}$$

where

$$\mu = \frac{A\,\delta}{\sqrt{2}}, \quad \delta = \frac{\beta}{\sqrt{1+\beta^2}},$$
$$q = \sqrt{1+\beta^2}\,\frac{\sigma_{\alpha,k}}{\sigma_{\alpha,k-1}}$$
$$G = \sqrt{\frac{2}{\pi}}\frac{C_{\alpha,k}}{B\,\sigma_{\alpha,k-1}\,C_{\alpha,k-1}}$$

and both $C_{\alpha,k} = \frac{\alpha\Gamma((k-1)/2)}{\Gamma((k-1)/\alpha)}$ and $\sigma_{\alpha,k}$ are according to FCM in (7.4). The factor $1/\beta$ is part of this tail approximation and should not be interpreted as a singularity of the exact PDF. When $\beta = 0$, (12.1) reduces directly to the symmetric density $L_{\alpha,k}(x;\beta) = L_{\alpha,k}(x)$.

The right tail ($x > 0$) is simply

$$2L_{\alpha,k}(x) - \widehat{L}_{\alpha,k}(-x;\beta) \tag{12.35}$$

where the second term $\widehat{L}_{\alpha,k}(-x;\beta)$ becomes much smaller than the first term as $x \to \infty$. △

PROOF. Assume $\beta > 0$ and $x < 0$. Since $\Phi_{\mathcal{N}}(z) = \frac{1}{2}[1 + \mathrm{erf}(z/\sqrt{2})]$, the skew-normal factor in (12.1) satisfies

$$2\Phi_{\mathcal{N}}(\beta x s) = 1 - \mathrm{erf}\left(-\frac{\beta x s}{\sqrt{2}}\right).$$

Set $u = -\beta x s/\sqrt{2}$, so that $u > 0$. Applying the approximation in the preceding display gives the factor $1/u = -\sqrt{2}/(\beta x s)$; the factor s in the integral then cancels, leaving $1/(\beta x)$. Thus,

$$\begin{aligned} L_{\alpha,k}(x;\beta) &\approx \int_0^\infty \mathcal{N}(xs)\,\frac{1}{B\sqrt{\pi}\,u}\left(1 - e^{-Au}\right)e^{-u^2}\overline{\chi}_{\alpha,k}(s)\,s\,ds \\ &= -\frac{1}{B\pi\beta x}\int_0^\infty e^{-(1+\beta^2)x^2s^2/2}\left(1 - e^{A\beta x s/\sqrt{2}}\right)\overline{\chi}_{\alpha,k}(s)\,ds. \end{aligned}$$

Now substitute

$$r = \frac{\sigma_{\alpha,k-1}}{\sigma_{\alpha,k}} s, \qquad y = qx = \sqrt{1+\beta^2}\,\frac{\sigma_{\alpha,k}}{\sigma_{\alpha,k-1}} x.$$

Using the FCM density in (7.4), this change of variables gives

$$\overline{\chi}_{\alpha,k}(s)\,ds = \frac{C_{\alpha,k}}{\sigma_{\alpha,k-1}C_{\alpha,k-1}}\,\overline{\chi}_{\alpha,k-1}(r)\,r\,dr.$$

Moreover, since

$$\delta = \frac{\beta}{\sqrt{1+\beta^2}}, \qquad \mu = \frac{A\delta}{\sqrt{2}},$$

we have

$$\mu y r = \frac{A\beta x s}{\sqrt{2}}.$$

Therefore,

$$\begin{aligned} L_{\alpha,k}(x;\beta) &\approx -\frac{1}{B\pi\beta x}\,\frac{C_{\alpha,k}}{\sigma_{\alpha,k-1}C_{\alpha,k-1}}\int_0^\infty e^{-(yr)^2/2}\,(1-e^{\mu y r})\,\overline{\chi}_{\alpha,k-1}(r)\,r\,dr \\ &= \frac{1}{\beta x}\sqrt{\frac{2}{\pi}}\,\frac{C_{\alpha,k}}{B\,\sigma_{\alpha,k-1}C_{\alpha,k-1}}\left[e^{\mu^2/2}L_{\alpha,k-1}(y||\mu) - L_{\alpha,k-1}(y)\right], \end{aligned}$$

where the last equality follows from the shifted-GSaS definition with k replaced by $k-1$. Since $y = qx$, this proves the stated left-tail approximation.

For the right tail, use the identity

$$2\Phi_{\mathcal{N}}(z) + 2\Phi_{\mathcal{N}}(-z) = 2.$$

Together with the symmetry of $\mathcal{N}(xs)$, this implies

$$L_{\alpha,k}(x;\beta) + L_{\alpha,k}(-x;\beta) = 2L_{\alpha,k}(x).$$

Replacing the left-tail term by the approximation just derived gives

$$L_{\alpha,k}(x;\beta) \approx 2L_{\alpha,k}(x) - \widehat{L}_{\alpha,k}(-x;\beta), \qquad x > 0.$$

□

REMARK 12.16. In the case of $\beta \to 0$, the apparent $1/\beta$ singularity in the approximation is removable at the level of the formal expansion. Indeed,

$$\mu = \frac{A\delta}{\sqrt{2}} = \frac{A\beta}{\sqrt{2}} + O(\beta^3),$$

and the shifted and unshifted terms in the bracket differ by

$$e^{\mu^2/2}L_{\alpha,k-1}(qx||\mu) - L_{\alpha,k-1}(qx) = O(\mu) = O(\beta).$$

This cancels the explicit $1/\beta$ factor. However, the approximation is not uniform as $\beta \to 0$. For fixed x, the error-function approximation has the local limit

$$\lim_{z\to 0}\frac{(1-e^{-Az})e^{-z^2}}{B\sqrt{\pi}\,z} = \frac{A}{B\sqrt{\pi}} \approx 0.984225,$$

whereas the exact complementary error-function factor equals 1 at the origin. Hence the formula is best viewed as a suppressed-tail approximation, effectively for $\beta|x| \gg 1$, rather than a formula to use near $\beta = 0$.

12.7. Maximum Skewness and Half GSaS

When $\beta \to \pm\infty$, a GAS-SN becomes a half-GSaS, which is a one-sided distribution with the notation of $L^{\pm}_{\alpha,k} := L_{\alpha,k}(\beta = \pm\infty)$. Its PDF is

$$L^{+}_{\alpha,k}(x) = \begin{cases} 2L_{\alpha,k}(x) & \text{if } x \geq 0, \\ 0 & \text{otherwise.} \end{cases} \tag{12.36}$$

It follows the reflection rule of $L^{-}_{\alpha,k}(x) = L^{+}_{\alpha,k}(-x)$. Hence, we only need to study the $+\infty$ case.

A half-GSaS possesses the maximum skewness that a GAS-SN family can achieve for a given pair of (α, k). In Section 10.5, it was mentioned that the maximum skewness of the SN family is only 0.9953. GAS-SN allows the skewness to reach infinity potentially.

From (12.7), the n-th moment is

$$\begin{aligned} \mathbb{E}(X^n | L^{+}_{\alpha,k}) &= 2L^{*}_{\alpha,k}(n+1) \\ \mathbb{E}(X^n | L^{-}_{\alpha,k}) &= 2L^{*}_{\alpha,k}(n+1)\,(-1)^n \end{aligned} \tag{12.37}$$

Therefore, it is straightforward to calculate the skewness.

The skewness of half-GSaS $L^{+}_{\alpha,k}$ is shown in Figure 12.8 in the (k, α) coordinate. There is a clear division of infinity by the line from $(2, 2)$ to $(4, 0)$.




FIGURE 12.8. The contour plot of skewness of the half-GSaS $L^{+}_{\alpha,k}$ by (k, α). This represents the maximum skewness that the GAS-SN family can achieve.

The contour plot of the skewness is shown in the (k, ϵ) coordinate in Figure 12.9. Each contour line approaches a straight line as k increases.




FIGURE 12.9. The contour plot of skewness of the half-GSaS $L^+_{\alpha,k}$ by (k, ϵ) where $\epsilon = 1/\alpha$. Each contour line approaches a straight line as k increases.

12.8. Fractional Skew Exponential Power Distribution

As shown in Definition 3.6 and Section 9 of [20], the negative k space is reserved for the fractional exponential power distribution, whose PDF is $\mathcal{E}_{\alpha,k}(x) := L_{\alpha,-k}(x)$. All it takes is to have $\overline{\chi}_{\alpha,k}(s)$ in (12.1) properly defined for negative k, which is done in (7.11).

It is natural to extend it with the skew-normal family such that its PDF becomes

$$\mathcal{E}_{\alpha,k}(x;\beta) = L_{\alpha,-k}(x;\beta). \tag{12.38}$$

Then we obtain another flexible skew distribution with a different type of tail behavior. Detailed analysis of this distribution is left for future research.

12.9. Quadratic Form

A squared GAS-SN variable Q is distributed as a fractional F distribution with $d = 1$. That is,

$$Q := \left(\frac{Y-\xi}{\omega}\right)^2 = Z^2 \sim F_{\alpha,1,k}, \quad \text{for all } \beta. \tag{12.39}$$

Notice that Q is based on the standard variable Z, which is invariant to the location and scale. See Chapter 8 for more detail.

12.10. Univariate MLE

In this section, we document how we fit the one-dimensional data with univariate GAS-SN. The main algorithm is *maximum likelihood estimation* (MLE), supplemented with several components of regularization.

We applied the MLE fitting on the daily returns of VIX and SPX from 1990 to mid-2025, each about 8900 samples. The MLE program is implemented in **python** and **scipy** on github at https://github.com/slihn/gas-impl/blob/main/gas_impl/mle_gas_sn.py.

In the univariate case, the hyperparameter space is $\Theta = \{\alpha, k, \beta, \omega, \xi\}$, where $\alpha \in (0, 2)$, $k \in (2, \infty)$, $\omega > 0$, and $\beta, \xi \in \mathbb{R}$. Assume there are N samples in the data set, $Y = \{y_i, i \in 1, 2, ..., N\}$, the main component of the objective function is the minus log-likelihood (MLLK):

$$\text{MLLK}(\Theta) = -\frac{1}{N}\sum_{i=1}^{N} \log(\phi(y_i; \Theta)) \tag{12.40}$$

where $\phi(y; \Theta)$ is the PDF of the univariate location scale family (12.4).

Additional components of regularization are added to the objective function $\ell(\Theta)$. Specifically, the L2 distances between the empirical and theoretical statistics are added as follows:

- Skewness: $|\Delta\gamma_1|^2 := |\Delta\text{skewness}(Y)|^2$. Section 12.4.
- Kurtosis: $|\Delta\gamma_2|^2 := |\Delta\text{kurtosis}(Y)|^2$. Section 12.4.
- The mean of the quadratic form: $\Delta\mu_Q^2 := |\Delta\text{mean}(Q)|^2$. Section 15.6.

The MLE seeks the optimal Θ that minimizes the objective function:

$$\hat{\Theta} = \text{argmin } \ell(\Theta) \tag{12.41}$$

$$\text{where } \ell(\Theta) = \text{MLLK}(\Theta) + |\Delta\gamma_1|^2 + |\Delta\gamma_2|^2 + \Delta\mu_Q^2 \tag{12.42}$$

A custom version of the stochastic descent (SD) algorithm is developed. Our experience shows that it is better to standardize the data set to one standard deviation, so that all parameters in Θ are approximately on the same scale.

It is also important to control the learning rate so that it does not take a too large step on α, empirically, no more than 0.01 per step. This ensures that the SD does not wander into the *undefined* regions for $\ell(\Theta)$. This is particularly important for the SPX fit below.

The SD algorithm calculates the gradients for each hyperparameter. And make a small move along the direction that is most likely to minimize $\ell(\Theta)$. The scale of the move is based on the learning rate, which can be dynamically adjusted. Some randomness is added to the small move. This allows the algorithm to explore the nearby region and increases its choices.

12.11. Examples of Univariate MLE Fits

12.11.1. VIX fit. Figure 12.10 shows the result of the MLE fit to the daily VIX returns from 1990 to mid-2025. Data are standardized to one standard deviation. This helps the SD algorithm to move correctly in all dimensions of Θ.

The VIX data are strongly right-skewed, with sample skewness about 2.0. The right tail is substantially extended by several *panic selling* episodes in which the VIX rises sharply within a single day. As a result, the sample kurtosis is approximately 17.

The top two graphs compare the histogram with the fitted theoretical density. The graph on the right shows the density on a logarithmic scale, which makes it easier to assess the tail fit down to the 10^{-3} level. In particular, the theoretical density does not fully capture the far right tail beyond about 7.

The fitted theoretical parameters are $\alpha \approx 0.77$, $k \approx 5.2$, and $\beta \approx 0.95$. This fitted point can be located in Figure 12.1.

The PP-plot in the bottom-left graph indicates that the overall fit is acceptable. The points remain close to the 45-degree line, although this plot is relatively insensitive to tail behavior.

The bottom-right graph shows the QQ-plot of the quadratic form, or squared variable. This diagnostic is particularly useful for assessing how well the distribution captures the combined tail behavior in absolute terms. The 45-degree line is OK below 20, but as the quantiles get larger, the

observed quantiles start to tilt upward. This means the top 0.5 percent of the combined tail is not properly captured by the distribution.




FIGURE 12.10. MLE fit of daily VIX returns from 1990 to mid-2025. The data are standardized to unit variance. Sample skewness is 2.0, and sample kurtosis is 16.5. $\hat{\Theta} = \{\alpha = 0.77, k = 5.2, \beta = 0.95\}$. Top left: density and histogram. Top right: log-density and histogram. Bottom left: PP-plot. Bottom right: QQ-plot of the quadratic form.

12.11.2. SPX fit. Figure 12.11 shows the result of the MLE fit to the daily SPX returns from 1990 to mid-2025. The data are also standardized to one standard deviation.

The SPX data are slightly left-skewed, with the sample skewness slightly below -0.1. The tails are extended by several large one-day market declines, producing sample kurtosis of approximately 11 while leaving the skewness relatively modest.

The top two graphs compare the histogram with the fitted theoretical density. The graph on the right shows the density on a logarithmic scale, making it easier to assess the tail fit down to the 10^{-3} level. The theoretical density does not fully capture the tails beyond about 7.

The fitted theoretical distribution has parameters $\alpha = 0.15$, $k = 19.9$, and $\beta = -0.11$. This MLLK fit is in the low-α, high-k region, marked as "SPX" in Figure 12.4.

The PP-plot in the bottom-left graph indicates that the overall fit is satisfactory, with most points lying on the 45-degree line.

In the QQ-plot of the quadratic form, the points remain close to the 45-degree line up to about 20. However, for larger quantiles, the observed quantiles begin to bend upward. In particular, the ten largest observations deviate materially from the theoretical curve.




FIGURE 12.11. MLE fit of daily SPX log returns from 1990 to mid-2025. The data are standardized to unit variance. Sample skewness is -0.14, and sample kurtosis is 10.6. $\hat{\Theta} = \{\alpha = 0.15, k = 19.9, \beta = -0.11\}$. Top left: density and histogram. Top right: log-density and histogram. Bottom left: PP-plot. Bottom right: QQ-plot of the quadratic form.

CHAPTER 13

Fractional Feller Square-Root Process

This chapter is copied from Section 11 of [20] for the generation of random variables for FG, FCM, and FCM2. Combining this with an SN variable provides a path to generate the random variable for GAS-SN and beyond.

For example, assuming that a sequence of random numbers $\{S_t > 0\}$ can be generated for FCM, it is straightforward to simulate random numbers $\{X_t\}$ for GAS-SN using the ratio of $X_t = Y_t/S_t$, where Y_t is a standard skew-normal variable $Y_t \sim SN(0, 1, \beta)$ in Chapter 12.

Instead of randomly generating $\{S_t\}$, we propose an innovative method based on *Feller square-root process*[11]. Given a user-specific volatility $\sigma_u > 0$ that describes how fast S_t should change, a scalar function $\mu(x)$, and a scale parameter $\theta_u > 0$ (default to 1), we assume that the random variable S_t should evolve according to the following generalized process:

$$dS_t = \sigma_u^2\, \mu\left(\frac{S_t}{\theta_u}\right) dt + \sigma_u \sqrt{S_t}\, dW_t \tag{13.1}$$

As $t \to \infty$, $\{S_t\}$ will be distributed as the equilibrium distribution for which $\mu(x)$ is designated.

13.0.1. The Fokker-Planck Equation. The $\mu(x)$ solution can be derived from the Fokker-Planck equation. We obtain the following beautiful relation:

LEMMA 13.1. *$\mu(x)$ is one half of the elasticity of the terminal density function $p(x)$ of S_t at $t \to \infty$ plus one half:*

$$\mu(x) = \frac{1}{2}\mathcal{L}\, p(x) + \frac{1}{2} \tag{13.2}$$

where $\mathcal{L}(\cdot) := x\frac{d}{dx}\log(\cdot)$ is the elasticity operator defined in Section 3.7.

△

PROOF. Assume $p(x, t)$ is the density function of (13.1) for S_t. It should satisfy the Fokker-Planck equation ($\theta_u = 1$):

$$\frac{\partial}{\partial t}p(x,t) = -\frac{\partial}{\partial x}\left[\sigma_u^2\, \mu(x)\, p(x,t)\right] + \frac{\partial^2}{\partial x^2}\left[\frac{1}{2}(\sigma_u\sqrt{x})^2\, p(x,t)\right]$$

As $t \to \infty$, $p(x, t)$ approaches the terminal density function $p(x)$. The time dependency is removed. σ_u^2 cancels out from both sides and is irrelevant to the solution. The ODE of $p(x)$ becomes

$$\frac{\partial}{\partial x}\left(\mu(x)\, p(x)\right) = \frac{1}{2}\frac{\partial^2}{\partial x^2}\left(x\, p(x)\right)$$

Apply $\int_x^\infty dx$ to both sides. Assuming that $\mu(x)p(x)$ at $x = \infty$ should be zero, we get

$$\mu(x)p(x) = \frac{1}{2}\frac{d}{dx}\left(x\, p(x)\right) = \frac{1}{2}\left(x\frac{d}{dx}p(x) + p(x)\right)$$

Moving $p(x)$ from LHS to RHS, we obtain (13.2).

□

13.0.2. Generation of Random Variables for FG.

LEMMA 13.2. The $\mu(x)$ solution for FG is obviously

$$\mu(x) = \frac{1}{2}\mathcal{L}\,\mathfrak{N}_\alpha(x;\sigma,d,p) + \frac{1}{2}$$

With Lemma 3.10, $\mu(x)$ is reduced to a function of $\mathcal{L}\,M_\alpha(x)$:

$$\mu(x) = \frac{p}{2}\left[\mathcal{L}\,M_\alpha\right]\left(\left(\frac{x}{\sigma}\right)^p\right) + \frac{d+p}{2}. \tag{13.3}$$

△

As an application, since $\mathcal{L}\,M_{1/2}(x) = -x^2/2$ from Section 3.7, we obtain a simple power-law solution at $\alpha = 1/2$:

$$\mu(x)|_{\mathfrak{N}_{1/2}} = -\frac{p}{4}\left(\frac{x}{\sigma}\right)^{2p} + \frac{d+p}{2} \tag{13.4}$$

Note that (13.4) at $p = 1/2$ subsumes the renown Cox–Ingersoll–Ross (CIR) model[6] since the $\mu(x)$ of the model is a linear $a(b-x)$ type, according to its stochastic process of $dS_t = a(b - S_t)\,dt + \sigma_u\sqrt{S_t}\,dW_t$.[1]

To prepare for the solution of FCM, we prefer to use $Q_\alpha(x)$ defined in (3.42):

$$Q_\alpha(x) := -\frac{W_{-\alpha,-1}(-x)}{W_{-\alpha,0}(-x)}$$

LEMMA 13.3. From (3.44), the $\mu(x)$ solution of a FG in terms of $Q_\alpha(x)$ is

$$\mu(x) = \frac{p}{2\alpha}Q_\alpha\left(\left(\frac{x}{\sigma}\right)^p\right) + \left(\frac{d}{2} - \frac{p}{2\alpha}\right) \tag{13.5}$$

Notice that p/α and d are just constant terms, and σ only affects the scale of x. Neither of them has any effect on the shape of $\mu(x)$.

△

13.1. Generation of Random Variables for FCM

Obviously, what really matters for GAS-SN and GSaS is the solution of FCM, The $\mu(x)$ solution for $\overline{\chi}_{\alpha,k}$ is denoted as $\mu_{\alpha,k}(x)$. Note that from this point on, $\alpha \in (0,2)$.

To further simplify the symbology for FCM, define

$$Q_\alpha^{(\chi)}(z) := Q_{\frac{\alpha}{2}}\left(z^\alpha\right),\ \text{where } \alpha \in (0,2).$$

Assuming $k > 0$, we set $\sigma = \sigma_{\alpha,k}, d = k-1, p/\alpha = 2$ and α replaced by $\alpha/2$ in (13.5). We get

$$\mu_{\alpha,k}(x) = Q_\alpha^{(\chi)}\left(\frac{x}{\sigma_{\alpha,k}}\right) + \left(\frac{k-3}{2}\right) \tag{13.6}$$

For validation, $\mu_{1,k}(x) = k(1-x^2)/2$ can be used to simulate Student's t. And $\mu_{\alpha,1}(x)$ provides a method to simulate an SaS $L_{\alpha,1}(x)$:

$$\mu_{\alpha,1}(x) = Q_\alpha^{(\chi)}\left(\sqrt{2}x\right) - 1$$

Fig. 13.1 shows a simulation of random variables based on the (α,k) parameter obtained from the fit of the S&P 500 daily log returns. The rest of the parameters are in the caption of the figure. First, as outlined above, $\mu_{\alpha,k}(s)$ is calculated analytically as shown in the right graph. Second, it enables

[1] It can also be subsumed by the FG at $\alpha = 0, p = 1$. But $\alpha = 0$ is a singular point and we prefer to avoid using it when possible.

the FG simulation $\{S_t\}$ as shown in the left graph. Third, GSaS $\{X_t\}$ is simulated via $X_t = \mathcal{N}/S_t$, where $\mathcal{N}$ is drawn from a standard normal variable.

The simulation is performed daily. The duration of the sampling is 200,000 years. The red areas are histograms of the simulated data. The blue lines are from the theoretical density functions. They match well.




FIGURE 13.1. Simulation of random variables based on the (α, k) parameters obtained from the fit of the S&P 500 daily log returns. The red areas are the histograms from simulated data. The blue lines are from theoretical formulas. The settings of the simulation are $\alpha = 0.813, k = 3.292, dt = 1/365, \sigma_u = 0.85$. Sampling duration is 200,000 years. The simulation takes 11 minutes in python. $\mu_{\alpha,k}(s)$ is discretized to 0.01 and cached to increase performance.

13.2. Generation of Random Variables for FCM2

LEMMA 13.4. The $\mu(x)$ solution for $\overline{\chi}^2_{\alpha,k}$ is

$$\mu^{(2)}_{\alpha,k}(x) = \frac{1}{2}\mu_{\alpha,k}\left(\sqrt{x}\right) \tag{13.7}$$

△

PROOF. From (7.60), we have

$$\overline{\chi}^2_{\alpha,k}(x) := \mathfrak{N}_{\alpha/2}(x; \sigma = \sigma^2_{\alpha,k}, d = (k-1)/2, p = \alpha/2) \qquad (k > 0)$$

Combined with (13.5), we obtain the solution for $\overline{\chi}^2_{\alpha,k}$ as

$$\begin{aligned}\mu^{(2)}_{\alpha,k}(x) &= \frac{1}{2}Q_{\alpha/2}\left(\left(\frac{\sqrt{x}}{\sigma_{\alpha,k}}\right)^{\alpha}\right) + \left(\frac{k-1}{4} - \frac{1}{2}\right) \\ &= \frac{1}{2}Q^{(\chi)}_{\alpha}\left(\frac{\sqrt{x}}{\sigma_{\alpha,k}}\right) + \frac{k-3}{4},\end{aligned}$$

which is just

$$\mu^{(2)}_{\alpha,k}(x) = \frac{1}{2}\mu_{\alpha,k}\left(\sqrt{x}\right).$$

□

This solution can be used to simulate the F distribution in Chapter 8. Let $U_1 \sim \chi^2_d/d = \bar{\chi}^2_{1,d}$ and $U_2 \sim \overline{\chi}^2_{\alpha,k}$, then $F_{\alpha,d,k} \sim U_1/U_2$ is a fractional F distribution.

Part 4

Multivariate Distributions

CHAPTER 14

Multivariate SN Distribution - Review

In this chapter, we start to explore the multivariate distributions. Data sets from the real world are often multidimensional. A flexible multivariate distribution framework with skewness and kurtosis can be very useful. That is what we aim to achieve in the next few chapters.

The foundation is the standard multivariate normal distribution $\mathcal{N}_d(0, \bar{\Omega})$, where d is the dimension of the random variable, and $\bar{\Omega}$ is a $d \times d$ correlation matrix[41].

In Chapter 5 of Azzalini, the skew normal distribution $SN_d(0,\bar{\Omega}, \boldsymbol{\beta})$ adds skewness to it from the skew parameter $\boldsymbol{\beta}$ [1]. In its Chapter 6, the skew-elliptical distribution is discussed. The multivariate skew-t distribution $ST_d(0, \bar{\Omega}, \boldsymbol{\beta}, k)$ is constructed by combining $SN_d(0,\bar{\Omega}, \boldsymbol{\beta})$ with $\chi_k/\sqrt{k}$ in a ratio distribution.

Our work builds on top of this concept of the skew-elliptical distribution. By expanding the denominator of $\chi_k/\sqrt{k}$ to the FCM $\overline{\chi}_{\alpha,k}$, the fractional dimension α is added to the shape parameters. This forms a super-distribution family called *multivariate GAS-SN elliptical distribution* with the notation $L_{\alpha,k}(0, \bar{\Omega}, \boldsymbol{\beta})$ for its standard distribution.

The multivariate skew-elliptical distribution has beautiful properties inherited from the multivariate elliptical distribution framework. However, its deficiency is obvious in real-world applications: The structure is multivariate, but the shape parameters α and k are scalars. All dimensions share the same (α, k). This restricts the kurtoses of 1D marginal distributions to a similar range. It even creates some strange phenomena that are hard to interpret in the SPX-VIX 2D fit (see Section 17.1.1).

To overcome such a restriction, we propose a more flexible framework called *multivariate adaptive distribution*, in which the shape parameters $(\boldsymbol{\alpha}, \boldsymbol{k})$ are d dimensional vectors, just like their skew counterpart $\boldsymbol{\beta}$.

The flexibility in shapes comes with an expensive computational cost. It is analogous to the *curse of dimensionality* problem. It becomes much harder to verify the results beyond the bivariate case for the adaptive distribution.

The study of quadratic form $Z^\intercal \bar{\Omega}^{-1} Z$ from the skew-elliptical distribution results in the fractional extension of the F distribution $F_{\alpha,d,k}$. The QQ-plot based on the quadratic form and the fractional F distribution is a powerful validation of the goodness of the fit.

14.1. Definition

We summarize the results of Chapter 5 of Azzalini[1]. On the one hand, we need to clarify the symbology here that is slightly different from that in his book. On the other hand, our multivariate distributions rely on many results from there, which are collected in this chapter.

DEFINITION 14.1. The PDF of a standard multivariate normal distribution $\mathcal{N}_d(0, \bar{\Omega})$ is defined as

$$\mathcal{N}_d(\boldsymbol{x}; \bar{\Omega}) := \frac{1}{(2\pi)^{d/2} \det(\bar{\Omega})^{1/2}} \exp\left(-\frac{1}{2}\boldsymbol{x}^\intercal \bar{\Omega}^{-1} \boldsymbol{x}\right), \quad \boldsymbol{x} \in \mathbb{R}^d. \tag{14.1}$$

where $\bar{\Omega}$ is a $d \times d$ correlation matrix[41]. That is, $\bar{\Omega}$ is positive definite and all its diagonal elements are equal to 1.

DEFINITION 14.2. A standard multivariate skew-normal variable is denoted as $Z \sim SN_d(0,\bar{\Omega},\boldsymbol{\beta})$, where $\boldsymbol{\beta} \in \mathbb{R}^d$ is the skew parameter (or the slant parameter). Its PDF is

$$\mathcal{N}_d(\boldsymbol{x};\bar{\Omega},\boldsymbol{\beta}) := \mathcal{N}_d(\boldsymbol{x};\,\bar{\Omega})\,\Phi_{\mathcal{N}}(\boldsymbol{\beta}^\intercal\boldsymbol{x}), \tag{14.2}$$

where $\Phi_{\mathcal{N}}(x)$ is the CDF of a standard normal distribution $\mathcal{N}(0,1)$.

Notice that this is a multivariate expansion of SN in Section 10.1. When $d = 1$, (14.2) becomes (10.2).

14.2. The Location-Scale Family

Its location-scale family is $Y = \boldsymbol{\xi} + \boldsymbol{\omega}Z \sim SN_d(\boldsymbol{\xi},\Omega,\boldsymbol{\beta})$, where $\boldsymbol{\xi} \in \mathbb{R}^d$ is the location parameter, $\boldsymbol{\omega} = \text{diag}(\omega_1,...,\omega_d)$ is a $d \times d$ diagonal scale matrix ($\omega_i > 0, \forall i$) and $\Omega = \boldsymbol{\omega}\bar{\Omega}\boldsymbol{\omega}$.

The PDF of Y becomes

$$f_Y(\boldsymbol{x}) = \det(\boldsymbol{\omega})^{-1}\,\mathcal{N}_d(\boldsymbol{z};\bar{\Omega},\boldsymbol{\beta}), \tag{14.3}$$

where $\boldsymbol{z} = \boldsymbol{\omega}^{-1}(\boldsymbol{x}-\boldsymbol{\xi})$.

The location-scale distribution is used for real-world applications. Internally, it has to be calculated via the standard distribution. The main reason is that $\boldsymbol{\beta}$ has to work with $\boldsymbol{z}$ and $\bar{\Omega}$, instead of $\boldsymbol{x}$ and $\boldsymbol{\omega}$.

14.3. Quadratic Form

DEFINITION 14.3. The quadratic form of a multivariate SN distribution (MSN) is defined as

$$Q := \frac{1}{d}(Y-\boldsymbol{\xi})^\intercal\Omega^{-1}(Y-\boldsymbol{\xi}) = \frac{1}{d}Z^\intercal\bar{\Omega}^{-1}Z. \tag{14.4}$$

Q distributes as $\chi^2_d/d = \bar{\chi}^2_{1,d}$ for all $\boldsymbol{\beta}$. The distribution of Q is independent of $\boldsymbol{\beta}$. This is an important property due to the rotational invariance of the elliptical distribution.

Notice that our definition of Q is slightly different from that of Azzalini. We prefer to have the distribution of Q tied to the FCM and the fractional F distribution directly without any constant adjustment. This will make things much simpler in Section 15.6.

To prove $Q \sim \chi^2_d/d$, we quote Corollary 5.9 from [1] below for a skew-normal distribution with 0 location:

LEMMA 14.4. If $Y \sim SN_d(0,\Omega,\boldsymbol{\beta})$ and A is a $d \times d$ symmetric matrix, then

$$Y^\intercal AY = X^\intercal AX$$

where $X \sim \mathcal{N}_d(0,\Omega)$. △

This lemma allows $\boldsymbol{\beta}$ to be removed from the statistics of Q. Hence, $Q \sim X^\intercal\Omega^{-1}X/d \sim \chi^2_d/d$.

14.4. Stochastic Representation

Assuming $X_0 \sim \mathcal{N}_d(0,\bar{\Omega})$ and $X_1 \sim \mathcal{N}(0,1)$, then the first representation of $Z \sim SN_d(0,\bar{\Omega},\boldsymbol{\beta})$ is

$$Z = \begin{cases} X_0 & \text{if } X_1 > \boldsymbol{\beta}^\intercal X_0, \\ -X_0 & \text{otherwise.} \end{cases} \tag{14.5}$$

This form of selective sampling is quite useful in generating random numbers for Z. It is essentially an extension of (10.1).

This scheme can be rephrased in a more interesting representation. First, define the multivariate version of δ as

$$\boldsymbol{\delta} = (1 + \boldsymbol{\beta}^\intercal \bar{\Omega} \boldsymbol{\beta})^{-1/2} \, \bar{\Omega} \boldsymbol{\beta}, \quad (\boldsymbol{\delta} \in \mathbb{R}^d) \tag{14.6}$$

which is used to construct a $(d+1) \times (d+1)$ correlation matrix

$$\Omega^* = \begin{pmatrix} \bar{\Omega} & \boldsymbol{\delta} \\ \boldsymbol{\delta}^\intercal & 1 \end{pmatrix}.$$

Ω^* is used to generate two marginals, $X_0 \in \mathbb{R}^d$ and $X_1 \in \mathbb{R}$, such that

$$\begin{pmatrix} X_0 \\ X_1 \end{pmatrix} \sim \mathcal{N}_{d+1}(0, \Omega^*),$$

which leads to the second representation

$$Z = \begin{cases} X_0 & \text{if } X_1 > 0, \\ -X_0 & \text{otherwise.} \end{cases} \tag{14.7}$$

This form resembles (10.7). It shows that the function of $\boldsymbol{\delta}$ is to add the correlation between X_0 and X_1 through Ω^* in the selective sampling. This makes (14.7) slightly different from (14.5).

14.5. Moments

The first two moments of Z have simple analytic forms. Its first moment is

$$\mu_z = \mathbb{E}(Z) = b \, \boldsymbol{\delta}, \quad \text{where } b = \sqrt{2/\pi}. \tag{14.8}$$

The second moment is simply $\bar{\Omega}$. Its variance is

$$\Sigma_z = \text{var}\{Z\} = \bar{\Omega} - b^2 \, \boldsymbol{\delta}\boldsymbol{\delta}^\intercal. \tag{14.9}$$

It is easy to obtain $\mathbb{E}\{Y \, Y^\intercal\} = \Omega$ for the location-scale variable Y.

Define the important invariant quantity for the skewness.

$$\beta_* = (\boldsymbol{\beta}^\intercal \bar{\Omega} \boldsymbol{\beta})^{1/2} \geq 0, \tag{14.10}$$

which is a nonnegative scalar quantity. It encapsulates the departure from normality for the distribution.

The quadratic form $\mu_z^\intercal \Sigma_z^{-1} \mu_z$ can be simplified to

$$\mu_z^\intercal \Sigma_z^{-1} \mu_z = \frac{b^2 \beta_*^2}{1 + (1 - b^2)\beta_*^2}. \tag{14.11}$$

A related quantity is

$$\delta_* = (\boldsymbol{\delta}^\intercal \bar{\Omega}^{-1} \boldsymbol{\delta})^{1/2} \tag{14.12}$$

where $\delta_* \in [\, 0, 1)$ has the scale of a positive correlation coefficient.

The two are connected by

$$\delta_*^2 = \frac{\beta_*^2}{1 + \beta_*^2}, \quad \beta_*^2 = \frac{\delta_*^2}{1 - \delta_*^2}.$$

Or in a trigonometric form, there exists an angle $\theta \in [\, 0, \frac{\pi}{2})$ such that $\tan\theta = \beta_*$ and $\sin\theta = \delta_*$. In such an expression, $\theta > 0$ captures the "degree" of departure from normality.

14.6. Canonical Form

The concept of a canonical form in SN is very important and fascinating. Due to the rotational symmetry, an MSN can be rotated and rescaled to an "identity" MSN with a scalar skew parameter.

By Proposition 5.12 of [1], there exists an affine transformation $Z^* = A_*(Y - \boldsymbol{\xi})$ such that $Z^* \sim SN_d(0, \boldsymbol{I}_d, \boldsymbol{\beta}_{Z^*})$, where $\boldsymbol{I}_d$ is a $d \times d$ identity matrix, and $\boldsymbol{\beta}_{Z^*} = (\beta_*, 0, ..., 0)^\intercal$. β_* is defined by (14.10), which is an invariant under transformation.

The variable Z^* is called *the canonical variable*. It is d-dimensional. But only one dimension is skew-normal, which is designated as the first dimension. All other dimensions are standard normal distributions. That is, the PDF of Z^* is

$$\begin{aligned}\mathcal{N}_*(\boldsymbol{x}; \beta_*) &= 2\Phi_{\mathcal{N}}(\beta_* x_1) \prod_{i=1}^{d} \mathcal{N}(x_i) \\ &= \mathcal{N}(x_1; \beta_*) \prod_{i=2}^{d} \mathcal{N}(x_i).\end{aligned}$$

This structure helps tremendously for the subsequent development of the elliptical distribution and adaptive distribution.

Proposition 5.13 in [1] describes how to find such A_*. Due to rotational symmetry, there are many choices of A_*. This is not a problem as long as we always look at the system in quadratic form.

LEMMA 14.5 (Affine Transformation). Let $C = \Omega^{1/2}$ be the unique positive definite symmetric square root of Ω. Define $M = C^{-1}\Sigma C^{-1}$, where $\Sigma = \text{var}\{Y\}$. Let $Q\Lambda Q^\intercal$ denote a spectral decomposition of M, where we assume that the diagonal elements in the eigenvalue matrix Λ are arranged in increasing order.

Let $H = C^{-1}Q$. Then H is the matrix operator to convert Y to Z^*,

$$Z^* = H^\intercal(Y - \boldsymbol{\xi}).$$

Since $\boldsymbol{\delta}_{Z^*} = H^\intercal \boldsymbol{\omega} \boldsymbol{\delta}$ and $\boldsymbol{\beta}_{Z^*} = \boldsymbol{\delta}_{Z^*}/(1 - \delta_*^2)$, the choice of H must make the first element of $\boldsymbol{\delta}_{Z^*}$ a nonnegative number, that is, $\delta_* \geq 0$. All other elements, except the first ones in $\boldsymbol{\delta}_{Z^*}$ and $\boldsymbol{\beta}_{Z^*}$, must be zero. △

REMARK 14.6. The significance of this lemma is that the skew-elliptical distributions derived from the SN framework can only have a single source of skewness. It might be mixed up and not easy to observe in real-world data. But there is only one source from the theoretical perspective. Everything else comes from the multivariate normal distribution.

If we want a more "sophisticated" distribution that provides multiple sources of skewness, we have to go beyond the skew-elliptical distributions.

14.7. 1D Marginal Distribution

We are particularly interested in the 1D marginal distribution, since this is what is actually observed in a data set. When we optimize a data fit, we can add the log-likelihood of the 1D marginal distributions to the objective function, so that the fitting of each dimension is properly addressed.

In fact, for the adaptive distribution, the full 2D likelihood is so compute-intensive that it is too slow to perform MLE on a desktop. The alternative is to compute the sum of the log-likelihoods of each 1D marginal distribution, in addition to the regularization on other statistical quantities, such as the correlation coefficient between each data pair.

We quote the results from Section 5..1.4 of [1] and adapt them to the 1D case.

LEMMA 14.7. (The marginal β) Assume that the marginal is on the first dimension. The correlation matrix is decomposed as

$$\bar{\Omega} = \begin{pmatrix} \bar{\Omega}_{11} & \bar{\Omega}_{12} \\ \bar{\Omega}_{21} & \bar{\Omega}_{22} \end{pmatrix}.$$

The formula can be simplified due to $\bar{\Omega}_{11} = 1$ in the 1D case.

The marginal distribution is $Y_1 \sim SN(\xi_1, \Omega_{11}, \beta_{1(2)})$. Its $\beta_{1(2)}$ is derived as

$$\beta_{1(2)} = \left(1 + \beta_2^{\intercal} \bar{\Omega}_{22.1} \beta_2\right)^{-1/2} \left(\beta_1 + \bar{\Omega}_{12} \beta_2\right) \quad \text{where} \quad \bar{\Omega}_{22.1} = \bar{\Omega}_{22} - \bar{\Omega}_{21} \bar{\Omega}_{12}. \tag{14.13}$$

△

LEMMA 14.8. (The marginals of a bivariate distribution) The bivariate case is quite simple:

$$\bar{\Omega} = \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix}. \tag{14.14}$$

Assume that we want to get the marginal β of the i-th dimension, $\beta_{i(j)}$, where j is the other dimension. Then

$$\beta_{i(j)} = \frac{\beta_i + \rho \beta_j}{\sqrt{1 + \beta_j^2 \, |\bar{\Omega}|}} \tag{14.15}$$

where $|\bar{\Omega}| = 1 - \rho^2$. Since Ω_{ii} is ω_i^2, the i-th marginal distribution is $Y_i \sim SN(\xi_i, \omega_i^2, \beta_{i(j)})$. The ξ_i and ω_i are the location and scale parameters in the i-th dimension that can be calculated directly from the data. △

We observe that ρ in the numerator describes how much β_j is mixed with β_i, while $|\bar{\Omega}|$ in the denominator describes how much β_j reduces the scale.

When $\rho = 0$, there is no mixing from the other dimension, only a reduction in total scale. That is, $\beta_{i(j)}|_{\rho=0} = \beta_i / \sqrt{1 + \beta_j^2}$.

CHAPTER 15

Multivariate GAS-SN Elliptical Distribution

15.1. Definition

This chapter follows the structure laid out in Chapter 6 of Azzalini (2013)[1]. We implemented the skew-elliptical distribution by our $\overline{\chi}_{\alpha,k}$, which fully extends his multivariate skew-t distribution.

DEFINITION 15.1. Assume $Z_0 \sim SN_d(0,\bar{\Omega},\boldsymbol{\beta})$ is a $d \times d$ standard multivariate skew-normal (SN) distribution, and $V \sim \overline{\chi}_{\alpha,k}$ is a standard FCM. $\bar{\Omega}$ is a correlation matrix.

Then $Z \sim Z_0/V$ is a $d \times d$ standard multivariate GAS-SN elliptical distribution. It is given the notation of $Z \sim L_{\alpha,k}(0,\bar{\Omega},\boldsymbol{\beta})$.

Equivalently, using the location-scale notation, $Z \sim SN_d(0,\Sigma,\boldsymbol{\beta})$ where $\Sigma = \bar{\Omega}/V^2$.

Assume $\mathcal{N}_d(\boldsymbol{x};\bar{\Omega},\boldsymbol{\beta})$ is the PDF of a standard multivariate normal distribution $\mathcal{N}_d(0,\bar{\Omega})$ [41]. $\Phi_{\mathcal{N}}(x)$ is the CDF of a standard normal distribution.

We expand on the construction of multivariate SN distribution in (14.1) and (14.2). And the PDF of $Z \sim L_{\alpha,k}(0,\bar{\Omega},\boldsymbol{\beta})$ is

$$\begin{aligned} L_{\alpha,k}(\boldsymbol{x};\bar{\Omega},\boldsymbol{\beta}) &= \int_0^\infty ds\, \overline{\chi}_{\alpha,k}(s)\, s^d\, \mathcal{N}_d(\boldsymbol{x}s;\bar{\Omega},\boldsymbol{\beta}) \\ &= 2\int_0^\infty ds\, \overline{\chi}_{\alpha,k}(s)\, s^d\, \mathcal{N}_d(\boldsymbol{x}s;\,\bar{\Omega})\, \Phi_{\mathcal{N}}(\boldsymbol{\beta}^\intercal \boldsymbol{x}\, s). \end{aligned} \tag{15.1}$$

The s^d term comes from $\det(s\boldsymbol{I}_d)$ where $\boldsymbol{I}_d$ is the $d \times d$ identity matrix. It is easy to see how it is reduced to a univariate GAS-SN distribution when $d = 1$.

15.1.1. Multivariate Skew-t Distribution. An important bridge between multivariate SN and GAS-SN is the multivariate skew-t distribution. It is documented in Section 6.2 of [1].

It is fully consistent with multivariate GAS-SN by setting $\alpha = 1$. That is, in his notation of skew-t: $ST_d(\Omega,\boldsymbol{\beta},k) \sim L_{1,k}(\Omega,\boldsymbol{\beta})$.

15.2. Location-Scale Family

The location-scale family follows the standard procedure: $Y = \boldsymbol{\xi} + \boldsymbol{\omega}Z$, which is denoted as $Y \sim L_{\alpha,k}(\boldsymbol{\xi},\Omega,\boldsymbol{\beta})$, where $\Omega := \boldsymbol{\omega}^\intercal\bar{\Omega}\boldsymbol{\omega}$ is the covariance matrix, and $\boldsymbol{\omega}$ is a $d \times d$ diagonal scale matrix.

The PDF of Y is

$$L_{\alpha,k}(\boldsymbol{x};\,\boldsymbol{\xi},\Omega,\boldsymbol{\beta}) := \det(\boldsymbol{\omega})^{-1} L_{\alpha,k}(\boldsymbol{z};\bar{\Omega},\boldsymbol{\beta}) \tag{15.2}$$

where $\boldsymbol{z} := \boldsymbol{\omega}^{-1}(\boldsymbol{x}-\boldsymbol{\xi})$. Notice that it has to be computed via the standard PDF.

15.3. Moments

The first moment of Z is $\mu_z := b\,\boldsymbol{\delta}$, where $b := \sqrt{2/\pi}\,\mathbb{E}(X^{-1}|\overline{\chi}_{\alpha,k})$.

The second moment of Z is $m_2\,\bar{\Omega}$, where $m_2 = \mathbb{E}(X^{-2}|\overline{\chi}_{\alpha,k})$. Hence, the covariance is

$$\mathrm{var}\{Z\} := \Sigma_z := m_2\,\bar{\Omega} - b^2\,\boldsymbol{\delta}\boldsymbol{\delta}^\intercal$$

The moments of Y follow the rule of the location-scale family. The first moment of Y is $\boldsymbol{\xi} + \boldsymbol{\omega}\,\mu_z$. The covariance of Y is $\boldsymbol{\omega}\,\Sigma_z\,\boldsymbol{\omega}$.

15.4. Canonical Form

The concept of canonical form in GAS-SN is extended from the multivariate SN in Section 14.6. There exists an affine transformation $Z^* = A_*(Y - \boldsymbol{\xi})$ such that $Z^* \sim L_{\alpha,k}(0, \boldsymbol{I}_d, \boldsymbol{\beta}_{Z*})$, where $\boldsymbol{\beta}_{Z*} = (\beta_*, 0, ..., 0)^\intercal$ and β_* is defined by (14.10). And the algorithm of finding A_* is exactly the same as in Section 14.6.

The variable Z^*, which is called *canonical variable*, comprises d independent components. Only one of them contains the skew component. That is, the PDF of Z^* is

$$L_{\alpha,k_*}(\boldsymbol{x}; \beta_*) = 2\int_0^\infty ds\,\overline{\chi}_{\alpha,k}(s)\,s^d \prod_{i=1}^{d} \mathcal{N}(x_i s)\Phi_{\mathcal{N}}(\beta_* x_1\, s). \tag{15.3}$$

It can be further simplified to an elegant univariate-style integral. When $|\boldsymbol{x}| \neq 0$, let $\beta_*(\boldsymbol{x}) := \beta_* x_1/|\boldsymbol{x}| \in \mathbb{R}$, and

$$L_{\alpha,k_*}(\boldsymbol{x}; \beta_*) = (2\pi)^{-(d-1)/2}\int_0^\infty ds\,\overline{\chi}_{\alpha,k}(s)\,s^d\,\mathcal{N}(|\boldsymbol{x}|s; \beta_*(\boldsymbol{x})). \tag{15.4}$$

When $|\boldsymbol{x}| = 0$, It is simply

$$L_{\alpha,k_*}(0; \beta_*) = (2\pi)^{-d/2}\,\mathbb{E}(X^d|\overline{\chi}_{\alpha,k}), \tag{15.5}$$

independent of β_*.

15.5. Marginal 1D Distribution

The construction of the 1D marginal distribution from Y extends directly from Section 14.7, where $\beta_{1(2)}$ is calculated.

Then the marginal distribution is an univariate GAS-SN: $Y_1 \sim L_{\alpha,k}(\xi_1, \Omega_{11}, \beta_{1(2)})$.

15.6. Quadratic Form

The quadratic form is

$$Q := \frac{1}{d}(Y - \boldsymbol{\xi})^\intercal \Omega^{-1}(Y - \boldsymbol{\xi}) = \frac{1}{d} Z^\intercal \bar{\Omega}^{-1} Z. \tag{15.6}$$

This leads to the fractional extension of the classic F distribution.

Q distributes like a fractional F distribution, $Q \sim F_{\alpha,d,k}$ for all β. The QQ-plot between the empirical data and theoretical values is used to evaluate the goodness of a fit. A perfect fit should produce a 45-degree line.

To prove, from Section 15.1, we have $Z \sim Z_0/V$, $Z_0 \sim SN_d(0, \bar{\Omega}, \boldsymbol{\beta})$, and $V \sim \overline{\chi}_{\alpha,k}$. Put them together,

$$Q = \frac{1}{d} Z^\intercal \bar{\Omega}^{-1} Z = \frac{Z_0^\intercal \bar{\Omega}^{-1} Z_0}{d\,V^2} \sim \left(\frac{X^2}{d}\right)/V^2$$

where $X \sim \mathcal{N}_d(0, \bar{\Omega})$, according to Lemma 14.4.

Since $X^2 \sim \chi^2_d$ and $V^2 \sim \overline{\chi}^2_{\alpha,k}$, this leads to $Q \sim F_{\alpha,d,k}$, according to Section 8.1.

Azzalini (2013) provided a point of validation from his multivariate skew-t distribution. From Section 6.2 of [1], Q of a skew-t variable distributes like the classic $F(d, k)$. This is a special case of our fractional F distribution at $\alpha = 1$. That is, $Q \sim F_{1,d,k}$.

15.7. Multivariate MLE

Maximum likelihood estimation for the bivariate elliptical model can be implemented with stochastic gradient descent (SGD). The objective is the minus-log likelihood (MLLK), computed by summing the negative log density over the observed data. At each iteration, the gradients with respect to the hyperparameters are used to update the parameter vector, with the step size controlled by a learning rate. Small random perturbations may also be added to improve local exploration of the objective surface.

For the bivariate optimization, the parameter space is

$$\Theta = \{\rho, \alpha, k, \beta_1, \beta_2, w_1, w_2, \xi_1, \xi_2\}$$

where $\alpha \in (0, 2)$, $k \in (2, \infty)$, $w_1 > 0$, $w_2 > 0$, and $\beta_1, \beta_2, \xi_1, \xi_2 \in \mathbb{R}$.

Since $\rho \in (-1, 1)$, the optimization uses an unconstrained parameter $\rho_\theta \in \mathbb{R}$ and maps it to the correlation coefficient by $\rho = (2/\pi) \arctan(\rho_\theta)$.

In practice, each dimension of the data set is normalized to unit standard deviation. This keeps the gradient components on comparable scales and improves the stability of the SGD updates.

Let Y denote a data set of size N, and let $L(Y_i; \Theta)$ be the PDF evaluated at observation Y_i. Then

$$\text{MLLK}(\Theta; Y) := -\sum_{i=1}^{N} \log\, L(Y_i; \Theta)$$

$$\text{Gradient}(\Theta) := \left\{ \frac{\partial\, \text{MLLK}(\Theta; Y)}{\partial \Theta_j}, \forall\, \Theta_j \in \Theta \right\}$$

When N is large, evaluating $L(Y_i; \Theta)$ at every observation may be computationally expensive. A histogram approximation can be used to compress the data into a smaller number of bins.

Additional regularization can be added to the MLLK. In the bivariate setting, useful penalties include the L2 distances between the empirical and model-implied skewness and kurtosis of the marginal one-dimensional distributions.

CHAPTER 16

Multivariate GAS-SN Adaptive Distribution (Experimental)

16.1. Definition

The adaptive distribution allows each dimension to have its own shape parameters (α_i, k_i). This is the main departure from the elliptical construction, where all dimensions share the same scalar pair (α, k).

Accordingly, $\boldsymbol{\alpha} = \{\alpha_i\}$ and $\boldsymbol{k} = \{k_i\}$ are d-dimensional parameter vectors. The mixture therefore uses a corresponding collection of standard FCM variables, $\{\overline{\chi}_{\alpha_i,k_i} : i = 1, \ldots, d\}$.

DEFINITION 16.1. Assume that Z_0 is a d-dimensional random variable from the standard multivariate skew-normal distribution $SN_d(0,\bar{\Omega},\boldsymbol{\beta})$, where $\bar{\Omega}$ is a correlation matrix.

Let Z be defined component-wise by the ratio representation $Z_i \sim (Z_0)_i/\overline{\chi}_{\alpha_i,k_i}$. Then $Z \sim \overrightarrow{L}_{\boldsymbol{\alpha},\boldsymbol{k}}(0,\bar{\Omega},\boldsymbol{\beta})$ is a standard multivariate GAS-SN adaptive distribution. The arrow notation emphasizes that the shape parameters $(\boldsymbol{\alpha}, \boldsymbol{k})$ are vectors.

Let $\mathcal{N}(x;\bar{\Omega})$ denote the PDF of the standard multivariate normal distribution $N(0,\bar{\Omega})$ [41], and let $\Phi_{\mathcal{N}}(x)$ denote the CDF of the standard normal distribution.

For $Z \sim \overrightarrow{L}_{\boldsymbol{\alpha},\boldsymbol{k}}(0,\bar{\Omega},\boldsymbol{\beta})$, the PDF is

$$\overrightarrow{L}_{\boldsymbol{\alpha},\boldsymbol{k}}(\boldsymbol{x};\bar{\Omega},\boldsymbol{\beta}) = 2\int\cdots\int_0^\infty \mathcal{N}\left(\boldsymbol{s}\,\boldsymbol{x};\bar{\Omega}\right)\,\Phi_{\mathcal{N}}\Big(\boldsymbol{\beta}^{\intercal}(\boldsymbol{s}\,\boldsymbol{x})\Big)\prod_{i=1}^{d} s_i ds_i\,\overline{\chi}_{\alpha_i,k_i}(s_i). \tag{16.1}$$

where $\boldsymbol{s} := \mathrm{diag}(s_1,\ldots,s_d)$ is the $d \times d$ diagonal matrix generated by the vector $\{s_i\}$. When $d = 1$, this construction reduces to the univariate GAS-SN distribution.

Compared with the elliptical PDF (15.1), the adaptive PDF in (16.1) requires a d-dimensional integral. This makes its evaluation substantially more computationally demanding.

16.2. Location-Scale Family

The location-scale family is obtained by setting $Y = \boldsymbol{\xi} + \boldsymbol{\omega}Z$, denoted by $Y \sim \overrightarrow{L}_{\boldsymbol{\alpha},\boldsymbol{k}}(\boldsymbol{\xi},\Omega,\boldsymbol{\beta})$. Here $\boldsymbol{\omega}$ is the $d \times d$ diagonal scale matrix, and $\Omega = \boldsymbol{\omega}^{\intercal}\bar{\Omega}\boldsymbol{\omega}$ is the corresponding covariance matrix.

The PDF of Y is

$$\overrightarrow{L}_{\boldsymbol{\alpha},\boldsymbol{k}}(\boldsymbol{x};\,\boldsymbol{\xi},\Omega,\boldsymbol{\beta}) := \det(\boldsymbol{\omega})^{-1}\,\overrightarrow{L}_{\boldsymbol{\alpha},\boldsymbol{k}}(\boldsymbol{z};\bar{\Omega},\boldsymbol{\beta}). \tag{16.2}$$

where $\boldsymbol{z} := \boldsymbol{\omega}^{-1}(\boldsymbol{x}-\boldsymbol{\xi})$. The density must be evaluated through the standard PDF because the mixing variables $\{s_i\}$ act on the standardized variable Z, not directly on the location-scale variable Y.

16.3. Moments

The first moment of Z is $\mu_z := \boldsymbol{b} \odot \boldsymbol{\delta}$, where $b_i := \sqrt{2/\pi}\,\mathbb{E}(X^{-1}|\overline{\chi}_{\alpha_i,k_i})$ and $\odot$ is the Hadamard product.

The (i,j) element of the second moment of Z is

$$\boldsymbol{m_2}(i,j) := \begin{cases} \mathbb{E}(X^{-2}|\overline{\chi}_{\alpha_i,k_i}) & \text{if } i=j, \\ \bar{\Omega}_{i,j}\mathbb{E}(X^{-1}|\overline{\chi}_{\alpha_i,k_i})\mathbb{E}(X^{-1}|\overline{\chi}_{\alpha_j,k_j}) & \text{if } i\neq j. \end{cases}$$

where $\bar{\Omega}_{i,i}=1$ is ignored in the first line. Hence, the covariance is

$$\mathrm{var}\{Z\} := \Sigma_z := \boldsymbol{m_2} - \mu_z\,\mu_z^\intercal$$

The moments of Y follow the rule of the location-scale family. The first moment of Y is $\boldsymbol{\xi}+\boldsymbol{\omega}\,\mu_z$. The covariance of Y is $\boldsymbol{\omega}\,\mathrm{var}\{Z\}\,\boldsymbol{\omega}$.

16.4. Canonical Form

The adaptive distribution does not have the rotational symmetry of an elliptical distribution. Consequently, its canonical form is not especially useful, because it is not connected to the other distributions in the family by a simple affine transformation.

Assume that the variable Z^* is a *canonical variable*. Then $Z^* \sim \overrightarrow{L}_{\boldsymbol{\alpha},\boldsymbol{k}}(0,\boldsymbol{I}_d,\boldsymbol{\beta}_{Z*})$, where $\boldsymbol{\beta}_{Z*} = (\beta_*,0,...,0)^\intercal$ and β_* are defined by (14.10).

The PDF of Z^* is

$$\overrightarrow{L}_{\boldsymbol{\alpha},\boldsymbol{k}_*}(\boldsymbol{x};\beta_*) = L_{\alpha_1,k_1}(x_1;\beta_*)\prod_{j=2}^{d} L_{\alpha_j,k_j}(x_j). \tag{16.3}$$

Thus, only the first component has the GAS-SN form, while the remaining components are GSaS components, each with its own shape pair (α,k).

Only the first component of μ_z is nonzero, namely $\sqrt{2/\pi}\,\delta_*\,\mathbb{E}(X^{-1}|\overline{\chi}_{\alpha_1,k_1})$. The matrix $\boldsymbol{m_2}$ is diagonal, with $\boldsymbol{m_2}(i,i) = \mathbb{E}(X^{-2}|\overline{\chi}_{\alpha_i,k_i})$.

16.5. Marginal 1D Distribution

The construction of the 1D marginal distribution from Y extends directly from Section 14.7, where $\beta_{1(2)}$ is calculated.

Then the marginal distribution is an univariate GAS-SN: $Y_1 \sim L_{\alpha_1,k_1}(\xi_1,\Omega_{11},\beta_{1(2)})$.

16.6. 2D Adaptive MLE

Maximum likelihood estimation can be implemented with a stochastic gradient descent (SGD) procedure, but the adaptive model requires several practical adjustments. In the bivariate case, the parameter space is

$$\Theta = \{\rho,\alpha_1,\alpha_2,k_1,k_2,\beta_1,\beta_2,w_1,w_2,\xi_1,\xi_2\}$$

where $\alpha_1,\alpha_2\in(0,2)$, $k_1,k_2\in(2,\infty)$, $w_1>0$, $w_2>0$, and $\beta_1,\beta_2,\xi_1,\xi_2\in\mathbb{R}$. Since $\rho\in(-1,1)$, the optimization uses an unconstrained parameter $\rho_\theta\in\mathbb{R}$ and maps it to the correlation coefficient by $\rho = (2/\pi)\arctan(\rho_\theta)$.

Direct evaluation of the adaptive PDF is computationally expensive, even in two dimensions. For this reason, the MLLK objective is evaluated through the two marginal one-dimensional distributions and supplemented with an L2 regularization term on the correlation coefficient.

In practice, each dimension of the data set is normalized to unit standard deviation. This keeps the gradient components on comparable scales and improves the stability of the SGD updates.

Let Y represent the data set of size N, and $L_m(Y_i;\Theta)$ is the marginal 1D PDF at dimension m $(m = 1...d)$, then

$$\text{MLLK}(\Theta;Y) := -\sum_{i=1}^{N}\sum_{m=1}^{d} \log\, L_m(Y_i;\Theta)$$

$$\text{Gradient}(\Theta) := \left\{ \frac{\partial\, \text{MLLK}(\Theta;Y)}{\partial \Theta_j}, \forall\, \Theta_j \in \Theta \right\}$$

Once the MLLK and gradients are calculated. The program makes a small move along the direction that is most likely to minimize the MLLK. The scale of the move is based on a learning rate, which can be adjusted dynamically. Some randomness can be added to the small move. This allows the algorithm to explore the nearby regions, and maybe get "unstuck" over unexpected edges.

When N is large, it may not be computationally feasible to evaluate every $L(Y_i;\Theta)$. One may use histogram to compress the data into smaller numbers of bins.

More regularization can be added to the MLLK. For instance, we find it makes a lot of sense to add the L2 distances of the skewness and kurtosis on the marginal 1D distributions.

We also regulate the mean of the quadratic form. But the exact distribution of the quadratic form is still under research.

CHAPTER 17

Fitting SPX-VIX Daily Returns with Bivariate Distributions

Two MLE fits are performed for the VIX/SPX daily log returns from 1990 to 2025. The first fit uses the bivariate elliptical GAS-SN distribution. The second fit uses the bivariate adaptive GAS-SN distribution.

The major difference is that the adaptive distribution allows each dimension to have its own (α, k) shape. However, it is much more compute-intensive, it requires alternative methods to work around. And it breaks the rotational symmetry that the elliptical distribution has. This requires a different approach to evaluate the quadratic form.

17.1. Elliptical Fit

The bivariate elliptical MLE program is similar to the univariate MLE program. But the hyper-parameter space is much larger:

$$\Theta = \{\rho, \alpha, k, \beta_1, \beta_2, w_1, w_2, \xi_1, \xi_2\}$$

where $\alpha \in (0, 2)$, $k \in (2, \infty)$, $\omega_1 > 0, \omega_2 > 0$, and $\beta_1, \beta_2, \xi_1, \xi_2 \in \mathbb{R}$.

ρ is the correlation coefficient. Since $\rho \in (-1, 1)$, the optimization uses an unconstrained parameter $\rho_\theta \in \mathbb{R}$ and maps it to the correlation coefficient by $\rho = (2/\pi)\arctan(\rho_\theta)$. In the program, ρ is converted to $\bar{\Omega}$ according to (14.14).

The bivariate MLE program is implemented in **python** and **scipy** on github at https://github.com/slihn/gas-impl/blob/main/gas_impl/mle_gas_sn_2d.py.

We run the MLE fitting on the daily returns of VIX and SPX from 1990 to mid-2025, about 8900 two-column samples. Each column in the data set is normalized to one standard deviation. This allows all gradients to have similar scales and helps the MLE to operate smoothly.

Assume there are N samples in the data set, $Y = \{\boldsymbol{y}_i, i \in 1, 2, ..., N\}$, the minus log-likelihood (MLLK) is

$$\text{MLLK}(\Theta) = -\frac{1}{N}\sum_{i=1}^{N} \log(L_{\alpha,k}(\boldsymbol{y}_i; \boldsymbol{\xi}, \Omega, \boldsymbol{\beta})) \tag{17.1}$$

where $L_{\alpha,k}(\boldsymbol{x}; \boldsymbol{\xi}, \Omega, \boldsymbol{\beta})$ is the multivariate PDF of the location scale family (15.2).

When N is large, it may not be computationally feasible to compute the PDF on every $\boldsymbol{y}_i$. A histogram may be used to compress the data into a grid of bins.

Two components of regularization are added to the objective function $\ell(\Theta)$. The L2 distances between the empirical and theoretical statistics are added as follows:

- Correlation: $|\Delta\rho(Y)|^2$.
- The mean of the quadratic form: $\Delta\mu_Q^2 := |\Delta\text{mean}(Q)|^2$. Section 15.6.

MLE seeks the optimal Θ that minimizes the objective function:

$$\hat{\Theta} = \text{argmin } \ell(\Theta) \tag{17.2}$$

$$\text{where } \ell(\Theta) = \text{MLLK}(\Theta) + |\Delta\rho(Y)|^2 + \Delta\mu_Q^2 \tag{17.3}$$

17.1.1. The VIX-SPX Bivariate Elliptical Fit. Figure 17.1 shows the results of the bivariate elliptical MLE fit on the VIX/SPX daily log returns. The top two graphs show the 2D scatter plot (left) and the contour plot (right) of the samples. Two overlapping lines are drawn to indicate the angles of the correlation, theoretical vs. empirical. The main accomplishment of this fit is that the correlation coefficient matches nicely at about -0.7.

The contour plot is compared to the theoretical elliptical contour plot in the middle left graph. We note that the sample contours look rectangular instead of elliptical. This is an important research topic left for the future.

The remaining three graphs are for the quadratic form Q in (15.6). The PP-plot in the middle right graph and the QQ-plot in log scale in the bottom right graph show very good match with a clear 45-degree line.

However, the QQ-plot in the bottom left graph is less ideal. The tail is tilted upward after 20. This indicates a poor fit on the outside of the contours. This is probably due to the fact that an elliptical distribution could not capture the rectangular nature of the contours.

17.1.2. The Issue in Marginal Distributions. One major issue with the fit is related to the 1-dimensional marginals. The bivariate MLE finds the best fit at $\alpha = 0.75, k = 4.5$. This is a strange place when we examine it in Figure 12.1. When the bivariate distribution is projected to the 1-dimensional marginal distributions according to Section 15.5, the univariate GAS-SN distributions are near the border of infinite kurtosis.

(The reader is reminded that the degrees of freedom need to be higher than 4 to have valid kurtosis in the Student's t distribution. $k = 4.5$ is in the neighborhood of that threshold.)

Although the implied kurtoses are substantially inaccurate, the marginal plots in Figures 17.2 and 17.3 appear visually acceptable except in the upper-right region. There, the theoretical peak of the VIX marginal density is higher than the observed peak, whereas the theoretical peak of the SPX marginal density is lower than the observed peak.

This discrepancy likely reflects the fact that VIX and SPX require different shape parameters (α, k) for their marginal behavior. The current elliptical construction does not allow separate choices of α and k for the two dimensions, and it remains an open problem to incorporate that additional flexibility within the model.

Overall, these results suggest that a single bivariate distribution is too restrictive to describe the full 35-year history of the SPX and VIX data. Further work is needed to obtain a more satisfactory joint model. A major step forward is to apply this distribution in regime-switching models, such as the Hidden Markov Model (HMM), statistical jump model[**38**], and mixture-VAE model[**29**].




FIGURE 17.1. Elliptical MLE fit of VIX/SPX daily log returns from 1990 to 2025 with the bivariate elliptical GAS-SN distribution. Data is standardized to one standard deviation on each axis. The optimal parameters are: $\hat{\Theta} = \{\rho_\theta = -2.12, \alpha = 0.75, k = 4.5, \beta_0 = 0.78, \beta_1 = 0.27, \omega_0 = 0.92, \omega_1 = 0.88, \xi_0 = -0.35, \xi_1 = 0.19\}$.




FIGURE 17.2. VIX Marginal from Elliptical MLE fit of VIX/SPX daily log returns from 1990 to 2025 with the bivariate elliptical GAS-SN distribution. Data is standardized to one standard deviation on each axis.




FIGURE 17.3. SPX Marginal from Elliptical MLE fit of VIX/SPX daily log returns from 1990 to 2025 with the bivariate elliptical GAS-SN distribution. Data is standardized to one standard deviation on each axis.

17.2. Adaptive Fit

The adaptive fit is obtained by maximizing the two marginal likelihoods with an additional regularization term, such as the L^2 distance between the empirical and theoretical correlations. This surrogate objective is used because direct bivariate maximum-likelihood estimation is computationally infeasible on the available hardware.

The resulting contour plot in Figure 17.4 exhibits moderately rectangular level sets, indicating that the adaptive specification captures part of the dependence structure. However, the theoretical correlation reaches only about -0.5, whereas the empirical correlation is closer to -0.7.

TODO need to redo the addaptive fit

Although the adaptive distribution allows each dimension to have its own shape parameters, this additional flexibility does not automatically produce a close bivariate fit. In practice, the correlation parameter interacts strongly with the skew parameters, which makes the joint optimization problem difficult. The remaining discrepancy is therefore mainly a dependence-calibration issue rather than a failure of the univariate SPX marginal fit.

Another practical issue is that the quadratic form requires an additional multiplier, or scale adjustment, to produce a satisfactory fit. The theoretical basis for this adjustment remains unclear. Moreover, the squared QQ plots of the marginals in Figures 17.5 and 17.6 show that the adaptive fit reproduces the tails less accurately than the elliptical fit.




FIGURE 17.4. MLE fit of VIX/SPX daily log returns from 1990 to 2025 with the bivariate adaptive distribution. Data is standardized to one standard deviation on each axis.




FIGURE 17.5. VIX Marginal from MLE fit of VIX/SPX daily log returns from 1990 to 2025 with the bivariate adaptive GAS-SN distribution. Data is standardized to one standard deviation on each axis.




FIGURE 17.6. SPX Marginal from MLE fit of VIX/SPX daily log returns from 1990 to 2025 with the bivariate adaptive GAS-SN distribution. Data is standardized to one standard deviation on each axis.

APPENDIX A

List of Useful Formula

A.1. Gamma Function

Gamma function is used extensively in this paper. First, note that $\Gamma(\frac{1}{2}) = \sqrt{\pi}$. Its **reflection formula** is

$$\Gamma(z)\Gamma(1-z) = \frac{\pi}{\sin(\pi z)} \tag{A.1}$$

And the **Legendre duplication formula** is

$$\Gamma(z)\Gamma(z+1/2) = 2^{1-2z}\sqrt{\pi}\,\Gamma(2z) \tag{A.2}$$

Gamma function Asymptotic: At $x \to 0$, gamma function becomes

$$\begin{aligned} \lim_{x\to 0}\Gamma(x) &\sim \frac{1}{x} \\ \lim_{x\to 0}\frac{\Gamma(ax)}{\Gamma(bx)} &= \frac{b}{a} \qquad (ab \neq 0) \end{aligned} \tag{A.3}$$

For a very large x, assume a, b are finite,

$$\lim_{x\to\infty}\frac{\Gamma(x+a)}{\Gamma(x+b)} \sim x^{a-b} \tag{A.4}$$

Sterling's formula is used to expand the kurtosis formula for a large k, which is:

$$\lim_{x\to\infty}\Gamma(x+1) \sim \sqrt{2\pi x}\Big(\frac{x}{e}\Big)^x, \tag{A.5}$$

$$\text{or } \lim_{x\to\infty}\Gamma(x) \sim \sqrt{2\pi}\,x^{x-1/2}e^{-x}. \tag{A.6}$$

A.2. Transformation

Laplace transform of cosine is[1]

$$\int_0^\infty dt\,\cos(xt)e^{-t/\nu} = \frac{\nu^{-1}}{x^2+\nu^{-2}} = \frac{\nu}{(\nu x)^2+1} \tag{A.7}$$

Gaussian transform of cosine is[2]

$$\begin{aligned} \int_0^\infty dt\,\cos(xt)\,e^{-t^2/2} &= \sqrt{\frac{\pi}{2}}\,e^{-x^2/2} \\ \text{Hence } \int_0^\infty dt\,\cos(xt)\,e^{-t^2/2s^2} &= \sqrt{\frac{\pi}{2}}\,s\,e^{-(sx)^2/2} \end{aligned} \tag{A.8}$$

[1] See https://proofwiki.org/wiki/Laplace_Transform_of_Cosine

[2] See https://www.wolframalpha.com/input?i=integrate+cos%28a+x%29+e%5E%28-x%5E2%2F2%29+dx+from+0+to+infty

A.3. Half-Normal Distribution

The moments of the half-normal distribution (HN)[3] are used several times. Its PDF is defined as

$$p_{HN}(x;\sigma) := \sqrt{\frac{2}{\pi}}\frac{1}{\sigma}\, e^{-x^2/(2\sigma^2)}\,,\ x>0 \tag{A.9}$$

which is a special case of GG with $d=1, p=2, a=\sqrt{2}\sigma$. Its moments are

$$E_{HN}(T^n) = \sigma^n \frac{2^{n/2}}{\sqrt{\pi}}\,\Gamma\left(\frac{n+1}{2}\right) \tag{A.10}$$

which are the same as those of a normal distribution.

[3] See https://en.wikipedia.org/wiki/Half-normal_distribution

Bibliography

[1] Adelchi Azzalini. *The Skew-Normal and Related Families*. Institute of Mathematical Statistics Monographs. Cambridge University Press, 2013.

[2] H. Bateman and A. Erdélyi. *Higher Transcendental Functions*, volume 3 of *California Institute of Technology Bateman Manuscript Project*. McGraw-Hill Book Company., 1955.

[3] Berndt Bruce C. *Ramanujan's Notebooks, Part I*. Springer New York, NY, 1985.

[4] Peter Carr, Hélyette Geman, Dilip B. Madan, and Marc Yor. Self-decomposability and option pricing. *Mathematical Finance*, 17(1):31–57, 2007.

[5] Peter K. Clark. A subordinated stochastic process model with finite variance for speculative prices. *Econometrica*, 41(1):135–155, 1973.

[6] John C. Cox, Jonathan E. Ingersoll, and Stephen A. Ross. A theory of the term structure of interest rates. *Econometrica*, 53(2):385–407, 1985.

[7] J.C. Crisanto-Neto, M.G.E. da Luz, E.P. Raposo, and G.M. Viswanathan. An efficient series approximation for the lévy α-stable symmetric distribution. *Physics Letters A*, 382(35):2408–2413, 2018.

[8] Luc Devroye. Random variate generation for exponentially and polynomially tilted stable distributions. *ACM Trans. Model. Comput. Simul.*, 19(4), November 2009.

[9] Luc Devroye and Lancelot F. James. On simulation and properties of the stable law. *Statistical Methods & Applications*, 23:307 – 343, 2014.

[10] *NIST Digital Library of Mathematical Functions*. https://dlmf.nist.gov/, Release 1.2.0 of 2024-03-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.

[11] William Feller. Two singular diffusion problems. *Annals of Mathematics*, 54(1):173–182, 1951.

[12] William Feller. On a generalization of marcel riesz' potentials and the semi-groups generated by them. *Meddelanden Lunds Universitets Matematiska Seminarium*, pages 73–81, 1952.

[13] William Feller. *An Introduction to Probability Theory and its Applications (2nd Edition)*, volume 2. John Wiley & Sons, Inc., 1971.

[14] Wikimedia Foundation. Inverse distribution, 2023. Retrieved December 1, 2023, from https://en.wikipedia.org/wiki/Inverse_distribution.

[15] Lancelot F. James. Generalized mittag leffler distributions arising as limits in preferential attachment models, 2015.

[16] George Karagiannidis and Athanasios Lioumpas. An improved approximation for the gaussian q-function. *IEEE Communications Letters*, 11:644–646, 08 2007.

[17] Paul Lévy. *Calcul des Probabilités*. Gauthier-Villars, 1925.

[18] Xiaoyue Li and John M. Mulvey. Optimal portfolio execution in a regime-switching market with non-linear impact costs: Combining dynamic program and neural network. *arXiv preprint*, 6 2023. Available at arXiv: https://arxiv.org/abs/2306.08809.

[19] Stephen H.T. Lihn. Stable count distribution for the volatility indices and space-time generalized stable characteristic function. *SSRN Working Paper*, 7 2020. Available at SSRN: https://ssrn.com/abstract=3659383.

[20] Stephen H.T. Lihn. Generalization of the alpha-stable distribution with the degree of freedom. *arXiv preprint*, 5 2024. Available at arXiv: https://arxiv.org/abs/2405.04693.

[21] F. Mainardi. *Fractional Calculus and Waves in Linear Viscoelasticity: An Introduction to Mathematical Models*. Imperial College Press, 2010. See Appendix F at https://appliedmath.brown.edu/sites/default/files/fractional/36%20TheWrightFunctions.pdf.

[22] Francesco Mainardi and Armando Consiglio. The wright functions of the second kind in mathematical physics. *Mathematics*, 8(6), 2020.

[23] Francesco Mainardi, Yuri Luchko, and Gianni Pagnini. The fundamental solution of the space-time fractional diffusion equation. *Frac. Calc. Appl. Anal.*, 4, 03 2007. See https://arxiv.org/abs/cond-mat/0702419.

[24] Francesco Mainardi and Gianni Pagnini. Mellin-barnes integrals for stable distributions and their convolutions. *Fractional Calculus and Applied Analysis*, 11:443–456, 2008.

[25] Till Massing. Parametric estimation of tempered stable laws. *ALEA, Latin American Journal of Probability and Mathematical Statistics*, 21:1567–1600, 2024.

[26] Arak Mathai and Hans Haubold. *Fractional and Multivariable Calculus: Model Building and Optimization Problems*. Springer Cham, 12 2017.

[27] Hjalmar Mellin. Zur theorie zweier allgemeinen klassen bestimmter integrale. *Acta Societatis Scientiarum Fennicæ*, page 1–75, 1897.

[28] The mpmath development team. *mpmath: a Python library for arbitrary-precision floating-point arithmetic (version 1.3.0)*, 2023. `http://mpmath.org/`.

[29] Yuqi Nie, John M. Mulvey, H. Vincent Poor, Chenyu Yu, and Hao Huang. Deep generative models meet statistical methods: A generalized framework for financial regime identification. *SSRN*, 2024. Available at SSRN: https://ssrn.com/abstract=5332057.

[30] John Nolan. *Univariate Stable Distributions, Models for Heavy Tailed Data*. Springer Cham, 01 2020.

[31] Donald B. Owen. Tables for Computing Bivariate Normal Probabilities. *The Annals of Mathematical Statistics*, 27(4):1075 – 1090, 1956.

[32] Richard B. Paris, Armando Consiglio, and Mainardi Francesco. On the asymptotics of wright functions of the second kind. *Fract. Calc. Appl. Anal.*, 24(1):54–72, 01 2021.

[33] Jim Pitman. Combinatorial stochastic processes. In *Ecole d'Eté de Probabilités de Saint-Flour XXXII - 2002*, page 66. Springer Berlin, Heidelberg, 2006.

[34] Jim Pitman and Marc Yor. The two-parameter poisson-dirichlet distribution derived from a stable subordinator. *The Annals of Probability*, 25(2):855–900, 1997.

[35] Dimiter Prodanov. Computation of the wright function from its integral representation. In *Advances in Nonlinear Dynamics*, volume 1, pages 421–431. Springer Nature Switzerland, 05 2024.

[36] Jan Rosiński. Tempering stable processes. *Stochastic Processes and their Applications*, 117(6):677–707, 2007.

[37] W. R. Schneider. Stable distributions: Fox function representation and generalization. In S. Albeverio, G. Casati, and D. Merlini, editors, *Stochastic Processes in Classical and Quantum Systems*, pages 497–511, Berlin, Heidelberg, 1986. Springer Berlin Heidelberg.

[38] Y. Shu, C. Yu, and J.M. Mulvey. Downside risk reduction using regime-switching signals: a statistical jump model approach. *Journal of Asset Management*, 25:493–507, 2024.

[39] E. W. Stacy. A generalization of the gamma distribution. *The Annals of Mathematical Statistics*, 33(3):1187 – 1192, 1962. See also https://en.wikipedia.org/wiki/Generalized_gamma_distribution.

[40] Student. The probable error of a mean. *Biometrika*, 6:1–25, 1908. See also https://en.wikipedia.org/wiki/Student%27s_t-distribution.

[41] Y.L. Tong. *The Multivariate Normal Distribution*. Research in Criminology. Springer New York, 1990.

[42] Pauli Virtanen, Ralf Gommers, Travis E. Oliphant, Matt Haberland, Tyler Reddy, David Cournapeau, Evgeni Burovski, Pearu Peterson, Warren Weckesser, Jonathan Bright, Stéfan J. van der Walt, Matthew Brett, Joshua Wilson, K. Jarrod Millman, Nikolay Mayorov, Andrew R. J. Nelson, Eric Jones, Robert Kern, Eric Larson, C J Carey, İlhan Polat, Yu Feng, Eric W. Moore, Jake VanderPlas, Denis Laxalde, Josef Perktold, Robert Cimrman, Ian Henriksen, E. A. Quintero, Charles R. Harris, Anne M. Archibald, Antônio H. Ribeiro, Fabian Pedregosa, Paul van Mulbregt, and SciPy 1.0 Contributors. SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. *Nature Methods*, 17:261–272, 2020.

[43] E. M. Wright. On the coefficients of power series having exponential singularities. *Journal London Math. Soc.*, s1-8, Issue 1 (1933):71–79, 1933.

[44] E. M. Wright. The asymptotic expansion of the generalized bessel function. *Proc. London Math. Soc.*, s2-38, Issue 1 (1935):257–270, 1935.