

1 **Introduction to the Fractional Distribution Families**

2 Stephen Horng-Twu Lihn

3 stevelihn@gmail.com

4 Version: Draft 2026-08

5 ORFE, Princeton University

6 Dedicated to Professor John M. Mulvey for his 80th birthday in 2026.

7 (as of October 4, 2026)

Contents

10	Chapter 1. Introduction	1
11	1.1. Motivation	1
12	1.2. Elements of Probability Distributions	2
13	1.3. Organization of This Book	3
14	1.4. Unresolved Issues	4
15	Part 1. Mathematical Functions	5
16	Chapter 2. Mellin Transform	7
17	2.1. One-Sided Distribution	8
18	2.2. Two-Sided Distribution	9
19	2.3. Ramanujan's Master Theorem	10
20	Chapter 3. The Wright Function and The Inverse Stable Law	13
21	3.1. Definition	13
22	3.2. Classic Results	13
23	3.3. The M-Wright Functions	14
24	3.4. The Inverse Stable Law	15
25	3.5. The Fractional Gamma-Star Function	16
26	3.6. The Elasticity Operator	17
27	3.7. The Elasticity of the M-Wright Functions	18
28	3.8. Numerical Methods of the M-Wright Functions	20
29	Chapter 4. The Alpha-Stable Distribution - Review	23
30	4.1. Classic Result	23
31	4.2. Extremal Distributions	24
32	4.3. Ratio Distribution Approach	24
33	4.4. SaS	26
34	Chapter 5. Fractional Hypergeometric Functions	27
35	5.1. Fractional Confluent Hypergeometric Function	27
36	5.2. Fractional Gauss Hypergeometric Function	29
37	Part 2. One-Sided Distributions	33
38	Chapter 6. FG: Fractional Gamma Distribution and Tilted Stable Law	35
39	6.1. Definition	35
40	6.2. Determination of C	35
41	6.3. FG Subsumes Generalized Gamma Distribution	36
42	6.4. Mellin Transform	36
43	6.5. The Inverse Tilted Stable Law	36

44	6.6. CDF and Fractional Incomplete Gamma Function	38
45	6.7. Inverse Expression of Several Fractional Distributions	39
46	6.8. Alternative Definition	40
47	Chapter 7. Fractional Chi Distributions	41
48	7.1. Introduction to Fractional Chi Distribution	41
49	7.2. FCM: Fractional Chi-Mean Distribution	41
50	7.3. FCM Moments	42
51	7.4. FCM Reflection Formula and Negative k	43
52	7.5. FCM: Lognormal Asymptotics at Infinite Degrees of Freedom	43
53	7.6. FCM2: Fractional Chi-Squared-Mean Distribution	45
54	7.7. FCM2 Mellin Transform	46
55	7.8. FCM2 Moments	46
56	7.9. FCM2 Increment of k	47
57	7.10. Sum of Two Chi-Squares with Correlation	47
58	Chapter 8. Fractional F Distribution	49
59	8.1. Definition	49
60	8.2. PDF at Zero	50
61	8.3. Mellin Transform	50
62	8.4. Moments	50
63	Chapter 9. The Framework of Continuous Gaussian Mixture	53
64	9.1. The Inverse Chi Distribution	53
65	9.2. The Characteristic Distributions	54
66	9.3. Summary of Gaussian Mixture	55
67	9.4. FCM Extensions	55
68	9.5. Application of FCM Variants	57
69	Part 3. Two-Sided Univariate Distributions	61
70	Chapter 10. SN: The Skew-Normal Distribution - Review	63
71	10.1. Definition	63
72	10.2. The Location-Scale Family	64
73	10.3. Invariant Quantities	64
74	10.4. Mellin Transform	65
75	10.5. Moments	66
76	Chapter 11. GAS: Generalized Alpha-Stable Distribution (Experimental)	69
77	11.1. Definition	69
78	11.2. Limitation	70
79	11.3. Workaround	71
80	11.4. Moments	73
81	Chapter 12. GAS-SN: Generalized Alpha-Stable Distribution with Skew-Normal	75
82	12.1. Definition	75
83	12.2. The Location-Scale Family	76
84	12.3. Mellin Transform	76
85	12.4. Moments	76
86	12.5. Tail Behavior	81
87	12.6. Maximum Skewness and Half GSaS	83

88	12.7. Fractional Skew Exponential Power Distribution	85
89	12.8. Quadratic Form	85
90	12.9. Univariate MLE	85
91	12.10. Examples of Univariate MLE Fits	86
92	Chapter 13. Fractional Feller Square-Root Process	89
93	13.1. Generation of Random Variables for FCM	90
94	13.2. Generation of Random Variables for FCM2	91
95	Part 4. Multivariate Distributions	93
96	Chapter 14. Multivariate SN Distribution - Review	95
97	14.1. Definition	95
98	14.2. The Location-Scale Family	96
99	14.3. Quadratic Form	96
100	14.4. Stochastic Representation	96
101	14.5. Moments	97
102	14.6. Canonical Form	98
103	14.7. 1D Marginal Distribution	98
104	Chapter 15. Multivariate GAS-SN Elliptical Distribution	101
105	15.1. Definition	101
106	15.2. Location-Scale Family	101
107	15.3. Moments	101
108	15.4. Canonical Form	102
109	15.5. Marginal 1D Distribution	102
110	15.6. Quadratic Form	102
111	15.7. Multivariate MLE	103
112	Chapter 16. Multivariate GAS-SN Adaptive Distribution (Experimental)	105
113	16.1. Definition	105
114	16.2. Location-Scale Family	105
115	16.3. Moments	105
116	16.4. Canonical Form	106
117	16.5. Marginal 1D Distribution	106
118	16.6. 2D Adaptive MLE	106
119	Chapter 17. Fitting SPX-VIX Daily Returns with Bivariate Distributions	109
120	17.1. Elliptical Fit	109
121	17.2. Adaptive Fit	114
122	Appendix A. List of Useful Formula	119
123	A.1. Gamma Function	119
124	A.2. Transformation	119
125	A.3. Half-Normal Distribution	120
126	Bibliography	121

CHAPTER 1

Introduction

1.1. Motivation

In quantitative finance, asset return data often exhibit substantial skewness and kurtosis. In portfolio optimization and market-regime modeling[17, 35, 27], a standard example is the joint behavior of the S&P 500 Index (SPX) and the CBOE Volatility Index (VIX), whose daily prices have been publicly available since 1990¹. Although such data sets are readily available, fitting them with existing parametric distributions remains difficult. Even with the many probability distributions implemented in modern statistical software, such as **scipy.stats**, the fit is often inadequate.

This book develops a multivariate elliptical distribution system based on the Wright function[40, 41, 2]. As shown in the next section, this subordinator approach is identical to the tilted stable law[7]. The construction combines and extends the α -stable distribution[16] and the multivariate skew-t distribution[1]. The resulting family is intended to model real-world data sets with pronounced tail behavior with more accuracy.

For example, the daily return distribution of VIX has kurtosis about 16 and skewness about 2.0. Its standardized peak density is approximately 0.55 (see Figure 12.7). For comparison, the excess kurtosis of the t distribution[37] is $6/(k-4)$ for $k > 4$, so matching this level of kurtosis would require k to be very close to 4. However, the corresponding standardized peak density is only 0.53 at $k = 4$. Thus, the VIX data lie near the boundary of what the t distribution can represent.

The daily return distribution of SPX is even more challenging (see Figure 12.8). In addition to kurtosis of about 11, its standardized peak density is approximately 0.65. A reasonable fit requires a t distribution with about 3 degrees of freedom ($k \approx 3$). However, from a theoretical standpoint, finite kurtosis does not exist for the t distribution until $k > 4$.

These two examples illustrate a basic difficulty in fitting existing parametric distributions. In particular, it is difficult to match both the kurtosis and the peak density simultaneously.

Our new multivariate distribution is designed to fit both data sets with satisfactory accuracy while matching empirical skewness, kurtosis, and peak density. Goodness of fit is evaluated not only through the density function but also through the extent to which the distribution captures tail behavior via the quadratic form. These fits are presented in Chapter 17.

¹SPX data: Courtesy of S&P Dow Jones Indices LLC, from <https://fred.stlouisfed.org/series/SP500>. VIX data: Courtesy of Chicago Board Options Exchange (CBOE), from <https://fred.stlouisfed.org/series/VIXCLS>. Retrieved from FRED, Federal Reserve Bank of St. Louis. Not for commercial use.

1.2. Elements of Probability Distributions

The word "fractional" can be understood as adding the Lévy stability index α to a known classic distribution. Depending on the context, α could be in $[0, 1]$ or $[0, 2]$.

For example, in the Mellin transform of the PDF of a distribution, $\Gamma(s + c)$ in the classic world becomes $\Gamma(\alpha s + c)$ or $\Gamma(s/\alpha + c)$ in the fractional world. When the coefficient of s is $\frac{1}{2}$, 1, or 2, the fractional distribution subsumes the classic distribution, since the Legendre duplication formula (A.2) becomes applicable. But where do these fractional $\Gamma(\alpha \dots)$ functions come from?

The most fundamental and elegant representation is based on the **Tilted Stable law** (TS). The tilted stable law emerged from the study of the Poisson-Dirichlet distribution by Pitman and Yor[32]. In the limit of infinite samplings, the normalized distribution of the size of the cluster approaches the tilted stable law: $T_{\alpha, \beta}^{-\alpha} = (T_{\alpha, \beta})^{-\alpha}$, where α is the stability index in $[0, 1]$ and $\beta \geq 0$ is the power of the polynomial tilt. T_α is the stable law without tilt $T_{\alpha, 0}$.

Take the Chinese restaurant process as an example (Section 3 of [31]). Let K_n be the number of tables occupied after n customers, and n_j be customers already in table j . The Pitman-Yor probability for the next customer to join the table j or to open a new table is

$$\begin{aligned} \Pr(\text{join table } j) &= \frac{n_j - \alpha}{\beta + n}, \\ \Pr(\text{new table} \mid K_n) &= \frac{\beta + \alpha K_n}{\beta + n}. \end{aligned}$$

Its α -diversity limit is $\frac{K_n}{n^\alpha} \rightarrow T_{\alpha, \beta}^{-\alpha}$. Hence, α plays a discount role as the larger α discounts the attractiveness of joining the existing tables. On the other hand, β is a baseline strength for creating a new table, independent of n and K_n .

In [7] and [8], the superscript is generalized to form a three-parameter distribution family $T_{\alpha, \beta}^{-\gamma}$ where $\gamma \in \mathbb{R}$. To avoid collisions of the greeks, we also use the text-based notation: $TS(\alpha, \beta, \gamma)$. When $\gamma > 0$, it represents a local time process. When $\gamma < 0$, it represents a volatility process.

This three-parameter distribution law is identical to our fractional gamma distribution with a different parameterization. In Table 1, we list all the one-sided distributions in this book that have direct mapping to $T_{\alpha, \beta}^{-\gamma}$.

Fractional Distribution	Section	Notation	$T_{\alpha, \beta}^{-\gamma}$ or $TS(\alpha, \beta, \gamma)$			
			α	β	γ	scale
One-sided stable	4.2	L_α (aka T_α)	α	0	-1	1
M-Wright (or Mittag-Leffler)	3.4	M_α (aka $T_\alpha^{-\alpha}$)	α	0	α	1
Fractional gamma	6.5	$\mathfrak{N}_\alpha(\sigma, d, p)$	α	$\alpha d/p$	α/p	σ
Fractional chi (FCM)	7.2.1	$\bar{\chi}_{\alpha, k}(k > 0)$	$\alpha/2$	$(k-1)/2$	1/2	$\sigma_{\alpha, k}$
Inverse FCM	9.4	$\bar{\chi}_{\alpha, k}^\dagger$	$\alpha/2$	$(k-1)/2$	-1/2	$\sigma_{\alpha, k}^{-1}$
Characteristic FCM	7.4.2	$\bar{\chi}_{\alpha, -k}$	$\alpha/2$	$k/2$	-1/2	$\sigma_{\alpha, k}^{-1}$
Fractional chi-squared (FCM2)	7.6.1	$\bar{\chi}_{\alpha, k}^2$	$\alpha/2$	$(k-1)/2$	1	$\sigma_{\alpha, k}^2$
Characteristic FCM2	7.6.1	$\bar{\chi}_{\alpha, -k}^2$	$\alpha/2$	$k/2$	-1	$\sigma_{\alpha, k}^{-2}$

TABLE 1. The tilted stable law for the fractional distributions in this book. k is the "degree of freedom" parameter as in Student's t and χ^2 .

Therefore, $T_{\alpha,\beta}$ is seen as **the atom** of all one-sided distributions. The polynomial tilt is expressed in its density function: $\frac{\Gamma(1+\beta)}{\Gamma(1+\beta/\alpha)} x^{-\beta} L_{\alpha}(x)$ ($x > 0$). [31, 7] And importantly, $\bar{\chi}_{\alpha,-k}^2$ is directly linked to $T_{\alpha/2,k/2}$ according to the last row of the table.

In addition to Table 1, two fractional distributions are formed by a ratio of two tilted stable laws.

- (1) Fractional F distribution in Section 8.1.2: It is $F_{\alpha,d,k} \sim \bar{\chi}_{1,d}^2 / \bar{\chi}_{\alpha,k}^2$
- (2) The α -stable distribution in Section 4.3.2: We have

$$L_{\alpha}^{\theta} \sim M_{\gamma} / T_{\gamma\alpha,0}^{-\gamma} = (L_{\gamma\alpha} / L_{\gamma})^{\gamma}, \text{ where } \gamma = \frac{\alpha - \theta}{2\alpha}, \alpha \in (0, 2].$$

This is for the γ portion of the population in $x > 0$. And (α, θ) is in Feller's parametrization (4.1) [11, 12]. (A similar formula was derived in [8].)

To construct the two-sided distributions, we need the **second atom, Selective Sampling**, to produce the skewness [1].

We take the well-known *multivariate normal distribution* $\mathcal{N}_d(0, \bar{\Omega})$, where d is the dimension, and $\bar{\Omega}$ is a $d \times d$ correlation matrix. The selective sampling works as follows (Section 14.4).

Assuming $X_0 \sim \mathcal{N}_d(0, \bar{\Omega})$, $X_1 \sim \mathcal{N}(0, 1)$, and $\beta \in \mathbb{R}^d$ is the skew parameter, they form the skew-normal distribution $Z \sim SN_d(0, \bar{\Omega}, \beta)$ such that (Section 2.1 of [1])

$$Z = \begin{cases} X_0 & \text{if } X_1 > \beta^{\top} X_0, \\ -X_0 & \text{otherwise.} \end{cases}$$

A two-sided stable distribution is built by properly assembling the ratio $Z/TS(\dots)$ or the product $Z \times TS(\dots)$.

The crown jewel is the skew multivariate elliptical distribution, based on the ratio:

$$SN_d(0, \bar{\Omega}, \beta) / \bar{\chi}_{\alpha,k} \quad (\text{Definition 15.1})$$

The most important chapters of the book are

- Chapter 12 on the univariate GAS-SN distribution; and
- Chapter 15 on the multivariate GAS-SN elliptical distribution.

The reader can think of the entire book as aiming at developing tools (atoms) to create these two distributions.

Our univariate GAS-SN distribution is supposed to be the most flexible two-sided distribution up to date for statisticians to fit a univariate data set, such as return distributions in finance.

Our multivariate GAS-SN elliptical distribution is intended to be the most flexible multivariate distribution to date that extends the multivariate skew-t and skew-normal distributions [1].

A reference implementation in **python** can be found on Github at: <https://github.com/slihn/gas-impl>. This provides a baseline if these distributions make their way to standard libraries, such as **scipy.stats** in the future.

1.3. Organization of This Book

This book is divided into three parts.

Part I describes the mathematical foundation needed for the construction of fractional distributions. It contains several higher transcendental functions. Several classic special functions are extended with a fractional parameter.

Each distribution has its density function (PDF) and distribution function (CDF). Its Mellin transform. The squared variable or quadratic forms. Therefore, new mathematical tools are needed to address them.

Part II contains the univariate one-sided fractional distributions that are invented. All of them have their classic counterparts. For example, the generalized gamma distribution (GG) is upgraded. All the χ and F related distributions are also upgraded.

Part III contains the two-sided univariate fractional distributions. The Azzalini (2013) book is used as the blueprint[1]. It is integrated with the symmetric distributions developed in my 2024 work[19].

This book can be viewed as an integration between the two works, literally going chapter-by-chapter. The consistency of such integration and harmony speaks volumes.

The fourth part introduces the multivariate fractional distributions. These families extend the univariate constructions in Part III and subsume the skew-normal and skew- t families discussed in Azzalini's book.

Their main advantage is the ability to represent a wide range of skewness, kurtosis, and peak density within a unified framework. This flexibility makes them suitable for data sets whose distributional features cannot be adequately captured by more classical parametric models.

Modern data sets are often large in both sample size and dimension, making tail behavior increasingly important to model accurately. In finance, this is especially true for the left tail, where risk is concentrated.

An adaptive version of the multivariate distribution is therefore developed so that each dimension may carry its own shape parameters. This construction is used to fit one of the most challenging financial data sets considered in this book: daily SPX and VIX returns since 1990. The corresponding methodology is presented in Part IV.

1.4. Unresolved Issues

Although the two multivariate distributions present new opportunities to fit the data sets that were thought impossible previously, the outcomes pose new challenges.

On the one hand, maximum-likelihood estimation (MLE) for the elliptical distribution is relatively straightforward to implement. The resulting fit (Figures 17.1, 17.2, 17.3) appears satisfactory at the visual level. However, the fitted values of (α, k) lie near a region of effectively infinite kurtosis when the bivariate distribution is projected onto its two one-dimensional marginals. This tension between visual fit and implied marginal behavior remains an important unresolved issue.

On the other hand, the adaptive distribution suffers from the curse of dimensionality, which makes direct MLE computationally prohibitive. Therefore, a modified fitting algorithm is used. The resulting fit (Figures 17.4, 17.5, 17.6) is qualitatively reasonable, but several limitations remain. In particular, the theoretical correlation coefficient remains smaller in absolute value than the empirical value of about -0.7 , and the quadratic form does not yet have a matching F distribution.

The broader mathematical and statistical questions raised by these constructions remain open.

Part 1

Mathematical Functions

Mellin Transform

255

256 This chapter begins with the Mellin-transform tools used throughout the book. Readers who
257 are mainly interested in the statistical models may treat this and the following chapter as technical
258 background, but the notation introduced here is used repeatedly in the later constructions.

259 The Mellin transform, introduced by Hjalmar Mellin in 1897 [25], is a central tool for analyzing
260 one-sided probability distributions. Once the Mellin transform of a density is known, the moments and
261 several derivative relations can often be obtained directly. This makes the transform especially useful
262 for the Wright function, the α -stable distribution, and the fractional χ distribution, whose connections
263 are most naturally expressed in Mellin-transform form.

264

265 DEFINITION 2.1. This chapter provides an overview of the Mellin transform. Following the notation
266 of [23], the Mellin transform of a function $f(x)$ properly defined for $x \geq 0$ is

$$(2.1) \quad f^*(s) := \mathcal{M}\{f(x); s\} = \int_0^\infty f(x) x^{s-1} dx, \quad c_1 < \Re(s) < c_2.$$

267 The role of c_1, c_2 will be explained in the following.

268 If $f^*(s)$ has analytic continuation on the complex plane, the inverse Mellin transform is

$$(2.2) \quad f(x) = \mathcal{M}^{-1}\{f^*(s); x\} = \frac{1}{2\pi i} \int_{C-i\infty}^{C+i\infty} f^*(s) x^{-s} ds, \quad c_1 < C < c_2.$$

269 Equation (2.1) shows the direct connection between the Mellin transform and distributional mo-
270 ments. If $f(x)$ is the PDF of a one-sided distribution, then its n -th moment is $\mathbb{E}(X^n|f) = f^*(n+1)$.
271 Thus, modifying the Mellin transform $f^*(s)$ provides a systematic way to construct new distributions
272 from existing ones.

273 For compactness, we use the juxtaposition notation $\xleftrightarrow{\mathcal{M}}$ and write $f(x) \xleftrightarrow{\mathcal{M}} f^*(s)$, with the validity
274 strip inherited from (2.1) and (2.2). The precise choice of the contour parameter C is needed for
275 numerical inversion in (2.2), but otherwise the notation mainly serves to keep the transform identities
276 concise.

277 LEMMA 2.2. The main rules of Mellin transform used in this paper are:

$$(2.3) \quad f(ax) \xleftrightarrow{\mathcal{M}} a^{-s} f^*(s), \quad a > 0$$

$$(2.4) \quad x^k f(x) \xleftrightarrow{\mathcal{M}} f^*(s+k),$$

$$(2.5) \quad f(x^p) \xleftrightarrow{\mathcal{M}} \frac{1}{p} f^*(s/p), \quad p \neq 0$$

and the following ones involving an integral,

$$(2.6) \quad h(x) = \int_0^\infty f(xs)g(s) s ds \xleftrightarrow{\mathcal{M}} h^*(s) = f^*(s)g^*(2-s), \quad (\text{ratio distribution})$$

$$(2.7) \quad \gamma_f(x) = \int_0^x f(x) dx \xleftrightarrow{\mathcal{M}} -s^{-1}f^*(s+1), \quad (\text{lower incomplete function})$$

$$(2.8) \quad \Gamma_f(x) = \int_x^\infty f(x) dx \xleftrightarrow{\mathcal{M}} s^{-1}f^*(s+1). \quad (\text{upper incomplete function})$$

The ratio distribution rule (2.6) is widely used in our fractional distribution system. Notice that the argument of $g^*(s)$ is transformed via $s \rightarrow 2-s$.

For (2.7) and (2.8), the valid range of C is decremented by one: $c_1 - 1 < C < c_2 - 1$.

△

EXAMPLE 2.3. A simple exercise is the Mellin transform of the standard normal distribution. It starts with

$$e^{-x} \xleftrightarrow{\mathcal{M}} \Gamma(s)$$

via the definition of the gamma function itself.

By applying (2.5) then (2.3), we get

$$(2.9) \quad \mathcal{N}(x) \xleftrightarrow{\mathcal{M}} \mathcal{N}^*(s) = \frac{2^{(s-3)/2}}{\sqrt{\pi}} \Gamma\left(\frac{s}{2}\right)$$

where $\mathcal{N}(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$ is our notation for the PDF of a standard normal distribution.

EXAMPLE 2.4. A slightly more complicated exercise is the Mellin transform of the PDF of the fractional gamma distribution (FG) in Chapter 6. But we only work out its skeleton here.

Assume we have a function $F_\alpha(x)$ whose Mellin transform is

$$F_\alpha(x) \xleftrightarrow{\mathcal{M}} \frac{\Gamma(s)}{\Gamma(\alpha s)}.$$

It undergoes the following transforms:

$$\begin{aligned} F_\alpha(x^p) &\xleftrightarrow{\mathcal{M}} \frac{1}{|p|} \frac{\Gamma(s/p)}{\Gamma(\alpha s/p)}, \\ x^{d-1} F_\alpha(x^p) &\xleftrightarrow{\mathcal{M}} \frac{1}{|p|} \frac{\Gamma((s+d-1)/p)}{\Gamma(\alpha(s+d-1)/p)}, \end{aligned}$$

which is the prototype of FG before further normalization.

2.1. One-Sided Distribution

When the subject matter is a probability distribution, the two rules of incomplete functions, (2.7) and (2.8), provide a clear path to obtain its distribution function (CDF) while $f(x)$ is associated with its density function (PDF).

For a one-sided distribution denoted f , it is straightforward to define it, since the domain of the distribution $x \geq 0$ is exactly the same as the domain of the Mellin transform. Hence, $f(x)$ is exactly its density function and $\gamma_f(x)$ is its CDF.

By assigning $s = n + 1$, its n -th moment is

$$(2.10) \quad \mathbb{E}(X^n | f) = f^*(n+1)$$

The zeroth moment is the total density that should be normalized to one. That is $f^*(1) = 1$.

The mean of the distribution is $\mathbb{E}(X | f) = f^*(2)$. For a distribution whose mean needs to be determined, for instance, the fractional χ distribution in Chapter 7, this formula is useful.

2.2. Two-Sided Distribution

When the distribution is two-sided, one more rule is needed.

DEFINITION 2.5 (The Reflection Rule). Assume the distribution is two sided, its domain of support is $x \in \mathbb{R}$. The reflection rule requires that its density function $f(x)$ is based on a skew parameter $\beta \in \mathbb{R}$ that satisfies

$$(2.11) \quad f(-x; \beta) := f(x; -\beta) \quad \text{for } x > 0.$$

Let $c_\beta < 1$ be the one-sided integral of

$$c_\beta = \int_0^\infty f(x; \beta) dx.$$

(2.11) leads to $c_{-\beta} + c_\beta = 1$ since $\int_{-\infty}^\infty f(x; \beta) dx = 1$.

2.2.1. Mellin Transform of a Two-sided CDF.

LEMMA 2.6. The Mellin transform of the CDF $\Phi(x)$ of a two-sided distribution has two parts. Both can be derived from its density function transform, $f(x; \beta) \xleftrightarrow{\mathcal{M}} f^*(s; \beta)$, in the positive domain.

From (2.7), let $\gamma_f(x; \beta) \xleftrightarrow{\mathcal{M}} \Phi^*(s; \beta) := -s^{-1} f^*(s+1; \beta)$. Then for $x > 0$, the Mellin transform of the CDF can be expressed as

$$\begin{aligned} \Phi(x) - \Phi(0) &\xleftrightarrow{\mathcal{M}} \Phi^*(s; \beta), \\ 1 - \Phi(0) - \Phi(-x) &\xleftrightarrow{\mathcal{M}} \Phi^*(s; -\beta). \end{aligned}$$

△

PROOF. Note that $\Phi(0) = c_{-\beta} = 1 - c_\beta$. When $x \geq 0$, its CDF is

$$\Phi(x) = \int_{-\infty}^x f(x; \beta) dx = c_{-\beta} + \int_0^x f(x; \beta) dx = \Phi(0) + \gamma_f(x; \beta).$$

In the negative domain, its CDF is

$$\begin{aligned} \Phi(-x) &= \int_{-\infty}^{-x} f(x; \beta) dx = \int_x^\infty f(x; -\beta) dx \\ &= 1 - \Phi(0) - \int_0^x f(x; -\beta) dx = 1 - \Phi(0) - \gamma_f(x; -\beta). \end{aligned}$$

□

The point is that, once the Mellin transform of either the PDF or CDF is known, the other one can be derived by simple algebraic rules.

2.2.2. From Mellin Transform to Moments. By assigning $s = n + 1$, it is easy to show that its n -th moment is

$$(2.12) \quad \mathbb{E}(X^n | f) = f^*(n+1; \beta) + (-1)^n f^*(n+1; -\beta)$$

$$(2.13) \quad = -n [\Phi^*(n; \beta) + (-1)^n \Phi^*(n; -\beta)]$$

The moment formula is tightly linked to $\Phi^*(n; \beta)$.

The Zeroth Moment. The total density can be regarded as the zeroth moment. Hence, c_β can be determined by

$$(2.14) \quad c_\beta = \int_0^\infty f(x; \beta) dx = f^*(1; \beta).$$

Its application is in (10.9).

2.3. Ramanujan's Master Theorem

In order to keep things simple, we describe all the distributions via the Mellin transform of their PDFs. Due to Ramanujan's master theorem[3], not only can the moments be obtained from the Mellin transform but also all the derivatives of the PDF at $x = 0$. We get its series representation "for free", so to speak.

LEMMA 2.7 (Ramanujan's master theorem). If $f(x)$ has an expansion of the form

$$(2.15) \quad f(x) = \sum_{n=0}^{\infty} \frac{\varphi(n)}{n!} (-x)^n$$

then its Mellin transform is given by

$$(2.16) \quad f(x) \xleftrightarrow{\mathcal{M}} f^*(s) = \Gamma(s) \varphi(-s)$$

△

Assume that $g^*(s) := f^*(s)/\Gamma(s)$ exists on the complex plane, $s \in \mathbb{C}$. Its connection to the derivatives of the PDF at $x = 0$ is as follow.

LEMMA 2.8. The Taylor series of $f(x)$ is

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n$$

where $f^{(n)}(0)$ is the n -th derivative of $f(x)$ at $x = 0$.

Then $f^{(n)}(0)$ can be obtained from $g^*(s)$ by

$$(2.17) \quad f^{(n)}(0) = (-1)^n g^*(-n)$$

At $x = 0$, we have $f(0) = g^*(0)$.

△

The power of the master theorem is that, once the Mellin transform is known, the Taylor series is also known immediately. We provide a contrived example from next chapter as a showcase.

EXAMPLE 2.9. The Mellin transform of the Wright function from (3.5) is $f(-x) \xleftrightarrow{\mathcal{M}} f^*(s) = \Gamma(s)/\Gamma(\delta - \lambda s)$. Then its $g^*(s) = 1/\Gamma(\delta - \lambda s)$.

According to Lemma 2.8, its Taylor series should be

$$f(-x) := \sum_{n=0}^{\infty} \frac{(-1)^n g^*(-n)}{n!} x^n = \sum_{n=0}^{\infty} \frac{g^*(-n)}{n!} (-x)^n$$

Replace $-x$ with z , and plug in $g^*(-n)$, we have

$$f(z) := \sum_{n=0}^{\infty} \frac{z^n}{n! \Gamma(\lambda n + \delta)}$$

This is the series representation (3.1) where we essentially "derived" it from the master theorem.

The major application in this book is in Chapter 11. In the experimental construction of the generalized α -stable distribution, the theorem is used to remedy the discontinuity of the PDF in $x = 0$.

354 **2.3.1. Distribution Function.** The form of the Mellin transform in (2.16) has an important
 355 implication when $f(x)$ is a density function.

356 LEMMA 2.10. Assume $x > 0$, its complimentary distribution function $\Gamma_f(x) := \int_x^\infty f(x) dx$ has
 357 the series representation of

$$(2.18) \quad \Gamma_f(x) = \sum_{n=0}^{\infty} \frac{\varphi(n-1)}{n!} (-x)^n$$

358

△

359 PROOF. From (2.8), the Mellin transform of $\Gamma_f(x)$ is

$$\Gamma_f(x) = \int_x^\infty f(x) dx \xleftrightarrow{\mathcal{M}} s^{-1} f^*(s+1)$$

360 which can be simplified to

$$\begin{aligned} s^{-1} f^*(s+1) &= s^{-1} \Gamma(s+1) \varphi(-s-1) \\ &= \Gamma(s) \varphi(-s-1). \end{aligned}$$

361 This is still in the form of (2.16), with a transformation rule of $s \rightarrow s+1$ in the function $\varphi(-s)$.

362 Applying the master theorem of (2.15), we get (2.18).

363

□

364 We use the CDF of the M-Wright function from (3.17) as an example.

365 LEMMA 2.11. The goal is to show

$$(2.19) \quad \int_x^\infty M_\alpha(t) dt = W_{-\alpha,1}(-x).$$

366

△

367 PROOF. We start with the Mellin transform of $M_\alpha(x)$ from (3.13),

$$M_\alpha(x) \xleftrightarrow{\mathcal{M}} M_\alpha^*(s) = \frac{\Gamma(s)}{\Gamma((1-\alpha) + \alpha s)}$$

368 which yields $\varphi(-s) = 1/\Gamma((1-\alpha) + \alpha s)$.

369 Therefore, its $\Gamma_f(x)$ should be

$$\Gamma_f(x) = \sum_{n=0}^{\infty} \frac{1}{n! \Gamma((1-\alpha) - \alpha(n-1))} (-x)^n = \sum_{n=0}^{\infty} \frac{1}{n! \Gamma(-\alpha n + 1)} (-x)^n$$

370 which is $W_{-\alpha,1}(-x)$ according to (3.1).

371

□

CHAPTER 3

The Wright Function and The Inverse Stable Law

3.1. Definition

The Wright function is a basic building block in the fractional-distribution system developed in this book. It was introduced by E. M. Wright in the 1930s[40, 41], and Bateman later catalogued it together with the Mittag-Leffler function[2].

Its importance became much clearer in the later literature, especially through the work of F. Mainardi, who introduced the M-Wright function $M_\alpha(x)$. In this framework, $M_\alpha(x)$ plays the role of a fractional analogue of the exponential function e^{-x} . This chapter collects the main facts about the Wright function needed in the remainder of the book.

DEFINITION 3.1. The series representation of the Wright function is

$$(3.1) \quad W_{\lambda,\delta}(z) := \sum_{n=0}^{\infty} \frac{z^n}{n! \Gamma(\lambda n + \delta)} \quad (\lambda \geq -1, z \in \mathbb{C})$$

Its shape parameters are pairs (λ, δ) . The apparent limit is $W_{0,1}(z) = e^z$.

Four variants of the Wright function are used repeatedly in this book. The first pair is

- $M_\alpha(z) := W_{-\alpha,1-\alpha}(-z)$
- $F_\alpha(z) := W_{-\alpha,0}(-z)$

with $\alpha \in [0, 1]$. They satisfy $M_\alpha(z) = F_\alpha(z)/(\alpha z)$.

The function $M_\alpha(z)$ is known as *the M-Wright function*, or equivalently *the Mainardi function*[20, 24, 21]. See Section 3.3 for further details. Its role as a fractional extension of classical exponential behavior is reflected in the two reference cases $M_0(z) = \exp(-z)$ and $M_{\frac{1}{2}}(z) = \frac{1}{\sqrt{\pi}} \exp(-z^2/4)$.

The second pair is

- $W_{-\alpha,-1}(-z)$
- $-W_{-\alpha,1-2\alpha}(-z)$

These functions arise naturally from derivatives of $F_\alpha(z)$ and $M_\alpha(z)$. They are useful in random-variable generation, as in (3.19) and Section 11 of [19], and in some cases they reduce to polynomial expressions.

3.2. Classic Results

The recurrence relations of the Wright function are (Chapter 18, Vol 3 of [2])

$$(3.2) \quad \lambda z W_{\lambda,\lambda+\mu}(z) = W_{\lambda,\mu-1}(z) + (1-\mu)W_{\lambda,\mu}(z)$$

$$(3.3) \quad \frac{d}{dz} W_{\lambda,\mu}(z) = W_{\lambda,\lambda+\mu}(z)$$

The moments of the Wright function are (See (1.4.28) of [24])

$$(3.4) \quad \mathbb{E}(X^{d-1}) = \int_0^\infty x^{d-1} W_{-\lambda,\delta}(-x) dx = \frac{\Gamma(d)}{\Gamma(d\lambda + \delta)}$$

400 The way it is written is in fact its Mellin transform:

$$(3.5) \quad W_{\lambda,\delta}(-x) \xleftrightarrow{\mathcal{M}} W_{\lambda,\delta}^*(s) = \frac{\Gamma(s)}{\Gamma(\delta - \lambda s)}$$

401 $W_{\lambda,\delta}(z)$ has the following Hankel integral representation:

$$(3.6) \quad W_{\lambda,\delta}(z) = \frac{1}{2\pi i} \int_H dt \frac{\exp(t + z t^{-\lambda})}{t^\delta}$$

402 Prodanov[33] derived an integral representation of the Wright function. We use the branch with
403 $\lambda < 0$ and $\delta \leq 1$ from Theorem 1, namely

$$(3.7) \quad W_{\lambda,\delta}(z) = \frac{1}{\pi} \int_0^\infty \frac{dr}{r^\delta} \sin(\sin(\lambda\pi)w + \delta\pi) e^{\cos(\lambda\pi)w-r}, \quad \text{where } w = z r^{-\lambda}.$$

404 This representation is well suited to numerical evaluation by tanh-sinh quadrature.

405 The four-parameter Wright function is defined as

$$(3.8) \quad W \begin{bmatrix} a, & b \\ \lambda, & \mu \end{bmatrix} (z) := \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(n+1)} \frac{\Gamma(an+b)}{\Gamma(\lambda n + \mu)}$$

406 This function is a higher-order Wright function. It was used seriously for the first time by the
407 author[19].

408 3.3. The M-Wright Functions

409 Mainardi has introduced two auxiliary functions of Wright type (see F.2 of [20]). Assume $\alpha \in [0, 1]$,

$$(3.9) \quad F_\alpha(z) := W_{-\alpha,0}(-z) \quad (z > 0)$$

$$(3.10) \quad M_\alpha(z) := W_{-\alpha,1-\alpha}(-z) = \frac{1}{\alpha z} F_\alpha(z) \quad (z > 0)$$

410 The relation between $M_\alpha(z)$ and $F_\alpha(z)$ in (3.10) is an application of (3.2) by setting $\lambda = -\alpha, \mu = 1$.

411 $F_\alpha(z)$ has the following Hankel integral representation:

$$(3.11) \quad F_\alpha(z) = \frac{1}{2\pi i} \int_H dt \exp(t - z t^\alpha)$$

412 Both functions have simple Mellin transforms from (3.5):

$$(3.12) \quad F_\alpha(x) \xleftrightarrow{\mathcal{M}} F_\alpha^*(s) = \frac{\Gamma(s)}{\Gamma(\alpha s)}$$

$$(3.13) \quad M_\alpha(x) \xleftrightarrow{\mathcal{M}} M_\alpha^*(s) = \frac{\Gamma(s)}{\Gamma((1-\alpha) + \alpha s)}$$

413 The function $F_\alpha(z)$ is used to define fractional one-sided distributions. However, its series repre-
414 sentation is not especially convenient for numerical computation, because many terms may be required
415 to reach a given precision.

416 By contrast, $M_\alpha(z)$ admits a series representation that is more computationally practical, espe-
417 cially for small values of α :

$$(3.14) \quad M_\alpha(z) = \sum_{n=0}^{\infty} \frac{(-z)^n}{n! \Gamma(-\alpha n + (1-\alpha))} = \frac{1}{\pi} \sum_{n=1}^{\infty} \frac{(-z)^{n-1}}{(n-1)!} \Gamma(\alpha n) \sin(\alpha n \pi) \quad (0 < \alpha < 1)$$

418 $M_\alpha(0) = 1/\Gamma(1-\alpha)$ monotonically decreases from 1 to 0 as α increases from 0 to 1.

419 The function $M_\alpha(z)$ also has simple closed-form expressions in the special cases $\alpha = 0$ and $\alpha = \frac{1}{2}$:

$$(3.15) \quad M_0(z) = \exp(-z) \text{ and } M_{\frac{1}{2}}(z) = \frac{1}{\sqrt{\pi}} \exp(-z^2/4).$$

420 $M_\alpha(z)$ can be computed to high accuracy when properly implemented with arbitrary-precision
 421 floating point library, such as the `mpmath` package[26]. In this regard, it is much more "useful" than
 422 $F_\alpha(z)$.

423 This is particularly important in working with large degrees of freedom and extreme values of
 424 α , mainly close to 0 and 1. The typical 64-bit floating-point algorithm suffers from overflow and/or
 425 underflow. See Section 3.8 for more details.

426 $M_\alpha(z)$ has the asymptotic representation in the *generalized gamma* (GG) style: (see F.20 of [20])

$$(3.16) \quad M_\alpha\left(\frac{x}{\alpha}\right) = A x^{d-1} e^{-B x^p}$$

where $p = 1/(1 - \alpha)$, $d = p/2$, $A = \sqrt{p/(2\pi)}$, $B = 1/(\alpha p)$.

427 Additional correction terms in the asymptotic expansion have been derived up to the order $x^{-6/(1-\alpha)}$ [30].
 428 This formula is important in guiding (3.14) to high precision for large x , where the series representation
 429 often fails to converge.

430 $M_\alpha(x)$ can be used as the density function of a one-sided distribution M_α [21], because $\int_0^\infty M_\alpha(x)dx =$
 431 1 and $M_\alpha(x)$ for $x \geq 0$. Its CDF is another Wright function:

$$(3.17) \quad \int_0^x M_\alpha(t)dt = 1 - W_{-\alpha,1}(-x).$$

432 This is proved in Lemma 2.11.

433 The absolute moments of $M_\alpha(x)$ in $\mathbb{R}^+$ are

$$(3.18) \quad \mathbb{E}(M_\alpha^n) = \int_0^\infty t^n M_\alpha(t)dt = \frac{\Gamma(n+1)}{\Gamma(n\alpha+1)}, \quad n > -1.$$

434 Hence, its mean is located at $1/\Gamma(\alpha+1)$, which is equal to 1 when $\alpha = 0, 1$. Its variance is $2/\Gamma(2\alpha +$
 435 $1) - 1/\Gamma(\alpha+1)^2$. The variance becomes zero when $\alpha = 1$, consistent with $M_1(x) = \delta(x-1)$.

436 Differentiating $M_\alpha(z)$, and from (3.14), we get

$$(3.19) \quad \frac{d}{dz} M_\alpha(z) = -W_{-\alpha,1-2\alpha}(-z) = \frac{-1}{\pi} \sum_{n=2}^{\infty} \frac{(-z)^{n-2}}{(n-2)!} \Gamma(\alpha n) \sin(\alpha n \pi)$$

437 Note that $\frac{d}{dz} M_\alpha(0) = -\frac{1}{\pi} \Gamma(2\alpha) \sin(2\alpha\pi)$. This also indicates that

$$(3.20) \quad \frac{d}{dz} F_\alpha(z) = \alpha \left(1 + z \frac{d}{dz}\right) M_\alpha(z)$$

438 which can be implemented from $M_\alpha(z)$ through (3.14) and (3.19). These differential forms lead to the
 439 concept of elasticity in Section 3.6 and below.

3.4. The Inverse Stable Law

440 M_α is said to follow the *inverse stable law*. In Section 4.2, the extremal stable distribution L_α will
 441 be introduced¹. The law is

$$(3.21) \quad M_\alpha \sim (L_\alpha)^{-\alpha}.$$

443 This can be proved by the moment formula. Let $X = M_\alpha, Y = L_\alpha$. Then $\mathbb{E}(Y^{-q}) = \frac{\Gamma(q/\alpha)}{\alpha\Gamma(q)}$. Set
 444 $q = \alpha n$. We get $Y^{-q} = (Y^{-\alpha})^n = X^n$. It leads to $\mathbb{E}(X^n) = \frac{\Gamma(n)}{\alpha\Gamma(\alpha n)} = \frac{\Gamma(n+1)}{\Gamma(\alpha n+1)}$. That is (3.18).

445 Since $M_{1/2}/\sqrt{2}$ has the half-normal density on $(0, \infty)$, it provides a simple reference case for the
 446 inverse stable law.

¹Historically, people also used the notation S_α because S is the first letter of "stable".

447 The calculation of M_α follows the framework in Theorem 1 of [7]. This sets the foundation for the
 448 tilted stable law in Section 6.5.

449 DEFINITION 3.2. The factor $A_\alpha(q)$ is often called *the Zolotarev function*. It is defined as

$$(3.22) \quad A_\alpha(q) = \frac{\sin(\pi\alpha q)}{\sin(\pi q)^{1/\alpha}} \sin(\pi(1-\alpha)q)^c, \quad c = \frac{1-\alpha}{\alpha}.$$

LEMMA 3.3. For the inverse stable law M_α , the sampling is performed via Kanter's method:

$$M_\alpha = A_\alpha(Q)^{-\alpha} G^{1-\alpha}.$$

- 450 (1) Q is from a uniform random variable in $(0, 1)$.
 451 (2) G is a gamma variable: $G \sim \text{Gamma}(1, \text{scale} = 1)$.

452

△

453 3.5. The Fractional Gamma-Star Function

454 The so-called γ^* function is documented in 8.2.6 and 8.2.7 of DLMF[9]. It is defined as follows:

$$\gamma^*(s, x) := \frac{1}{\Gamma(s)} \int_0^1 t^{s-1} e^{-xt} dt = \frac{x^{-s}}{\Gamma(s)} \gamma(s, x)$$

455 The finite integral in $t \in [0, 1]$ is transformed from the incomplete gamma function, which takes the
 456 form of $\gamma(s, x) = \int_0^x t^{s-1} e^{-t} dt$.

457 $\gamma^*(s, x)$ can be extended fractionally in a straightforward manner. It is used to calculate the CDF
 458 of the FG in Chapter 6. See (6.13) for details.

459 DEFINITION 3.4 (The fractional γ^* function). It is defined by replacing e^{-xt} with $M_\alpha(xt)$ such
 460 that

$$(3.23) \quad \gamma_\alpha^*(s, x) := \frac{\Gamma((1-\alpha) + \alpha s)}{\Gamma(s)} \int_0^1 dt t^{s-1} M_\alpha(xt)$$

461 The $\alpha \rightarrow 0$ limit of $\gamma_\alpha^*(s, x)$ subsumes the classic γ^* function, that is, $\gamma_0^*(s, x) = \gamma^*(s, x)$. This is
 462 reflected in the simple fact that $M_0(xt) = \exp(-xt)$.

463 The γ^* function is a subset of the fractional confluent hypergeometric function in Lemma 5.4.

464

465 The supplementary γ_α^* function is defined as

$$(3.24) \quad \Gamma_\alpha^*(s, x) := \frac{\Gamma((1-\alpha) + \alpha s)}{\Gamma(s)} \int_1^\infty dt t^{s-1} M_\alpha(xt)$$

466 LEMMA 3.5. It has an analytic relation such as

$$(3.25) \quad \Gamma_\alpha^*(s, x) = x^{-s} - \gamma_\alpha^*(s, x)$$

467

△

468 PROOF. The total integral of $t \in \mathbb{R}$ has an analytic solution due to (3.18):

$$\begin{aligned} \Gamma_\alpha^*(s) &:= \frac{\Gamma((1-\alpha) + \alpha s)}{\Gamma(s)} \int_0^\infty dt t^{s-1} M_\alpha(xt) \\ &= \frac{\Gamma((1-\alpha) + \alpha s)}{\Gamma(s) x^s} \int_0^\infty dz z^{s-1} M_\alpha(z) \\ &= x^{-s}. \end{aligned}$$

469 Split the integral into two parts: $\int_0^1$ and $\int_1^\infty$. We obtain (3.25). □

470 The identity formula is

$$(3.26) \quad x^s \Gamma_\alpha^*(s, x) + x^s \gamma_\alpha^*(s, x) = 1$$

471 This identity is critical in understanding how to use γ_α^* versus Γ_α^* when calculating the CDF of a FG
472 distribution.

473 Numerically, it is better to use γ_α^* when $x \leq 1$ and to use Γ_α^* when $x > 1$.

474 3.6. The Elasticity Operator

475 In (3.19) and (3.20), we encountered an important mathematical structure called "elasticity"
476 which will be used in Chapter 13. It provides an elegant view of the inner structure of the FG density
477 functions.

478 DEFINITION 3.6 (The elasticity operator). Assume $f(x)$ is differentiable for $x \in \mathbb{R}$. The elasticity
479 of $f(x)$ is defined as

$$(3.27) \quad \mathcal{L} f(x) := \frac{x}{f(x)} \frac{d}{dx} f(x)$$

$$(3.28) \quad = \frac{d \log f(x)}{d \log x}, \quad \text{when } x > 0 \text{ and } f(x) > 0.$$

480 The second line can be interpreted as the percentage change of $f(x)$ over a percentage change of x .
481 This is often used in statistics and economics. (It is an extension of the Euler dilation operator, $x \frac{d}{dx}$.)

482 To illustrate its property, if $f(x) \sim x^k$ locally, then $\mathcal{L} f(x) \approx k$. It informs *local degree of homo-*
483 *geneity* in the scaling analysis.

484 More generally, some algebraic rules of $\mathcal{L}$ are

- 485 • $\mathcal{L}[f(x)g(x)] = \mathcal{L}f(x) + \mathcal{L}g(x)$; multiplication becomes addition.
- 486 • $\mathcal{L}[f(g(x))] = \mathcal{L}g(x) \times [\mathcal{L}f](g(x))$; composition becomes multiplication.
- 487 • $\mathcal{L}(x^k) = k$; the trivial case is $\mathcal{L}(x) = 1$.
- 488 • $\mathcal{L}(e^{-x}) = -x$;
- 489 • $\mathcal{L}(\text{constant})$ is zero;

490 As an application, it is a good exercise to derive $\mathcal{L}[f((x/\sigma)^p)] = p[\mathcal{L}f]((x/\sigma)^p)$.

491
492 The recurrence relations of the Wright function, (3.2) and (3.3), can be rewritten using the $\mathcal{L}$ operator.
493 They become two expressions of the elasticity of the Wright function.

494 Define the ratio of two Wright functions as

$$(3.29) \quad Q_{\lambda, \mu, \delta}(z) = \frac{W_{\lambda, \mu + \delta}(z)}{W_{\lambda, \mu}(z)}.$$

495 It follows immediately that (3.2) becomes

$$(3.30) \quad \lambda z Q_{\lambda, \mu, \lambda}(z) = Q_{\lambda, \mu, -1}(z) + 1 - \mu.$$

496 LEMMA 3.7. The elasticity of the Wright function is expressed by the following ratios:

$$(3.31) \quad \mathcal{L} W_{\lambda, \mu}(z) = \frac{1}{\lambda} Q_{\lambda, \mu, -1}(z) + \frac{1 - \mu}{\lambda},$$

$$(3.32) \quad \mathcal{L} W_{\lambda, \mu}(z) = z Q_{\lambda, \mu, \lambda}(z).$$

497 △

498 PROOF. The second line is straightforward from (3.3). The first line is derived from the second
499 line by replacing the $z Q_{\lambda, \mu, \lambda}(z)$ term on the RHS with (3.30). □

3.7. The Elasticity of the M-Wright Functions

What we are most interested in is the elasticity of $M_\alpha(x)$:

$$(3.33) \quad \mathcal{L} M_\alpha(x) = [\mathcal{L} W_{-\alpha, 1-\alpha}](-x)$$

which is from (3.10). Note that $\mathcal{L} F_\alpha(x)$ is trivial if $\mathcal{L} M_\alpha(x)$ is known. This is due to (3.20), we have

$$(3.34) \quad \mathcal{L} F_\alpha(x) = \mathcal{L} M_\alpha(x) + 1.$$

However, $\mathcal{L} F_\alpha(x)$ has a representation that is more friendly to FCM. From (3.31),

$$(3.35) \quad \mathcal{L} F_\alpha(x) = [\mathcal{L} W_{-\alpha, 0}](-x) = \frac{1}{\alpha} Q_\alpha(x) - \frac{1}{\alpha},$$

$$(3.36) \quad \text{where } Q_\alpha(x) := -Q_{-\alpha, 0, -1}(-x) = -\frac{W_{-\alpha, -1}(-x)}{W_{-\alpha, 0}(-x)}.$$

It follows that $Q_\alpha(x) = \alpha \mathcal{L} M_\alpha(x) + (1 + \alpha)$.

The following lemma converts the elasticity of the FG PDF to either $\mathcal{L} M_\alpha(x)$ or $Q_\alpha(x)$.

LEMMA 3.8. Let $\mathfrak{N}(x)$ represent the functional form of FG PDF (6.1) where $\mathfrak{N}(x) = x^{d-1} F_\alpha\left(\left(\frac{x}{\sigma}\right)^p\right)$ (apart from a constant multiplier). The elasticity of $\mathfrak{N}(x)$ is

$$(3.37) \quad \mathcal{L} \mathfrak{N}(x) = p [\mathcal{L} M_\alpha]\left((x/\sigma)^p\right) + (d + p - 1).$$

Alternatively, a useful ratio form for the FCM where p/α is a constant is

$$(3.38) \quad \mathcal{L} \mathfrak{N}(x) = \frac{p}{\alpha} Q_\alpha\left(\left(\frac{x}{\sigma}\right)^p\right) - \frac{p}{\alpha} + (d - 1).$$

We observe that the role of the degrees of freedom d is very simple in $\mathcal{L} \mathfrak{N}(x)$. It shifts the constant level, but it does not affect the shape of $\mathcal{L} \mathfrak{N}(x)$.

△

$\mathcal{L} M_\alpha(x)$ has simple behaviors in a few cases. For example,

$$\begin{aligned} \mathcal{L} M_0(x) &= -x; \\ \mathcal{L} M_{1/2}(x) &= -x^2/2. \end{aligned}$$

When $x \rightarrow 0$,

$$(3.39) \quad \mathcal{L} M_\alpha(x) \sim -b_1 x, \quad \text{where } b_1 := \frac{\Gamma(1 - \alpha)}{\Gamma(1 - 2\alpha)}.$$

When $\alpha \in [0, 1/2]$, $\mathcal{L} M_\alpha(x) < 0$ for all $x > 0$. It is a monotonically decreasing function for $x \in [0, \infty)$.

When $x \rightarrow \infty$, the GG-style asymptotic form in (3.16) leads to

$$(3.40) \quad \mathcal{L} M_\alpha(x) \sim -\alpha^{\alpha/(1-\alpha)} x^{1/(1-\alpha)} + \frac{\alpha - 1/2}{1 - \alpha},$$

in which the first term is dominant. It leads to the asymptotic limit of second-order elasticity:

$$(3.41) \quad \lim_{x \rightarrow \infty} \mathcal{L}[-\mathcal{L} M_\alpha](x) \rightarrow \frac{1}{1 - \alpha}.$$

It follows immediately from (3.32) that (with $z \rightarrow -x$)

$$(3.42) \quad \mathcal{L} M_\alpha(x) = -x \frac{W_{-\alpha, 1-2\alpha}(-x)}{W_{-\alpha, 1-\alpha}(-x)} = -x Q_{-\alpha, 1-\alpha, -\alpha}(-x)$$

where the series form of the numerator is in (3.19). We can compute the numerator and denominator individually, then take the ratio. Or we can derive its series representation as follows.

LEMMA 3.9. The series representation of $\mathcal{L} M_\alpha(x) = -x Q_{-\alpha, 1-\alpha, -\alpha}(-x)$ is

$$\mathcal{L} M_\alpha(x) = \sum_{k=1}^{\infty} c_k x^k$$

522 where

$$(3.43) \quad c_k = \frac{(-1)^k}{(k-1)!} b_k + \sum_{j=1}^{k-1} \frac{(-1)^{(j+1)}}{j!} b_j c_{k-j}, \quad k \geq 1;$$

$$(3.44) \quad b_n = \frac{\Gamma(1-\alpha)}{\Gamma(1-\alpha(n+1))}, \quad n \geq 1.$$

523

$\triangle$

PROOF. From (3.14), we have

$$M_\alpha(x) = \sum_{n=0}^{\infty} a_n x^n, \quad a_n = \frac{(-1)^n}{n! \Gamma(1-\alpha(n+1))}.$$

Then (3.19) can be written as

$$\frac{d}{dx} M_\alpha(x) = -W_{-\alpha, 1-2\alpha}(-x) = \sum_{n=1}^{\infty} n a_n x^{n-1}.$$

And

$$x \frac{M'_\alpha(x)}{M_\alpha(x)} = \frac{\sum_{n \geq 1} n a_n x^n}{\sum_{n \geq 0} a_n x^n}.$$

The coefficients satisfy the standard recurrence of series divisions, which becomes

$$c_k = \frac{1}{a_0} \left(k a_k - \sum_{j=1}^{k-1} a_j c_{k-j} \right), \quad k \geq 1.$$

524 With $a_0 = \frac{1}{\Gamma(1-\alpha)}$, and $\frac{a_n}{a_0} = \frac{(-1)^n}{n!} b_n$, it leads to (3.43) and (3.44).

525

$\square$

REMARK 3.10. The first three coefficients are explicitly derived as follows.

$$c_1 = -b_1 = -\frac{\Gamma(1-\alpha)}{\Gamma(1-2\alpha)},$$

$$c_2 = b_2 - b_1^2 = \frac{\Gamma(1-\alpha)}{\Gamma(1-3\alpha)} - \left(\frac{\Gamma(1-\alpha)}{\Gamma(1-2\alpha)} \right)^2,$$

$$c_3 = -\frac{1}{2} b_3 + \frac{3}{2} b_2 b_1 - b_1^3 = -\frac{\Gamma(1-\alpha)}{2 \Gamma(1-4\alpha)} + \frac{3}{2} \frac{\Gamma(1-\alpha)^2}{\Gamma(1-2\alpha) \Gamma(1-3\alpha)} - \left(\frac{\Gamma(1-\alpha)}{\Gamma(1-2\alpha)} \right)^3.$$

The small- x expansion up to the x^3 term is

$$\mathcal{L} M_\alpha(x) = \left[-\frac{\Gamma(1-\alpha)}{\Gamma(1-2\alpha)} \right] x + c_2 x^2 + c_3 x^3 + O(x^4).$$

3.8. Numerical Methods of the M-Wright Functions

To properly compute the subsequent special functions and distributions in this book, we need a very robust numerical implementation of $F_\alpha(x)$ and $M_\alpha(x)$ for *the entire range* of $\alpha \in [0, 1]$ and $x \geq 0$. Since $F_\alpha(x) = \alpha x M_\alpha(x)$, we can easily compute one from the other in most cases. It is a matter of which approach is faster, more convenient, and precise.

3.8.1. Handling alpha for zero and one. When $\alpha = 0$, we should use $M_0(x) = e^{-x}$. $\lim_{\alpha \rightarrow 0} F_\alpha(x)$ should be handled carefully in the fractional gamma distribution.

When $\alpha = 1$, we could use a normal distribution to simulate the delta function: $M_1(x) = \mathcal{N}(x; 1, \sigma^2)$ where $\sigma = 0.001$. This is to ensure that $\int_0^\infty M_1(x) dx = 1$.

3.8.2. Using `scipy.stats.levy-stable`. Both functions can be derived from the one-sided α -stable distribution $L_\alpha(x)$ of Section 4.2, which is implemented in `scipy.stats.levy-stable` package[39].

For example, due to (3.21), $M_\alpha(x)$ can be computed using $L_\alpha(x) = \alpha x^{-\alpha-1} M_\alpha(x^{-\alpha})$ where $x > 0$. On the other hand, for $\beta > 1/2$, we can also use $M_\beta(x) = \alpha L_\alpha^{\alpha-2}(x)$ where $\beta = 1/\alpha$.

These two numerical methods are good for the bulk of α and x . However, they begin to lose precision for small $\alpha < 0.08$ and large $\alpha > 0.99$. They are also not good enough for small $x < 0.01$.

3.8.3. Using the series sum in `numpy`. Based on (3.14), we define the sum of the series of finite terms as

$$(3.45) \quad M_\alpha^{(m)}(x) = \frac{1}{\pi} \sum_{n=1}^m \frac{(-x)^{n-1}}{(n-1)!} \Gamma(\alpha n) \sin(\alpha n \pi). \quad (0 < \alpha < 1)$$

This method implemented in `numpy` and `scipy` is good for several scenarios. First, to cover the small x area ($x < 0.01$), use $M_\alpha^{(7)}(x)$ if $\alpha < 0.9$.

Otherwise, we could use $M_\alpha^{(80)}(x)$ for $\alpha \leq 0.998$ and $x < 0.85$. The sum of 80 terms takes more time to compute. But it is a necessary path when `scipy.stats.levy-stable` approach loses precision.

3.8.4. Using the series sum in `mpmath`. $M_\alpha^{(m)}(x)$ implemented in `mpmath` is our de facto implementation to calibrate the precision of other approaches. In order to make it a good baseline implementation, we must carefully choose `mp.prec` and m to use.

After rigorous testing, it was found that `mp.prec` ≥ 64 provides sufficient precision. Therefore, `mp.prec` = 128 is more than abundant up to three decimal points in α . `mpmath` is smart about handling summing many small terms, especially with large amount of cancellation due to the $\sin(\alpha n \pi)$ factor in (3.45). This feature can be used during the calibration phase.

The more important numerical choice is the truncation level m . In practice, $m = 40,000$ is sufficient for $\alpha < 0.9$, whereas a much larger value, $m = 80,000$, is needed when α is very close to 1, such as $\alpha = 0.998$. Larger values of m substantially increase the computational cost of the series summation. This approach is therefore appropriate during calibration, but it is too expensive for the final NumPy-style implementation.

This will be elaborated on in the next section.

3.8.5. Using the asymptotic approximation. Paris et al. [30] derived a more refined asymptotic formula, where (3.16) is simply its first term. Theorem 2.2 of that paper is recaptured in the following.

LEMMA 3.11.

$$(3.46) \quad M_\alpha(x) \sim \frac{A(\alpha)}{2\pi} X^{\alpha-1/2} e^{-X} \sum_{n=0}^{\infty} c_j(\alpha) (-X)^{-j}. \quad (0 < \alpha < 1)$$

563 where $c_j(\alpha)$ is in its (2.4) up to $j = 6$. Other parameters are $A(\alpha) = \sqrt{\frac{2\pi}{\alpha}} \left(\frac{\alpha}{\kappa}\right)^\alpha$ and $X = \kappa(hx)^{1/\kappa}$
 564 with $\kappa = 1 - \alpha$ and $h = \alpha^\alpha$.
 565 $\triangle$

566 When $M_\alpha(x)$ is small, (3.46) could be very precise with an error as small as 10^{-5} . Our strategy
 567 is to use other implementations to get $M_\alpha(x)$ to a small number, e.g. 10^{-6} in most cases, and at least
 568 10^{-3} in some difficult cases. Then use (3.46) for larger x up to infinity (the maximum 64-bit float).

569 This right-tail strategy works for the bulk of α from 0.1 to 0.9. The transition interval (defined as
 570 $M_\alpha(x) \in [10^{-5}, 10^{-6}]$) could be precomputed by the faster (3.16).

571 For α from 0.9 to 0.99, the `mpmath` version of $M_\alpha^{(m)}(x)$ is more precise to determine the transition
 572 interval.

573 For α from 0.001 to 0.1, the asymptotic form is adjusted to

$$(3.47) \quad M_\alpha(x) \sim A'(\alpha) e^{-B'(\alpha)X'}, \quad \text{where } X' = x^{1/\kappa}.$$

574 $A'(\alpha)$ and $B'(\alpha)$ are obtained from a linear regression in the transition interval: $\log M_\alpha(x) \sim \log A'(\alpha) -$
 575 $B'(\alpha)X'$.

576 **3.8.6. Using the integral form.** For α in the range from 0.99 to 0.998, the `scipy` implementa-
 577 tion of $M_\alpha(x)$ loses precision rapidly in the right tail. In this regime, we use (3.7) to supplement the
 578 computation as long as $M_\alpha(x) > 10^{-3}$.

579 The main numerical difficulty arises when the target value of $M_\alpha(x)$ is very small. In that case,
 580 the integrand in (3.7) becomes highly oscillatory and is effectively nonzero only over a very narrow
 581 range of r^δ . Standard integration algorithms have difficulty identifying this small interval, resolving
 582 the oscillations, and performing the required cancellation accurately. A more sophisticated quadrature
 583 method would be needed, and this is left for future research.

584 For $M_\alpha(x) < 10^{-3}$ at large x , we still use (3.46) asymptotically. This equation is fine for large α ,
 585 as long as the numeric overflow is handled properly.

The Alpha-Stable Distribution - Review

The two-sided distributions developed in this book are built on the α -stable distribution, introduced in Paul Lévy's 1925 monograph[16]. Among its parameters, the central one is the *stability index* $\alpha \in (0, 2]$, which we interpret throughout the book as the principal *fractional* parameter.

This chapter reviews the α -stable distribution from the Mellin transform perspective. That viewpoint is used repeatedly in the later construction of fractional extensions. The ratio-distribution representation of the density in Section 4.3 is due to the author.

4.1. Classic Result

The α -stable distribution has two shape parameters. There are many parameterizations that have been studied (see p.5 of [28]). We are primarily concerned with Feller's (α, θ) parameterization [11, 12], where α is called the stability index with a range of $0 < \alpha \leq 2$, and θ is an angle that injects skewness into the distribution when it is not zero¹

An innovative approach is to study its Mellin transform. This presentation is used because it is *simpler* and provides great insight into its structure.

LEMMA 4.1. The Mellin transform of its PDF is

$$(4.2) \quad L_{\alpha}^{\theta}(x) \xleftrightarrow{\mathcal{M}} \epsilon \frac{\Gamma(s)\Gamma(\epsilon(1-s))}{\Gamma(\gamma(1-s))\Gamma(1-\gamma+\gamma s)}$$

$$\text{where } \epsilon = \frac{1}{\alpha}, \gamma = \frac{\alpha - \theta}{2\alpha}.$$

where $0 < C < 1$ implicitly. This is defined for $x \geq 0$. The reflection rule is used for $x < 0$ such that $L_{\alpha}^{\theta}(x) := L_{\alpha}^{-\theta}(-x)$.

△

This formula was first derived by Schneider in 1986[34], rediscovered by Mainardi et al. in 2001[22], and later summarized by Mainardi and Pagnini in (2.8) of [23].

Equation (4.2) is written not directly in (α, θ) , but in the auxiliary variables (ϵ, γ) . This reparameterization is especially convenient in Mellin space, where the transform takes a more symmetric form.

In the Feller parameterization, the admissible range $|\theta| \leq \min\{\alpha, 2 - \alpha\}$ is known as the Feller-Takayasu diamond. In the (ϵ, γ) variables, the same constraint becomes (a) $0 \leq \gamma \leq 1$ when $\epsilon > 1$, and (b) $1 - \epsilon \leq \gamma \leq \epsilon$ when $\epsilon \leq 1$.²

¹The conversion from/to the S1 parameterization is summarized as

$$(4.1) \quad \beta \tan\left(\frac{\alpha\pi}{2}\right) = -\tan\left(\frac{\theta\pi}{2}\right), \quad \text{scale} = \cos^{1/\alpha}\left(\frac{\theta\pi}{2}\right).$$

where $\beta \in [-1, 1]$ is the standard skew parameter implemented in `scipy.stats.levy_stable`.

²Conversely, if γ is fixed, condition (b) implies $\alpha \leq \min\{1/\gamma, 1/(1-\gamma)\}$.

4.1.1. The Reflection Rule. Note that the reflection of $\theta \rightarrow -\theta$ in the (α, θ) parametrization is equivalent to the reflection of $\gamma \rightarrow 1 - \gamma$ in the (ϵ, γ) parametrization.

Since we often mingle the two parameterizations, this alternative view can be very helpful in certain scenarios. For example, the total density in the positive domain is $\int_0^\infty L_\alpha^\theta(x) = \gamma$. By the reflection rule, $\int_0^\infty L_\alpha^{-\theta}(x) = 1 - \gamma$. Hence, the total density $\int_{-\infty}^\infty L_\alpha^\theta(x) = \gamma + (1 - \gamma) = 1$.

4.2. Extremal Distributions

There are two extremal regimes in which the skew parameter θ reaches the boundary of its admissible range. These cases are important because the M-Wright functions $F_\alpha(x)$ and $M_\alpha(x)$ from Section 3.3 arise naturally from them.

The first extremal regime corresponds to $\gamma = 0$ or $\gamma = 1$, or equivalently $\theta = \pm\alpha \leq 1$. By the reflection rule, it is sufficient to study the case $\theta = -\alpha$, or $\gamma = 1$.

This yields the one-sided α -stable distribution:

$$L_\alpha(x) := L_\alpha^{-\alpha}(x) \xleftrightarrow{\mathcal{M}} \epsilon \frac{\Gamma(\epsilon(1-s))}{\Gamma(1-s)}$$

In the literature [7, 8], this distribution is often characterized by the Laplace transform

$$\mathbb{E}[e^{-tL_\alpha}] = e^{-t^\alpha}, \quad t \geq 0.$$

REMARK 4.2. Starting from (3.12) where $F_\alpha(x) \xleftrightarrow{\mathcal{M}} F_\alpha^*(s) = \Gamma(s)/\Gamma(\alpha s)$, and applying three Mellin-transform operations, namely $x \rightarrow x^\alpha$, multiplication by x , and then $x \rightarrow x^{-1}$, we recover the classical identity

$$(4.3) \quad L_\alpha(x) = x^{-1} F_\alpha(x^{-\alpha}) \quad (x \geq 0 \text{ and } 0 < \alpha \leq 1)$$

with the endpoint case $L_1(x) = \delta(x - 1)$. This leads to the important random variable relation of $M_\alpha \sim (L_\alpha)^{-\alpha}$ in (3.21).

REMARK 4.3. $L_\alpha(x)$ can be computed with `scipy.stats.levy_stable`[39] in the S1 parameterization by setting `beta=1` and `scale=cos(\alpha\pi/2)^{1/\alpha}` for $0 < \alpha < 1$.³ This provides a practical numerical starting point for developing the new fractional distributions and testing the constructions in proof-of-concept form.

The second extremal regime occurs when $\theta = \alpha - 2$. In this case $\epsilon = \gamma = 1/\alpha$, and

$$L_\alpha^{\alpha-2}(x) \xleftrightarrow{\mathcal{M}} \epsilon \frac{\Gamma(s)}{\Gamma(1 - \epsilon + \epsilon s)}$$

Comparing this expression with (3.13) gives the classical identity: (See, for example, formula (F.49) in [20])

$$(4.4) \quad L_\alpha^{\alpha-2}(x) = \frac{1}{\alpha} M_{1/\alpha}(x) \quad (x \in \mathbb{R} \text{ and } 1 < \alpha \leq 2)$$

This identity extends the M-Wright function to $x < 0$, because $L_\alpha^{\alpha-2}(x)$ is two-sided.

4.3. Ratio Distribution Approach

Important insight can be obtained by interpreting (4.2) as a ratio distribution (2.6). We split (4.2) into two components:

$$(4.5) \quad L_\alpha^\theta(x) \xleftrightarrow{\mathcal{M}} \epsilon \left[\frac{\Gamma(s)}{\Gamma(1 - \gamma + \gamma s)} \right] \left[\frac{\Gamma(\epsilon(1-s))}{\Gamma(\gamma(1-s))} \right]$$

The first bracket is the Mellin transform of the M-Wright function (3.13).

³See Chapter 1 of [28] for further discussion of stable-law parameterizations. We do not pursue that issue here.

643 The second bracket is related to the Mellin transform of the fractional χ -mean distribution (FCM)
 644 at $k = 1$:

$$(4.6) \quad \begin{aligned} \bar{\chi}_{\alpha,1}^{\theta}(x) &\stackrel{\mathcal{M}}{\longleftrightarrow} \bar{\chi}_{\alpha,1}^{\theta,*}(s) \\ &= \epsilon \gamma^{\gamma(s-1)-1} \frac{\Gamma(\epsilon(s-1))}{\Gamma(\gamma(s-1))} \end{aligned}$$

645 For a ratio distribution, the Mellin rule requires replacing s with $2 - s$ in $\bar{\chi}_{\alpha,1}^{\theta,*}(s)$. Consequently,
 646 the factor $s - 1$ in the second line of (4.6) becomes $1 - s$ in the second bracket of (4.5).

647 **4.3.1. Rescaled M-Wright Function.** Additionally, a small nuance here is to deal with scaling
 648 factors. Define the rescaled M-Wright function

$$(4.7) \quad \tilde{M}_{\gamma}(x) := \gamma^{1-\gamma} M_{\gamma}(x/\gamma^{\gamma}), \quad x \geq 0$$

$$(4.8) \quad \tilde{M}_{\gamma}(-x) := \tilde{M}_{1-\gamma}(x) \quad (\text{reflection rule}).$$

649 Since $\int_0^{\infty} \tilde{M}_{\gamma}(x) dx = \gamma$ and $\int_{-\infty}^0 \tilde{M}_{\gamma}(x) dx = \int_0^{\infty} \tilde{M}_{1-\gamma}(x) dx = 1 - \gamma$, we have $\int_0^{\infty} M_{\gamma}(x) = 1$.
 650 Hence, $\tilde{M}_{\gamma}(x)$ is a valid two-sided density function.

651 The scale is adjusted so that it matches the standard normal distribution: $\tilde{M}_{1/2}(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$
 652 of $\mathcal{N}(0, 1)$.

653 According to (2.3), the rescaling of PDF modifies the Mellin transform from (3.13) to

$$(4.9) \quad \begin{aligned} \tilde{M}_{\gamma}(x) &\stackrel{\mathcal{M}}{\longleftrightarrow} \tilde{M}_{\gamma}^{*}(s) \\ &= \gamma^{1-\gamma+\gamma s} \frac{\Gamma(s)}{\Gamma(1-\gamma+\gamma s)} \end{aligned}$$

654 from which the $\gamma^{1-\gamma+\gamma s}$ term cancels out its counterpart in $\bar{\chi}_{\alpha,1}^{\theta,*}(2-s)$ nicely.

655 Therefore, we find a new method to construct the α -stable distribution using the following integral.

656 **LEMMA 4.4** (The ratio-distribution representation of the α -stable distribution). The Mellin trans-
 657 form of the PDF (4.2) becomes

$$(4.10) \quad L_{\alpha}^{\theta}(x) \stackrel{\mathcal{M}}{\longleftrightarrow} \tilde{M}_{\gamma}^{*}(s) \bar{\chi}_{\alpha,1}^{\theta,*}(2-s)$$

658 from which the PDF can be written in a ratio distribution form of

$$(4.11) \quad L_{\alpha}^{\theta}(x) := \int_0^{\infty} \tilde{M}_{\gamma}(xs) \bar{\chi}_{\alpha,1}^{\theta}(s) s ds \quad (x \geq 0)$$

659 Since the Mellin integral is only valid for $x > 0$, it is supplemented with *the reflection rule*:

$$(4.12) \quad L_{\alpha}^{\theta}(-x) := L_{\alpha}^{-\theta}(x)$$

660 △

661 **4.3.2. Ratio of Stable Law Approach.** It can be expressed elegantly in terms of the stable
 662 law. The normalization term γ^{γ} cancels out between $\tilde{M}_{\gamma}$ and $\bar{\chi}_{\gamma\alpha,1}^{\theta}$ (See Section 7.2.1). For $x \geq 0$, we
 663 obtain the following law of ratio:

$$(4.13) \quad L_{\alpha}^{\theta} \sim M_{\gamma} / T_{\gamma\alpha,0}^{-\gamma} = (L_{\gamma\alpha} / L_{\gamma})^{\gamma}$$

664 for the γ portion of the population. For $x < 0$, the law is modified by replacing γ with $1 - \gamma$ for the
 665 $(1 - \gamma)$ portion of the population due to the reflection rule.

666

This construction places $\bar{\chi}_{\alpha,1}^\theta$ in the central role. We define it at one degree of freedom $k = 1$. In Chapter 7, we will add *degrees of freedom* k to it and make it $\bar{\chi}_{\alpha,k}^\theta$, which is the fractional extension of the classic χ distribution.

Subsequently, in Chapter 11, we will add *degrees of freedom* k to the α -stable distribution and merge it with Student's t distribution.

4.4. SaS

Note that $\theta = 0$ is equivalent to $\gamma = 1/2$. The distribution is symmetric, with the nickname of "SaS", which stands for "Symmetric α -Stable".

Its Mellin transform is simplified to

$$\begin{aligned} L_\alpha^0(x) &\stackrel{\mathcal{M}}{\longleftrightarrow} \epsilon \left[\frac{\Gamma(s)}{\Gamma((1+s)/2)} \right] \left[\frac{\Gamma(\epsilon(1-s))}{\Gamma((1-s)/2)} \right] \\ (4.14) \quad &= \epsilon \left[\frac{2^{s-1}}{\sqrt{\pi}} \Gamma\left(\frac{s}{2}\right) \right] \left[\frac{\Gamma(\epsilon(1-s))}{\Gamma((1-s)/2)} \right]. \end{aligned}$$

The first bracket is the Mellin transform of a normal distribution (2.9) with a scale. The second bracket is $\bar{\chi}_{\alpha,1}^0 \ast (2-s)$ from above.

Hence, the PDF of SaS is

$$(4.15) \quad L_\alpha^0(x) = \int_0^\infty \mathcal{N}(xs) \bar{\chi}_{\alpha,1}^0(s) s ds.$$

This is one of the foundations of GAS-SN in (12.1).

4.4.1. Method of Normal Mixture. SaS in (4.15) will be generalized to GSaS in (12.3) in Chapter 12. Both integrals are in the normal mixture structure (9.1) that enjoys several nice properties described in Chapter 9.

The classic exponential power distribution (Section 3.11.1 of [28]) is the characteristic function transform in Lemma 9.2.

Fractional Hypergeometric Functions

This chapter develops fractional versions of two classical hypergeometric functions: the confluent hypergeometric function ${}_1F_1(a, b; x)$, also denoted by $M(a, b; x)$ (Chapter 13 of DLMF[9]), and the Gauss hypergeometric function ${}_2F_1(a, b, c; x)$ (Chapter 15 of DLMF).

The confluent case appears in the CDFs of the FG and FCM distributions, whereas the Gauss case appears in the CDFs of the GSaS and F distributions. We therefore first record the relevant DLMF formulas and then convert them to the Mellin-transform convention used in (2.2).

From DLMF 13.2.4 and 13.4.16, the Mellin transform of the Kummer function is

$$M(a, b; -x) = \frac{\Gamma(b)}{2\pi i \Gamma(a)} \int_{C-i\infty}^{C+i\infty} \frac{\Gamma(a-s)\Gamma(s)}{\Gamma(b-s)} x^{-s} ds,$$

where $a \neq 0, -1, -2, \dots$

From DLMF 15.1.2 and 15.6.6, the Mellin transform of the Gauss hypergeometric function is

$${}_2F_1(a, b, c; -x) = \frac{\Gamma(c)}{2\pi i \Gamma(a)\Gamma(b)} \int_{C-i\infty}^{C+i\infty} \frac{\Gamma(a-s)\Gamma(b-s)\Gamma(s)}{\Gamma(c-s)} x^{-s} ds,$$

where $a, b \neq 0, -1, -2, \dots$

In the Mellin-transform notation used in this book, these formulas become

$$(5.1) \quad M(a, b; -x) \xleftrightarrow{\mathcal{M}} M^*(a, b; s) = \frac{\Gamma(b)}{\Gamma(a)} \frac{\Gamma(a-s)\Gamma(s)}{\Gamma(b-s)},$$

$$(5.2) \quad {}_2F_1(a, b, c; -x) \xleftrightarrow{\mathcal{M}} {}_2F_1^*(a, b, c; s) = \frac{\Gamma(c)}{\Gamma(a)\Gamma(b)} \frac{\Gamma(a-s)\Gamma(b-s)\Gamma(s)}{\Gamma(c-s)}.$$

We now introduce the corresponding fractional extensions.

5.1. Fractional Confluent Hypergeometric Function

The fractional confluent hypergeometric function (FCHF) is the union of the Kummer function and the Wright function. It allows us to extend many classic functions to their fractional forms.

We start with its Mellin transform. And we follow with the integral and series representations.

DEFINITION 5.1. The Mellin transform of the FCHF is

$$(5.3) \quad M_{\lambda, \delta}(a, b; -x) \xleftrightarrow{\mathcal{M}} M_{\lambda, \delta}^*(a, b; s) = \frac{\Gamma(b)}{\Gamma(a)} \frac{\Gamma(a-s)\Gamma(s)}{\Gamma(\delta - \lambda s)\Gamma(b-s)}$$

where the $\Gamma(\delta - \lambda s)$ term is from the Wright function (3.5).

LEMMA 5.2. The integral representation from DLMF 13.4.1 is extended to

$$(5.4) \quad M_{\lambda, \delta}(a, b; z) = \frac{\Gamma(b)}{\Gamma(a)\Gamma(b-a)} \int_0^1 W_{\lambda, \delta}(zt) t^{a-1} (1-t)^{b-a-1} dt$$

The obvious limit $W_{0,1}(zt) = e^{zt}$ restores it to the classic DLMF formula.

△

PROOF. Replace the Wright function in (5.4) with its Hankel integral (3.6),

$$M_{\lambda,\delta}(a, b; z) = \frac{\Gamma(b)}{2\pi i \Gamma(a)} \int_0^1 \int_{Ha} \left(\frac{e^{s+zt} s^{-\lambda}}{s^\delta} ds \right) t^{a-1} (1-t)^{b-a-1} dt$$

which can be simplified to

$$M_{\lambda,\delta}(a, b; z) = \frac{1}{2\pi i} \int_{Ha} (s^{-\delta} e^s ds) M(a, b; -z s^{-\lambda})$$

Substitute the Mellin integral from (5.1) to it,

$$\begin{aligned} M_{\lambda,\delta}(b, c; -z) &= \frac{1}{2\pi i} \int_{Ha} (s^{-\delta} e^s ds) \frac{\Gamma(b)}{2\pi i \Gamma(a)} \int_{C-i\infty}^{C+i\infty} \frac{\Gamma(b-t)\Gamma(t)}{\Gamma(c-t)} (z s^{-\lambda})^{-t} dt \\ &= \frac{\Gamma(b)}{2\pi i \Gamma(a)} \int_{C-i\infty}^{C+i\infty} \left[\frac{1}{2\pi i} \int_{Ha} s^{\lambda t - \delta} e^s ds \right] \frac{\Gamma(b-t)\Gamma(t)}{\Gamma(c-t)} z^{-t} dt \\ &= \frac{\Gamma(b)}{2\pi i \Gamma(a)} \int_{C-i\infty}^{C+i\infty} \left[\frac{1}{\Gamma(\delta - \lambda t)} \right] \frac{\Gamma(b-t)\Gamma(t)}{\Gamma(c-t)} z^{-t} dt \end{aligned}$$

which is the Mellin transform in (5.3).

From the second line to the third line, we use the well-known Hankel integral of the reciprocal gamma function:

$$\frac{1}{2\pi i} \int_{Ha} s^{-z} e^s ds = \frac{1}{\Gamma(z)}$$

□

LEMMA 5.3. The series representation is

$$(5.5) \quad M_{\lambda,\delta}(a, b; z) := \sum_{n=0}^{\infty} \left[\frac{(a)_n}{(b)_n \Gamma(\lambda n + \delta)} \right] \frac{z^n}{n!}$$

where $(a)_n, (b)_n$ are Pochhammer symbols.

△

PROOF. Take (5.3) and apply Ramanujan's master theorem from Section 2.3. This produces $(M_{\lambda,\delta}^*(a, b; s)/\Gamma(s))|_{s=-n}$, which is equal to the bracket term, since $(x)_n = \Gamma(x+n)/\Gamma(x)$. □

5.1.1. FCHF Subsumes the Kummer Function. It is obvious that $M_{0,1}(a, b; x) = M(a, b; x)$.

5.1.2. FCHF Subsumes the M-Wright Function. By using the same setting from (3.10), we get

$$M_\alpha(z) = M_{-\alpha, 1-\alpha}(c, c; -z) \quad (c \neq 0)$$

5.1.3. FCHF Subsumes Fractional Gamma-Star Function. An important variant of FCHF is the fractionalization of the incomplete gamma function. The reader is referred to Sections 8 and 13 of DLMF[9] and Wikipedia for background information.

We are mainly concerned with the following setup:

$$M_{-\alpha, 1-\alpha}(c, c+1; -x) = c \int_0^1 M_\alpha(xt) t^{c-1} dt$$

This integral is found in (3.23). Hence, we obtain -

LEMMA 5.4. The fractional γ^* function (3.23) has the following FCHF representation:

$$(5.6) \quad \gamma_\alpha^*(s, x) = \frac{\Gamma(\alpha s - \alpha + 1)}{\Gamma(s + 1)} M_{-\alpha, 1-\alpha}(s, s + 1; -x)$$

△

The fractional γ^* function is the basis for expressing the CDF of the fractional gamma distribution in Section 6.6. In fact, this was the main motivation to enrich the classic confluent hypergeometric function.

5.2. Fractional Gauss Hypergeometric Function

The fractional Gauss hypergeometric function (FGHF) arises from the ratio distribution between an elementary function and FCM2 ($\hat{\chi}_{\alpha, k}^2$) in Section 7.6.

When $\alpha = 1$, the Mellin transform of FCM2 is reduced from a fractional form to a classic form in (7.27). The ratio distribution is reduced to a Gauss hypergeometric function ${}_2F_1$. Hence, we consider the general form of such a ratio distribution as fractional ${}_2F_1$.

We start by modifying the Mellin transform from (5.2) (DLMF 15.6.6). Then we derive the integral and series representations from it.

DEFINITION 5.5. The Mellin transform of the fractional Gauss hypergeometric function is

$$(5.7) \quad \begin{aligned} {}_2F_1(a, b, c, \epsilon; -x) &\stackrel{\mathcal{M}}{\longleftrightarrow} {}_2F_1^*(a, b, c, \epsilon; s) \\ &= \left[M^*(a, c; s) \right] \left[\frac{B(k/2, 1/2)}{\Gamma(1/2)} \hat{\chi}_{\alpha, k}^{2*}(3/2 - s) \right] \end{aligned}$$

where $\epsilon = 1/\alpha$ is the convention from (4.2), and $b = (k + 1)/2$. $M^*(a, c; s)$ is from (5.1), and $\hat{\chi}_{\alpha, k}^{2*}(s)$ is from (7.26) (we jump ahead). And $B(x, y)$ is the beta function.

This structure is a fractional form of the generalized hypergeometric function ${}_3F_2$ (DLMF 16.5.1, replace s with $-s$). To see this, expand (5.7) and we get

$$(5.8) \quad {}_2F_1^*(a, b, c, \epsilon; s) = \left[\frac{\Gamma(c)}{\Gamma(a)} \frac{\Gamma(a-s)\Gamma(s)}{\Gamma(c-s)} \right] \left[2^{2s-1} \frac{B(k/2, 1/2)}{\Gamma(1/2)} \frac{\Gamma((k-1)/2)}{\Gamma(\epsilon(k-1))} \frac{\Gamma(2\epsilon(k/2-s))}{\Gamma(k/2-s)} \right].$$

There are five gamma functions that contain s : three in the numerator, two in the denominator. And the $\Gamma(2\epsilon(k/2-s))$ term is fractional.

5.2.1. FGHF Subsumes the Gauss Hypergeometric Function.

LEMMA 5.6. When $\epsilon = 1$,

$${}_2F_1^*(a, b, c, \epsilon = 1; s) = {}_2F_1^*(a, b, c; s)$$

△

PROOF. Let $\epsilon = 1$, the second bracket becomes

$$(5.9) \quad \frac{B(k/2, 1/2)}{\Gamma(1/2)} \frac{\Gamma(k/2 + 1/2 - s)}{\Gamma(k/2)} = \frac{\Gamma(k/2 + 1/2 - s)}{\Gamma(k/2 + 1/2)} = \frac{\Gamma(b-s)}{\Gamma(b)}.$$

Hence, (5.7) is reduced to the classic limit of ${}_2F_1^*(a, b, c; s)$ in (5.2). □

5.2.2. The Integral Form.

LEMMA 5.7. The integral form of FGHF is

$$(5.10) \quad {}_2F_1(a, b, c, \epsilon; -x) := \frac{B(k/2, 1/2)}{\Gamma(1/2)} \int_0^\infty M(a, c; -x\nu) \hat{\chi}_{\alpha, k}^2(\nu) \sqrt{\nu} d\nu$$

where $\epsilon = 1/\alpha$ and $b = (k+1)/2$. $M(a, c; x)$ is the Kummer function (Chapter 13, DLMF). $\hat{\chi}_{\alpha, k}^2(x)$ is from (7.18). △

PROOF. We use the generalized convolution formula:

$$h(x) = \int_0^\infty f(xs)g(s) s^p ds \xleftrightarrow{\mathcal{M}} h^*(s) = f^*(s)g^*(1+p-s),$$

Clearly f is M , and g is $\hat{\chi}_{\alpha, k}^2$. Substitute $p = 1/2$ due to the $\sqrt{\nu}$ term. The Mellin transform of (5.10) is

$${}_2F_1(a, b, c, \epsilon; -x) \xleftrightarrow{\mathcal{M}} \frac{B(k/2, 1/2)}{\Gamma(1/2)} M^*(a, c; s) \hat{\chi}_{\alpha, k}^{2*}(3/2 - s)$$

This is exactly (5.7). □

5.2.3. Relation between FGHF and Real-World Usage. This section explains how FGHF connects to the later FCM, GAS, and GAS-SN constructions. The connection is most transparent at the level of ratio-distribution integrals: many of the univariate PDFs used later can be viewed as specializations of the FGHF integral representation.

To isolate this structure, suppress the parameters that are not needed for the present comparison. Define $F(-x) := {}_2F_1(a, b, c, \epsilon; -x)$ and $M(-x) := M^*(a, c; -x)$ for $x \geq 0$. Then (5.10) becomes

$$(5.11) \quad F(-x) := B \int_0^\infty M(-x\nu) \bar{\chi}_{\alpha, k}^2(\nu; \sigma = 1/4) \sqrt{\nu} d\nu$$

where we employ the notation $\bar{\chi}_{\alpha, k}^2(x) = \bar{\chi}_{\alpha, k}^2(x; \sigma = \frac{1}{4})$ from (7.18), and $B := B(\frac{k}{2}, \frac{1}{2})/\Gamma(\frac{1}{2})$.

LEMMA 5.8. Let $F'(-x)$ denote the scaled FGHF used in the later ratio-distribution applications. The following ratio-distribution integrals can be expressed in terms of F' :

$$(5.12) \quad \left\{ \begin{array}{l} \int_0^\infty M(-xs) \bar{\chi}_{\alpha, k}^2(s) \sqrt{s} ds \\ \int_0^\infty M(-xs^2) \bar{\chi}_{\alpha, k}(s) s ds \end{array} \right\} = F'(-x) := \frac{2\sqrt{\pi} \sigma_{\alpha, k}}{B(\frac{k}{2}, \frac{1}{2})} F(-4\sigma_{\alpha, k}^2 x)$$

Equivalently, in the full FGHF notation,

$$(5.13) \quad \left\{ \begin{array}{l} \int_0^\infty M(a, c; -xs) \bar{\chi}_{\alpha, k}^2(s) \sqrt{s} ds \\ \int_0^\infty M(a, c; -xs^2) \bar{\chi}_{\alpha, k}(s) s ds \end{array} \right\} = F'_{\alpha, k}(a, c; -x) := \frac{2\sqrt{\pi} \sigma_{\alpha, k}}{B(\frac{k}{2}, \frac{1}{2})} {}_2F_1(a, b, c, \epsilon; -4\sigma_{\alpha, k}^2 x)$$

where $\epsilon = 1/\alpha$ and $b = (k+1)/2$ on the RHS. △

PROOF. Let Q be the scale that we want to solve. (5.11) is rewritten to $F'(-x)$ such that

$$F'(-x) := \frac{\sqrt{Q}}{B} F(-Qx) = \sqrt{Q} \int_0^\infty M(-Qx\nu) \bar{\chi}_{\alpha, k}^2(\nu; \sigma = 1/4) \sqrt{\nu} d\nu.$$

Let $s = Q\nu$,

$$\begin{aligned} F'(-x) &= \int_0^\infty M(-xs) \bar{\chi}_{\alpha,k}^2(s/Q; \sigma = 1/4)/Q \sqrt{s} ds \\ &= \int_0^\infty M(-xs) \bar{\chi}_{\alpha,k}^2(s; \sigma = Q/4) \sqrt{s} ds \end{aligned}$$

Let $Q = 4\sigma_{\alpha,k}^2$, we obtain the integral form in terms of FCM2,

$$F'(-x) = \int_0^\infty M(-xs) \bar{\chi}_{\alpha,k}^2(s) \sqrt{s} ds$$

This is the first line of (5.12). Then apply (7.20) and (7.21) to get the second line. And on the FGHF side, we have

$$F'(-x) = \frac{\sqrt{Q}}{B} F(-Qx) = \frac{2\sqrt{\pi} \sigma_{\alpha,k}}{B(\frac{k}{2}, \frac{1}{2})} F(-4\sigma_{\alpha,k}^2 x)$$

781

□

5.2.4. Example 1: GSaS. In Lemma 8.3 of [19], a fractional extension was explored for the CDF of GSaS. We formalized it further here. However, we note that the $M(-x)$ function needed to describe GAS-SN is more complicated than a Kummer function. See (10.2) and (10.3).

LEMMA 5.9. Assume $\Phi[L_{\alpha,k}](x)$ is the CDF of a GSaS, which is (12.2) with $\beta = 0$. It can be expressed by the scaled FGHF via

$$(5.14) \quad \Phi[L_{\alpha,k}](x) = \frac{1}{2} + \frac{x}{\sqrt{2\pi}} F'_{\alpha,k} \left(\frac{1}{2}, \frac{3}{2}; -\frac{x^2}{2} \right).$$

787

△

PROOF. From Lemma 8.3 of [19],

$$\Phi[L_{\alpha,k}](x) = \frac{1}{2} + \frac{x}{\sqrt{k}} M_{\alpha,k} \left(a, c; -\frac{x^2}{k} \right),$$

where $a = \frac{1}{2}, c = \frac{3}{2}$ and

$$M_{\alpha,k}(a, c; x) := \sqrt{\frac{k}{2\pi}} \int_0^\infty s ds M \left(a, c; \frac{xks^2}{2} \right) \bar{\chi}_{\alpha,k}(s).$$

This pattern fits right in with the second line of (5.13). It is immediately clear that its $M_{\alpha,k}(a, c; x)$ is our $\sqrt{k/2\pi} F'_{\alpha,k}(a, c; kx/2)$. Therefore,

$$\Phi[L_{\alpha,k}](x) = \frac{1}{2} + \frac{x}{\sqrt{2\pi}} F'_{\alpha,k} \left(a, c; -\frac{x^2}{2} \right),$$

where $a = \frac{1}{2}, c = \frac{3}{2}$.

793

□

Notice that this formula is much cleaner, without the cluttering of k in the previous attempt in [19].

795

5.2.5. Example 2: Fractional F.

LEMMA 5.10. From (8.2), the standard CDF of a fractional F distribution $F_{\alpha,d,k}$ is

$$\Phi[F_{\alpha,d,k}](x) = \frac{1}{\Gamma(\frac{d}{2})} \int_0^\infty ds \gamma\left(\frac{d}{2}, \frac{dxs}{2}\right) \bar{\chi}_{\alpha,k}^2(s).$$

It can be expressed by the scaled FGHF via

$$(5.15) \quad \Phi[F_{\alpha,d,k}](x) = \left[C_{\alpha,d,k} \frac{(dx/2)^{d/2}}{\Gamma(\frac{d}{2}+1)} \right] F'_{\alpha,k+d-1}\left(\frac{d}{2}, \frac{d}{2}+1; -\frac{dx}{2\Sigma}\right).$$

where $C_{\alpha,d,k}$ is defined in (5.16) and $\Sigma := \sigma_{\alpha,k+d-1}^2 / \sigma_{\alpha,k}^2$. △

PROOF. Note that

$$\frac{1}{\Gamma(\frac{d}{2})} \gamma\left(\frac{d}{2}, \frac{x}{2}\right) = \frac{(x/2)^{d/2}}{\Gamma(\frac{d}{2}+1)} M\left(\frac{d}{2}, \frac{d}{2}+1; -\frac{x}{2}\right).$$

Then

$$\begin{aligned} \Phi[F_{\alpha,d,k}](x) &= \int_0^\infty \left[\frac{(dxs/2)^{d/2}}{\Gamma(\frac{d}{2}+1)} M\left(\frac{d}{2}, \frac{d}{2}+1; -\frac{dxs}{2}\right) \right] \bar{\chi}_{\alpha,k}^2(s) ds \\ &= \frac{(dx/2)^{d/2}}{\Gamma(\frac{d}{2}+1)} \int_0^\infty M\left(\frac{d}{2}, \frac{d}{2}+1; -\frac{dxs}{2}\right) s^{(d-1)/2} \bar{\chi}_{\alpha,k}^2(s) \sqrt{s} ds. \end{aligned}$$

When $d = 1$, it fits right in with FGHF. When $d > 1$, it needs more work.

From (7.7), let $m = (d-1)/2$, then $k+2m = k+d-1$ and

$$\Phi[F_{\alpha,d,k}](x) = C_{\alpha,d,k} \frac{(dx/2)^{d/2}}{\Gamma(\frac{d}{2}+1)} \int_0^\infty M\left(\frac{d}{2}, \frac{d}{2}+1; -\frac{dxy}{2\Sigma}\right) \bar{\chi}_{\alpha,k+d-1}^2(y) \sqrt{y} dy,$$

where $\Sigma := \sigma_{\alpha,k+d-1}^2 / \sigma_{\alpha,k}^2$ and $y = \Sigma s$, and

$$(5.16) \quad C_{\alpha,d,k} := \frac{\sigma_{\alpha,k}^{d-1}}{\sqrt{\Sigma}} \frac{C_{\alpha,k}}{C_{\alpha,k+d-1}} = \frac{\sigma_{\alpha,k}^d}{\sigma_{\alpha,k+d-1}} \frac{C_{\alpha,k}}{C_{\alpha,k+d-1}}.$$

The integral matches the FGHF pattern in Lemma 5.12, and we get (5.15). □

REMARK 5.11. One final note. There is a connection between (5.14) and (5.15). When $d = 1$, $\Sigma = 1$ and $C_{\alpha,d,k} = 1$. Then

$$(5.17) \quad \Phi[F_{\alpha,1,k}](x^2) = \frac{2x}{\sqrt{2\pi}} F'_{\alpha,k}\left(\frac{1}{2}, \frac{3}{2}; -\frac{x^2}{2}\right)$$

which is $2\Phi[L_{\alpha,k}](x) - 1$ in (5.14).

This is a reflection of Lemma 8.3. If the variable X distributes as a GSaS $L_{\alpha,k}$, then X^2 distributes as a one-dimensional F, aka $F_{\alpha,1,k}$. It is particularly easy to see this relation in the FGHF form above.

Part 2

One-Sided Distributions

FG: Fractional Gamma Distribution and Tilted Stable Law

The fractional gamma distribution (FG) is one of the central one-sided families used throughout this book. It contains the fractional χ -mean distribution (FCM) as a special case and provides a fractional extension of the generalized gamma distribution.

The same family was called *the generalized stable count distribution* in [19], extending the "stable count distribution" terminology introduced in [18]. In the present formulation, the name *fractional gamma distribution* is more natural because the construction is organized around a fractional analogue of the gamma family.

A key result of this chapter is that FG is exactly a reparameterized inverse tilted stable law. This identification provides the probabilistic foundation for FG and for the distributional constructions built from it in later chapters.

6.1. Definition

DEFINITION 6.1 (Fractional Gamma distribution (FG)). FG is a four-parameter one-sided distribution family, whose PDF is defined as

$$(6.1) \quad \mathfrak{N}_\alpha(x; \sigma, d, p) := C \left(\frac{x}{\sigma} \right)^{d-1} F_\alpha \left(\left(\frac{x}{\sigma} \right)^p \right) \quad (x \geq 0)$$

where $F_\alpha(x) = W_{-\alpha,0}(-x) = \alpha x M_\alpha(x)$ from (3.9) and $\alpha \in [0, 1]$ controls the shape of the Wright function; σ is the scale parameter; p is also the shape parameter controlling the tail behavior ($p \neq 0, dp \geq 0$); d is the *degree of freedom* parameter. When $\alpha \rightarrow 1$, the PDF becomes a Dirac delta function: $\delta(x - \sigma)$ assuming σ is finite. When $d \geq 1$, all the moments of the FG exist and have closed forms.

6.2. Determination of C

The normalization constant C is:

$$(6.2) \quad C = \begin{cases} \frac{|p|}{\sigma} \frac{\Gamma(\alpha d/p)}{\Gamma(d/p)} & , \text{ for } \alpha \neq 0, d \neq 0. \\ \frac{|p|}{\sigma \alpha} & , \text{ for } \alpha \neq 0, d = 0. \end{cases}$$

It is important to note that d and p are allowed to be negative, as long as $dp \geq 0$.

PROOF. The normalization constant C in (6.1) is obtained from the requirement that the integral of the PDF must be 1:

$$\int_0^\infty \mathfrak{N}_\alpha(x; \sigma, d, p) dx = \frac{C \sigma}{|p|} \frac{\Gamma(\frac{d}{p})}{\Gamma(\frac{d}{p} \alpha)} = 1$$

where the integral is carried out by the moment formula of the Wright function.

We typically constrain $dp \geq 0$ and p is typically positive. However, it becomes negative in the inverse distribution and/or characteristic distribution types. So we need $|p|$ to ensure that C is positive.

For the case of $\alpha \neq 0$ and $d \rightarrow 0$, due to (A.3), we have

$$C = \frac{|p|}{\sigma\alpha} \quad (\alpha \neq 0, d = 0)$$

These two cases are combined to form (6.2). $\square$

6.3. FG Subsumes Generalized Gamma Distribution

Since the Wright function extends an exponential function to the fractional space, FG is the fractional extension of the generalized gamma (GG) distribution[36], whose PDF is defined as:

$$(6.3) \quad f_{\text{GG}}(x; a, d, p) = \frac{|p|}{a\Gamma(d/p)} \left(\frac{x}{a}\right)^{d-1} e^{-(x/a)^p}.$$

The parallel use of parameters is obvious, except that a in GG is replaced by σ in FG to avoid confusion with α .

GG is subsumed to FG in two ways:

$$(6.4) \quad f_{\text{GG}}(x; \sigma, d, p) := \begin{cases} \mathfrak{N}_0(x; \sigma, d = d - p, p) & , \text{ at } \alpha = 0. \\ \mathfrak{N}_{\frac{1}{2}}(x; \sigma = \frac{\sigma}{2^{2/p}}, d = d - \frac{p}{2}, p = \frac{p}{2}) & , \text{ at } \alpha = \frac{1}{2}. \end{cases}$$

The first line is treated as the definition of FG at $\alpha = 0$. The proof is given in [19].

Although the first line is more obvious, it is the second line that leads to the fractional extension of the χ distribution.

6.4. Mellin Transform

From Example 2.4, we add σ and C . The Mellin transform of the PDF of the fractional gamma distribution is

$$(6.5) \quad \begin{aligned} \mathfrak{N}_\alpha(x; \sigma, d, p) &\stackrel{\mathcal{M}}{\longleftrightarrow} \frac{C \sigma^s}{|p|} \frac{\Gamma((s + d - 1)/p)}{\Gamma(\alpha(s + d - 1)/p)} \\ &= \sigma^{s-1} \frac{\Gamma(\alpha d/p)}{\Gamma(d/p)} \frac{\Gamma((s + d - 1)/p)}{\Gamma(\alpha(s + d - 1)/p)}, \end{aligned}$$

where C is from Section 6.2. The typical limiting case for the gamma functions shall be taken care in each scenario.

Substituting $s = n + 1$ into (6.5), we obtain the moment formula of the FG distribution,

$$(6.6) \quad \mathbb{E}[X^n] = \sigma^n \frac{\Gamma(\alpha d/p)}{\Gamma(d/p)} \frac{\Gamma((n + d)/p)}{\Gamma(\alpha(n + d)/p)}.$$

FG is often used in a ratio distribution, such as the role of $g^*(s)$ in (2.6), where $s \rightarrow 2 - s$. The term $s + d - 1$ becomes $d + 1 - s$. Furthermore, in the FCM case, since $d = k - 1$, it becomes the elegant $k - s$ term.

6.5. The Inverse Tilted Stable Law

FG is the foundation for all the distributions that follow in this book. It is beautifully rooted in the polynomially tilted inverse stable distribution[7, 14]. We will show that the FG random variable is exactly the re-parametrized *inverse tilted stable variable*.

We use the T notation from [7] where $T_{\alpha,\beta}$ is the polynomially tilted stable variable. Its density function is $\propto x^{-\beta} L_\alpha(x), x > 0$. β is the *tilted angle* due to polynomial bias.¹

It is generalized by adding the third parameter γ so that $T_{\alpha,\beta}^{-\gamma} = (T_{\alpha,\beta})^{-\gamma}$. The inverse power of γ transforms the density function to a polynomially tilted M-Wright function, which will be proved

¹It will be shown in Section 7.6 that $\chi_{\alpha,-k}^2$ is equal to $T_{\alpha/2,k/2}$.

in the Lemma 6.3. We might use the notion $TS(\alpha, \beta, \gamma)$ to avoid collisions between greeks since β, γ might have different meanings in other contexts later in the book.

Let $X \sim \mathfrak{N}_\alpha(\sigma, d, p)$ be the FG random variable. The conclusion is $\beta = \alpha d/p, \gamma = \alpha/p$. That is,

$$(6.7) \quad X = \sigma T_{\alpha, \alpha d/p}^{-\alpha/p}.$$

873

The calculation of $T_{\alpha, \beta}$ is carried out in Theorem 1 of [7]. We reproduce the formula for the more familiar $U = T_{\alpha, \beta}^{-\alpha}$ in the following way.

875

LEMMA 6.2. For the tilted stable law $T_{\alpha, \beta}$, the sampling is performed via the modified Kanter's method,

$$T_{\alpha, \beta} = A_\alpha(Q_\beta) G_\beta^{-c}, \quad c = \frac{1 - \alpha}{\alpha}.$$

876

- (1) The factor $A_\alpha(q)$ is the Zolotarev function from (3.22).
- (2) Q_β is the sampling of A_α with angle β , whose density is

$$f_{Q_\beta}(q) = I_\beta A_\alpha(q)^{-\beta}, \quad \text{where } I_\beta = \frac{\Gamma(1 + \beta)\Gamma(1 + c\beta)}{\Gamma(1 + \beta/\alpha)}, \quad 0 < q < 1.$$

877

When $\beta = 0$, Q_0 is uniformly distributed on $(0, 1)$.

- (3) G_β is a gamma variable

$$G_\beta \sim \text{Gamma}(1 + c\beta, \text{scale} = 1).$$

878

Then

$$(6.8) \quad U = T_{\alpha, \beta}^{-\alpha} = A_\alpha(Q_\beta)^{-\alpha} G_\beta^{1-\alpha}.$$

879

△

The case of $\beta = 0$ is added because it is an interesting special case: $T_{\alpha, 0}^{-\alpha} = M_\alpha$ where M_α represents the M-Wright function treated as a one-sided distribution.

880

Moving from U to X is straightforward,

$$(6.9) \quad X = \sigma U^{1/p}$$

883

LEMMA 6.3. For FG, the corresponding tilted stable parameters are $\beta = \alpha d/p$ and $\gamma = \alpha/p$.

884

△

PROOF. It follows immediately from (6.7) that $\gamma = \alpha/p$. From [7, 14], $U = T_{\alpha, \beta}^{-\alpha}$ has density

$$f_U(u) = J_\beta u^{\beta/\alpha} M_\alpha(u) = \frac{J_\beta}{\alpha} u^{\beta/\alpha - 1} F_\alpha(u), \quad \text{where } J_\beta = \frac{\Gamma(\beta + 1)}{\Gamma(\beta/\alpha + 1)}.$$

Let

$$u = \left(\frac{x}{\sigma}\right)^p.$$

Then

$$\left|\frac{du}{dx}\right| = \frac{|p|}{\sigma} \left(\frac{x}{\sigma}\right)^{p-1}.$$

885

Hence the density of $X = \sigma U^{1/p}$ is

$$\begin{aligned} f_X(x) &= f_U(u) \left|\frac{du}{dx}\right| \\ &= \frac{|p|J_\beta}{\alpha\sigma} \left(\frac{x}{\sigma}\right)^{p\beta/\alpha - 1} F_\alpha\left(\left(\frac{x}{\sigma}\right)^p\right). \end{aligned}$$

Comparing this with (6.1), we obtain

$$\frac{p\beta}{\alpha} - 1 = d - 1,$$

hence

$$\beta = \frac{\alpha d}{p}.$$

886 The normalization constant is also consistent, because

$$\begin{aligned} \frac{|p|J_\beta}{\alpha\sigma} &= \frac{|p|}{\sigma} \frac{\Gamma(\beta)}{\Gamma(\beta/\alpha)} \\ &= \frac{|p|}{\sigma} \frac{\Gamma(\alpha d/p)}{\Gamma(d/p)}, \end{aligned}$$

887 which agrees with (6.2). The case $d = 0$ follows by the same limiting argument used in Section 6.2.

888

□

889 The numeric sampler implements the logarithmic form:

$$(6.10) \quad \log U = -\alpha \log T_{\alpha,\beta} = (1 - \alpha) \log G_\beta - \alpha \log A_\alpha(Q_\beta).$$

890 Then

$$(6.11) \quad \begin{aligned} \log X &= -\gamma \log T_{\alpha,\beta} = \log \sigma + c\gamma \log G_\beta - \gamma \log A_\alpha(Q_\beta) \\ \text{where } \beta &= \frac{\alpha d}{p}, \gamma = \frac{\alpha}{p}, c = \frac{1 - \alpha}{\alpha}. \end{aligned}$$

891 6.6. CDF and Fractional Incomplete Gamma Function

892 The CDF of FG is

$$(6.12) \quad \Phi(x) := \int_0^x \mathfrak{N}_\alpha(s; \sigma, d, p) ds \quad (x \geq 0).$$

893 This integral leads to fractionalization of the incomplete gamma function in Section 3.5.

894 LEMMA 6.4. The CDF of FG can be represented by γ_α^* in (3.23) or Γ_α^* in (3.24) as

$$(6.13) \quad \Phi(x) = \begin{cases} z^{d+p} \gamma_\alpha^*(d/p + 1, z^p) & , \text{ when } p > 0. \\ z^{d+p} \Gamma_\alpha^*(d/p + 1, z^p) & , \text{ when } p < 0. \end{cases}$$

895 where $z = x/\sigma$ is the standardized variable.

896 This could be viewed as one form of fractional extension to the regularized lower incomplete gamma
897 function, $\gamma(s, z)/\Gamma(s)$, which is the CDF of GG mentioned above.

898 Due to this result, this distribution is called the *fractional gamma distribution* (FG).

899

△

900 PROOF. The CDF of FG is

$$\begin{aligned} \Phi(x) &= \int_0^x \mathfrak{N}_\alpha(s; \sigma, d, p) ds \\ &= C \int_0^x ds \left(\frac{s}{\sigma}\right)^{d-1} W_{-\alpha,0} \left(-\left(\frac{s}{\sigma}\right)^p\right). \\ &= C \int_0^x ds \left(\frac{s}{\sigma}\right)^{d-1} F_\alpha \left(\left(\frac{s}{\sigma}\right)^p\right). \end{aligned}$$

901 Since $F_\alpha(x) = \alpha x M_\alpha(x)$ from (3.9), and let $u = s/x$, then $u \in [0, 1]$ and

$$\begin{aligned}\Phi(x) &= \alpha C \int_0^x ds \left(\frac{s}{\sigma}\right)^{d+p-1} M_\alpha\left(\left(\frac{s}{\sigma}\right)^p\right) \\ &= \alpha C x \int_0^1 du \left(\frac{xu}{\sigma}\right)^{d+p-1} M_\alpha\left(\left(\frac{xu}{\sigma}\right)^p\right)\end{aligned}$$

902 Recognize that $u^p \in [0, 1]$ when $p > 0$. Let $t = u^p$, and $dt/t = p du/u$,

$$\Phi(x) = \frac{\alpha \sigma C}{p} z^{d+p} \int_0^1 dt t^{d/p} M_\alpha(z^p t)$$

903 Compare the last line with γ_α^* in (3.23), and we get

$$\Phi(x) = \frac{\alpha \sigma C}{p} \frac{\Gamma(\frac{d}{p} + 1)}{\Gamma((1 - \alpha) + \alpha(\frac{d}{p} + 1))} z^{d+p} \gamma_\alpha^*(d/p + 1, z^p)$$

904 Using the case of $\alpha \neq 0, d \neq 0$ for C , it can be shown that the constant part is just 1. Hence,

$$\Phi(x) = z^{d+p} \gamma_\alpha^*(d/p + 1, z^p), \quad \text{for } p > 0.$$

905 On the other hand, $u^p \in [1, \infty]$ when $p < 0$. The range of the integral changes to $\int_1^\infty$. It leads to
906 the use of the supplementary function (3.24):

$$\Phi(x) = z^{d+p} \Gamma_\alpha^*(d/p + 1, z^p).$$

907

□

908 Numerically, it is better to use γ_α^* when $z^p \leq 1$ and to use Γ_α^* when $z^p > 1$, as long as the identity
909 relation (3.25) is preserved.

910 6.7. Inverse Expression of Several Fractional Distributions

911 Several known fractional distributions could be expressed in the FG in Table 1. This shows that
912 the FG is the super set of the one-sided fractional distribution system. Its parametrization provides
913 immense flexibility to express other formerly known one-sided distributions.

Distribution (PDF)	Wright Equiv.	FG: $\mathfrak{N}_\alpha(x; \sigma, d, p)$			
		α	σ	d	p
One-sided stable: $L_\alpha(x)$	$x^{-1} W_{-\alpha, 0}(-x^{-\alpha})$	α	1	0	$-\alpha$
Stable Count: $\mathfrak{N}_\alpha(x)$		α	1	1	α
Stable Vol: $V_\alpha(x)$		$\frac{\alpha}{2}$	$\frac{1}{\sqrt{2}}$	1	α
M-Wright: $M_\alpha(x)$	$\frac{1}{\alpha x} W_{-\alpha, 0}(-x)$	α	1	0	1
M-Wright II: $\Gamma(\alpha) F_\alpha(x)$	$\Gamma(\alpha) W_{-\alpha, 0}(-x)$	α	1	1	1

TABLE 1. FG mapping of several known fractional distributions in the literature. $\mathfrak{N}_\alpha(x)$ and $V_\alpha(x)$ first appeared in [18], which led to this work.

6.8. Alternative Definition

DEFINITION 6.5. It is reasonable to argue that the PDF of FG can be defined via the M-Wright function directly, such that

$$(6.14) \quad \mathfrak{N}'_{\alpha}(x; \sigma, d', p) := C' \left(\frac{x}{\sigma} \right)^{d'-1} M_{\alpha} \left(\left(\frac{x}{\sigma} \right)^p \right). \quad (x \geq 0)$$

However, since $F_{\alpha}(z) = \alpha z M_{\alpha}(z)$, it is easy to see that

$$\mathfrak{N}'_{\alpha}(x; \sigma, d', p) = \alpha C' \left(\frac{x}{\sigma} \right)^{d'+p-1} F_{\alpha} \left(\left(\frac{x}{\sigma} \right)^p \right).$$

Therefore, this is merely a reparameterization of $d = d' + p$. This definition will encounter some issues in FCM later due to the assignment of $d \rightarrow k - 1$, $\alpha \rightarrow \alpha/2$ and $p \rightarrow \alpha$ (see (7.4)). We learn from Figures 12.1 and 12.2 that there is a natural linear relation between k and $\epsilon = 1/\alpha$. Mixing the role of d with α from p is not a good idea.

Fractional Chi Distributions

7.1. Introduction to Fractional Chi Distribution

Chapter 4 showed that the Mellin transform of the α -stable density can be interpreted as the transform of a ratio distribution. This observation motivates the fractional χ construction. The two factors are

$$L_{\alpha}^{\theta}(x) \xleftrightarrow{\mathcal{M}} \epsilon \left[\frac{\Gamma(s)}{\Gamma(1 - \gamma + \gamma s)} \right] \left[\frac{\Gamma(\epsilon(1 - s))}{\Gamma(\gamma(1 - s))} \right]$$

$$\text{where } \epsilon = \frac{1}{\alpha}, \gamma = \frac{\alpha - \theta}{2\alpha}.$$

The first bracket is the Mellin transform of the M-Wright function. The second bracket is interpreted as the Mellin transform of the fractional χ -mean distribution (FCM) at $k = 1$:

$$\bar{\chi}_{\alpha,1}^{\theta}(x) \xleftrightarrow{\mathcal{M}} \bar{\chi}_{\alpha,1}^{\theta *} (s) \propto \frac{\Gamma(\epsilon(s - 1))}{\Gamma(\gamma(s - 1))},$$

up to the normalization constant and scale in the PDF.

The connection follows from the Mellin-transform rule for ratio distributions: the transform of the denominator is evaluated at $2 - s$. Thus the term $s - 1$ in $\bar{\chi}_{\alpha,1}^{\theta *} (s)$ becomes $1 - s$ in the stable-density factor above. In this chapter, the degrees-of-freedom parameter k is introduced by replacing $s - 1$ with $s + k - 2$, giving

$$(7.1) \quad \bar{\chi}_{\alpha,k}^{\theta}(x) \xleftrightarrow{\mathcal{M}} \bar{\chi}_{\alpha,k}^{\theta *} (s) \propto \frac{\Gamma(\epsilon(s + k - 2))}{\Gamma(\gamma(s + k - 2))}.$$

This prototype is the starting point for the full definition of FCM.

7.2. FCM: Fractional Chi-Mean Distribution

There are two ways to define FCM. The first approach is to define it via Mellin transform. The second approach is to define the shape of its PDF.

DEFINITION 7.1 (Fractional χ -mean distribution (FCM) via Mellin Transform). The Mellin transform of FCM's PDF is enriched from (7.1) to

$$(7.2) \quad \bar{\chi}_{\alpha,k}^{\theta}(x) \xleftrightarrow{\mathcal{M}} \bar{\chi}_{\alpha,k}^{\theta *} (s)$$

$$= (\sigma_{\alpha,k}^{\theta})^{s-1} \frac{\Gamma(\gamma(k - 1))}{\Gamma(\epsilon(k - 1))} \frac{\Gamma(\epsilon(s + k - 2))}{\Gamma(\gamma(s + k - 2))},$$

$$\text{where } \sigma_{\alpha,k}^{\theta} := \gamma^{\gamma} k^{\gamma - \epsilon}.$$

The main differences are (1) to address the normalization of the total density, and (2) to have a proper scale $\sigma_{\alpha,k}^{\theta}$ such that it is consistent with the classic χ distribution and α -stable distribution.

For positive k , the PDF of an FCM is

$$(7.3) \quad \bar{\chi}_{\alpha,k}^{\theta}(x) := \mathfrak{N}_{\gamma\alpha}(x; \sigma = \sigma_{\alpha,k}^{\theta}, d = k-1, p = \alpha) \quad (x \geq 0)$$

$$= \frac{\Gamma(\gamma(k-1))}{\epsilon\Gamma(\epsilon(k-1))} (\sigma_{\alpha,k}^{\theta})^{1-k} x^{k-2} F_{\gamma\alpha} \left(\left(\frac{x}{\sigma_{\alpha,k}^{\theta}} \right)^{\alpha} \right),$$

where $\mathfrak{N}_{\lambda}(x; \sigma, d, p)$ is FG (6.1), and $F_{\lambda}(x) := W_{-\lambda,0}(-x) = \lambda x M_{\lambda}(x)$ is the Wright function of the second kind (3.9).

Notice the appearances of γ that replaces all the $1/2$ in Section 7.6 of [19]. That is how θ comes into play in the upgraded FCM. This full representation is used in Chapter 11.

However, for GAS-SN in Chapter 12 and beyond, such θ upgrade is unnecessary. The skew-normal framework is based on modulation of normal distributions. It is required to have $\theta = 0$, aka $\gamma = 1/2$.

Hence, we recite the original definition of FCM PDF ($k > 0$):

$$(7.4) \quad \bar{\chi}_{\alpha,k}(x) = \bar{\chi}_{\alpha,k}^0(x) := \mathfrak{N}_{\alpha/2}(x; \sigma = \sigma_{\alpha,k}, d = k-1, p = \alpha) \quad (x \geq 0)$$

$$= (C_{\alpha,k}) (\sigma_{\alpha,k})^{1-k} x^{k-2} F_{\frac{\alpha}{2}} \left(\left(\frac{x}{\sigma_{\alpha,k}} \right)^{\alpha} \right),$$

where

$$(7.5) \quad C_{\alpha,k} := \frac{\alpha\Gamma((k-1)/2)}{\Gamma((k-1)/\alpha)}$$

$$(7.6) \quad \sigma_{\alpha,k} := \frac{|k|^{1/2-1/\alpha}}{\sqrt{2}}.$$

Note that the difference between (7.3) and (7.4) is very small: Just replace $\mathfrak{N}_{\gamma\alpha}(\dots)$ to $\mathfrak{N}_{\alpha/2}(\dots)$.

7.2.1. Inverse Tilted Stable Law. According to Section 6.5, (7.4) implies $\bar{\chi}_{\alpha,k} = \sigma_{\alpha,k} T_{\alpha/2, (k-1)/2}^{-1/2}$. The special case for stable distribution decomposition (4.11) is $\bar{\chi}_{\alpha,1}^{\theta} = \gamma^{\gamma} T_{\gamma\alpha,0}^{-\gamma}$ and $\bar{\chi}_{\alpha,1} = T_{\alpha/2,0}^{-1/2} / \sqrt{2}$.

7.2.2. FCM CDF. Extending directly from Lemma 6.4, we have

LEMMA 7.2. The CDF of FCM can be represented by γ_{α}^* in (3.23) as

$$(7.7) \quad \Phi[\bar{\chi}_{\alpha,k}](x) = z^{k-1+\alpha} \gamma_{\alpha/2}^* \left(\frac{k-1+\alpha}{\alpha}, z^{\alpha} \right), \quad (k > 0, \alpha \in [0, 2])$$

where $z = x/\sigma_{\alpha,k}$.

△

7.2.3. FCM Subsumes Chi. It is easy to show that $\bar{\chi}_{1,k}$ is χ_k with a scale of $1/\sqrt{k}$, due to (3.15).

7.3. FCM Moments

By letting $s = n+1$ and $\theta = 0$ in (7.2), its n -th moment is

$$(7.8) \quad \mathbb{E}(X^n | \bar{\chi}_{\alpha,k}) = (\sigma_{\alpha,k})^n \frac{\Gamma((k-1)/2)}{\Gamma((k-1)/\alpha)} \frac{\Gamma((n+k-1)/\alpha)}{\Gamma((n+k-1)/2)}, \quad (k > 0, \alpha > 0)$$

which requires $k > 1$ and $n+k > 1$ to avoid the singularity of the gamma functions (See Section 7.6 of [19]).

966 The explicit form of the first moment is

$$(7.9) \quad \mathbb{E}(X|\bar{\chi}_{\alpha,k}) = \sigma_{\alpha,k} \frac{\Gamma((k-1)/2)}{\Gamma((k-1)/\alpha)} \frac{\Gamma(k/\alpha)}{\Gamma(k/2)} = \sigma_{\alpha,k} \frac{C_{\alpha,k}}{C_{\alpha,k+1}}. \quad (k > 0, \alpha > 0)$$

967 Notice that it can be used as a bridge connecting the coefficient C between k and $k+1$.

968 The moment formula of FCM is fundamental to all the fractional distributions built on top of it.
969 However, ironically, due to the nature of a ratio distribution, it is often evaluated as negative moments
970 $n < 0$. Hence, n is restricted in the range of $1 - k < n < 0$.

971 This results in non-existing moments when k is not "large enough", which happens to be a core
972 feature of the α -stable distribution and Student's t distribution. Our two-dimensional parameter space
973 (α, k) adds more complexity to it.

974 **7.3.1. FCM at Infinite Degrees of Freedom.** The choice of $\sigma_{\alpha,k}$ is intentional, such that

$$(7.10) \quad \lim_{k \rightarrow \infty} \mathbb{E}(X^n|\bar{\chi}_{\alpha,k}) = \alpha^{-n/\alpha}. \quad (k > 0, \alpha > 0)$$

975 Under such conditions, its variance is zero. That is, FCM becomes a delta function, $\delta(x - \alpha^{-1/\alpha})$,
976 as $k \rightarrow \infty$.

977 7.4. FCM Reflection Formula and Negative k

978 **7.4.1. FCM for Negative k .** We quote Definition 3.2 of [19] for FCM in the negative k space.
979 Its PDF defined by FG is

$$(7.11) \quad \bar{\chi}_{\alpha,-k}(x) := \mathfrak{N}_{\alpha/2}(x; \sigma = (\sigma_{\alpha,k})^{-1}, d = -k, p = -\alpha). \quad (x \geq 0, k > 0)$$

980 This is defined as the *characteristic FCM* discussed in Chapter 9. That is, $[\bar{\chi}_\phi]_{\alpha,k} := \bar{\chi}_{\alpha,-k}$ in Lemma
981 9.6. It is derived from the properties of the α -stable characteristic function.

982 **7.4.2. Tilted Stable Law.** According to Section 6.5, (7.11) implies $\bar{\chi}_{\alpha,-k} = (\sigma_{\alpha,k})^{-1} T_{\alpha/2, k/2}^{1/2}$.
983 Notice that the superscript $(-\gamma = 1/2)$ is positive. The distribution is based on L_α , instead of M_α .

984 The explicit PDF derivation is documented in Section 9.4. We quote the inverse relation:

$$(7.12) \quad \bar{\chi}_{\alpha,-k}(x) = \frac{x^{-3}}{\mathbb{E}(X|\bar{\chi}_{\alpha,k})} \bar{\chi}_{\alpha,k}\left(\frac{1}{x}\right). \quad (x \geq 0, k > 0)$$

985 which is proved in (9.18) in Section 9.4.

986 $\bar{\chi}_{\alpha,-k}$ is used to define the fractional exponential power distribution within the GSaS (and GAS-
987 SN) nomenclature, which subsumes the exponential power distribution (Section 3.11.1 of [28]). See
988 Section 12.7. Readers interested in more details are referred to the FCM sections in [19].

989 We quote the FCM reflection formula from Section 7 of [19] to summarize the relation:

$$(7.13) \quad \mathbb{E}(X^n|\bar{\chi}_{\alpha,-k}) = \frac{\mathbb{E}(X^{-n+1}|\bar{\chi}_{\alpha,k})}{\mathbb{E}(X|\bar{\chi}_{\alpha,k})}, \quad k > 0.$$

990 This indicates an elegant relation for the first moment that $\mathbb{E}(X|\bar{\chi}_{\alpha,-k}) = 1/\mathbb{E}(X|\bar{\chi}_{\alpha,k})$.

991 7.5. FCM: Lognormal Asymptotics at Infinite Degrees of Freedom

992 **DEFINITION 7.3** (The Lognormal Distribution). Let $Z \sim \mathcal{N}(0, \sigma^2)$, then $X \sim e^Z$ is the lognormal
993 distribution family, denoted as Lognormal(0, σ^2), where σ is the shape parameter.

994 **LEMMA 7.4.** Let $V \sim \bar{\chi}_{\alpha,k}$, $\epsilon = 1/\alpha$, and

$$T := \frac{\epsilon - \frac{1}{2}}{z}, \quad \text{where } z := \begin{cases} k - 1, & k > 0, \\ |k|, & k < 0. \end{cases}$$

995 Let $\alpha \rightarrow 0$ and $|k| \rightarrow \infty$ with T fixed, i.e. $\alpha \approx 1/(T|k|)$. Then

$$\log V - \mathbb{E}[\log V] \xrightarrow{d} \mathcal{N}(0, T), \quad \text{equivalently} \quad \frac{V}{e^{\mathbb{E}[\log V]}} \xrightarrow{d} \text{Lognormal}(0, T).$$

996 The fractional χ family contains the lognormal *in its closure, not as a member*. Both branches of
 997 k reach the same limit. $\triangle$

998 PROOF. From (7.2), with $\theta = 0$ and hence $\gamma = 1/2$, write the log moment generating function of
 999 $\log V$ as

$$\log \mathbb{E}(V^s) = \zeta s \log \sigma_{\alpha, k} + A(z + \zeta s) - A(z), \quad A(x) = \log \Gamma(\epsilon x) - \log \Gamma(x/2),$$

1000 where $\zeta = 1$ on the $k > 0$ branch and $\zeta = -1$ on the $k < 0$ branch. If $W := \log V - \mathbb{E}[\log V]$, then its
 1001 log moment generating function is

$$\log \mathbb{E}(e^{sW}) = \log \mathbb{E}(V^s) - s \mathbb{E}[\log V].$$

1002 Since $\mathbb{E}[\log V]$ is the derivative of $\log \mathbb{E}(V^s)$ at $s = 0$, this subtraction removes the first-order term in
 1003 s . Thus only the cumulants of order $n \geq 2$ remain.

1004 For $n \geq 2$, the cumulants of the centered variable $\log V - \mathbb{E}[\log V]$ are

$$\kappa_n = \zeta^n \left[\epsilon^n \psi^{(n-1)}(\epsilon z) - 2^{-n} \psi^{(n-1)}(z/2) \right].$$

1005 Here $\psi^{(m)}$ denotes the m -th polygamma function, i.e. the m -th derivative of $\psi(x) = \frac{d}{dx} \log \Gamma(x)$. Along
 1006 the path $\epsilon = 1/2 + Tz$, the standard polygamma expansion gives

$$\kappa_2 = T + O(z^{-2}), \quad \kappa_n = O(z^{2-n}) \quad (n \geq 3).$$

1007 Thus all higher cumulants vanish while the variance converges to T , so

$$\log V - \mathbb{E}[\log V] \xrightarrow{d} \mathcal{N}(0, T).$$

1008 Exponentiating gives the stated lognormal limit by the continuous mapping theorem. At finite z , the
 1009 third cumulant remains nonzero, with standardized log-skewness $\mp 1/(\sqrt{T} z) + O(z^{-2})$, so the lognormal
 1010 appears only as a limiting member of the closure. $\square$

1012 Centering term. The centering constant is explicit. With the same branch sign ζ as in the proof,

$$\mathbb{E}[\log V] = \zeta \left[\log \sigma_{\alpha, k} + \epsilon \psi(\epsilon z) - \frac{1}{2} \psi(z/2) \right], \quad \log \sigma_{\alpha, k} = \left(\frac{1}{2} - \epsilon \right) \log |k| - \frac{1}{2} \log 2,$$

1013 where ψ is the digamma function. Along the lognormal path $\epsilon = \frac{1}{2} + Tz$, this gives

$$\mathbb{E}[\log V] = \zeta \epsilon \log \epsilon + O(1).$$

1014 Thus the mean of $\log V$ drifts to $+\infty$ on the positive- k branch and to $-\infty$ on the negative- k branch,
 1015 which is why the lognormal limit is stated after centering by $\mathbb{E}[\log V]$.

7.6. FCM2: Fractional Chi-Squared-Mean Distribution

If $Z \sim \bar{\chi}_{\alpha,k}$, then $X \sim Z^2$ is FCM2, denoted as $X \sim \bar{\chi}_{\alpha,k}^2$. This is the fractional extension of the classic χ_k^2/k , which is subsumed by it at $\alpha = 1$.

$\bar{\chi}_{\alpha,k}^2$ is used in the fractional F distribution in the area of the squared variable and the quadratic form in the multivariate elliptical distribution.

DEFINITION 7.5. The PDF of FCM2 is

$$(7.14) \quad \bar{\chi}_{\alpha,k}^2(x) = \frac{1}{2\sqrt{x}} \bar{\chi}_{\alpha,k}(\sqrt{x}) \quad (x \geq 0, \alpha \in [0, 2])$$

Expressed in FG and (7.4), it is

$$(7.15) \quad \begin{aligned} \bar{\chi}_{\alpha,k}^2(x) &:= \mathfrak{N}_{\alpha/2}(x; \sigma = \sigma_{\alpha,k}^2, d = (k-1)/2, p = \alpha/2) \\ &= \frac{C_{\alpha,k}}{2\sigma_{\alpha,k}^2} \left(\frac{x}{\sigma_{\alpha,k}^2} \right)^{k/2-3/2} F_{\frac{\alpha}{2}} \left(\left(\frac{x}{\sigma_{\alpha,k}^2} \right)^{\alpha/2} \right). \end{aligned} \quad (k > 0)$$

Or for the negative degree of freedom,

$$(7.16) \quad \bar{\chi}_{\alpha,-k}^2(x) := \mathfrak{N}_{\alpha/2}(x; \sigma = \sigma_{\alpha,k}^{-2}, d = -k/2, p = -\alpha/2)$$

7.6.1. Inverse Tilted Stable Law. According to Section 6.5, (7.15) implies $\bar{\chi}_{\alpha,k}^2 = \sigma_{\alpha,k}^2 T_{\alpha/2, (k-1)/2}^{-1}$ ($k > 0$). The inverse power is one.

For the negative degree of freedom, we have $\bar{\chi}_{\alpha,-k}^2 = \sigma_{\alpha,k}^{-2} T_{\alpha/2, k/2}$. That is, without interference of the scale, the characteristic FCM2 is equivalent to the tilted stable variable: $\chi_{\alpha,-k}^2 = T_{\alpha/2, k/2}$.

When dealing with the fractional Gauss hypergeometric function (FGHF) in Section 5.2, we need two more variations from FCM2. The first allows an FCM2 to take a different scale:

$$(7.17) \quad \bar{\chi}_{\alpha,k}^2(x; \sigma) := \mathfrak{N}_{\alpha/2}(x; \sigma = \sigma, d = (k-1)/2, p = \alpha/2) \quad (k > 0)$$

from which the constant-scale variant is defined by replacing $\sigma_{\alpha,k}$ with $1/2$,

$$(7.18) \quad \hat{\chi}_{\alpha,k}^2(x) := \bar{\chi}_{\alpha,k}^2(x; \sigma = 1/4) = \mathfrak{N}_{\alpha/2}(x; \sigma = 1/4, d = (k-1)/2, p = \alpha/2) \quad (k > 0)$$

Notice the hat symbol replaces the bar symbol.

7.6.2. FCM2 CDF. Extending directly from Lemma 6.4, we have:

LEMMA 7.6. The CDF of FCM2 can be represented by γ_{α}^* as

$$(7.19) \quad \Phi[\bar{\chi}_{\alpha,k}^2](x) = z^{(k-1+\alpha)/2} \gamma_{\alpha/2}^* \left(\frac{k-1+\alpha}{\alpha}, z^{\alpha/2} \right) \quad (k > 0, \alpha \in [0, 2])$$

where $z = x/\sigma_{\alpha,k}^2$. △

7.6.3. FCM2 Subsumes Chi-squared. It is easy to show that $\bar{\chi}_{1,k}^2$ is χ_k^2 with a scale of $1/k$, due to (3.15).

7.6.4. Representing FCM by FCM2. In (7.14), let $s = \sqrt{x}$, we get the inverse relation:

$$(7.20) \quad \bar{\chi}_{\alpha,k}(s) = 2s \bar{\chi}_{\alpha,k}^2(s^2) \quad (s \geq 0)$$

Many ratio distribution integrals involving FCM can be rewritten in terms of FCM2, such that

$$(7.21) \quad \begin{aligned} f(x) &:= \int_0^\infty g(xs) \bar{\chi}_{\alpha,k}(s) s ds \\ &= \int_0^\infty g(x\sqrt{\nu}) \bar{\chi}_{\alpha,k}^2(\nu) \sqrt{\nu} d\nu \end{aligned}$$

For the CDF case, the incomplete integral can be transformed as

$$\begin{aligned} F(x) &:= \int_0^x f(x) dx = \int_0^\infty G(xs) \bar{\chi}_{\alpha,k}(s) ds \\ &= \int_0^\infty G(x\sqrt{\nu}) \bar{\chi}_{\alpha,k}^2(\nu) d\nu \end{aligned} \quad (7.22)$$

where $G(x) := \int_0^x g(x) dx$. The lower bound of the incomplete integrals can be $-\infty$ such as $\int_{-\infty}^x dx$ too.

7.6.5. Universal Expression. Assume $x \geq 0$, let $M(x^2) := G(x)/x$ in (7.22) or $g(x)$ in (7.21), we get the universal expression of

$$F(x) = x \int_0^\infty M(x^2\nu) \bar{\chi}_{\alpha,k}^2(\nu) \sqrt{\nu} d\nu \quad (7.23)$$

$$f(x) = \int_0^\infty M(x^2\nu) \bar{\chi}_{\alpha,k}^2(\nu) \sqrt{\nu} d\nu \quad (7.24)$$

Most of the univariate PDFs and CDFs in subsequent chapters can be understood in such framework. It is just a matter of what $M(x)$ is.

When $M(x)$ can be expressed by a Kummer function (apart from a negative sign), these integrals are members of the FGHF in Section 5.2.

7.7. FCM2 Mellin Transform

From (6.5), the Mellin transform of FCM2's PDF is

$$\begin{aligned} \bar{\chi}_{\alpha,k}^2(x) &\xleftrightarrow{\mathcal{M}} \bar{\chi}_{\alpha,k}^{2*}(s) \\ &= (\sigma_{\alpha,k})^{2s-2} \frac{\Gamma((k-1)/2) \Gamma((s+k/2-3/2) \times 2/\alpha)}{\Gamma((k-1)/\alpha) \Gamma(s+k/2-3/2)}. \end{aligned} \quad (k > 0) \quad (7.25)$$

Likewise, for the constant-scale variant, it becomes

$$\begin{aligned} \hat{\chi}_{\alpha,k}^2(x) &\xleftrightarrow{\mathcal{M}} \hat{\chi}_{\alpha,k}^{2*}(s) \\ &= 2^{2-2s} \frac{\Gamma((k-1)/2) \Gamma((s+k/2-3/2) \times 2/\alpha)}{\Gamma((k-1)/\alpha) \Gamma(s+k/2-3/2)}, \end{aligned} \quad (k > 0) \quad (7.26)$$

whose most important special case is $\alpha = 1$,

$$\hat{\chi}_{1,k}^2(x) \xleftrightarrow{\mathcal{M}} \hat{\chi}_{1,k}^{2*}(s) = \frac{\Gamma(s+k/2-1)}{\Gamma(k/2)} \quad (7.27)$$

$\Gamma(s+k/2-1)$ in $\hat{\chi}_{1,k}^{2*}(s)$ is just an ordinary gamma function without a fractional coefficient in front of s . This property is the basis that connects the fractional Gauss hypergeometric function to its classic form in Section 5.2.

7.8. FCM2 Moments

From the Mellin transform by $s = n + 1$, its n -th moment is

$$\begin{aligned} \mathbb{E}(X^n | \bar{\chi}_{\alpha,k}^2) &= \mathbb{E}(X^{2n} | \bar{\chi}_{\alpha,k}) \\ &= (\sigma_{\alpha,k})^{2n} \frac{\Gamma((k-1)/2) \Gamma((n+k/2-1/2) \times 2/\alpha)}{\Gamma((k-1)/\alpha) \Gamma(n+k/2-1/2)}. \end{aligned} \quad (k > 0) \quad (7.28)$$

As mentioned in Section 7.3, due to the nature of a ratio distribution, it is often evaluated as negative moments, $n < 0$. Hence, n is confined in the range of $1/2 - k/2 < n < 0$.

This puts stricter constraint on non-existing moments than FCM when k is not "large enough". For instance, in the case of fractional F distribution in Section 8.4, $k \approx 3$ is in the neighborhood where its second moment barely exists. This makes it rather hard for the statistics of the SPX daily return data set, since its k is just slightly larger than 3 while α is slightly below 1.

7.9. FCM2 Increment of k

LEMMA 7.7. When x^m is multiplied to $\bar{\chi}_{\alpha,k}^2(x)$, it follows a scaling rule where k is incremented to $k + 2m$ in the parametrization.

$$(7.29) \quad x^m \bar{\chi}_{\alpha,k}^2(x) = \sigma_{\alpha,k}^{2m} Q \frac{C_{\alpha,k}}{C_{\alpha,k+2m}} \bar{\chi}_{\alpha,k+2m}^2(y).$$

where $Q := \sigma_{\alpha,k+2m}^2 / \sigma_{\alpha,k}^2$ and $y = Qx$. △

PROOF. From (7.15),

$$x^m \bar{\chi}_{\alpha,k}^2(x) = \sigma_{\alpha,k}^{2m} \frac{C_{\alpha,k}}{2\sigma_{\alpha,k}^2} \left(\frac{x}{\sigma_{\alpha,k}^2} \right)^{(k+2m)/2-3/2} F_{\frac{\alpha}{2}} \left(\left(\frac{x}{\sigma_{\alpha,k}^2} \right)^{\alpha/2} \right).$$

We see that $\bar{\chi}_{\alpha,k}^2$ should become $\bar{\chi}_{\alpha,k+2m}^2$ according to the power in the $x^{(k+2m)/2-3/2}$ term, but other parts of the formula need to be adjusted too.

Since

$$\bar{\chi}_{\alpha,k+2m}^2(y) = \frac{C_{\alpha,k+2m}}{2\sigma_{\alpha,k+2m}^2} \left(\frac{y}{\sigma_{\alpha,k+2m}^2} \right)^{(k+2m)/2-3/2} F_{\frac{\alpha}{2}} \left(\left(\frac{y}{\sigma_{\alpha,k+2m}^2} \right)^{\alpha/2} \right),$$

we obtain $y = x \sigma_{\alpha,k+2m}^2 / \sigma_{\alpha,k}^2$ in order to match the two structurally.

Then take the ratio of $x^m \bar{\chi}_{\alpha,k}^2(x) / \bar{\chi}_{\alpha,k+2m}^2(y)$ to determine the needed constant, we arrive at (7.29). □

7.10. Sum of Two Chi-Squares with Correlation

The sum of bivariate variables is studied here.

LEMMA 7.8. Let $Z = Z_1/s_1 + Z_2/s_2$ where Z_1, Z_2 are two independent χ_1^2 variables. The PDF of Z is

$$\begin{aligned} \chi_{11}^2(z, s_1, s_2) &= \frac{\sqrt{s_1 s_2}}{2} e^{-s_2 z/2} {}_1F_1 \left(\frac{1}{2}, 1; \frac{(s_2 - s_1)z}{2} \right) \\ &= \frac{\sqrt{s_1 s_2}}{2} e^{-(s_1 + s_2)z/4} I_0(|s_2 - s_1|z/4) \end{aligned}$$

We apply DLMF 12.6.9 to get the second line, where the symmetry of a, b is explicit since $I_0(x)$ is symmetric. For $x \gg 1$, $I_0(x) \approx e^x / \sqrt{2\pi x}$ (DLMF 10.40.5). △

When $Z_1 = U_1^2$, $Z_2 = U_2^2$, and U_1, U_2 has correlation ρ , then s_1, s_2 must be modified by the eigenvalue solution of $\bar{\Omega}^{-1} \text{diag}(\mathbf{s})$ such that

$$\chi_{11}^2(z, s_1, s_2, \rho) = \chi_{11}^2(z, s'_1, s'_2)$$

$$\text{where } (s'_1, s'_2) = \frac{(s_1 + s_2) \pm \sqrt{(s_1 - s_2)^2 - 4\rho^2 s_1 s_2}}{2(1 - \rho^2)}$$

Fractional F Distribution

The classic F distribution comes from the ratio of two χ^2 distributions. Assume $U_1 \sim \chi_d^2/d$ and $U_2 \sim \chi_k^2/k$, then $F \sim U_1/U_2$ is an F distribution, $F_{d,k}$.

Two use cases were mentioned in Azzalini (2013)[1]. In Section 4.3 there, the squared variable of a univariate skew-t with k degrees of freedom is distributed as $F_{1,k}$.

In Section 6.2 there, the quadratic form from a $d \times d$ multivariate skew-t with k degrees of freedom is distributed as $F_{d,k}$.

Thus, the meaning of d and k is quite clear in such a context: d is the dimension of the multivariate skew-normal process; k is the degree of freedom in the denominator of the ratio distribution. This chapter extends it fractionally.

8.1. Definition

DEFINITION 8.1. Assume $U_1 \sim \chi_d^2/d$ and $U_2 \sim \bar{\chi}_{\alpha,k}^2$, then $F \sim U_1/U_2$ is a fractional F distribution. We use the notation $F \sim F_{\alpha,d,k}$.

The standard PDF of $F_{\alpha,d,k}$ is

$$(8.1) \quad F_{\alpha,d,k}(x) = \int_0^\infty s \, ds \, [d \chi_d^2(dxs)] \bar{\chi}_{\alpha,k}^2(s)$$

and note that the classic term in the integrand, $d \chi_d^2(dz)$, is equivalent to our $\bar{\chi}_{1,d}^2(z)$.

The reader should be aware of the subtlety that "ds" in "s ds" is the calculus notation, while d in $[d \chi_d^2(dxs)]$ is the constant from $F_{\alpha,d,k}$.

The standard CDF of $F_{\alpha,d,k}$ is

$$(8.2) \quad \Phi[F_{\alpha,d,k}](x) = \int_0^x F_{\alpha,d,k}(x) \, dx$$

$$(8.3) \quad = \int_0^\infty \left[\frac{1}{\Gamma(\frac{d}{2})} \gamma\left(\frac{d}{2}, \frac{dxs}{2}\right) \right] \bar{\chi}_{\alpha,k}^2(s) \, ds$$

since the CDF of a χ_d^2 is the regularized lower incomplete gamma function of $\gamma(\frac{d}{2}, \frac{x}{2})/\Gamma(\frac{d}{2})$.

It can also be represented by a fractional Gauss hypergeometric function. See Section 5.2.5.

8.1.1. The Origin of Fractional F. $F_{\alpha,d,k}$ is connected to the quadratic form of a d -dimensional multivariate GAS-SN distribution, $L_{\alpha,k}(0, \bar{\Omega}, \beta)$. Indeed, its three parameters, α, d, k , are designated such that the symbols convey the same meanings. However, $\bar{\Omega}$ and β doesn't affect the outcome of $F_{\alpha,d,k}$.

To elaborate from Section 15.6, assume Z is a $d \times d$ multivariate skew-normal (SN) distribution $SN(0, \bar{\Omega}, \beta)$, and $\bar{\chi}_{\alpha,k}$ is a standard FCM. Then $X = Z/\bar{\chi}_{\alpha,k}$ is an $L_{\alpha,k}(0, \bar{\Omega}, \beta)$.

The quadratic form of X is $Q = \frac{1}{d} X^\top \bar{\Omega}^{-1} X$. And $Q \sim F_{\alpha,d,k}$ is a fractional F distribution.

8.1.2. Ratio of Titled Stable Law. It is obvious that Fractional F is a ratio of tilted stable law.

$$\begin{aligned} F_{\alpha,d,k} &\sim \bar{\chi}_{1,d}^2 / \bar{\chi}_{\alpha,k}^2 \\ &= \frac{\sigma_{\alpha,k}^2}{d} \frac{T_{\alpha/2,(k-1)/2}}{T_{1/2,(d-1)/2}} \end{aligned}$$

8.1.3. Fractional F Subsumes F.

LEMMA 8.2. When $\alpha = 1$, it becomes a classic F. That is, $F_{1,d,k} = F_{d,k}$. $\triangle$

8.1.4. Fractional F Subsumes GSaS-Squared and GAS-SN-Squared. The following cases are for $d = 1$:

LEMMA 8.3. If $X_1 \sim L_{\alpha,k}$, then $X_1^2 \sim F_{\alpha,1,k}$. $\triangle$

LEMMA 8.4. If $X_2 \sim L_{\alpha,k}(\beta)$, then $X_2^2 \sim F_{\alpha,1,k}$, independent of β . $\triangle$

They will be discussed in Chapter 12.

8.2. PDF at Zero

The PDF of an F distribution is singular as $x \rightarrow 0$ when $d < 2$. We can see that from

$$\begin{aligned} F_{\alpha,d,k}(x) &\approx \frac{(d/2)^{d/2}}{\Gamma(d/2)} x^{d/2-1} \int_0^\infty s^{d/2} ds \bar{\chi}_{\alpha,k}^2(s) \\ (8.4) \quad &= \frac{(d/2)^{d/2}}{\Gamma(d/2)} \mathbb{E}(X^d | \bar{\chi}_{\alpha,k}) x^{d/2-1} \end{aligned}$$

for very small x .

When $d = 1$, the peak is divergent as $F_{\alpha,1,k}(x) \approx \frac{1}{\sqrt{2\pi}} \mathbb{E}(X | \bar{\chi}_{\alpha,k}) \sqrt{x}^{-1}$. But its CDF $\propto \sqrt{x}$.

When $d = 2$, this peak is finite. $F_{\alpha,2,k}(0) = \mathbb{E}(X^2 | \bar{\chi}_{\alpha,k})$.

When $d > 2$, $F_{\alpha,d,k}(x)$ drops to zero at $x = 0$. This strange phenomenon seems to indicate that the bivariate system is the lowest dimension to have stable quadratic statistics. And a three dimension system is likely more stable. But we only analyze the bivariate case in this book.

8.3. Mellin Transform

From (7.25), and note that $\bar{\chi}_d^2 = \bar{\chi}_{1,d}^2$, the Mellin transform of Fractional F's PDF is

$$(8.5) \quad F_{\alpha,d,k}(x) \xleftrightarrow{\mathcal{M}} (\bar{\chi}_{1,d}^2)^*(s) (\bar{\chi}_{\alpha,k}^2)^*(2-s) \quad (d > 0, k > 0)$$

$$\begin{aligned} (8.6) \quad &= \left(\sqrt{2d} \sigma_{\alpha,k} \right)^{2-2s} \left[\frac{\Gamma((d-1)/2)}{\Gamma(d-1)} \frac{\Gamma((k-1)/2)}{\Gamma((k-1)/\alpha)} \right] \left[\frac{\Gamma(2p(s))}{\Gamma(p(s))} \frac{\Gamma(2q(s)/\alpha)}{\Gamma(q(s))} \right], \\ &\text{where } p(s) := s + d/2 - 3/2, \quad q(s) := 1/2 + k/2 - s. \end{aligned}$$

The number of gamma functions can be reduced via the Legendre duplication formula (A.2).

8.4. Moments

Its n -th moment is

$$\begin{aligned} \mathbb{E}(X^n | F_{\alpha,d,k}) &= d^{-n} \mathbb{E}(X^n | \chi_d^2) \mathbb{E}(X^{-n} | \bar{\chi}_{\alpha,k}^2) \\ (8.7) \quad &= \left(\frac{2}{d} \right)^n (d/2)_n \mathbb{E}(X^{-n} | \bar{\chi}_{\alpha,k}^2). \end{aligned}$$

where $(d/2)_n$ is the Pochhammer symbol, $(a)_n := \Gamma(a+n)/\Gamma(a)$.

Its first moment is $\mathbb{E}(X^{-1} | \bar{\chi}_{\alpha,k}^2) = \mathbb{E}(X^{-2} | \bar{\chi}_{\alpha,k}^2)$, independent of d . This is due to $\mathbb{E}(X | \chi_d^2) = d$.

Note that this first moment is also the second moment of an univariate GAS-SN in (12.9), or simply the variance of the corresponding GSaS.

Its second moment is $(1 + 2/d) \mathbb{E}(X^{-2}|\bar{\chi}_{\alpha,k}^2)$. Hence, its variance is

$$\begin{aligned} \text{var}\{F_{\alpha,d,k}\} &= (1 + 2/d) \mathbb{E}(X^{-2}|\bar{\chi}_{\alpha,k}^2) - \mathbb{E}(X^{-1}|\bar{\chi}_{\alpha,k}^2)^2 \\ &= (1 + 2/d) \mathbb{E}(X^{-4}|\bar{\chi}_{\alpha,k}^2) - \mathbb{E}(X^{-2}|\bar{\chi}_{\alpha,k}^2)^2. \end{aligned} \tag{8.8}$$

8.4.1. Stability Issue of the Second Moment. The moment formula appears to be straightforward. But the devil is in the detail.

The stability of moments symbolizes the challenge of stability in the α -stable distribution. Even the second moment has dramatic behavior when k is smaller than 4.

First, we recognize that the first moment of F is actually the second moment of the underlying two-sided distribution because the variable of F is squared. Having a finite and stable first moment in F is quite meaningful. But it is much harder to make sense of the variance when k is too small.

Notice that, when $d \rightarrow \infty$, the variance is independent of d ,

$$\text{var}\{F_{\alpha,\infty,k}\} = \mathbb{E}(X^{-2}|\bar{\chi}_{\alpha,k}^2) - \mathbb{E}(X^{-1}|\bar{\chi}_{\alpha,k}^2)^2$$

This is the most relevant quantity if it exists. Other variance of finite d could be expressed as the following:

$$\text{var}\{F_{\alpha,d,k}\} - \text{var}\{F_{\alpha,\infty,k}\} = \frac{2}{d} \mathbb{E}(X^{-2}|\bar{\chi}_{\alpha,k}^2).$$

The Framework of Continuous Gaussian Mixture

The construction of a symmetric two-sided distribution is in the form of a continuous Gaussian mixture. Both the ratio and product distribution methods could be used.

In the case of the symmetric α -stable distribution (SaS)[6], the exponential power distribution comes from its characteristic function (CF)[28]. In our framework, both distributions will be expanded by adding the parameter for the degrees of freedom (k).

In addition, we use positive k for the former and negative k for the latter. Hence, the " $k \in \mathbb{R}$ " domain will consolidate both distributions into one.

We would like to present a unified framework and familiarize the reader with the notations, which would otherwise be subtle and confusing.

The results of this chapter are the following. First, the symmetric two-sided distribution can be enriched using the characteristic function transform.

Second, the features of the two-sided distribution are transferred to the one-sided distribution because of the Gaussian mixture.

Third, that one-sided distribution could be transformed to the inverse distribution that represents the marginal distribution of a volatility process. This is more meaningful in applications, such as quantitative finance.

9.1. The Inverse Chi Distribution

Assume the PDF of a two-sided symmetric distribution is $L(x)$ where $x \in \mathbb{R}$. It has zero mean, $\mathbb{E}(X|L) = 0$. Assume the PDF of a one-sided distribution is $\chi(x)$ ($x > 0$) such that

$$(9.1) \quad L(x) := \int_0^\infty s \, ds \, \mathcal{N}(xs) \chi(s)$$

This is not new. It is the definition of a ratio distribution with a standard normal variable $\mathcal{N}$. This is the first form of the Gaussian mixture: $L \sim \mathcal{N}/\chi$. A typical example is that L is a Student's t distribution when χ is $\sqrt{\chi_k^2}$.

The skewness is added by replacing the normal distribution $\mathcal{N}$ with its skew-normal counterpart $\mathcal{N}(\beta)$. See next chapter for more details.

It has the equivalent expression in terms of a product distribution using *the inverse distribution* $\chi^\dagger$ such that $L \sim \mathcal{N}\chi^\dagger$ [13]. This is the second form of the Gaussian mixture.

The advantage of this expression is that $\chi^\dagger$ is closer to our typical understanding of the marginal distribution of a volatility process. For example, when the Brownian motion process $dX_t = \sigma_t dW_t$ is measured in a particular time interval Δt , we have $\Delta X_t \sim L$ and $\sigma_t \sim \chi^\dagger$.

Since χ in the ratio form is more natural in the expression of the α -stable distribution, we are more inclined to use the ratio distribution. The reader should keep this subtlety in mind.

LEMMA 9.1. (Inverse distribution) Let the inverse distribution of χ be $\chi^\dagger$, the relation between their density functions is

$$(9.2) \quad \chi^\dagger(s) := s^{-2} \chi\left(\frac{1}{s}\right)$$

1182 such that

$$(9.3) \quad \int_0^\infty s \, ds \, \mathcal{N}(xs) \chi(s) = \int_0^\infty \frac{ds}{s} \mathcal{N}\left(\frac{x}{s}\right) \chi^\dagger(s) \quad (x \in \mathbb{R})$$

$$(9.4) \quad \int_0^\infty s \, ds \, \mathcal{N}(xs) \chi^\dagger(s) = \int_0^\infty \frac{ds}{s} \mathcal{N}\left(\frac{x}{s}\right) \chi(s)$$

1183 The proof is straightforward by a change of variable $t = 1/s$. You can move between LHS and RHS
1184 easily.

1185 $\triangle$

1186

1187 9.2. The Characteristic Distributions

1188 The characteristic transform of a symmetric α -stable distribution (SaS) leads to an exponential
1189 power distribution. We would like to generalize this concept in this section.

1190 When the PDF of a distribution is represented by a Gaussian mixture, the features of a charac-
1191 teristic transform are transferred to its χ distribution counterpart.

1192 We use the notation $\text{CF}\{g\}(t) = \mathbb{E}(e^{itX}|g)$ to represent the characteristic function transform of
1193 the PDF $g(x)$. Note that $\mathcal{N}$ has a special property that its CF is still itself: $\text{CF}\{\mathcal{N}\}(t) = \sqrt{2\pi} \mathcal{N}(t)$.

1194 LEMMA 9.2. (Characteristic function transform of L) Let $\phi(t)$ be the CF of L such that $\phi(t) :=$
1195 $\text{CF}\{L\}(t) = \int_{-\infty}^\infty dx \exp(itx) L(x)$. (9.1) is transformed to

$$(9.5) \quad \phi(t) = \sqrt{2\pi} \int_0^\infty ds \, \mathcal{N}\left(\frac{t}{s}\right) \chi(s) \quad (t \in \mathbb{R})$$

1196 The proof is straightforward from the fact that $\text{CF}\{\mathcal{N}(0, \sigma^2)\}(t) = \sqrt{2\pi} \mathcal{N}(\sigma t)$.

1197 $\triangle$

1198 (9.5) is quite similar to the RHS of (9.4). It inspires us to define a new distribution pair, L_ϕ and
1199 $\chi_\phi^\dagger$, in terms of a product distribution, such that

$$(9.6) \quad L_\phi(x) := \int_0^\infty \frac{ds}{s} \mathcal{N}\left(\frac{x}{s}\right) \chi_\phi^\dagger(s) \quad (x \in \mathbb{R})$$

$$(9.7) \quad \chi_\phi^\dagger(s) := \frac{s \chi(s)}{\mathbb{E}(X|\chi)}$$

1200 where $\mathbb{E}(X|\chi)$ is the first moment of χ . Here $\chi_\phi^\dagger$ is the inverse distribution of χ_ϕ , which can be
1201 reverse-engineered according to (9.2),

$$(9.8) \quad \chi_\phi(s) := \frac{s^{-3}}{\mathbb{E}(X|\chi)} \chi\left(\frac{1}{s}\right)$$

1202

1203 We are in an interesting place: We start with a one-sided distribution χ , we derive two variants
1204 from it: χ_ϕ and $\chi_\phi^\dagger$. We also obtain two two-sided distributions: L and L_ϕ .

1205 We shall call χ_ϕ *the characteristic distribution* of χ as it facilitates $L_\phi \sim \mathcal{N}/\chi_\phi$.

9.3. Summary of Gaussian Mixture

In summary, considering the inverse distribution and characteristic transform, we obtain four Gaussian mixture relations. The first two parallel relations in terms of ratio distribution:

$$L \sim \mathcal{N}/\chi;$$

$$L_\phi \sim \mathcal{N}/\chi_\phi, \quad \chi_\phi(s) := \frac{s^{-3}}{\mathbb{E}(X|\chi)} \chi\left(\frac{1}{s}\right).$$

In terms of production distribution, they become

$$L \sim \mathcal{N} \chi^\dagger, \quad \chi^\dagger(s) := s^{-2} \chi\left(\frac{1}{s}\right) = \frac{s}{\mathbb{E}(X|\chi_\phi)} \chi_\phi(s);$$

$$L_\phi \sim \mathcal{N} \chi_\phi^\dagger, \quad \chi_\phi^\dagger(s) := \frac{s}{\mathbb{E}(X|\chi)} \chi(s).$$

Both $\chi^\dagger$ and $\chi_\phi^\dagger$ are the volatility processes that generate L and L_ϕ .

Next, the ϕ suffix in $\chi_\phi(s)$ will be replaced by the *negation* (sign change) of the degree of freedom ($k \rightarrow -k$).

The multiplication of $\frac{s}{\mathbb{E}(X|f)}$ is closely related to an increase in the degree of freedom ($k \rightarrow k+1$) of the underlying FG, which is equivalent to the scaling of χ_ϕ and χ .

$$g(s) = \frac{s}{\mathbb{E}(X|\chi)} f_k(s; \sigma_k)$$

$$= \frac{1}{\sigma} f_{k+1}\left(\frac{s}{\sigma}; \sigma_{k+1}\right)$$

9.4. FCM Extensions

In this section, we explicitly define the three extensions of $\bar{\chi}_{\alpha,k}$ and prove their connections. They are sub-families of FG. Hence, their density functions should be expressed by $\mathfrak{N}_{\alpha/2}(x; \dots)$ in (6.1).

First, we quote the PDF of $\bar{\chi}_{\alpha,k}$ from (7.4) below. For $x \geq 0$ and $k > 0$,

$$(9.9) \quad \bar{\chi}_{\alpha,k}(x) := \mathfrak{N}_{\alpha/2}(x; \sigma = \sigma_{\alpha,k}, d = k-1, p = \alpha) \quad (x \geq 0, k > 0)$$

$$= (C_{\alpha,k}) (\sigma_{\alpha,k})^{1-k} x^{k-2} F_{\frac{\alpha}{2}} \left(\left(\frac{x}{\sigma_{\alpha,k}} \right)^\alpha \right).$$

The characteristic FCM $[\bar{\chi}_\phi]_{\alpha,k}$ uses the $-k$ index of $\bar{\chi}_{\alpha,k}$. Its PDF is

$$(9.10) \quad [\bar{\chi}_\phi]_{\alpha,k}(x) = \bar{\chi}_{\alpha,-k}(x) := \mathfrak{N}_{\alpha/2}(x; \sigma = (\sigma_{\alpha,k})^{-1}, d = -k, p = -\alpha).$$

$$= (C_{\alpha,k+1}) (\sigma_{\alpha,k})^{-k} x^{-k-1} F_{\frac{\alpha}{2}} \left((\sigma_{\alpha,k} x)^{-\alpha} \right).$$

DEFINITION 9.3. The density functions of the two inverse distributions are as follows.

$$(9.11) \quad \bar{\chi}_{\alpha,k}^\dagger(x) := \mathfrak{N}_{\alpha/2}(x; \sigma = (\sigma_{\alpha,k})^{-1}, d = -k+1, p = -\alpha).$$

$$= (C_{\alpha,k}) (\sigma_{\alpha,k})^{-k+1} x^{-k} F_{\frac{\alpha}{2}} \left((\sigma_{\alpha,k} x)^{-\alpha} \right).$$

And

$$(9.12) \quad [\bar{\chi}_\phi^\dagger]_{\alpha,k}(x) = \bar{\chi}_{\alpha,-k}^\dagger(x) := \mathfrak{N}_{\alpha/2}(x; \sigma = \sigma_{\alpha,k}, d = k, p = \alpha).$$

$$= (C_{\alpha,k+1}) (\sigma_{\alpha,k})^{-k} x^{k-1} F_{\frac{\alpha}{2}} \left(\left(\frac{x}{\sigma_{\alpha,k}} \right)^\alpha \right).$$

REMARK 9.4. In the following subsections, we will show that only the characteristic FCM is a new distribution family, which we use $-k$ to align its parameter such that $[\bar{\chi}_\phi]_{\alpha,k}(x) := \bar{\chi}_{\alpha,-k}(x)$ in (9.14).

The other two variations are just rescaled $k \pm 1$ of FCM. Since we will add location and scale to a distribution family, they don't add new explanatory power to the data in real-world applications.

9.4.1. Expressed via Alpha-Stable Extremal Distribution. From (4.3), we have $L_\alpha(x) = x^{-1}F_\alpha(x^{-\alpha})$. The two density functions, $\bar{\chi}_{\alpha,k}^\dagger(x)$ and $[\bar{\chi}_\phi]_{\alpha,k}(x)$, can be expressed by $L_\alpha(x)$ more elegantly since

$$(9.13) \quad F_{\frac{\alpha}{2}}(x^{-\alpha}) = x^2 L_{\frac{\alpha}{2}}(x^2).$$

Therefore, the PDF of the characteristic distribution becomes

$$(9.14) \quad [\bar{\chi}_\phi]_{\alpha,k}(x) := \bar{\chi}_{\alpha,-k}(x) = (C_{\alpha,k+1}) (\sigma_{\alpha,k})^{-k+2} x^{-k+1} L_{\frac{\alpha}{2}}((\sigma_{\alpha,k} x)^2).$$

which is a FG-type manipulation in the α -stable extremal distribution.

Likewise, the PDF of the inverse distribution becomes

$$(9.15) \quad \bar{\chi}_{\alpha,k}^\dagger(x) = (C_{\alpha,k}) (\sigma_{\alpha,k})^{-k+3} x^{-k+2} L_{\frac{\alpha}{2}}((\sigma_{\alpha,k} x)^2).$$

LEMMA 9.5. Show that

$$(9.16) \quad \bar{\chi}_{\alpha,k}^\dagger(x) = \frac{x}{\mathbb{E}(X|[\bar{\chi}_\phi]_{\alpha,k})} [\bar{\chi}_\phi]_{\alpha,k}(x)$$

△

PROOF. Divide (9.15) by (9.14). The result is $\bar{\chi}_{\alpha,k}^\dagger(x)/[\bar{\chi}_\phi]_{\alpha,k}(x) = x \sigma_{\alpha,k} C_{\alpha,k}/C_{\alpha,k+1}$, which is $x \mathbb{E}(X|\bar{\chi}_{\alpha,k})$ according to (7.9). And $\mathbb{E}(X|\bar{\chi}_{\alpha,k}) = 1/\mathbb{E}(X|[\bar{\chi}_\phi]_{\alpha,k})$. We arrive at the desired result. □

Furthermore, this indicates the following.

LEMMA 9.6. $\bar{\chi}_{\alpha,k}^\dagger(x)$ is a rescaled $[\bar{\chi}_\phi]_{\alpha,k-1}(x)$, which is $\bar{\chi}_{\alpha,-k+1}(x)$. That is,

$$(9.17) \quad \bar{\chi}_{\alpha,k}^\dagger(x) = \frac{1}{\sigma} [\bar{\chi}_\phi]_{\alpha,k-1}\left(\frac{x}{\sigma}\right), \quad \text{where } \sigma = \frac{\sigma_{\alpha,k-1}}{\sigma_{\alpha,k}}.$$

△

PROOF. From (9.14), we have

$$[\bar{\chi}_\phi]_{\alpha,k-1}(x) = (C_{\alpha,k}) (\sigma_{\alpha,k-1})^{-k+3} x^{-k+2} L_{\frac{\alpha}{2}}((\sigma_{\alpha,k-1} x)^2).$$

Compare the argument in $L_{\frac{\alpha}{2}}()$ to (9.15) after the substitution x/σ . We obtain $\sigma_{\alpha,k-1}^2/\sigma^2 = \sigma_{\alpha,k}^2$ and arrive at the desired result for σ . □

9.4.2. Proofs of Other Relations.

LEMMA 9.7. Show that (9.8) is true in the FCM implementation.

$$(9.18) \quad [\bar{\chi}_\phi]_{\alpha,k}(x) = \frac{x^{-3}}{\mathbb{E}(X|\bar{\chi}_{\alpha,k})} \bar{\chi}_{\alpha,k}\left(\frac{1}{x}\right)$$

△

PROOF. The proof is straightforward. First, replace x with $1/x$ in (9.9). Second, divide it by (9.10). The result is $[\bar{\chi}_\phi]_{\alpha,k}(x)/[\bar{\chi}_\phi]_{\alpha,k}(x) = \sigma_{\alpha,k} x^3 C_{\alpha,k}/C_{\alpha,k+1}$, which is $x^3 \mathbb{E}(X|\bar{\chi}_{\alpha,k})$ according to (7.9). We arrive at the desired result. □

LEMMA 9.8. Show that

$$(9.19) \quad \left[\bar{\chi}_\phi^\dagger \right]_{\alpha,k}(x) = \frac{x}{\mathbb{E}(X|\bar{\chi}_{\alpha,k})} \bar{\chi}_{\alpha,k}(x)$$

This indicates that $\left[\bar{\chi}_\phi^\dagger \right]_{\alpha,k}(x)$ is a rescaled $\bar{\chi}_{\alpha,k+1}(x)$. This makes it a trivial case. $\triangle$

PROOF. Divide (9.9) by (9.12). The result is $\bar{\chi}_{\alpha,k}(x) / \left[\bar{\chi}_\phi^\dagger \right]_{\alpha,k}(x) = \sigma_{\alpha,k} x^{-1} C_{\alpha,k} / C_{\alpha,k+1}$, which is $x^{-1} \mathbb{E}(X|\bar{\chi}_{\alpha,k})$ according to (7.9). We arrive at the desired result. $\square$

These relations from different angles ensure that all the formulas are consistent with each other.

9.5. Application of FCM Variants

I have hypothesized that the distribution of the daily VIX indexes can be fitted by the FCM-type distributions[18]. Now we have a more sophisticated parameterization system. We can attempt to fit the data and study how the FCM variants behave.

The data set is the daily VIX data from 1/1990 to 4/2026. The data is converted to a histogram with about 230 bins. Each VIX cycle could stretch over more than a decade. And we only have less than three decades of data. The shape of the data could be somewhat twisted, depending on where the cutoff is in the current cycle.

9.5.1. The FCM Fit. We fit the data by two methods. The first fit uses the FCM itself with the location parameter, whose PDF is

$$(9.20) \quad f_1(x) = \frac{1}{\omega} \bar{\chi}_{\alpha,k} \left(\frac{x - \varepsilon}{\omega} \right)$$

as shown in Figure 9.1.

The distribution fits the peak nicely. But the peak of the data before the lower bound is sharper than the distribution can explain.

The distribution follows the shape of the right tail in the data until $x = 7$. In the far right tail of $x > 7$, the data have a fatter tail than the distribution.

9.5.2. The Characteristic FCM Fit. The second fit uses the characteristic FCM without the location parameter, whose PDF is

$$(9.21) \quad f_2(x) = \frac{1}{\omega} [\bar{\chi}_\phi]_{\alpha,k} \left(\frac{x}{\omega} \right)$$

as shown in Figure 9.2.

It is intentional to force the location parameter to zero. We want to examine the behavior of the form $x^{-k} L_{\frac{\alpha}{2}}(x^2)$, which suppresses the distribution to zero when x is small. This provides an equivalent of a location parameter that shifts the density to the right.

The distribution fits the peak nicely. However, the data reach their peak at a location lower than that of the distribution. We also observe that the cutoff point of the data at the lower bound is sharper than that of the distribution.

An interesting result of the fit is $\alpha \sim 1$ which is an exponential tail like the classic χ distribution.

The distribution follows the shape of the right tail in the data until $x = 9$. In the far right tail of $x > 9$, the distribution has a fatter tail than the data.

REMARK 9.9. The fatter tail of the $[\bar{\chi}_\phi]_{\alpha,k}$ distribution is interesting. In the real world, an extreme VIX reading indicates a collapse of the equity market, which is catastrophic. This can not happen often, or even once if it is extremely large. Indeed, after every large market crash, the U.S. regulator implemented mechanisms to slow the market panic.




FIGURE 9.1. FCM fit of VIX daily data from 1990 to 4/2026. Data is standardized to one standard deviation. The optimal parameters are: $\{\alpha = 0.645, k = 2.39, \varepsilon = 1.19, \omega = 0.86\}$. The top left chart is the PDF vs x . The top right chart is the PDF in the log scale vs x . The bottom left chart is the complimentary CDF vs x . The bottom right chart is the complimentary CDF in the log scale vs x .

1286 It appears that the FCM distribution is too conservative in the right tail, while the characteristic
 1287 FCM is too aggressive. The right tail behavior of real-world data lies between the two, which is likely
 1288 determined empirically and could evolve over time.




FIGURE 9.2. Characteristic FCM fit of VIX daily data from 1990 to 4/2026. Data is standardized to one standard deviation. This fit forces the location parameter to zero. The optimal parameters are: $\{\alpha = 1.035, k = 4.45, \omega = 2.33\}$. The top left chart is the PDF vs x . The top right chart is the PDF in the log scale vs x . The bottom left chart is the complimentary CDF vs x . The bottom right chart is the complimentary CDF in the log scale vs x .

Part 3

Two-Sided Univariate Distributions

SN: The Skew-Normal Distribution - Review

10.1. Definition

The skew-normal distribution family is well documented in A. Azzalini's 2013 monograph[1]. We recap the results and clarify the symbology. My contribution is to incorporate the skew-normal methodology into the fractional distributions wherever suitable. The enhanced distributions are flexible and can adapt to many different shapes and tails with high skewness and kurtosis.

10.1.1. The Selective Sampling. The *selective sampling* method is used to inject skewness into the stochastic system, which is otherwise symmetric. This mechanism is fairly common in an applied context, for example, in social sciences, where a variable X_0 is observed only when a correlated variable X_1 , which is usually unobserved, satisfies a certain condition (p.128 of [1]).

In quantitative finance, the condition could be market regimes. In a two-regime model, a market index such as the S&P 500 index (SPX) is classified into the growth regime or the crash regime at a given time. It is well known that the volatility of the market behaves differently in each regime. In the growth regime, volatility tends to be low, and the market is trending upward. In the crash regime, volatility tends to be high, and the market is trending downward.

A univariate random variable $Z \sim SN(0, 1, \beta)$ is a standard skew-normal variable with skew parameter $\beta \in \mathbb{R}$ (Section 2.1 of [1]). The sign of β determines the sign of its skewness (10.14).

One of its stochastic representations is

$$(10.1) \quad Z = \begin{cases} X_0 & \text{if } X_1 < \beta X_0, \\ -X_0 & \text{otherwise.} \end{cases}$$

where X_0, X_1 are independent $\mathcal{N}(0, 1)$ variables.

An alternative representation uses filtering, or rejection, such that $Z = (X_0 | X_1 < \beta X_0)$. That is, X_0 is accepted as Z only when the condition $X_1 < \beta X_0$ is satisfied. Otherwise, it is discarded.

10.1.2. The PDF and CDF. The standard PDF is

$$(10.2) \quad \mathcal{N}(x; \beta) := 2\mathcal{N}(x)\Phi_{\mathcal{N}}(\beta x), \quad (x \in \mathbb{R})$$

where $\mathcal{N}(x)$ and $\Phi_{\mathcal{N}}(x)$ are the PDF and CDF of $\mathcal{N}(0, 1)$.

Its extremal distribution occurs at $\beta \rightarrow \infty$, where $\Phi_{\mathcal{N}}(\beta x)$ becomes a step function. The PDF becomes that of a half-normal distribution.

The standard CDF is

$$(10.3) \quad \Phi_{SN}(x; \beta) := \Phi_{\mathcal{N}}(x) - 2T(x, \beta)$$

where $T(h, a)$ is called the Owen's T function[29]. Its numerical methods are widely implemented in modern software packages.

Several important properties are quoted from Proposition 2.1 of [1]:

- $\mathcal{N}(0; 0) = 1/\sqrt{2\pi}$. Universal anchor at $x = 0, \beta = 0$.
- $\mathcal{N}(x; 0) = \mathcal{N}(x)$. Continuity at $\beta = 0$.
- $\mathcal{N}(-x; \beta) = \mathcal{N}(x; -\beta)$. This is the reflection rule.

- $Z^2 \sim \chi_1^2$, irrespective of β .

Notice that Z^2 is independent of β . This is an important property, but may not be intuitive for new students. This is due to the fact that the squares of X_0 and $-X_0$ are the same in (10.1). This property is carried into the quadratic form of the multivariate elliptical distribution.

10.2. The Location-Scale Family

Its location-scale family is $Y = \xi + \omega Z \sim SN(\xi, \omega^2, \beta)$, where $\xi \in \mathbb{R}$ and $\omega > 0$. Its PDF becomes

$$(10.4) \quad \frac{1}{\omega} \mathcal{N}\left(\frac{x - \xi}{\omega}; \beta\right).$$

10.3. Invariant Quantities

The following quantity plays an important role in the selective sampling concept of SN:

$$(10.5) \quad \delta = \frac{\beta}{\sqrt{1 + \beta^2}}, \quad \delta \in (-1, 1).$$

It can be thought of as some kind of correlation in the following. Inversely, β can be calculated from

$$(10.6) \quad \beta = \frac{\delta}{\sqrt{1 - \delta^2}}.$$

These two quantities will appear in many places in the ensuing chapters. They are invariants in the context of the multivariate elliptical distribution, called the Canonical Form.

In a trigonometry representation, one can think of δ as $\sin(\theta)$ of a right triangle, where one leg is 1, the other leg is β , and θ is the angle facing β .

Three representations use δ as the correlation coefficient to generate SN. (Section 2.1.3 of [1]) First, designate the correlation matrix as

$$\bar{\Omega} = \begin{pmatrix} 1 & \delta \\ \delta & 1 \end{pmatrix}.$$

The Cholesky factor of $\bar{\Omega}$ is

$$L = \begin{pmatrix} 1 & 0 \\ \delta & \sqrt{1 - \delta^2} \end{pmatrix},$$

so that $L L^T = \bar{\Omega}$.

Assume U_0 and U_1 are two independent $\mathcal{N}(0, 1)$ variates. The first representation of $Z \sim SN(0, 1, \beta)$ is

$$(10.7) \quad Z = \begin{cases} X_0 & \text{if } X_1 > 0, \\ -X_0 & \text{otherwise.} \end{cases}$$

where X_0, X_1 are marginals of a standard correlated normal bivariate with $\text{cor}\{X_0, X_1\} = \delta$ such that

$$\begin{pmatrix} X_0 \\ X_1 \end{pmatrix} = L \begin{pmatrix} U_0 \\ U_1 \end{pmatrix}.$$

The second representation is from

$$\begin{pmatrix} - \\ Z \end{pmatrix} = L \begin{pmatrix} U_0 \\ |U_1| \end{pmatrix}$$

such that $Z = \sqrt{1 - \rho^2} U_0 + \delta |U_1| \sim SN(0, 1, \beta)$.

The third representation is $Z = \max\{X_0, X_1\} \sim SN(0, 1, \beta)$, where X_0, X_1 are marginals of a standard correlated bivariate such that

$$\begin{pmatrix} X_0 \\ X_1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ \rho & \sqrt{1 - \rho^2} \end{pmatrix} \begin{pmatrix} U_0 \\ U_1 \end{pmatrix}.$$

1340 and $\text{cor}\{X_0, X_1\} = \rho = 1 - 2\delta^2$.

1341

10.4. Mellin Transform

1342 The following result is elegant, but also peculiar. It is discovered by the author.

1343 LEMMA 10.1. The Mellin transform of the SN PDF is

$$(10.8) \quad \mathcal{N}(x; \beta) \xleftrightarrow{\mathcal{M}} \mathcal{N}^*(s; \beta) := 2\mathcal{N}^*(s) \Phi[t_s](\beta\sqrt{s}),$$

$$\text{where } \mathcal{N}^*(s) = \frac{1}{2} \frac{2^{s/2}}{\sqrt{2\pi}} \Gamma\left(\frac{s}{2}\right)$$

1344 is the Mellin transform of the PDF of $\mathcal{N}(0, 1)$ in (2.9). And $\Phi[t_k](x)$ is the CDF of a Student's t
 1345 distribution with k degrees of freedom. But it is used in a strange way, where s substitutes k and goes
 1346 into x at the same time.

1347

△

1348 PROOF. We prove (10.8) via the CDF of GSaS with $\alpha = 1$. By definition,

$$\mathcal{N}^*(s; \beta) = \int_0^\infty x^{s-1} [2\mathcal{N}(x)\Phi_{\mathcal{N}}(\beta x)] dx.$$

1349 We use the known result from $\bar{\chi}_{1,k}$ where

$$x^{k-1}\mathcal{N}(x) = \frac{2^{k/2-1}\Gamma(k/2)}{\sqrt{2\pi k}} \bar{\chi}_{1,k}(x/\sqrt{k}) = \frac{1}{\sqrt{k}} \mathcal{N}^*(k) \bar{\chi}_{1,k}(x/\sqrt{k}).$$

1350 Then

$$\begin{aligned} \mathcal{N}^*(s; \beta) &= \frac{2\mathcal{N}^*(s)}{\sqrt{s}} \int_0^\infty \Phi_{\mathcal{N}}(\beta x) \bar{\chi}_{1,s}(x/\sqrt{s}) dx \\ &= 2\mathcal{N}^*(s) \int_0^\infty \Phi_{\mathcal{N}}(\beta\sqrt{s}t) \bar{\chi}_{1,s}(t) dt \quad \text{via } t = x/\sqrt{s}. \end{aligned}$$

1351 The integral is exactly the CDF of a GSaS, $L_{1,s}$, with the argument $\beta\sqrt{s}$. That is, $\mathcal{N}^*(s; \beta) =$
 1352 $2\mathcal{N}^*(s) \Phi[L_{1,s}](\beta\sqrt{s})$.

1353 When $\alpha = 1$, $L_{1,s}$ becomes t_s . Therefore, $\mathcal{N}^*(s; \beta) = 2\mathcal{N}^*(s) \Phi[t_s](\beta\sqrt{s})$.

1354

□

1355 The beauty of this lemma is that $\mathcal{N}^*(s; \beta)$ is the multiplication of a symmetric component and a
 1356 skew component, just like its PDF counterpart.

1357 From (2.14), we also obtain that

$$(10.9) \quad \Phi_{SN}(0; \beta) = 1 - \mathcal{N}^*(1; \beta) = \frac{1}{2} - \frac{1}{\pi} \arctan(\beta).$$

1358 This is due to $\mathcal{N}^*(1) = \frac{1}{2}$ and $\Phi[t_1](\beta) = \frac{1}{2} + \frac{1}{\pi} \arctan(\beta)$. This result is stated in Proposition 2.7 of
 1359 [1], and is proved here via the Mellin transform.

10.4.1. Mellin Transform of Owen's T Function. Another peculiar result from the Mellin transform is

LEMMA 10.2.

$$(10.10) \quad T(x, \beta) \xleftrightarrow{\mathcal{M}} s^{-1} \mathcal{N}^*(s+1) \left[\Phi[t_{s+1}](\beta\sqrt{s+1}) - \frac{1}{2} \right]$$

△

PROOF. Define the upper incomplete integral as

$$\begin{aligned} \Gamma_f(x) &:= \int_x^\infty \mathcal{N}(x; \beta) dx = 1 - \Phi_{SN}(x; \beta) \\ &= 1 - \Phi_{\mathcal{N}}(x) + 2T(x, \beta) \end{aligned}$$

According to Lemma 2.6, its Mellin transform is

$$\begin{aligned} \Gamma_f(x) &\xleftrightarrow{\mathcal{M}} s^{-1} \mathcal{N}^*(s+1; \beta) \\ &= 2s^{-1} \mathcal{N}^*(s+1) \Phi[t_{s+1}](\beta\sqrt{s+1}) \end{aligned}$$

Combining the two results above, we obtain

$$T(x, \beta) = \frac{\Gamma_f(x) - (1 - \Phi_{\mathcal{N}}(x))}{2} \xleftrightarrow{\mathcal{M}} s^{-1} \mathcal{N}^*(s+1) \left[\Phi[t_{s+1}](\beta\sqrt{s+1}) - \frac{1}{2} \right]$$

where $1 - \Phi_{\mathcal{N}}(x) \xleftrightarrow{\mathcal{M}} s^{-1} \mathcal{N}^*(s+1)$.

□

10.5. Moments

LEMMA 10.3. According to Section 2.2.2, by assigning $s = n+1$, the Mellin transform is converted to the moment formula. It is easy to show that the n -th moment of Z is

$$(10.11) \quad \begin{aligned} \mathbb{E}(Z^n) &= \mathbb{E}(X^n | \mathcal{N}(\beta)) = \mathcal{N}^*(n+1; \beta) + (-1)^n \mathcal{N}^*(n+1; -\beta) \\ &= 2\mathcal{N}^*(n+1) \times \begin{cases} 1, & \text{when } n \text{ is even,} \\ 1 - 2\Phi[t_{n+1}](-\beta\sqrt{n+1}), & \text{when } n \text{ is odd.} \end{cases} \end{aligned}$$

The even moments are identical to those of $\mathcal{N}(0, 1)$. It is the odd moments that make the difference when $\beta \neq 0$.

△

The first four moments of Z have simple analytic forms. Its first moment is

$$(10.12) \quad \mu_z = b\delta, \quad \text{where } b = \sqrt{2/\pi}.$$

The second moment is simply 1. Its variance is

$$(10.13) \quad \sigma_z^2 = 1 - (b\delta)^2.$$

The third moment is $b\delta(3 - \delta^2)$. Its skewness is

$$(10.14) \quad \gamma_1\{Z\} = \frac{4 - \pi}{2} \frac{\mu_z^3}{\sigma_z^3}.$$

The fourth moment is 3. Its kurtosis is

$$(10.15) \quad \gamma_2\{Z\} = 2(\pi - 3) \frac{\mu_z^4}{\sigma_z^4}.$$

The maximum skewness of SN is approximately 0.9953 and the maximum kurtosis is 0.8692. They are not very interesting, since the extremal distribution is just a half-normal distribution.

1380 However, these analytical forms are useful when SN is extended to GAS-SN. Both skewness and
1381 kurtosis are extended to much wider ranges, or even infinity!

CHAPTER 11

GAS: Generalized Alpha-Stable Distribution (Experimental)

This chapter studies how a degrees-of-freedom parameter k can be added to the α -stable distribution L_α^θ through the Mellin transform framework. The construction leads to the generalized α -stable distribution (GAS), denoted by $L_{\alpha,k}^\theta$.

This experiment is an early attempt and one of the cleanest approaches to understanding how k interacts with skewness. It is a valuable lesson on the mathematical structure of the α -stable distribution. Therefore, it is documented in this chapter.

The GAS construction is useful because it exposes how the new parameter k interacts with skewness in the stable-law setting. At the same time, the resulting distribution has a discontinuity associated with the reflection rule, which limits its direct use as a final statistical model.

Readers who are mainly interested in the final skew-normal-based construction may treat this chapter as optional background.

Thus, the main role of this chapter is structural. It explains how the fractional χ distribution and GSaS arise from the stable-law framework, and it motivates the later shift to the skew-normal approach, where skewness can be incorporated while preserving continuity of the PDF.

11.1. Definition

First, we recap the Mellin transform (4.5) of the PDF of the α -stable distribution from Section 4.3,

$$L_\alpha^\theta(x) \xleftrightarrow{\mathcal{M}} \epsilon \left[\frac{\Gamma(s)}{\Gamma(1-\gamma+\gamma s)} \right] \left[\frac{\Gamma(\epsilon(1-s))}{\Gamma(\gamma(1-s))} \right].$$

It is interpreted in Lemma 4.4 as a multiplication of two components,

$$L_\alpha^\theta(x) \xleftrightarrow{\mathcal{M}} \tilde{M}_\gamma^*(s) \bar{\chi}_{\alpha,1}^\theta * (2-s).$$

The PDF of the second term $\bar{\chi}_{\alpha,1}$ is defined as

$$\bar{\chi}_{\alpha,1}^\theta(x) \xleftrightarrow{\mathcal{M}} \bar{\chi}_{\alpha,1}^\theta * (s) \propto \frac{\Gamma(\epsilon(s-1))}{\Gamma(\gamma(s-1))},$$

apart from the normalization constant and scale in the PDF. It is interpreted as the FCM of "one degree of freedom" in Section 7.1.

In (7.1) it is shown that the "degrees of freedom" parameter k is added to the FCM by replacing $s-1$ with $s+k-2$ such that

$$\bar{\chi}_{\alpha,k}^\theta(x) \xleftrightarrow{\mathcal{M}} \bar{\chi}_{\alpha,k}^\theta * (s) \propto \frac{\Gamma(\epsilon(s+k-1))}{\Gamma(\gamma(s+k-1))}.$$

Next, it is natural to use $\bar{\chi}_{\alpha,k}^\theta * (s)$ in the Mellin space to extend L_α^θ as follows.

DEFINITION 11.1 (The ratio-distribution representation of (unadjusted) GAS). The Mellin transform of the PDF of (unadjusted) GAS is defined as

$$(11.1) \quad \tilde{L}_{\alpha,k}^\theta(x) \xleftrightarrow{\mathcal{M}} \tilde{M}_\gamma^*(s) \bar{\chi}_{\alpha,k}^\theta * (2-s)$$

Based on the Mellin transform, its PDF can be written in a ratio distribution form,

$$(11.2) \quad \tilde{L}_{\alpha,k}^{\theta}(x) := \int_0^{\infty} \tilde{M}_{\gamma}(xs) \bar{\chi}_{\alpha,k}^{\theta}(s) s ds \quad (x \geq 0)$$

Since the Mellin integral is only valid for $x \geq 0$, it is supplemented with *the reflection rule*:

$$(11.3) \quad \tilde{L}_{\alpha}^{\theta}(-x) := \tilde{L}_{\alpha}^{-\theta}(x)$$

Thus, we have constructed a version of GAS for $x \in \mathbb{R}$, which produces fat tails and skewness -

- (1) $\tilde{L}_{\alpha,k}^{\theta}$ subsumes the α -stable distribution L_{α}^{θ} .
- (2) $\tilde{L}_{\alpha,k}^{\theta}$ subsumes Student's t distribution t_k .
- (3) $\tilde{L}_{\alpha,k}^{\theta}$ subsumes the power-exponential distribution, with the proper definition of negative k in FCM.

What is wrong with it? The problem is that the PDF and its derivatives are discontinuous at $x = \pm 0$ when $k \neq 1$ and $\theta \neq 0$.

The remaining sections of this chapter will explain this problem and provide a remediation. The reader who just wants to explore the skew-normal implementation can safely skip the rest of this chapter. The conclusion is that such discontinuity makes the PDF far from mathematical elegance, which motivates the author to explore other alternatives. The answer is to abandon the M-Wright kernel for skewness ($\tilde{M}_{\gamma}(xs)$ in (11.2)), and integrate with the skew-normal distribution, outlined in the next chapter.

11.2. Limitation

The issue of discontinuity of the PDF $\tilde{L}_{\alpha,k}^{\theta}(x)$ at $x = 0$ is encountered when $k \neq 1$. We lay out a generic framework to understand and address it.

Assume that the unadjusted two-sided density function is $\tilde{f}(x) := \tilde{L}_{\alpha,k}^{\theta}(x)$, which is discontinuous at $x = 0$. It also must satisfy the reflection rule, where, for $x > 0$, $\tilde{f}(x) := \tilde{f}^{+}(x)$ and $\tilde{f}(-x) := \tilde{f}^{-}(-x)$. $\tilde{f}(x)$ can be expanded at $x = 0$ in terms of x by

$$(11.4) \quad \tilde{f}^{\pm}(x) := \tilde{L}_{\alpha,k}^{\pm\theta}(x) = \tilde{f}_0^{\pm} + \tilde{f}_1^{\pm} x + \dots$$

where $\tilde{f}_0^{\pm}$ are the densities at $x = 0$, and $\tilde{f}_1^{\pm}$ are the respective slopes (aka the first derivatives).

The series expansion can be achieved via either (11.2), or (11.1) in conjunction with Ramanujan's master theorem in Section 2.3, such that

$$(11.5) \quad \tilde{f}_0^{+} = \frac{\gamma^{1-\gamma}}{\Gamma(1-\gamma)} E(X|\bar{\chi}_{\alpha,k}^{\theta}),$$

$$(11.6) \quad \tilde{f}_1^{+} = \frac{-\gamma^{1-2\gamma}}{\Gamma(1-2\gamma)} E(X^2|\bar{\chi}_{\alpha,k}^{\theta}).$$

Notice that they are based on the first and second moments of $\bar{\chi}_{\alpha,k}^{\theta}$. ($\tilde{f}_0^{-}, \tilde{f}_1^{-}$) are obtained by applying the reflection rule from $(\tilde{f}_0^{+}, \tilde{f}_1^{+})$. That is, θ is replaced with $-\theta$, and γ with $1-\gamma$ in every occurrence of the formula.

Furthermore, it is known that

$$(11.7) \quad \int_0^{\infty} \tilde{f}^{+}(x) dx = \gamma, \quad \int_0^{\infty} \tilde{f}^{-}(x) dx = 1 - \gamma.$$

These two are the only conditions required for $\tilde{f}^{\pm}(x)$.

The discontinuity occurs because $\tilde{f}_0^{+} \neq \tilde{f}_0^{-}$ and $\tilde{f}_1^{+} \neq \tilde{f}_1^{-}$ when $k \neq 1$ and $\theta \neq 0$. In fact, this is true for all orders of derivatives $\tilde{f}_n^{+} \neq \tilde{f}_n^{-}$ in the n -th term, $\tilde{f}_n^{\pm} x^n$.

Obviously, when $\theta = 0$, the density function is symmetric by definition: $\tilde{f}^+(x) = \tilde{f}^-(x)$. There is no issue here. So the issue is specific to the injection of skewness from $\theta \neq 0$.

On the other hand, when $k = 1$, the density function is continuous under the reflection rule, regardless the value of θ . This is the original α -stable distribution. It is perfectly fine. So the issue is specific to our attempt of adding degrees of freedom $k \neq 1$.

Either one of θ or k are fine, but when we try to do both, the distribution is broken, so to speak. That is the limitation. The dilemma is that adding θ and k is exactly what we try to achieve.

11.3. Workaround

An adjustment algorithm is proposed such that the PDF and its first derivative are continuous.

DEFINITION 11.2 (The adjusted GAS). The PDF of the adjusted GAS is defined as

$$(11.8) \quad L_{\alpha,k}^{\pm\theta}(x) := \frac{1}{A^{\pm}\sigma^{\pm}} \tilde{f}^{\pm}(x) \left(\frac{x}{\sigma^{\pm}} \right) \quad (x \geq 0)$$

It is required that (a) the new density function satisfies the reflection rule of $L_{\alpha,k}^{\theta}(-x) := L_{\alpha,k}^{-\theta}(x)$; (b) $A^{\pm}, \sigma^{\pm}$ are constrained by the continuity conditions that, at $x = 0$, both its density is continuous: $L_{\alpha,k}^{\theta}(0) = L_{\alpha,k}^{-\theta}(0)$; and its slope is continuous: $\frac{d}{dx} L_{\alpha,k}^{\theta}(0) = -\frac{d}{dx} L_{\alpha,k}^{-\theta}(0)$.

With such definition, we proceed to find the solutions of $A^{\pm}, \sigma^{\pm}$. The solutions form a distribution family. There is a canonical solution, simple and elegant, from which all other solutions are derived as a member of the location-scale family.

A member in the location-scale family shares the same "shapes" such as the skewness and kurtosis. Apart from the location and scale, it brings nothing new to the table. Hence, we can focus on analyzing the canonical distribution.

DEFINITION 11.3 (Two essential quantities for the canonical distribution). We define two essential quantities:

$$(11.9) \quad \Sigma := -\frac{\tilde{f}_0^+ \tilde{f}_1^-}{\tilde{f}_0^- \tilde{f}_1^+}$$

$$(11.10) \quad \Psi := \Sigma \frac{\tilde{f}_0^+}{\tilde{f}_0^-} = -\left(\frac{\tilde{f}_0^+}{\tilde{f}_0^-} \right)^2 \frac{\tilde{f}_1^-}{\tilde{f}_1^+}$$

Notice that $\tilde{f}_0^+/\tilde{f}_0^-$ is the ratio of the original densities from two sides of $x = 0$. And $\tilde{f}_1^-/\tilde{f}_1^+$ is the ratio of the slopes of the two sides. Since $\tilde{f}_1^-, \tilde{f}_1^+$ always have the opposite signs, Σ is a positive quantity.

Note that Σ is singular when $\gamma = 1/2$. Both $\tilde{f}_1^-, \tilde{f}_1^+$ approach zero at the same speed. Hence, $\Sigma \rightarrow 1$ and $\Psi \rightarrow 1$.

The most important contribution is the discovery of the canonical distribution.

DEFINITION 11.4 (The canonical GAS). The canonical GAS distribution is defined according to $\sigma^+ = 1$ and $\sigma^- = \Sigma$. Hence, its PDF for $x \geq 0$ is (with the hat symbol)

$$(11.11) \quad \hat{L}_{\alpha,k}^{\theta}(x) := \frac{1}{A^+} \tilde{f}^+(x)$$

$$(11.12) \quad \hat{L}_{\alpha,k}^{-\theta}(x) := \frac{1}{A^-\Sigma} \tilde{f}^-\left(\frac{x}{\Sigma}\right)$$

where $A^+ = \gamma + \Psi(1 - \gamma)$ and $A^- = A^+/\Psi$ from Lemma 11.7.

The reflection rule applies: $\hat{L}_{\alpha,k}^{\theta}(-x) := \hat{L}_{\alpha,k}^{-\theta}(x)$.

11.3.1. The Location-scale Family. The following lemmas show that all other solutions must obey $\sigma^-/\sigma^+ = \Sigma$. They are just the location-scale family of the canonical distribution.

Briefly, all other solutions are defined by a choice of scale $\sigma^+ > 0$, such that

$$(11.13) \quad L_{\alpha,k}^\theta(x) := \frac{1}{\sigma^+} \widehat{L}_{\alpha,k}^\theta\left(\frac{x}{\sigma^+}\right)$$

For instance, we found that $\sigma^+ = \Sigma^\gamma$ to be a very good alternative. In the remark of Definition 11.9, we show that the n -th moment of $L_{\alpha,k}^\theta$ is just that of $\widehat{L}_{\alpha,k}$ multiplied by its scale $(\sigma^+)^n$.

LEMMA 11.5. The requirement that the density and slope of the *adjusted* density function should be smooth at $x = 0$ leads to

$$(11.14) \quad \frac{1}{A^+\sigma^+} \tilde{f}_0^+ = \frac{1}{A^-\sigma^-} \tilde{f}_0^-$$

$$(11.15) \quad \frac{1}{A^+(\sigma^+)^2} \tilde{f}_1^+ = -\frac{1}{A^-(\sigma^-)^2} \tilde{f}_1^-$$

△

PROOF. To solve $A^\pm$ and $\sigma^\pm$, take (11.8) and carry out the series expansions from (11.4):

$$(11.16) \quad L_{\alpha,k}^{\pm\theta}(x) = \frac{\tilde{f}_0^\pm}{A^\pm\sigma^\pm} + \frac{\tilde{f}_1^\pm}{A^\pm(\sigma^\pm)^2} x + \dots$$

(11.14) is straightforward from requiring $L_{\alpha,k}^\theta(0) = L_{\alpha,k}^{-\theta}(0)$ in (11.16). Likewise, (11.15) is the result of $\frac{d}{dx}L_{\alpha,k}^\theta(0) = \frac{d}{dx}L_{\alpha,k}^{-\theta}(0)$ from (11.16). □

LEMMA 11.6. The equations in Lemma 11.5 lead to the following invariant:

$$(11.17) \quad \frac{\sigma^-}{\sigma^+} = \Sigma$$

△

PROOF. Divide the LHS and RHS of (11.14) by those of (11.15) respectively,

$$\sigma^+ \frac{\tilde{f}_0^+}{\tilde{f}_1^+} = -\sigma^- \frac{\tilde{f}_0^-}{\tilde{f}_1^-}$$

Rearrange the items and we obtain (11.17). □

LEMMA 11.7. The solution for $A^\pm$ are

$$(11.18) \quad A^+ = \gamma + \Psi(1 - \gamma)$$

$$(11.19) \quad A^+/A^- = \Psi$$

△

PROOF. (11.19) is derived by rearranging the items in (11.14) and following the definition of Ψ .

(11.18) is derived from the fact that the total density of the adjusted distribution should be equal to 1, that is, $\int_{-\infty}^{\infty} f(x)dx = 1$. Hence,

$$\int_0^{\infty} f^+(x)dx + \int_0^{\infty} f^-(x)dx = \frac{1}{A^+} \int_0^{\infty} \tilde{f}^+(x)dx + \frac{1}{A^-} \int_0^{\infty} \tilde{f}^-(x)dx = 1$$

Apply (11.7), we get $\frac{\gamma}{A^+} + \frac{1-\gamma}{A^-} = 1$. Multiply it by A^+ on both sides, we obtain (11.18). □

We've shown that $A^\pm$ are well-defined constants based on (α, k, θ) , while $\sigma^\pm$ is a choice of parametrization, constrained by (11.17).

11.4. Moments

The structure of the *moments* reveals critical information about the adjusted distribution. We show the moment formula of the canonical distribution, and how the location-scale family relates to it.

To simplify the notations below, let

- $f^\pm = L_{\alpha,k}^{\pm\theta}$ be the adjusted distribution family,
- $\hat{f}^\pm = \hat{L}_{\alpha,k}^{\pm\theta}$ be the canonical distribution,
- $\tilde{f}^\pm = \tilde{L}_{\alpha,k}^{\pm\theta}$ be the original (unadjusted) distribution.

First, the n -th one-sided moments of the adjusted distribution are ($x > 0$)

$$(11.20) \quad E(X^n|f^\pm) = \frac{1}{A^\pm \sigma^\pm} \int_0^\infty x^n \tilde{f}^\pm(x/\sigma^\pm) dx = \frac{(\sigma^\pm)^n}{A^\pm} E(X^n|\tilde{f}^\pm)$$

where $E(X^n|\tilde{f}^\pm)$ are the original n -th one-sided moments. They can be obtained from the Mellin transform (11.1).

The n -th total moment, given the notation of m_n , is the sum of $E(X^n|f^+)$ and $(-1)^n E(X^n|f^-)$. We show the following.

LEMMA 11.8. The n -th total moment of the adjusted distribution is based on the original one-sided moments such as

$$(11.21) \quad m_n := E(X^n|f) = \frac{(\sigma^+)^n}{A^+} \left[E(X^n|\tilde{f}^+) + \Psi(-\Sigma)^n E(X^n|\tilde{f}^-) \right]$$

△

PROOF. By definition, we have

$$\begin{aligned} m_n := E(X^n|f) &= \int_{-\infty}^\infty x^n f(x) dx = \int_0^\infty x^n f^+(x) dx + (-1)^n \int_0^\infty x^n f^-(x) dx \\ &= E(X^n|f^+) + (-1)^n E(X^n|f^-) \end{aligned}$$

Apply (11.20), we get

$$m_n = \frac{(\sigma^+)^n}{A^+} E(X^n|\tilde{f}^+) + \frac{(-\sigma^-)^n}{A^-} E(X^n|\tilde{f}^-)$$

Factor out $\frac{(\sigma^+)^n}{A^+}$, apply $\sigma^-/\sigma^+ = \Sigma$ from Lemma 11.6, and $A^+/A^- = \Psi$ from 11.7, we obtain (11.21). □

LEMMA 11.9 (The moments of the canonical distribution). The n -th moment of the canonical distribution is

$$(11.22) \quad \hat{m}_n := E(X^n|\hat{f}) = \frac{1}{A^+} \left[E(X^n|\tilde{f}^+) + \Psi(-\Sigma)^n E(X^n|\tilde{f}^-) \right]$$

△

PROOF. Lemma 11.8 shows that the canonical distribution $\hat{f}$ is obtained by letting $\sigma^+ = 1$ and $\sigma^- = \Sigma$. Put them to (11.21), we obtain (11.22). □

Lastly, compare (11.21) with (11.22). We reach $m_n = (\sigma^+)^n \hat{m}_n$. That is, all other members in the adjusted distribution family are rescaled canonical distributions.

GAS-SN: Generalized Alpha-Stable Distribution with Skew-Normal

This fractional univariate distribution combines the features from a classic skew-normal distribution that provides skewness and a fractional distribution that provides fatter tails. The resulting distribution is analytically tractable. The PDF and all of its derivatives are continuous everywhere in $\mathbb{R}$.

12.1. Definition

DEFINITION 12.1. Assume $Z_0 \sim SN(0, 1, \beta)$ is a skew-normal variable and $V \sim \bar{\chi}_{\alpha,k}$ is an FCM variable.

Then $Z \sim Z_0/V$ is a variable with a GAS-SN distribution. We use the notation $Z \sim L_{\alpha,k}(\beta)$ for this standard distribution.

Assume $\mathcal{N}(x)$ and $\Phi_{\mathcal{N}}(x)$ are the PDF and CDF of $N(0, 1)$. The PDF of Z is

$$(12.1) \quad L_{\alpha,k}(x; \beta) = 2 \int_0^\infty \mathcal{N}(xs) \Phi_{\mathcal{N}}(\beta xs) \bar{\chi}_{\alpha,k}(s) s ds.$$

This is the fractional extension of (10.2).

Its CDF is

$$(12.2) \quad \begin{aligned} \Phi[L_{\alpha,k}(\beta)](x) &:= \int_0^\infty \Phi_{SN}(xs; \beta) \bar{\chi}_{\alpha,k}(s) ds. \\ &= \int_0^\infty [\Phi_{\mathcal{N}}(xs) - 2T(xs, \beta)] \bar{\chi}_{\alpha,k}(s) ds. \end{aligned}$$

where $\Phi_{SN}(xs; \beta)$ is the CDF of $SN(0, 1, \beta)$ in (10.3), and $T(h, a)$ is the Owen's T function.

We can clearly see that the CDF has two components: One from the symmetric part, and the other skew. The second component vanishes due to $T(h, 0) = 0$.

12.1.1. GAS-SN Subsumes GSaS.

LEMMA 12.2. When $\beta = 0$, it becomes a symmetric distribution, previously called GSaS. The notation of $L_{\alpha,k}$ is given in [19].

The PDF of a GSaS is

$$(12.3) \quad L_{\alpha,k}(x) = \int_0^\infty \mathcal{N}(xs) \bar{\chi}_{\alpha,k}(s) s ds.$$

When $\alpha \rightarrow 2$ or $k \rightarrow \infty$, the symmetric distribution approaches a normal distribution $N(0, \alpha^{2/\alpha})$ (Section 8.2 of [19]).

△

This integral is a normal mixture (9.1) that enjoys several nice properties outlined in Chapter 9.

In particular, the generalized exponential power distribution can be obtained via the characteristic function transform in Lemma 9.2 (Section 9 of [19]). We point out that the skew extension is straightforward, but leave the detailed description to future research.

12.1.2. GAS-SN Subsumes Skew-t Distribution. An important bridge between SN and GAS-SN is the skew-t (ST) distribution. It is documented in Section 4.3 of [1].

ST is fully consistent with GAS-SN by setting $\alpha = 1$. That is, in his notation, $T(\beta, k) = L_{1,k}(\beta)$.

12.2. The Location-Scale Family

Its location scale family is $Y = \xi + \omega Z \sim L_{\alpha,k}(\xi, \omega^2, \beta)$. Its PDF becomes

$$(12.4) \quad \phi(x) = \frac{1}{\omega} L_{\alpha,k} \left(\frac{x - \xi}{\omega}; \beta \right). \quad (x \in \mathbb{R})$$

In real-world applications, this PDF is used for optimization, e.g. in the maximum likelihood estimation (MLE). See Section 12.9.

12.3. Mellin Transform

The Mellin transform of the PDF follows the rule of the ratio distribution. From (10.8) and (7.2), we have

$$(12.5) \quad L_{\alpha,k}(x; \beta) \xleftrightarrow{\mathcal{M}} L_{\alpha,k}^*(s; \beta) \\ = \mathcal{N}^*(s; \beta) \bar{\chi}_{\alpha,k}^*(2 - s) \\ (12.6) \quad = [2 \Phi[t_s](\beta \sqrt{s})] \times [\mathcal{N}^*(s) \bar{\chi}_{\alpha,k}^*(2 - s)]$$

Notice that the contribution for the skewness is $2 \Phi[t_s](\beta \sqrt{s})$ in the first bracket, which becomes one if $\beta = 0$.

The second bracket is the Mellin transform of the GSaS PDF. From (2.9) and (7.2), it is

$$(12.7) \quad L_{\alpha,k}(x) \xleftrightarrow{\mathcal{M}} L_{\alpha,k}^*(s) = \mathcal{N}^*(s) \bar{\chi}_{\alpha,k}^*(2 - s) \\ = \frac{1}{2\sqrt{\pi}} \left(\frac{2}{\sigma} \right)^{s-1} \Gamma\left(\frac{s}{2}\right) \frac{\Gamma((k-1)/2)}{\Gamma((k-1)/\alpha)} \frac{\Gamma((k-s)/\alpha)}{\Gamma((k-s)/2)},$$

where $\sigma := k^{1/2-1/\alpha}$ and $k > 0$ is assumed.

12.4. Moments

Based on $\mathbb{E}(X^n | \mathcal{N}(\beta))$ from (10.11), the n -th moment of Z is

$$(12.8) \quad \mathbb{E}(X^n | L_{\alpha,k}(\beta)) := \mathbb{E}(X^n | \mathcal{N}(\beta)) \mathbb{E}(X^{-n} | \bar{\chi}_{\alpha,k}) \\ = 2 \mathcal{N}^*(n+1) \mathbb{E}(X^{-n} | \bar{\chi}_{\alpha,k}) \\ \times \begin{cases} 1, & \text{when } n \text{ is even,} \\ 1 - 2\Phi[t_{n+1}](-\beta \sqrt{n+1}), & \text{when } n \text{ is odd.} \end{cases}$$

Its first moment is $\mu_z = b \delta$, where $b = \sqrt{2/\pi} \mathbb{E}(X^{-1} | \bar{\chi}_{\alpha,k})$.

The second moment is $\mathbb{E}(X^{-2} | \bar{\chi}_{\alpha,k})$. Its variance is

$$(12.9) \quad \sigma_z^2 = \mathbb{E}(X^{-2} | \bar{\chi}_{\alpha,k}) - (b \delta)^2.$$

To simplify the symbology, let $q_n := \mathbb{E}(X^{-n} | \bar{\chi}_{\alpha,k})$. The third moment is $\delta_3 q_3$, where $\delta_3 =$

$\sqrt{\frac{2}{\pi}} \delta(3 - \delta^2)$. The fourth moment is $3 q_4$. To carry out the skewness γ_1 and excess kurtosis γ_2 ,

$$\gamma_1 \times \sigma_z^{3/2} = \delta_3 q_3 - 3 \mu_z q_2 + 2 \mu_z^3, \\ \gamma_2 \times \sigma_z^4 = 3(q_4 - q_2^2) - 4 \mu_z (\gamma_1 \times \sigma_z^{3/2}) + 2 \mu_z^4.$$

The maximum skewness and kurtosis can be infinite. Since $\delta = \sin \theta$, where $\beta = \tan \theta$, we have $\delta \in [-1, 1]$. Infinity has to come from q_3 and q_4 .

A typical example is the skew-t distribution at $\alpha = 1$. It is well known that kurtosis approaches infinity when k approaches 4 from above, and the skewness approaches infinity when k approaches 3 from above.

12.4.1. Excess Kurtosis of GSaS. It is important to understand the behavior of excess kurtosis γ_2 . However, the presence of skewness adds more complexity to γ_2 . Consider the symmetric case where $\beta = 0$, and we quote the result from [19] below.

The excess kurtosis of GSaS is plotted in Figure 12.1 in the (k, α) coordinate. Notice that a major division occurs along the line of $k = 5 - \alpha$. In the region where $0 < k \leq 5 - \alpha$, there are complicated patterns caused by the infinities of the gamma function. Only small pockets of valid kurtosis exist.

LEMMA 12.3. In the region where $k > 5 - \alpha$, the excess kurtosis of GSaS is a continuous function with positive values. At large k 's, the closed form of the moments can be expanded by Sterling's formula. The excess kurtosis γ_2 becomes part of a linear equation:

$$(12.10) \quad \left(\epsilon - \frac{1}{2}\right) = \left(\frac{k-3}{4}\right) \log\left(1 + \frac{\gamma_2}{3}\right), \quad \text{where } \epsilon = 1/\alpha$$

This equation shows how GSaS works under the **Central Limit Theorem**. GSaS approaches a normal distribution when γ_2 becomes zero. This can happen from two directions: when $\alpha \rightarrow 2$ or when $k \rightarrow \infty$.

△

The contour plot of excess kurtosis is shown in the (k, ϵ) coordinate in Figure 12.2. It is visually amusing. Notice the singular point at $\epsilon = 1/2, k = 3$.




FIGURE 12.1. The contour plot of excess kurtosis in GSaS by (k, α) .




FIGURE 12.2. The contour plot of excess kurtosis in GSaS by (k, ϵ) where $\epsilon = 1/\alpha$. This best describes the linearity in (12.10) for large k 's.

1592

1593 **12.4.2. Standardized Peak Density of GSaS.** For the symmetric case $\beta = 0$, the GAS-SN
 1594 density reduces to the GSaS density in (12.3). Its mode is located at the center. Under the location-
 1595 scale transformation

$$Y = \xi + \omega Z, \quad \phi_Y(x) = \frac{1}{\omega} L_{\alpha,k} \left(\frac{x - \xi}{\omega} \right),$$

1596 the peak height changes by the factor $1/\omega$. Hence the location-scale invariant peak height is obtained
 1597 by multiplying the peak density by the standard deviation.

1598 **DEFINITION 12.4.** Assume $Z \sim L_{\alpha,k}$ and that $\sigma_Z^2 < \infty$. The standardized peak density of GSaS
 1599 is defined as

$$(12.11) \quad \text{SPD}_{\alpha,k} := \sigma_Z L_{\alpha,k}(0) = \frac{1}{\sqrt{2\pi}} \sqrt{\mathbb{E}(X^{-2}|\bar{\chi}_{\alpha,k})} \mathbb{E}(X|\bar{\chi}_{\alpha,k}).$$

1600 Equivalently, for any location-scale version $Y = \xi + \omega Z$, it is

$$(12.12) \quad \text{SPD}_{\alpha,k} = \sigma_Y \phi_Y(\xi),$$

1601 so the quantity is independent of ξ and ω .

1602 **PROOF.** From (12.3), since $\mathcal{N}(0) = 1/\sqrt{2\pi}$,

$$L_{\alpha,k}(0) = \frac{1}{\sqrt{2\pi}} \mathbb{E}(X|\bar{\chi}_{\alpha,k}).$$

1603 In the symmetric case $\beta = 0$, (12.9) gives

$$\sigma_Z^2 = \mathbb{E}(X^{-2}|\bar{\chi}_{\alpha,k}).$$

1604 Therefore,

$$\text{SPD}_{\alpha,k} = \frac{1}{\sqrt{2\pi}} \sqrt{\mathbb{E}(X^{-2}|\bar{\chi}_{\alpha,k}) \mathbb{E}(X|\bar{\chi}_{\alpha,k})}.$$

1605

□

1606 Using (7.8), the same expression can be written as

$$(12.13) \quad \text{SPD}_{\alpha,k} = \frac{1}{\sqrt{2\pi}} \left[\frac{\Gamma((k-1)/2)}{\Gamma((k-1)/\alpha)} \right]^{3/2} \frac{\Gamma(k/\alpha)}{\Gamma(k/2)} \left[\frac{\Gamma((k-3)/\alpha)}{\Gamma((k-3)/2)} \right]^{1/2}.$$

1607 As a probabilistic standardized peak density, this formula requires the underlying gamma functions
1608 to render a positive number. The analytic continuation of (12.13) has a more complicated finite-real
1609 region in the (k, α) plane, as shown below.

1610 The contour plot of standardized peak density is shown in the (k, ϵ) coordinate in Figure 12.3.
1611 This view highlights the approximate linearity in (12.14) for large k 's.




FIGURE 12.3. The contour plot of standardized peak density in GSaS by (k, ϵ) , where $\epsilon = 1/\alpha$.

1612 LEMMA 12.5. At large k 's, the closed form (12.13) can be expanded by Stirling's formula. The
1613 standardized peak density becomes part of a linear equation:

$$(12.14) \quad \left(\epsilon - \frac{1}{2} \right) = \frac{2}{3} \left(k - \frac{4}{3} \right) \log \left(\sqrt{2\pi} \text{SPD}_{\alpha,k} \right), \quad \text{where } \epsilon = 1/\alpha.$$

This equation shows how the standardized peak density approaches the normal value under the **Central Limit Theorem**. The normal reference is $\text{SPD}_{2,\infty} = 1/\sqrt{2\pi} \approx 0.398942$, for which the logarithmic term vanishes.

△

12.4.3. Lognormal-Normal Asymptotics of GSaS. Combining (12.10) and (12.14), and assuming $\epsilon \neq 1/2$, eliminates the common factor $\epsilon - 1/2$. This gives

$$(12.15) \quad 1 + \frac{\gamma_2}{3} = \left(\sqrt{2\pi} \text{SPD}_{\alpha,k} \right)^H, \quad \text{where } H = \frac{8}{3} \left(\frac{1 - 4/(3k)}{1 - 3/k} \right).$$

As $k \rightarrow \infty$, $H \rightarrow 8/3$. The same exponent appears in the lognormal-normal limit below.

DEFINITION 12.6 (The Lognormal-Normal Distribution (LNN)). Let $Z \sim \mathcal{N}(0, 1)$ and $Y \sim \mathcal{N}(0, \sigma^2)$ be independent. The random variable $X = Ze^Y$ belongs to the lognormal-normal distribution family, denoted by $\text{LNN}(\sigma^2)$, where σ is the shape parameter[4].¹

The LNN has exact even moments

$$\mathbb{E}(X^{2m}) = (2m - 1)!! e^{2m^2 \sigma^2}.$$

Therefore, the variance is $e^{2\sigma^2}$ and

$$1 + \frac{\gamma_2}{3} = e^{4\sigma^2},$$

$$\sqrt{2\pi} \text{SPD}_{\text{LNN}} = e^{3\sigma^2/2}.$$

Eliminating σ^2 gives

$$(12.16) \quad 1 + \frac{\gamma_2}{3} = \left(\sqrt{2\pi} \text{SPD}_{\text{LNN}} \right)^{8/3}.$$

The LNN limit of GSaS is explained by Lemma 7.4. From Definition 12.1, the GSaS can be written as

$$X = \frac{Z}{V}, \quad Z \sim \mathcal{N}(0, 1), \quad V \sim \bar{\chi}_{\alpha,k}.$$

When $k \rightarrow \infty$ and $T = 1/(\alpha|k|)$ is kept at a nonzero finite value, the centered scale variable $V/e^{\mathbb{E}[\log V]}$ approaches e^Y , where $Y \sim \mathcal{N}(0, T)$. Hence, up to the deterministic scale factor $e^{-\mathbb{E}[\log V]}$,

$$\frac{Z}{V} \approx Ze^{-Y}.$$

Since $-Y$ has the same distribution as Y , this limiting shape is equivalently Ze^Y , the LNN form. Because both γ_2 and SPD are invariant under location-scale transformations, the deterministic scale factor does not affect the curve in the (γ_2, SPD) plane. Hence the terminal GSaS relation is the LNN relation (12.16).

This gives a practical visualization of GSaS for data analysis. Combining Figures 12.2 and 12.3 produces the contour plot in the γ_2 -SPD plane shown in Figure 12.4.

For a data set such as daily SPX log returns from 1990 to mid-2025 (see Figure 12.8), one can compute the sample kurtosis and estimate SPD, then locate the corresponding point on this chart. The chart gives an approximate GSaS starting value, which can then be refined by maximum likelihood estimation (MLLK).

At $\sigma = 0$, the LNN reduces to the standard normal distribution; as $\sigma \rightarrow 0$, it approaches this normal limit. This is the **Central Limit Theorem** direction, corresponding to the lower-left endpoint of Figure 12.4.

¹The author has studied LNN in a 2008 working paper. See <https://ssrn.com/abstract=1537686>. However, the role of LNN was rediscovered in 2026, when I attempted to fit the bear state of the global regime in the financial market. See https://github.com/slihn/Mixture-VAE/blob/main/slides/frac_slides.md.




FIGURE 12.4. The relation between excess kurtosis γ_2 and the standardized peak density (SPD) in GSaS, indicated by representative k 's. We added three points from the SPX global regime analysis to illustrate the usage (see footnote). "All data" is the statistics from the full dataset. "Bull data" is the statistics from the bull state, which lies in the negative k branch. "Bear data" is the statistics from the bear state, which lies on the terminal curve.

12.5. Tail Behavior

The tail behavior of GAS-SN is a "modified GSaS" type. Hence, it is well within what was known. Without losing generality, assume $\beta > 0$, that the decay of the left tail is more pronounced than that of the right tail. But it still follows the same power law of x^{-k} as in a $L_{\alpha,k}$.

It takes a small tweak to GSaS to capture that behavior.

DEFINITION 12.7. The shifted GSaS is defined as

$$(12.17) \quad L_{\alpha,k}(x|\mu) = \int_0^\infty \mathcal{N}(xs - \mu) \bar{\chi}_{\alpha,k}(s) s ds$$

Note that the shift μ is not a location parameter that shifts x . It is a shift inside the argument of $\mathcal{N}(\cdot)$. When $\mu = 0$, it is restored to the PDF of GSaS, $L_{\alpha,k}(x)$.

We use the following approximation of the erf function in (12.1)[15]

$$(12.18) \quad 1 - \text{erf}(x) \approx \frac{1}{B\sqrt{\pi}x} (1 - e^{-Ax}) e^{-x^2} \quad (x \geq 0)$$

where $A = 1.98$ and $B = 1.135$. It is much better than the first-order expansion of $e^{-x^2}/(\sqrt{\pi}x)$ for the entire range of $x \in [0, \infty)$.

LEMMA 12.8. For $\beta > 0$, the left tail ($x < 0$) of the PDF in (12.1) can be approximated by

$$(12.19) \quad \widehat{L}_{\alpha,k}(x; \beta) = \frac{G}{\beta x} \left[e^{\mu^2/2} L_{\alpha,k-1}(qx||\mu) - L_{\alpha,k-1}(qx) \right]$$

where

$$\begin{aligned} \mu &= \frac{A\delta}{\sqrt{2}}, \quad \delta = \frac{\beta}{\sqrt{1+\beta^2}}, \\ q &= \sqrt{1+\beta^2} \frac{\sigma_{\alpha,k}}{\sigma_{\alpha,k-1}} \\ G &= \sqrt{\frac{2}{\pi}} \frac{C_{\alpha,k}}{B \sigma_{\alpha,k-1} C_{\alpha,k-1}} \end{aligned}$$

and both $C_{\alpha,k} = \frac{\alpha \Gamma((k-1)/2)}{\Gamma((k-1)/\alpha)}$ and $\sigma_{\alpha,k}$ are according to FCM in (7.4). The factor $1/\beta$ is part of this tail approximation and should not be interpreted as a singularity of the exact PDF. When $\beta = 0$, (12.1) reduces directly to the symmetric density $L_{\alpha,k}(x; \beta) = L_{\alpha,k}(x)$.

The right tail ($x > 0$) is simply

$$(12.20) \quad 2L_{\alpha,k}(x) - \widehat{L}_{\alpha,k}(-x; \beta)$$

where the second term $\widehat{L}_{\alpha,k}(-x; \beta)$ becomes much smaller than the first term as $x \rightarrow \infty$. $\triangle$

PROOF. Assume $\beta > 0$ and $x < 0$. Since $\Phi_{\mathcal{N}}(z) = \frac{1}{2}[1 + \text{erf}(z/\sqrt{2})]$, the skew-normal factor in (12.1) satisfies

$$2\Phi_{\mathcal{N}}(\beta xs) = 1 - \text{erf}\left(-\frac{\beta xs}{\sqrt{2}}\right).$$

Set $u = -\beta xs/\sqrt{2}$, so that $u > 0$. Applying the approximation in the preceding display gives the factor $1/u = -\sqrt{2}/(\beta xs)$; the factor s in the integral then cancels, leaving $1/(\beta x)$. Thus,

$$\begin{aligned} L_{\alpha,k}(x; \beta) &\approx \int_0^\infty \mathcal{N}(xs) \frac{1}{B\sqrt{\pi}u} (1 - e^{-Au}) e^{-u^2} \bar{\chi}_{\alpha,k}(s) s ds \\ &= -\frac{1}{B\pi\beta x} \int_0^\infty e^{-(1+\beta^2)x^2 s^2/2} \left(1 - e^{A\beta xs/\sqrt{2}}\right) \bar{\chi}_{\alpha,k}(s) ds. \end{aligned}$$

Now substitute

$$r = \frac{\sigma_{\alpha,k-1}}{\sigma_{\alpha,k}} s, \quad y = qx = \sqrt{1+\beta^2} \frac{\sigma_{\alpha,k}}{\sigma_{\alpha,k-1}} x.$$

Using the FCM density in (7.4), this change of variables gives

$$\bar{\chi}_{\alpha,k}(s) ds = \frac{C_{\alpha,k}}{\sigma_{\alpha,k-1} C_{\alpha,k-1}} \bar{\chi}_{\alpha,k-1}(r) r dr.$$

Moreover, since

$$\delta = \frac{\beta}{\sqrt{1+\beta^2}}, \quad \mu = \frac{A\delta}{\sqrt{2}},$$

we have

$$\mu y r = \frac{A\beta xs}{\sqrt{2}}.$$

Therefore,

$$\begin{aligned} L_{\alpha,k}(x; \beta) &\approx -\frac{1}{B\pi\beta x} \frac{C_{\alpha,k}}{\sigma_{\alpha,k-1} C_{\alpha,k-1}} \int_0^\infty e^{-(yr)^2/2} (1 - e^{\mu y r}) \bar{\chi}_{\alpha,k-1}(r) r dr \\ &= \frac{1}{\beta x} \sqrt{\frac{2}{\pi}} \frac{C_{\alpha,k}}{B \sigma_{\alpha,k-1} C_{\alpha,k-1}} \left[e^{\mu^2/2} L_{\alpha,k-1}(y||\mu) - L_{\alpha,k-1}(y) \right], \end{aligned}$$

where the last equality follows from the shifted-GSaS definition with k replaced by $k-1$. Since $y = qx$, this proves the stated left-tail approximation.

For the right tail, use the identity

$$2\Phi_{\mathcal{N}}(z) + 2\Phi_{\mathcal{N}}(-z) = 2.$$

Together with the symmetry of $\mathcal{N}(xs)$, this implies

$$L_{\alpha,k}(x; \beta) + L_{\alpha,k}(-x; \beta) = 2L_{\alpha,k}(x).$$

Replacing the left-tail term by the approximation just derived gives

$$L_{\alpha,k}(x; \beta) \approx 2L_{\alpha,k}(x) - \hat{L}_{\alpha,k}(-x; \beta), \quad x > 0.$$

□

REMARK 12.9. In the case of $\beta \rightarrow 0$, the apparent $1/\beta$ singularity in the approximation is removable at the level of the formal expansion. Indeed,

$$\mu = \frac{A\delta}{\sqrt{2}} = \frac{A\beta}{\sqrt{2}} + O(\beta^3),$$

and the shifted and unshifted terms in the bracket differ by

$$e^{\mu^2/2} L_{\alpha,k-1}(qx||\mu) - L_{\alpha,k-1}(qx) = O(\mu) = O(\beta).$$

This cancels the explicit $1/\beta$ factor. However, the approximation is not uniform as $\beta \rightarrow 0$. For fixed x , the error-function approximation has the local limit

$$\lim_{z \rightarrow 0} \frac{(1 - e^{-Az})e^{-z^2}}{B\sqrt{\pi}z} = \frac{A}{B\sqrt{\pi}} \approx 0.984225,$$

whereas the exact complementary error-function factor equals 1 at the origin. Hence the formula is best viewed as a suppressed-tail approximation, effectively for $\beta|x| \gg 1$, rather than a formula to use near $\beta = 0$.

12.6. Maximum Skewness and Half GSaS

When $\beta \rightarrow \pm\infty$, a GAS-SN becomes a half-GSaS, which is a one-sided distribution with the notation of $L_{\alpha,k}^{\pm} := L_{\alpha,k}(\beta = \pm\infty)$. Its PDF is

$$(12.21) \quad L_{\alpha,k}^+(x) = \begin{cases} 2L_{\alpha,k}(x) & \text{if } x \geq 0, \\ 0 & \text{otherwise.} \end{cases}$$

It follows the reflection rule of $L_{\alpha,k}^-(x) = L_{\alpha,k}^+(-x)$. Hence, we only need to study the $+\infty$ case.

A half-GSaS possesses the maximum skewness that a GAS-SN family can achieve for a given pair of (α, k) . In Section 10.5, it was mentioned that the maximum skewness of the SN family is only 0.9953. GAS-SN allows the skewness to reach infinity potentially.

From (12.7), the n -th moment is

$$(12.22) \quad \begin{aligned} \mathbb{E}(X^n | L_{\alpha,k}^+) &= 2L_{\alpha,k}^*(n+1) \\ \mathbb{E}(X^n | L_{\alpha,k}^-) &= 2L_{\alpha,k}^*(n+1)(-1)^n \end{aligned}$$

Therefore, it is straightforward to calculate the skewness.

The skewness of half-GSaS $L_{\alpha,k}^+$ is shown in Figure 12.5 in the (k, α) coordinate. There is a clear division of infinity by the line from $(2, 2)$ to $(4, 0)$.

The contour plot of the skewness is shown in the (k, ϵ) coordinate in Figure 12.6. Each contour line approaches a straight line as k increases.




FIGURE 12.5. The contour plot of skewness of the half-GSaS $L_{\alpha,k}^+$ by (k, α) . This represents the maximum skewness that the GAS-SN family can achieve.




FIGURE 12.6. The contour plot of skewness of the half-GSaS $L_{\alpha,k}^+$ by (k, ϵ) where $\epsilon = 1/\alpha$. Each contour line approaches a straight line as k increases.

12.7. Fractional Skew Exponential Power Distribution

As shown in Definition 3.6 and Section 9 of [19], the negative k space is reserved for the fractional exponential power distribution, whose PDF is $\mathcal{E}_{\alpha,k}(x) := L_{\alpha,-k}(x)$. All it takes is to have $\bar{\chi}_{\alpha,k}(s)$ in (12.1) properly defined for negative k , which is done in (7.11).

It is natural to extend it with the skew-normal family such that its PDF becomes

$$(12.23) \quad \mathcal{E}_{\alpha,k}(x; \beta) = L_{\alpha,-k}(x; \beta).$$

Then we obtain another flexible skew distribution with a different type of tail behavior. Detailed analysis of this distribution is left for future research.

12.8. Quadratic Form

A squared GAS-SN variable Q is distributed as a fractional F distribution with $d = 1$. That is,

$$(12.24) \quad Q := \left(\frac{Y - \xi}{\omega} \right)^2 = Z^2 \sim F_{\alpha,1,k}, \quad \text{for all } \beta.$$

Notice that Q is based on the standard variable Z , which is invariant to the location and scale. See Chapter 8 for more detail.

12.9. Univariate MLE

In this section, we document how we fit the one-dimensional data with univariate GAS-SN. The main algorithm is *maximum likelihood estimation* (MLE), supplemented with several components of regularization.

We applied the MLE fitting on the daily returns of VIX and SPX from 1990 to mid-2025, each about 8900 samples. The MLE program is implemented in **python** and **scipy** on github at https://github.com/slihn/gas-impl/blob/main/gas_impl/mle_gas_sn.py.

In the univariate case, the hyperparameter space is $\Theta = \{\alpha, k, \beta, \omega, \xi\}$, where $\alpha \in (0, 2)$, $k \in (2, \infty)$, $\omega > 0$, and $\beta, \xi \in \mathbb{R}$. Assume there are N samples in the data set, $Y = \{y_i, i \in 1, 2, \dots, N\}$, the main component of the objective function is the minus log-likelihood (MLLK):

$$(12.25) \quad \text{MLLK}(\Theta) = -\frac{1}{N} \sum_{i=1}^N \log(\phi(y_i; \Theta))$$

where $\phi(y; \Theta)$ is the PDF of the univariate location scale family (12.4).

Additional components of regularization are added to the objective function $\ell(\Theta)$. Specifically, the L2 distances between the empirical and theoretical statistics are added as follows:

- Skewness: $|\Delta\gamma_1|^2 := |\Delta\text{skewness}(Y)|^2$. Section 12.4.
- Kurtosis: $|\Delta\gamma_2|^2 := |\Delta\text{kurtosis}(Y)|^2$. Section 12.4.
- The mean of the quadratic form: $\Delta\mu_Q^2 := |\Delta\text{mean}(Q)|^2$. Section 15.6.

The MLE seeks the optimal Θ that minimizes the objective function:

$$(12.26) \quad \hat{\Theta} = \text{argmin } \ell(\Theta)$$

$$(12.27) \quad \text{where } \ell(\Theta) = \text{MLLK}(\Theta) + |\Delta\gamma_1|^2 + |\Delta\gamma_2|^2 + \Delta\mu_Q^2$$

A custom version of the stochastic descent (SD) algorithm is developed. Our experience shows that it is better to standardize the data set to one standard deviation, so that all parameters in Θ are approximately on the same scale.

It is also important to control the learning rate so that it does not take a too large step on α , empirically, no more than 0.01 per step. This ensures that the SD does not wander into the *undefined* regions for $\ell(\Theta)$. This is particularly important for the SPX fit below.

The SD algorithm calculates the gradients for each hyperparameter. And make a small move along the direction that is most likely to minimize $\ell(\Theta)$. The scale of the move is based on the learning rate, which can be dynamically adjusted. Some randomness is added to the small move. This allows the algorithm to explore the nearby region and increases its choices.

12.10. Examples of Univariate MLE Fits

12.10.1. VIX fit. Figure 12.7 shows the result of the MLE fit to the daily VIX returns from 1990 to mid-2025. Data are standardized to one standard deviation. This helps the SD algorithm to move correctly in all dimensions of Θ .

The VIX data are strongly right-skewed, with sample skewness about 2.0. The right tail is substantially extended by several *panic selling* episodes in which the VIX rises sharply within a single day. As a result, the sample kurtosis is approximately 17.

The top two graphs compare the histogram with the fitted theoretical density. The graph on the right shows the density on a logarithmic scale, which makes it easier to assess the tail fit down to the 10^{-3} level. In particular, the theoretical density does not fully capture the far right tail beyond about 7.

The fitted theoretical parameters are $\alpha \approx 0.77$, $k \approx 5.2$, and $\beta \approx 0.95$. This fitted point can be located in Figure 12.1.

The PP-plot in the bottom-left graph indicates that the overall fit is acceptable. The points remain close to the 45-degree line, although this plot is relatively insensitive to tail behavior.

The bottom-right graph shows the QQ-plot of the quadratic form, or squared variable. This diagnostic is particularly useful for assessing how well the distribution captures the combined tail behavior in absolute terms. The 45-degree line is OK below 20, but as the quantiles get larger, the observed quantiles start to tilt upward. This means the top 0.5 percent of the combined tail is not properly captured by the distribution.

12.10.2. SPX fit. Figure 12.8 shows the result of the MLE fit to the daily SPX returns from 1990 to mid-2025. The data are also standardized to one standard deviation.

The SPX data are slightly left-skewed, with the sample skewness slightly below -0.1 . The tails are extended by several large one-day market declines, producing sample kurtosis of approximately 11 while leaving the skewness relatively modest.

The top two graphs compare the histogram with the fitted theoretical density. The graph on the right shows the density on a logarithmic scale, making it easier to assess the tail fit down to the 10^{-3} level. The theoretical density does not fully capture the tails beyond about 7.

The fitted theoretical distribution has parameters $\alpha = 0.15$, $k = 19.9$, and $\beta = -0.11$. This MLLK fit is in the low- α , high- k region, marked as "all data" in Figure 12.4.

The PP-plot in the bottom-left graph indicates that the overall fit is satisfactory, with most points lying on the 45-degree line.

In the QQ-plot of the quadratic form, the points remain close to the 45-degree line up to about 20. However, for larger quantiles, the observed quantiles begin to bend upward. In particular, the ten largest observations deviate materially from the theoretical curve.




FIGURE 12.7. MLE fit of daily VIX returns from 1990 to mid-2025. The data are standardized to unit variance. Sample skewness is 2.0, and sample kurtosis is 16.5. $\hat{\Theta} = \{\alpha = 0.77, k = 5.2, \beta = 0.95\}$. Top left: density and histogram. Top right: log-density and histogram. Bottom left: PP-plot. Bottom right: QQ-plot of the quadratic form.




FIGURE 12.8. MLE fit of daily SPX log returns from 1990 to mid-2025. The data are standardized to unit variance. Sample skewness is -0.14 , and sample kurtosis is 10.6 . $\hat{\Theta} = \{\alpha = 0.15, k = 19.9, \beta = -0.11\}$. Top left: density and histogram. Top right: log-density and histogram. Bottom left: PP-plot. Bottom right: QQ-plot of the quadratic form.

Fractional Feller Square-Root Process

This chapter is copied from Section 11 of [19] for the generation of random variables for FG, FCM, and FCM2. Combining this with an SN variable provides a path to generate the random variable for GAS-SN and beyond.

For example, assuming that a sequence of random numbers $\{S_t > 0\}$ can be generated for FCM, it is straightforward to simulate random numbers $\{X_t\}$ for GAS-SN using the ratio of $X_t = Y_t/S_t$, where Y_t is a standard skew-normal variable $Y_t \sim SN(0, 1, \beta)$ in Chapter 12.

Instead of randomly generating $\{S_t\}$, we propose an innovative method based on *Feller square-root process*[10]. Given a user-specific volatility $\sigma_u > 0$ that describes how fast S_t should change, a scalar function $\mu(x)$, and a scale parameter $\theta_u > 0$ (default to 1), we assume that the random variable S_t should evolve according to the following generalized process:

$$(13.1) \quad dS_t = \sigma_u^2 \mu\left(\frac{S_t}{\theta_u}\right) dt + \sigma_u \sqrt{S_t} dW_t$$

As $t \rightarrow \infty$, $\{S_t\}$ will be distributed as the equilibrium distribution for which $\mu(x)$ is designated.

13.0.1. The Fokker-Planck Equation. The $\mu(x)$ solution can be derived from the Fokker-Planck equation. We obtain the following beautiful relation:

LEMMA 13.1. $\mu(x)$ is one half of the elasticity of the terminal density function $p(x)$ of S_t at $t \rightarrow \infty$ plus one half:

$$(13.2) \quad \mu(x) = \frac{1}{2} \mathcal{L} p(x) + \frac{1}{2}$$

where $\mathcal{L}(\cdot) := x \frac{d}{dx} \log(\cdot)$ is the elasticity operator defined in Section 3.7.

△

PROOF. Assume $p(x, t)$ is the density function of (13.1) for S_t . It should satisfy the Fokker-Planck equation ($\theta_u = 1$):

$$\frac{\partial}{\partial t} p(x, t) = -\frac{\partial}{\partial x} [\sigma_u^2 \mu(x) p(x, t)] + \frac{\partial^2}{\partial x^2} \left[\frac{1}{2} (\sigma_u \sqrt{x})^2 p(x, t) \right]$$

As $t \rightarrow \infty$, $p(x, t)$ approaches the terminal density function $p(x)$. The time dependency is removed. σ_u^2 cancels out from both sides and is irrelevant to the solution. The ODE of $p(x)$ becomes

$$\frac{\partial}{\partial x} (\mu(x) p(x)) = \frac{1}{2} \frac{\partial^2}{\partial x^2} (x p(x))$$

Apply $\int_x^\infty dx$ to both sides. Assuming that $\mu(x)p(x)$ at $x = \infty$ should be zero, we get

$$\mu(x)p(x) = \frac{1}{2} \frac{d}{dx} (x p(x)) = \frac{1}{2} \left(x \frac{d}{dx} p(x) + p(x) \right)$$

Moving $p(x)$ from LHS to RHS, we obtain (13.2).

□

13.0.2. Generation of Random Variables for FG.

LEMMA 13.2. The $\mu(x)$ solution for FG is obviously

$$\mu(x) = \frac{1}{2} \mathcal{L} \mathfrak{N}_\alpha(x; \sigma, d, p) + \frac{1}{2}$$

With Lemma 3.8, $\mu(x)$ is reduced to a function of $\mathcal{L} M_\alpha(x)$:

$$(13.3) \quad \mu(x) = \frac{p}{2} [\mathcal{L} M_\alpha] \left(\left(\frac{x}{\sigma} \right)^p \right) + \frac{d+p}{2}.$$

△

As an application, since $\mathcal{L} M_{1/2}(x) = -x^2/2$ from Section 3.7, we obtain a simple power-law solution at $\alpha = 1/2$:

$$(13.4) \quad \mu(x)|_{\mathfrak{N}_{1/2}} = -\frac{p}{4} \left(\frac{x}{\sigma} \right)^{2p} + \frac{d+p}{2}$$

Note that (13.4) at $p = 1/2$ subsumes the renown Cox–Ingersoll–Ross (CIR) model[5] since the $\mu(x)$ of the model is a linear $a(b-x)$ type, according to its stochastic process of $dS_t = a(b-S_t)dt + \sigma_u \sqrt{S_t} dW_t$.¹

To prepare for the solution of FCM, we prefer to use $Q_\alpha(x)$ defined in (3.36):

$$Q_\alpha(x) := -\frac{W_{-\alpha,-1}(-x)}{W_{-\alpha,0}(-x)}$$

LEMMA 13.3. From (3.38), the $\mu(x)$ solution of a FG in terms of $Q_\alpha(x)$ is

$$(13.5) \quad \mu(x) = \frac{p}{2\alpha} Q_\alpha \left(\left(\frac{x}{\sigma} \right)^p \right) + \left(\frac{d}{2} - \frac{p}{2\alpha} \right)$$

Notice that p/α and d are just constant terms, and σ only affects the scale of x . Neither of them has any effect on the shape of $\mu(x)$.

△

13.1. Generation of Random Variables for FCM

Obviously, what really matters for GAS-SN and GSaS is the solution of FCM, The $\mu(x)$ solution for $\bar{\chi}_{\alpha,k}$ is denoted as $\mu_{\alpha,k}(x)$. Note that from this point on, $\alpha \in (0, 2)$.

To further simplify the symbology for FCM, define

$$Q_\alpha^{(\chi)}(z) := Q_{\frac{\alpha}{2}}(z^\alpha), \text{ where } \alpha \in (0, 2).$$

Assuming $k > 0$, we set $\sigma = \sigma_{\alpha,k}$, $d = k-1$, $p/\alpha = 2$ and α replaced by $\alpha/2$ in (13.5). We get

$$(13.6) \quad \mu_{\alpha,k}(x) = Q_\alpha^{(\chi)} \left(\frac{x}{\sigma_{\alpha,k}} \right) + \left(\frac{k-3}{2} \right)$$

For validation, $\mu_{1,k}(x) = k(1-x^2)/2$ can be used to simulate Student's t. And $\mu_{\alpha,1}(x)$ provides a method to simulate an SaS $L_{\alpha,1}(x)$:

$$\mu_{\alpha,1}(x) = Q_\alpha^{(\chi)}(\sqrt{2}x) - 1$$

Fig. 13.1 shows a simulation of random variables based on the (α, k) parameter obtained from the fit of the S&P 500 daily log returns. The rest of the parameters are in the caption of the figure. First, as outlined above, $\mu_{\alpha,k}(s)$ is calculated analytically as shown in the right graph. Second, it enables

¹It can also be subsumed by the FG at $\alpha = 0, p = 1$. But $\alpha = 0$ is a singular point and we prefer to avoid using it when possible.

the FG simulation $\{S_t\}$ as shown in the left graph. Third, GSaS $\{X_t\}$ is simulated via $X_t = \mathcal{N}/S_t$, where $\mathcal{N}$ is drawn from a standard normal variable.

The simulation is performed daily. The duration of the sampling is 200,000 years. The red areas are histograms of the simulated data. The blue lines are from the theoretical density functions. They match well.




FIGURE 13.1. Simulation of random variables based on the (α, k) parameters obtained from the fit of the S&P 500 daily log returns. The red areas are the histograms from simulated data. The blue lines are from theoretical formulas. The settings of the simulation are $\alpha = 0.813, k = 3.292, dt = 1/365, \sigma_u = 0.85$. Sampling duration is 200,000 years. The simulation takes 11 minutes in python. $\mu_{\alpha,k}(s)$ is discretized to 0.01 and cached to increase performance.

13.2. Generation of Random Variables for FCM2

LEMMA 13.4. The $\mu(x)$ solution for $\bar{\chi}_{\alpha,k}^2$ is

$$(13.7) \quad \mu_{\alpha,k}^{(2)}(x) = \frac{1}{2} \mu_{\alpha,k}(\sqrt{x})$$

△

PROOF. From (7.15), we have

$$\bar{\chi}_{\alpha,k}^2(x) := \mathfrak{N}_{\alpha/2}(x; \sigma = \sigma_{\alpha,k}^2, d = (k-1)/2, p = \alpha/2) \quad (k > 0)$$

Combined with (13.5), we obtain the solution for $\bar{\chi}_{\alpha,k}^2$ as

$$\begin{aligned} \mu_{\alpha,k}^{(2)}(x) &= \frac{1}{2} Q_{\alpha/2} \left(\left(\frac{\sqrt{x}}{\sigma_{\alpha,k}} \right)^\alpha \right) + \left(\frac{k-1}{4} - \frac{1}{2} \right) \\ &= \frac{1}{2} Q_{\alpha}^{(\chi)} \left(\frac{\sqrt{x}}{\sigma_{\alpha,k}} \right) + \frac{k-3}{4}, \end{aligned}$$

which is just

$$\mu_{\alpha,k}^{(2)}(x) = \frac{1}{2} \mu_{\alpha,k}(\sqrt{x}).$$

□

1829 This solution can be used to simulate the F distribution in Chapter 8. Let $U_1 \sim \chi_d^2/d = \bar{\chi}_{1,d}^2$ and
 1830 $U_2 \sim \bar{\chi}_{\alpha,k}^2$, then $F_{\alpha,d,k} \sim U_1/U_2$ is a fractional F distribution.

1831

Part 4

1832

Multivariate Distributions

Multivariate SN Distribution - Review

In this chapter, we start to explore the multivariate distributions. Data sets from the real world are often multidimensional. A flexible multivariate distribution framework with skewness and kurtosis can be very useful. That is what we aim to achieve in the next few chapters.

The foundation is the standard multivariate normal distribution $\mathcal{N}_d(0, \bar{\Omega})$, where d is the dimension of the random variable, and $\bar{\Omega}$ is a $d \times d$ correlation matrix[38].

In Chapter 5 of Azzalini, the skew normal distribution $SN_d(0, \bar{\Omega}, \beta)$ adds skewness to it from the skew parameter β [1]. In its Chapter 6, the skew-elliptical distribution is discussed. The multivariate skew-t distribution $ST_d(0, \bar{\Omega}, \beta, k)$ is constructed by combining $SN_d(0, \bar{\Omega}, \beta)$ with $\chi_k/\sqrt{k}$ in a ratio distribution.

Our work builds on top of this concept of the skew-elliptical distribution. By expanding the denominator of $\chi_k/\sqrt{k}$ to the FCM $\bar{\chi}_{\alpha, k}$, the fractional dimension α is added to the shape parameters. This forms a super-distribution family called *multivariate GAS-SN elliptical distribution* with the notation $L_{\alpha, k}(0, \bar{\Omega}, \beta)$ for its standard distribution.

The multivariate skew-elliptical distribution has beautiful properties inherited from the multivariate elliptical distribution framework. However, its deficiency is obvious in real-world applications: The structure is multivariate, but the shape parameters α and k are scalars. All dimensions share the same (α, k) . This restricts the kurtoses of 1D marginal distributions to a similar range. It even creates some strange phenomena that are hard to interpret in the SPX-VIX 2D fit (see Section 17.1.1).

To overcome such a restriction, we propose a more flexible framework called *multivariate adaptive distribution*, in which the shape parameters (α, k) are d dimensional vectors, just like their skew counterpart β .

The flexibility in shapes comes with an expensive computational cost. It is analogous to the *curse of dimensionality* problem. It becomes much harder to verify the results beyond the bivariate case for the adaptive distribution.

The study of quadratic form $Z^\top \bar{\Omega}^{-1} Z$ from the skew-elliptical distribution results in the fractional extension of the F distribution $F_{\alpha, d, k}$. The QQ-plot based on the quadratic form and the fractional F distribution is a powerful validation of the goodness of the fit.

14.1. Definition

We summarize the results of Chapter 5 of Azzalini[1]. On the one hand, we need to clarify the symbology here that is slightly different from that in his book. On the other hand, our multivariate distributions rely on many results from there, which are collected in this chapter.

DEFINITION 14.1. The PDF of a standard multivariate normal distribution $\mathcal{N}_d(0, \bar{\Omega})$ is defined as

$$(14.1) \quad \mathcal{N}_d(\mathbf{x}; \bar{\Omega}) := \frac{1}{(2\pi)^{d/2} \det(\bar{\Omega})^{1/2}} \exp\left(-\frac{1}{2} \mathbf{x}^\top \bar{\Omega}^{-1} \mathbf{x}\right), \quad \mathbf{x} \in \mathbb{R}^d.$$

where $\bar{\Omega}$ is a $d \times d$ correlation matrix[38]. That is, $\bar{\Omega}$ is positive definite and all its diagonal elements are equal to 1.

DEFINITION 14.2. A standard multivariate skew-normal variable is denoted as $Z \sim SN_d(0, \bar{\Omega}, \beta)$, where $\beta \in \mathbb{R}^d$ is the skew parameter (or the slant parameter). Its PDF is

$$(14.2) \quad \mathcal{N}_d(\mathbf{x}; \bar{\Omega}, \beta) := \mathcal{N}_d(\mathbf{x}; \bar{\Omega}) \Phi_{\mathcal{N}}(\beta^\top \mathbf{x}),$$

where $\Phi_{\mathcal{N}}(x)$ is the CDF of a standard normal distribution $\mathcal{N}(0, 1)$.

Notice that this is a multivariate expansion of SN in Section 10.1. When $d = 1$, (14.2) becomes (10.2).

14.2. The Location-Scale Family

Its location-scale family is $Y = \xi + \omega Z \sim SN_d(\xi, \Omega, \beta)$, where $\xi \in \mathbb{R}^d$ is the location parameter, $\omega = \text{diag}(\omega_1, \dots, \omega_d)$ is a $d \times d$ diagonal scale matrix ($\omega_i > 0, \forall i$) and $\Omega = \omega \bar{\Omega} \omega$.

The PDF of Y becomes

$$(14.3) \quad f_Y(\mathbf{x}) = \det(\omega)^{-1} \mathcal{N}_d(\mathbf{z}; \bar{\Omega}, \beta),$$

where $\mathbf{z} = \omega^{-1}(\mathbf{x} - \xi)$.

The location-scale distribution is used for real-world applications. Internally, it has to be calculated via the standard distribution. The main reason is that β has to work with $\mathbf{z}$ and $\bar{\Omega}$, instead of $\mathbf{x}$ and ω .

14.3. Quadratic Form

DEFINITION 14.3. The quadratic form of a multivariate SN distribution (MSN) is defined as

$$(14.4) \quad Q := \frac{1}{d}(Y - \xi)^\top \Omega^{-1}(Y - \xi) = \frac{1}{d}Z^\top \bar{\Omega}^{-1}Z.$$

Q distributes as $\chi_d^2/d = \bar{\chi}_{1,d}^2$ for all β . The distribution of Q is independent of β . This is an important property due to the rotational invariance of the elliptical distribution.

Notice that our definition of Q is slightly different from that of Azzalini. We prefer to have the distribution of Q tied to the FCM and the fractional F distribution directly without any constant adjustment. This will make things much simpler in Section 15.6.

To prove $Q \sim \chi_d^2/d$, we quote Corollary 5.9 from [1] below for a skew-normal distribution with 0 location:

LEMMA 14.4. If $Y \sim SN_d(0, \Omega, \beta)$ and A is a $d \times d$ symmetric matrix, then

$$Y^\top A Y = X^\top A X$$

where $X \sim \mathcal{N}_d(0, \Omega)$. △

This lemma allows β to be removed from the statistics of Q . Hence, $Q \sim X^\top \Omega^{-1} X / d \sim \chi_d^2 / d$.

14.4. Stochastic Representation

Assuming $X_0 \sim \mathcal{N}_d(0, \bar{\Omega})$ and $X_1 \sim \mathcal{N}(0, 1)$, then the first representation of $Z \sim SN_d(0, \bar{\Omega}, \beta)$ is

$$(14.5) \quad Z = \begin{cases} X_0 & \text{if } X_1 > \beta^\top X_0, \\ -X_0 & \text{otherwise.} \end{cases}$$

This form of selective sampling is quite useful in generating random numbers for Z . It is essentially an extension of (10.1).

This scheme can be rephrased in a more interesting representation. First, define the multivariate version of δ as

$$(14.6) \quad \boldsymbol{\delta} = (1 + \boldsymbol{\beta}^\top \bar{\Omega} \boldsymbol{\beta})^{-1/2} \bar{\Omega} \boldsymbol{\beta}, \quad (\boldsymbol{\delta} \in \mathbb{R}^d)$$

which is used to construct a $(d+1) \times (d+1)$ correlation matrix

$$\Omega^* = \begin{pmatrix} \bar{\Omega} & \boldsymbol{\delta} \\ \boldsymbol{\delta}^\top & 1 \end{pmatrix}.$$

Ω^* is used to generate two marginals, $X_0 \in \mathbb{R}^d$ and $X_1 \in \mathbb{R}$, such that

$$\begin{pmatrix} X_0 \\ X_1 \end{pmatrix} \sim \mathcal{N}_{d+1}(0, \Omega^*),$$

which leads to the second representation

$$(14.7) \quad Z = \begin{cases} X_0 & \text{if } X_1 > 0, \\ -X_0 & \text{otherwise.} \end{cases}$$

This form resembles (10.7). It shows that the function of $\boldsymbol{\delta}$ is to add the correlation between X_0 and X_1 through Ω^* in the selective sampling. This makes (14.7) slightly different from (14.5).

14.5. Moments

The first two moments of Z have simple analytic forms. Its first moment is

$$(14.8) \quad \mu_z = \mathbb{E}(Z) = b \boldsymbol{\delta}, \quad \text{where } b = \sqrt{2/\pi}.$$

The second moment is simply $\bar{\Omega}$. Its variance is

$$(14.9) \quad \Sigma_z = \text{var}\{Z\} = \bar{\Omega} - b^2 \boldsymbol{\delta} \boldsymbol{\delta}^\top.$$

It is easy to obtain $\mathbb{E}\{Y Y^\top\} = \Omega$ for the location-scale variable Y .

Define the important invariant quantity for the skewness.

$$(14.10) \quad \beta_* = (\boldsymbol{\beta}^\top \bar{\Omega} \boldsymbol{\beta})^{1/2} \geq 0,$$

which is a nonnegative scalar quantity. It encapsulates the departure from normality for the distribution.

The quadratic form $\mu_z^\top \Sigma_z^{-1} \mu_z$ can be simplified to

$$(14.11) \quad \mu_z^\top \Sigma_z^{-1} \mu_z = \frac{b^2 \beta_*^2}{1 + (1 - b^2) \beta_*^2}.$$

A related quantity is

$$(14.12) \quad \delta_* = (\boldsymbol{\delta}^\top \bar{\Omega}^{-1} \boldsymbol{\delta})^{1/2}$$

where $\delta_* \in [0, 1)$ has the scale of a positive correlation coefficient.

The two are connected by

$$\delta_*^2 = \frac{\beta_*^2}{1 + \beta_*^2}, \quad \beta_*^2 = \frac{\delta_*^2}{1 - \delta_*^2}.$$

Or in a trigonometric form, there exists an angle $\theta \in [0, \frac{\pi}{2})$ such that $\tan \theta = \beta_*$ and $\sin \theta = \delta_*$. In such an expression, $\theta > 0$ captures the "degree" of departure from normality.

14.6. Canonical Form

The concept of a canonical form in SN is very important and fascinating. Due to the rotational symmetry, an MSN can be rotated and rescaled to an "identity" MSN with a scalar skew parameter.

By Proposition 5.12 of [1], there exists an affine transformation $Z^* = A_*(Y - \xi)$ such that $Z^* \sim SN_d(0, \mathbf{I}_d, \beta_{Z^*})$, where $\mathbf{I}_d$ is a $d \times d$ identity matrix, and $\beta_{Z^*} = (\beta_*, 0, \dots, 0)^\top$. β_* is defined by (14.10), which is an invariant under transformation.

The variable Z^* is called *the canonical variable*. It is d -dimensional. But only one dimension is skew-normal, which is designated as the first dimension. All other dimensions are standard normal distributions. That is, the PDF of Z^* is

$$\begin{aligned} \mathcal{N}_*(\mathbf{x}; \beta_*) &= 2\Phi_{\mathcal{N}}(\beta_* x_1) \prod_{i=1}^d \mathcal{N}(x_i) \\ &= \mathcal{N}(x_1; \beta_*) \prod_{i=2}^d \mathcal{N}(x_i). \end{aligned}$$

This structure helps tremendously for the subsequent development of the elliptical distribution and adaptive distribution.

Proposition 5.13 in [1] describes how to find such A_* . Due to rotational symmetry, there are many choices of A_* . This is not a problem as long as we always look at the system in quadratic form.

LEMMA 14.5 (Affine Transformation). Let $C = \Omega^{1/2}$ be the unique positive definite symmetric square root of Ω . Define $M = C^{-1}\Sigma C^{-1}$, where $\Sigma = \text{var}\{Y\}$. Let $Q\Lambda Q^\top$ denote a spectral decomposition of M , where we assume that the diagonal elements in the eigenvalue matrix Λ are arranged in increasing order.

Let $H = C^{-1}Q$. Then H is the matrix operator to convert Y to Z^* ,

$$Z^* = H^\top(Y - \xi).$$

Since $\delta_{Z^*} = H^\top \omega \delta$ and $\beta_{Z^*} = \delta_{Z^*} / (1 - \delta_*^2)$, the choice of H must make the first element of δ_{Z^*} a nonnegative number, that is, $\delta_* \geq 0$. All other elements, except the first ones in δ_{Z^*} and β_{Z^*} , must be zero. $\triangle$

REMARK 14.6. The significance of this lemma is that the skew-elliptical distributions derived from the SN framework can only have a single source of skewness. It might be mixed up and not easy to observe in real-world data. But there is only one source from the theoretical perspective. Everything else comes from the multivariate normal distribution.

If we want a more "sophisticated" distribution that provides multiple sources of skewness, we have to go beyond the skew-elliptical distributions.

14.7. 1D Marginal Distribution

We are particularly interested in the 1D marginal distribution, since this is what is actually observed in a data set. When we optimize a data fit, we can add the log-likelihood of the 1D marginal distributions to the objective function, so that the fitting of each dimension is properly addressed.

In fact, for the adaptive distribution, the full 2D likelihood is so compute-intensive that it is too slow to perform MLE on a desktop. The alternative is to compute the sum of the log-likelihoods of each 1D marginal distribution, in addition to the regularization on other statistical quantities, such as the correlation coefficient between each data pair.

We quote the results from Section 5.1.4 of [1] and adapt them to the 1D case.

LEMMA 14.7. (The marginal β) Assume that the marginal is on the first dimension. The correlation matrix is decomposed as

$$\bar{\Omega} = \begin{pmatrix} \bar{\Omega}_{11} & \bar{\Omega}_{12} \\ \bar{\Omega}_{21} & \bar{\Omega}_{22} \end{pmatrix}.$$

1954 The formula can be simplified due to $\bar{\Omega}_{11} = 1$ in the 1D case.

1955 The marginal distribution is $Y_1 \sim SN(\xi_1, \Omega_{11}, \beta_{1(2)})$. Its $\beta_{1(2)}$ is derived as

$$(14.13) \quad \beta_{1(2)} = (1 + \beta_2^T \bar{\Omega}_{22.1} \beta_2)^{-1/2} (\beta_1 + \bar{\Omega}_{12} \beta_2)$$

where $\bar{\Omega}_{22.1} = \bar{\Omega}_{22} - \bar{\Omega}_{21} \bar{\Omega}_{12}$.

1956

△

1957 LEMMA 14.8. (The marginals of a bivariate distribution) The bivariate case is quite simple:

$$(14.14) \quad \bar{\Omega} = \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix}.$$

1958 Assume that we want to get the marginal β of the i -th dimension, $\beta_{i(j)}$, where j is the other
1959 dimension. Then

$$(14.15) \quad \beta_{i(j)} = \frac{\beta_i + \rho \beta_j}{\sqrt{1 + \beta_j^2 |\bar{\Omega}|}}$$

1960 where $|\bar{\Omega}| = 1 - \rho^2$. Since Ω_{ii} is ω_i^2 , the i -th marginal distribution is $Y_i \sim SN(\xi_i, \omega_i^2, \beta_{i(j)})$. The ξ_i and
1961 ω_i are the location and scale parameters in the i -th dimension that can be calculated directly from the
1962 data. △

1963 We observe that ρ in the numerator describes how much β_j is mixed with β_i , while $|\bar{\Omega}|$ in the
1964 denominator describes how much β_j reduces the scale.

1965 When $\rho = 0$, there is no mixing from the other dimension, only a reduction in total scale. That
1966 is, $\beta_{i(j)}|_{\rho=0} = \beta_i / \sqrt{1 + \beta_j^2}$.

Multivariate GAS-SN Elliptical Distribution

15.1. Definition

This chapter follows the structure laid out in Chapter 6 of Azzalini (2013)[1]. We implemented the skew-elliptical distribution by our $\bar{\chi}_{\alpha,k}$, which fully extends his multivariate skew-t distribution.

DEFINITION 15.1. Assume $Z_0 \sim SN_d(0, \bar{\Omega}, \beta)$ is a $d \times d$ standard multivariate skew-normal (SN) distribution, and $V \sim \bar{\chi}_{\alpha,k}$ is a standard FCM. $\bar{\Omega}$ is a correlation matrix.

Then $Z \sim Z_0/V$ is a $d \times d$ standard multivariate GAS-SN elliptical distribution. It is given the notation of $Z \sim L_{\alpha,k}(0, \bar{\Omega}, \beta)$.

Equivalently, using the location-scale notation, $Z \sim SN_d(0, \Sigma, \beta)$ where $\Sigma = \bar{\Omega}/V^2$.

Assume $\mathcal{N}_d(\mathbf{x}; \bar{\Omega}, \beta)$ is the PDF of a standard multivariate normal distribution $\mathcal{N}_d(0, \bar{\Omega})$ [38]. $\Phi_{\mathcal{N}}(x)$ is the CDF of a standard normal distribution.

We expand on the construction of multivariate SN distribution in (14.1) and (14.2). And the PDF of $Z \sim L_{\alpha,k}(0, \bar{\Omega}, \beta)$ is

$$\begin{aligned} L_{\alpha,k}(\mathbf{x}; \bar{\Omega}, \beta) &= \int_0^\infty ds \bar{\chi}_{\alpha,k}(s) s^d \mathcal{N}_d(\mathbf{x}s; \bar{\Omega}, \beta) \\ (15.1) \qquad \qquad \qquad &= 2 \int_0^\infty ds \bar{\chi}_{\alpha,k}(s) s^d \mathcal{N}_d(\mathbf{x}s; \bar{\Omega}) \Phi_{\mathcal{N}}(\beta^\top \mathbf{x} s). \end{aligned}$$

The s^d term comes from $\det(s\mathbf{I}_d)$ where $\mathbf{I}_d$ is the $d \times d$ identity matrix. It is easy to see how it is reduced to a univariate GAS-SN distribution when $d = 1$.

15.1.1. Multivariate Skew-t Distribution. An important bridge between multivariate SN and GAS-SN is the multivariate skew-t distribution. It is documented in Section 6.2 of [1].

It is fully consistent with multivariate GAS-SN by setting $\alpha = 1$. That is, in his notation of skew-t: $ST_d(\Omega, \beta, k) \sim L_{1,k}(\Omega, \beta)$.

15.2. Location-Scale Family

The location-scale family follows the standard procedure: $Y = \xi + \omega Z$, which is denoted as $Y \sim L_{\alpha,k}(\xi, \Omega, \beta)$, where $\Omega := \omega^\top \bar{\Omega} \omega$ is the covariance matrix, and ω is a $d \times d$ diagonal scale matrix.

The PDF of Y is

$$(15.2) \qquad \qquad \qquad L_{\alpha,k}(\mathbf{x}; \xi, \Omega, \beta) := \det(\omega)^{-1} L_{\alpha,k}(\mathbf{z}; \bar{\Omega}, \beta)$$

where $\mathbf{z} := \omega^{-1}(\mathbf{x} - \xi)$. Notice that it has to be computed via the standard PDF.

15.3. Moments

The first moment of Z is $\mu_z := b \delta$, where $b := \sqrt{2/\pi} \mathbb{E}(X^{-1} | \bar{\chi}_{\alpha,k})$.

The second moment of Z is $m_2 \bar{\Omega}$, where $m_2 = \mathbb{E}(X^{-2} | \bar{\chi}_{\alpha,k})$. Hence, the covariance is

$$\text{var}\{Z\} := \Sigma_z := m_2 \bar{\Omega} - b^2 \delta \delta^\top$$

The moments of Y follow the rule of the location-scale family. The first moment of Y is $\boldsymbol{\xi} + \boldsymbol{\omega} \mu_z$. The covariance of Y is $\boldsymbol{\omega} \Sigma_z \boldsymbol{\omega}$.

15.4. Canonical Form

The concept of canonical form in GAS-SN is extended from the multivariate SN in Section 14.6. There exists an affine transformation $Z^* = A_*(Y - \boldsymbol{\xi})$ such that $Z^* \sim L_{\alpha,k}(0, \mathbf{I}_d, \boldsymbol{\beta}_{Z^*})$, where $\boldsymbol{\beta}_{Z^*} = (\beta_*, 0, \dots, 0)^\top$ and β_* is defined by (14.10). And the algorithm of finding A_* is exactly the same as in Section 14.6.

The variable Z^* , which is called *canonical variable*, comprises d independent components. Only one of them contains the skew component. That is, the PDF of Z^* is

$$(15.3) \quad L_{\alpha,k_*}(\mathbf{x}; \beta_*) = 2 \int_0^\infty ds \bar{\chi}_{\alpha,k}(s) s^d \prod_{i=1}^d \mathcal{N}(x_i s) \Phi_{\mathcal{N}}(\beta_* x_1 s).$$

It can be further simplified to an elegant univariate-style integral. When $|\mathbf{x}| \neq 0$, let $\beta_*(\mathbf{x}) := \beta_* x_1 / |\mathbf{x}| \in \mathbb{R}$, and

$$(15.4) \quad L_{\alpha,k_*}(\mathbf{x}; \beta_*) = (2\pi)^{-(d-1)/2} \int_0^\infty ds \bar{\chi}_{\alpha,k}(s) s^d \mathcal{N}(|\mathbf{x}|s; \beta_*(\mathbf{x})).$$

When $|\mathbf{x}| = 0$, It is simply

$$(15.5) \quad L_{\alpha,k_*}(0; \beta_*) = (2\pi)^{-d/2} \mathbb{E}(X^d | \bar{\chi}_{\alpha,k}),$$

independent of β_* .

15.5. Marginal 1D Distribution

The construction of the 1D marginal distribution from Y extends directly from Section 14.7, where $\beta_{1(2)}$ is calculated.

Then the marginal distribution is an univariate GAS-SN: $Y_1 \sim L_{\alpha,k}(\xi_1, \Omega_{11}, \beta_{1(2)})$.

15.6. Quadratic Form

The quadratic form is

$$(15.6) \quad Q := \frac{1}{d} (Y - \boldsymbol{\xi})^\top \Omega^{-1} (Y - \boldsymbol{\xi}) = \frac{1}{d} Z^\top \bar{\Omega}^{-1} Z.$$

This leads to the fractional extension of the classic F distribution.

Q distributes like a fractional F distribution, $Q \sim F_{\alpha,d,k}$ for all β . The QQ-plot between the empirical data and theoretical values is used to evaluate the goodness of a fit. A perfect fit should produce a 45-degree line.

To prove, from Section 15.1, we have $Z \sim Z_0/V$, $Z_0 \sim SN_d(0, \bar{\Omega}, \boldsymbol{\beta})$, and $V \sim \bar{\chi}_{\alpha,k}$. Put them together,

$$Q = \frac{1}{d} Z^\top \bar{\Omega}^{-1} Z = \frac{Z_0^\top \bar{\Omega}^{-1} Z_0}{d V^2} \sim \left(\frac{X^2}{d} \right) / V^2$$

where $X \sim \mathcal{N}_d(0, \bar{\Omega})$, according to Lemma 14.4.

Since $X^2 \sim \chi_d^2$ and $V^2 \sim \bar{\chi}_{\alpha,k}^2$, this leads to $Q \sim F_{\alpha,d,k}$, according to Section 8.1.

Azzalini (2013) provided a point of validation from his multivariate skew-t distribution. From Section 6.2 of [1], Q of a skew-t variable distributes like the classic $F(d, k)$. This is a special case of our fractional F distribution at $\alpha = 1$. That is, $Q \sim F_{1,d,k}$.

15.7. Multivariate MLE

Maximum likelihood estimation for the bivariate elliptical model can be implemented with stochastic gradient descent (SGD). The objective is the minus-log likelihood (MLLK), computed by summing the negative log density over the observed data. At each iteration, the gradients with respect to the hyperparameters are used to update the parameter vector, with the step size controlled by a learning rate. Small random perturbations may also be added to improve local exploration of the objective surface.

For the bivariate optimization, the parameter space is

$$\Theta = \{\rho, \alpha, k, \beta_1, \beta_2, w_1, w_2, \xi_1, \xi_2\}$$

where $\alpha \in (0, 2)$, $k \in (2, \infty)$, $w_1 > 0$, $w_2 > 0$, and $\beta_1, \beta_2, \xi_1, \xi_2 \in \mathbb{R}$.

Since $\rho \in (-1, 1)$, the optimization uses an unconstrained parameter $\rho_\theta \in \mathbb{R}$ and maps it to the correlation coefficient by $\rho = (2/\pi) \arctan(\rho_\theta)$.

In practice, each dimension of the data set is normalized to unit standard deviation. This keeps the gradient components on comparable scales and improves the stability of the SGD updates.

Let Y denote a data set of size N , and let $L(Y_i; \Theta)$ be the PDF evaluated at observation Y_i . Then

$$\begin{aligned} \text{MLLK}(\Theta; Y) &:= - \sum_{i=1}^N \log L(Y_i; \Theta) \\ \text{Gradient}(\Theta) &:= \left\{ \frac{\partial \text{MLLK}(\Theta; Y)}{\partial \Theta_j}, \forall \Theta_j \in \Theta \right\} \end{aligned}$$

When N is large, evaluating $L(Y_i; \Theta)$ at every observation may be computationally expensive. A histogram approximation can be used to compress the data into a smaller number of bins.

Additional regularization can be added to the MLLK. In the bivariate setting, useful penalties include the L2 distances between the empirical and model-implied skewness and kurtosis of the marginal one-dimensional distributions.

Multivariate GAS-SN Adaptive Distribution (Experimental)

16.1. Definition

The adaptive distribution allows each dimension to have its own shape parameters (α_i, k_i) . This is the main departure from the elliptical construction, where all dimensions share the same scalar pair (α, k) .

Accordingly, $\boldsymbol{\alpha} = \{\alpha_i\}$ and $\mathbf{k} = \{k_i\}$ are d -dimensional parameter vectors. The mixture therefore uses a corresponding collection of standard FCM variables, $\{\bar{\chi}_{\alpha_i, k_i} : i = 1, \dots, d\}$.

DEFINITION 16.1. Assume that Z_0 is a d -dimensional random variable from the standard multivariate skew-normal distribution $SN_d(0, \bar{\Omega}, \boldsymbol{\beta})$, where $\bar{\Omega}$ is a correlation matrix.

Let Z be defined component-wise by the ratio representation $Z_i \sim (Z_0)_i / \bar{\chi}_{\alpha_i, k_i}$. Then $Z \sim \vec{L}_{\boldsymbol{\alpha}, \mathbf{k}}(0, \bar{\Omega}, \boldsymbol{\beta})$ is a standard multivariate GAS-SN adaptive distribution. The arrow notation emphasizes that the shape parameters $(\boldsymbol{\alpha}, \mathbf{k})$ are vectors.

Let $\mathcal{N}(x; \bar{\Omega})$ denote the PDF of the standard multivariate normal distribution $N(0, \bar{\Omega})$ [38], and let $\Phi_{\mathcal{N}}(x)$ denote the CDF of the standard normal distribution.

For $Z \sim \vec{L}_{\boldsymbol{\alpha}, \mathbf{k}}(0, \bar{\Omega}, \boldsymbol{\beta})$, the PDF is

$$(16.1) \quad \vec{L}_{\boldsymbol{\alpha}, \mathbf{k}}(\mathbf{x}; \bar{\Omega}, \boldsymbol{\beta}) = 2 \int \cdots \int_0^\infty \mathcal{N}(\mathbf{s} \mathbf{x}; \bar{\Omega}) \Phi_{\mathcal{N}}(\boldsymbol{\beta}^\top(\mathbf{s} \mathbf{x})) \prod_{i=1}^d s_i ds_i \bar{\chi}_{\alpha_i, k_i}(s_i).$$

where $\mathbf{s} := \text{diag}(s_1, \dots, s_d)$ is the $d \times d$ diagonal matrix generated by the vector $\{s_i\}$. When $d = 1$, this construction reduces to the univariate GAS-SN distribution.

Compared with the elliptical PDF (15.1), the adaptive PDF in (16.1) requires a d -dimensional integral. This makes its evaluation substantially more computationally demanding.

16.2. Location-Scale Family

The location-scale family is obtained by setting $Y = \boldsymbol{\xi} + \boldsymbol{\omega} Z$, denoted by $Y \sim \vec{L}_{\boldsymbol{\alpha}, \mathbf{k}}(\boldsymbol{\xi}, \Omega, \boldsymbol{\beta})$. Here $\boldsymbol{\omega}$ is the $d \times d$ diagonal scale matrix, and $\Omega = \boldsymbol{\omega}^\top \bar{\Omega} \boldsymbol{\omega}$ is the corresponding covariance matrix.

The PDF of Y is

$$(16.2) \quad \vec{L}_{\boldsymbol{\alpha}, \mathbf{k}}(\mathbf{x}; \boldsymbol{\xi}, \Omega, \boldsymbol{\beta}) := \det(\boldsymbol{\omega})^{-1} \vec{L}_{\boldsymbol{\alpha}, \mathbf{k}}(\mathbf{z}; \bar{\Omega}, \boldsymbol{\beta}).$$

where $\mathbf{z} := \boldsymbol{\omega}^{-1}(\mathbf{x} - \boldsymbol{\xi})$. The density must be evaluated through the standard PDF because the mixing variables $\{s_i\}$ act on the standardized variable Z , not directly on the location-scale variable Y .

16.3. Moments

The first moment of Z is $\mu_z := \mathbf{b} \odot \boldsymbol{\delta}$, where $b_i := \sqrt{2/\pi} \mathbb{E}(X^{-1} | \bar{\chi}_{\alpha_i, k_i})$ and $\odot$ is the Hadamard product.

The (i, j) element of the second moment of Z is

$$\mathbf{m}_2(i, j) := \begin{cases} \mathbb{E}(X^{-2} | \bar{\chi}_{\alpha_i, k_i}) & \text{if } i = j, \\ \bar{\Omega}_{i, j} \mathbb{E}(X^{-1} | \bar{\chi}_{\alpha_i, k_i}) \mathbb{E}(X^{-1} | \bar{\chi}_{\alpha_j, k_j}) & \text{if } i \neq j. \end{cases}$$

where $\bar{\Omega}_{i, i} = 1$ is ignored in the first line. Hence, the covariance is

$$\text{var}\{Z\} := \Sigma_z := \mathbf{m}_2 - \mu_z \mu_z^\top$$

The moments of Y follow the rule of the location-scale family. The first moment of Y is $\xi + \omega \mu_z$. The covariance of Y is $\omega \text{var}\{Z\} \omega$.

16.4. Canonical Form

The adaptive distribution does not have the rotational symmetry of an elliptical distribution. Consequently, its canonical form is not especially useful, because it is not connected to the other distributions in the family by a simple affine transformation.

Assume that the variable Z^* is a *canonical variable*. Then $Z^* \sim \vec{L}_{\alpha, \mathbf{k}}(0, \mathbf{I}_d, \beta_{Z^*})$, where $\beta_{Z^*} = (\beta_*, 0, \dots, 0)^\top$ and β_* are defined by (14.10).

The PDF of Z^* is

$$(16.3) \quad \vec{L}_{\alpha, \mathbf{k}_*}(\mathbf{x}; \beta_*) = L_{\alpha_1, k_1}(x_1; \beta_*) \prod_{j=2}^d L_{\alpha_j, k_j}(x_j).$$

Thus, only the first component has the GAS-SN form, while the remaining components are GSaS components, each with its own shape pair (α, k) .

Only the first component of μ_z is nonzero, namely $\sqrt{2/\pi} \delta_* \mathbb{E}(X^{-1} | \bar{\chi}_{\alpha_1, k_1})$. The matrix $\mathbf{m}_2$ is diagonal, with $\mathbf{m}_2(i, i) = \mathbb{E}(X^{-2} | \bar{\chi}_{\alpha_i, k_i})$.

16.5. Marginal 1D Distribution

The construction of the 1D marginal distribution from Y extends directly from Section 14.7, where $\beta_{1(2)}$ is calculated.

Then the marginal distribution is an univariate GAS-SN: $Y_1 \sim L_{\alpha_1, k_1}(\xi_1, \Omega_{11}, \beta_{1(2)})$.

16.6. 2D Adaptive MLE

Maximum likelihood estimation can be implemented with a stochastic gradient descent (SGD) procedure, but the adaptive model requires several practical adjustments. In the bivariate case, the parameter space is

$$\Theta = \{\rho, \alpha_1, \alpha_2, k_1, k_2, \beta_1, \beta_2, w_1, w_2, \xi_1, \xi_2\}$$

where $\alpha_1, \alpha_2 \in (0, 2)$, $k_1, k_2 \in (2, \infty)$, $w_1 > 0$, $w_2 > 0$, and $\beta_1, \beta_2, \xi_1, \xi_2 \in \mathbb{R}$. Since $\rho \in (-1, 1)$, the optimization uses an unconstrained parameter $\rho_\theta \in \mathbb{R}$ and maps it to the correlation coefficient by $\rho = (2/\pi) \arctan(\rho_\theta)$.

Direct evaluation of the adaptive PDF is computationally expensive, even in two dimensions. For this reason, the MLLK objective is evaluated through the two marginal one-dimensional distributions and supplemented with an L2 regularization term on the correlation coefficient.

In practice, each dimension of the data set is normalized to unit standard deviation. This keeps the gradient components on comparable scales and improves the stability of the SGD updates.

2104 Let Y represent the data set of size N , and $L_m(Y_i; \Theta)$ is the marginal 1D PDF at dimension m
 2105 ($m = 1 \dots d$), then

$$\begin{aligned} \text{MLLK}(\Theta; Y) &:= - \sum_{i=1}^N \sum_{m=1}^d \log L_m(Y_i; \Theta) \\ \text{Gradient}(\Theta) &:= \left\{ \frac{\partial \text{MLLK}(\Theta; Y)}{\partial \Theta_j}, \forall \Theta_j \in \Theta \right\} \end{aligned}$$

2106 Once the MLLK and gradients are calculated. The program makes a small move along the direction
 2107 that is most likely to minimize the MLLK. The scale of the move is based on a learning rate, which
 2108 can be adjusted dynamically. Some randomness can be added to the small move. This allows the
 2109 algorithm to explore the nearby regions, and maybe get "unstuck" over unexpected edges.

2110 When N is large, it may not be computationally feasible to evaluate every $L(Y_i; \Theta)$. One may use
 2111 histogram to compress the data into smaller numbers of bins.

2112 More regularization can be added to the MLLK. For instance, we find it makes a lot of sense to
 2113 add the L2 distances of the skewness and kurtosis on the marginal 1D distributions.

2114 We also regulate the mean of the quadratic form. But the exact distribution of the quadratic form
 2115 is still under research.

Fitting SPX-VIX Daily Returns with Bivariate Distributions

Two MLE fits are performed for the VIX/SPX daily log returns from 1990 to 2025. The first fit uses the bivariate elliptical GAS-SN distribution. The second fit uses the bivariate adaptive GAS-SN distribution.

The major difference is that the adaptive distribution allows each dimension to have its own (α, k) shape. However, it is much more compute-intensive, it requires alternative methods to work around. And it breaks the rotational symmetry that the elliptical distribution has. This requires a different approach to evaluate the quadratic form.

17.1. Elliptical Fit

The bivariate elliptical MLE program is similar to the univariate MLE program. But the hyper-parameter space is much larger:

$$\Theta = \{\rho, \alpha, k, \beta_1, \beta_2, w_1, w_2, \xi_1, \xi_2\}$$

where $\alpha \in (0, 2)$, $k \in (2, \infty)$, $\omega_1 > 0, \omega_2 > 0$, and $\beta_1, \beta_2, \xi_1, \xi_2 \in \mathbb{R}$.

ρ is the correlation coefficient. Since $\rho \in (-1, 1)$, the optimization uses an unconstrained parameter $\rho_\theta \in \mathbb{R}$ and maps it to the correlation coefficient by $\rho = (2/\pi) \arctan(\rho_\theta)$. In the program, ρ is converted to $\bar{\Omega}$ according to (14.14).

The bivariate MLE program is implemented in **python** and **scipy** on github at https://github.com/slihn/gas-impl/blob/main/gas_impl/mle_gas_sn_2d.py.

We run the MLE fitting on the daily returns of VIX and SPX from 1990 to mid-2025, about 8900 two-column samples. Each column in the data set is normalized to one standard deviation. This allows all gradients to have similar scales and helps the MLE to operate smoothly.

Assume there are N samples in the data set, $Y = \{\mathbf{y}_i, i \in 1, 2, \dots, N\}$, the minus log-likelihood (MLLK) is

$$(17.1) \quad \text{MLLK}(\Theta) = -\frac{1}{N} \sum_{i=1}^N \log(L_{\alpha,k}(\mathbf{y}_i; \boldsymbol{\xi}, \Omega, \boldsymbol{\beta}))$$

where $L_{\alpha,k}(\mathbf{x}; \boldsymbol{\xi}, \Omega, \boldsymbol{\beta})$ is the multivariate PDF of the location scale family (15.2).

When N is large, it may not be computationally feasible to compute the PDF on every $\mathbf{y}_i$. A histogram may be used to compress the data into a grid of bins.

Two components of regularization are added to the objective function $\ell(\Theta)$. The L2 distances between the empirical and theoretical statistics are added as follows:

- Correlation: $|\Delta\rho(Y)|^2$.
- The mean of the quadratic form: $\Delta\mu_Q^2 := |\Delta\text{mean}(Q)|^2$. Section 15.6.

MLE seeks the optimal Θ that minimizes the objective function:

$$(17.2) \quad \hat{\Theta} = \text{argmin } \ell(\Theta)$$

$$(17.3) \quad \text{where } \ell(\Theta) = \text{MLLK}(\Theta) + |\Delta\rho(Y)|^2 + \Delta\mu_Q^2$$

17.1.1. The VIX-SPX Bivariate Elliptical Fit. Figure 17.1 shows the results of the bivariate elliptical MLE fit on the VIX/SPX daily log returns. The top two graphs show the 2D scatter plot (left) and the contour plot (right) of the samples. Two overlapping lines are drawn to indicate the angles of the correlation, theoretical vs. empirical. The main accomplishment of this fit is that the correlation coefficient matches nicely at about -0.7.

The contour plot is compared to the theoretical elliptical contour plot in the middle left graph. We note that the sample contours look rectangular instead of elliptical. This is an important research topic left for the future.

The remaining three graphs are for the quadratic form Q in (15.6). The PP-plot in the middle right graph and the QQ-plot in log scale in the bottom right graph show very good match with a clear 45-degree line.

However, the QQ-plot in the bottom left graph is less ideal. The tail is tilted upward after 20. This indicates a poor fit on the outside of the contours. This is probably due to the fact that an elliptical distribution could not capture the rectangular nature of the contours.

17.1.2. The Issue in Marginal Distributions. One major issue with the fit is related to the 1-dimensional marginals. The bivariate MLE finds the best fit at $\alpha = 0.75, k = 4.5$. This is a strange place when we examine it in Figure 12.1. When the bivariate distribution is projected to the 1-dimensional marginal distributions according to Section 15.5, the univariate GAS-SN distributions are near the border of infinite kurtosis.

(The reader is reminded that the degrees of freedom need to be higher than 4 to have valid kurtosis in the Student's t distribution. $k = 4.5$ is in the neighborhood of that threshold.)

Although the implied kurtoses are substantially inaccurate, the marginal plots in Figures 17.2 and 17.3 appear visually acceptable except in the upper-right region. There, the theoretical peak of the VIX marginal density is higher than the observed peak, whereas the theoretical peak of the SPX marginal density is lower than the observed peak.

This discrepancy likely reflects the fact that VIX and SPX require different shape parameters (α, k) for their marginal behavior. The current elliptical construction does not allow separate choices of α and k for the two dimensions, and it remains an open problem to incorporate that additional flexibility within the model.

Overall, these results suggest that a single bivariate distribution is too restrictive to describe the full 35-year history of the SPX and VIX data. Further work is needed to obtain a more satisfactory joint model. A major step forward is to apply this distribution in regime-switching models, such as the Hidden Markov Model (HMM), statistical jump model[35], and mixture-VAE model[27].




FIGURE 17.1. Elliptical MLE fit of VIX/SPX daily log returns from 1990 to 2025 with the bivariate elliptical GAS-SN distribution. Data is standardized to one standard deviation on each axis. The optimal parameters are: $\hat{\Theta} = \{\rho_\theta = -2.12, \alpha = 0.75, k = 4.5, \beta_0 = 0.78, \beta_1 = 0.27, \omega_0 = 0.92, \omega_1 = 0.88, \xi_0 = -0.35, \xi_1 = 0.19\}$.




FIGURE 17.2. VIX Marginal from Elliptical MLE fit of VIX/SPX daily log returns from 1990 to 2025 with the bivariate elliptical GAS-SN distribution. Data is standardized to one standard deviation on each axis.




FIGURE 17.3. SPX Marginal from Elliptical MLE fit of VIX/SPX daily log returns from 1990 to 2025 with the bivariate elliptical GAS-SN distribution. Data is standardized to one standard deviation on each axis.

17.2. Adaptive Fit

The adaptive fit is obtained by maximizing the two marginal likelihoods with an additional regularization term, such as the L^2 distance between the empirical and theoretical correlations. This surrogate objective is used because direct bivariate maximum-likelihood estimation is computationally infeasible on the available hardware.

The resulting contour plot in Figure 17.4 exhibits moderately rectangular level sets, indicating that the adaptive specification captures part of the dependence structure. However, the theoretical correlation reaches only about -0.5 , whereas the empirical correlation is closer to -0.7 .

TODO need to redo the adaptive fit

Although the adaptive distribution allows each dimension to have its own shape parameters, this additional flexibility does not automatically produce a close bivariate fit. In practice, the correlation parameter interacts strongly with the skew parameters, which makes the joint optimization problem difficult. The remaining discrepancy is therefore mainly a dependence-calibration issue rather than a failure of the univariate SPX marginal fit.

Another practical issue is that the quadratic form requires an additional multiplier, or scale adjustment, to produce a satisfactory fit. The theoretical basis for this adjustment remains unclear. Moreover, the squared QQ plots of the marginals in Figures 17.5 and 17.6 show that the adaptive fit reproduces the tails less accurately than the elliptical fit.




FIGURE 17.4. MLE fit of VIX/SPX daily log returns from 1990 to 2025 with the bivariate adaptive distribution. Data is standardized to one standard deviation on each axis.




FIGURE 17.5. VIX Marginal from MLE fit of VIX/SPX daily log returns from 1990 to 2025 with the bivariate adaptive GAS-SN distribution. Data is standardized to one standard deviation on each axis.




FIGURE 17.6. SPX Marginal from MLE fit of VIX/SPX daily log returns from 1990 to 2025 with the bivariate adaptive GAS-SN distribution. Data is standardized to one standard deviation on each axis.

APPENDIX A

List of Useful Formula

A.1. Gamma Function

Gamma function is used extensively in this paper. First, note that $\Gamma(\frac{1}{2}) = \sqrt{\pi}$. Its **reflection formula** is

$$(A.1) \quad \Gamma(z)\Gamma(1-z) = \frac{\pi}{\sin(\pi z)}$$

And the **Legendre duplication formula** is

$$(A.2) \quad \Gamma(z)\Gamma(z+1/2) = 2^{1-2z}\sqrt{\pi}\Gamma(2z)$$

Gamma function Asymptotic: At $x \rightarrow 0$, gamma function becomes

$$(A.3) \quad \lim_{x \rightarrow 0} \Gamma(x) \sim \frac{1}{x}$$

$$\lim_{x \rightarrow 0} \frac{\Gamma(ax)}{\Gamma(bx)} = \frac{b}{a} \quad (ab \neq 0)$$

For a very large x , assume a, b are finite,

$$(A.4) \quad \lim_{x \rightarrow \infty} \frac{\Gamma(x+a)}{\Gamma(x+b)} \sim x^{a-b}$$

Sterling's formula is used to expand the kurtosis formula for a large k , which is:

$$(A.5) \quad \lim_{x \rightarrow \infty} \Gamma(x+1) \sim \sqrt{2\pi x} \left(\frac{x}{e}\right)^x,$$

$$(A.6) \quad \text{or } \lim_{x \rightarrow \infty} \Gamma(x) \sim \sqrt{2\pi} x^{x-1/2} e^{-x}.$$

A.2. Transformation

Laplace transform of cosine is¹

$$(A.7) \quad \int_0^\infty dt \cos(xt) e^{-t/\nu} = \frac{\nu^{-1}}{x^2 + \nu^{-2}} = \frac{\nu}{(\nu x)^2 + 1}$$

Gaussian transform of cosine is²

$$(A.8) \quad \int_0^\infty dt \cos(xt) e^{-t^2/2} = \sqrt{\frac{\pi}{2}} e^{-x^2/2}$$

Hence $\int_0^\infty dt \cos(xt) e^{-t^2/2s^2} = \sqrt{\frac{\pi}{2}} s e^{-(sx)^2/2}$

¹See https://proofwiki.org/wiki/Laplace_Transform_of_Cosine

²See <https://www.wolframalpha.com/input?i=integrate+cos%28a+x%29+e%5E%28-x%5E2%2F2%29+dx+from+0+to+infity>

A.3. Half-Normal Distribution

2208

2209 The moments of the half-normal distribution (HN)³ are used several times. Its PDF is defined as

$$(A.9) \quad p_{HN}(x; \sigma) := \sqrt{\frac{2}{\pi}} \frac{1}{\sigma} e^{-x^2/(2\sigma^2)}, \quad x > 0$$

2210 which is a special case of GG with $d = 1, p = 2, a = \sqrt{2}\sigma$. Its moments are

$$(A.10) \quad E_{HN}(T^n) = \sigma^n \frac{2^{n/2}}{\sqrt{\pi}} \Gamma\left(\frac{n+1}{2}\right)$$

2211 which are the same as those of a normal distribution.

³See https://en.wikipedia.org/wiki/Half-normal_distribution

Bibliography

- 2213 [1] Adelchi Azzalini. *The Skew-Normal and Related Families*. Institute of Mathematical Statistics Monographs. Cam-
2214 bridge University Press, 2013.
- 2215 [2] H. Bateman and A. Erdélyi. *Higher Transcendental Functions*, volume 3 of *California Institute of Technology*
2216 *Bateman Manuscript Project*. McGraw-Hill Book Company., 1955.
- 2217 [3] Berndt Bruce C. *Ramanujan's Notebooks, Part I*. Springer New York, NY, 1985.
- 2218 [4] Peter K. Clark. A subordinated stochastic process model with finite variance for speculative prices. *Econometrica*,
2219 41(1):135–155, 1973.
- 2220 [5] John C. Cox, Jonathan E. Ingersoll, and Stephen A. Ross. A theory of the term structure of interest rates. *Econo-*
2221 *metrica*, 53(2):385–407, 1985.
- 2222 [6] J.C. Crisanto-Neto, M.G.E. da Luz, E.P. Raposo, and G.M. Viswanathan. An efficient series approximation for the
2223 lévy α -stable symmetric distribution. *Physics Letters A*, 382(35):2408–2413, 2018.
- 2224 [7] Luc Devroye. Random variate generation for exponentially and polynomially tilted stable distributions. *ACM Trans.*
2225 *Model. Comput. Simul.*, 19(4), November 2009.
- 2226 [8] Luc Devroye and Lancelot F. James. On simulation and properties of the stable law. *Statistical Methods & Appli-*
2227 *cations*, 23:307 – 343, 2014.
- 2228 [9] *NIST Digital Library of Mathematical Functions*. <https://dlmf.nist.gov/>, Release 1.2.0 of 2024-03-15. F. W. J.
2229 Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders,
2230 H. S. Cohl, and M. A. McClain, eds.
- 2231 [10] William Feller. Two singular diffusion problems. *Annals of Mathematics*, 54(1):173–182, 1951.
- 2232 [11] William Feller. On a generalization of marcel riesz' potentials and the semi-groups generated by them. *Meddelanden*
2233 *Lunds Universitets Matematiska Seminarium*, pages 73–81, 1952.
- 2234 [12] William Feller. *An Introduction to Probability Theory and its Applications (2nd Edition)*, volume 2. John Wiley &
2235 Sons, Inc., 1971.
- 2236 [13] Wikimedia Foundation. Inverse distribution, 2023. Retrieved December 1, 2023, from [https://en.wikipedia.org/](https://en.wikipedia.org/wiki/Inverse_distribution)
2237 [wiki/Inverse_distribution](https://en.wikipedia.org/wiki/Inverse_distribution).
- 2238 [14] Lancelot F. James. Generalized mittag leffler distributions arising as limits in preferential attachment models, 2015.
- 2239 [15] George Karagiannidis and Athanasios Lioumpas. An improved approximation for the gaussian q-function. *IEEE*
2240 *Communications Letters*, 11:644–646, 08 2007.
- 2241 [16] Paul Lévy. *Calcul des Probabilités*. Gauthier-Villars, 1925.
- 2242 [17] Xiaoyue Li and John M. Mulvey. Optimal portfolio execution in a regime-switching market with non-linear impact
2243 costs: Combining dynamic program and neural network. *arXiv preprint*, 6 2023. Available at arXiv: [https://](https://arxiv.org/abs/2306.08809)
2244 arxiv.org/abs/2306.08809.
- 2245 [18] Stephen H.T. Lihn. Stable count distribution for the volatility indices and space-time generalized stable characteristic
2246 function. *SSRN Working Paper*, 7 2020. Available at SSRN: <https://ssrn.com/abstract=3659383>.
- 2247 [19] Stephen H.T. Lihn. Generalization of the alpha-stable distribution with the degree of freedom. *arXiv preprint*, 5
2248 2024. Available at arXiv: <https://arxiv.org/abs/2405.04693>.
- 2249 [20] F. Mainardi. *Fractional Calculus and Waves in Linear Viscoelasticity: An Introduction to Mathematical Models*. Im-
2250 perial College Press, 2010. See Appendix F at [https://appliedmath.brown.edu/sites/default/files/fractional/](https://appliedmath.brown.edu/sites/default/files/fractional/36%20TheWrightFunctions.pdf)
2251 [36%20TheWrightFunctions.pdf](https://appliedmath.brown.edu/sites/default/files/fractional/36%20TheWrightFunctions.pdf).
- 2252 [21] Francesco Mainardi and Armando Consiglio. The wright functions of the second kind in mathematical physics.
2253 *Mathematics*, 8(6), 2020.
- 2254 [22] Francesco Mainardi, Yuri Luchko, and Gianni Pagnini. The fundamental solution of the space-time fractional dif-
2255 fusion equation. *Frac. Calc. Appl. Anal.*, 4, 03 2007. See <https://arxiv.org/abs/cond-mat/0702419>.
- 2256 [23] Francesco Mainardi and Gianni Pagnini. Mellin-barnes integrals for stable distributions and their convolutions.
2257 *Fractional Calculus and Applied Analysis*, 11:443–456, 2008.
- 2258 [24] Arak Mathai and Hans Haubold. *Fractional and Multivariable Calculus: Model Building and Optimization Prob-*
2259 *lems*. Springer Cham, 12 2017.

- [25] Hjalmar Mellin. Zur theorie zweier allgemeinen klassen bestimmter integrale. *Acta Societatis Scientiarum Fennicæ*, page 1–75, 1897.
- [26] The mpmath development team. *mpmath: a Python library for arbitrary-precision floating-point arithmetic (version 1.3.0)*, 2023. <http://mpmath.org/>.
- [27] Yuqi Nie, John M. Mulvey, H. Vincent Poor, Chenyu Yu, and Hao Huang. Deep generative models meet statistical methods: A generalized framework for financial regime identification. *SSRN*, 2024. Available at SSRN: <https://ssrn.com/abstract=5332057>.
- [28] John Nolan. *Univariate Stable Distributions, Models for Heavy Tailed Data*. Springer Cham, 01 2020.
- [29] Donald B. Owen. Tables for Computing Bivariate Normal Probabilities. *The Annals of Mathematical Statistics*, 27(4):1075 – 1090, 1956.
- [30] Richard B. Paris, Armando Consiglio, and Mainardi Francesco. On the asymptotics of wright functions of the second kind. *Fract. Calc. Appl. Anal.*, 24(1):54–72, 01 2021.
- [31] Jim Pitman. Combinatorial stochastic processes. In *Ecole d’Eté de Probabilités de Saint-Flour XXXII - 2002*, page 66. Springer Berlin, Heidelberg, 2006.
- [32] Jim Pitman and Marc Yor. The two-parameter poisson-dirichlet distribution derived from a stable subordinator. *The Annals of Probability*, 25(2):855–900, 1997.
- [33] Dimiter Prodanov. Computation of the wright function from its integral representation. In *Advances in Nonlinear Dynamics*, volume 1, pages 421–431. Springer Nature Switzerland, 05 2024.
- [34] W. R. Schneider. Stable distributions: Fox function representation and generalization. In S. Albeverio, G. Casati, and D. Merlini, editors, *Stochastic Processes in Classical and Quantum Systems*, pages 497–511, Berlin, Heidelberg, 1986. Springer Berlin Heidelberg.
- [35] Y. Shu, C. Yu, and J.M. Mulvey. Downside risk reduction using regime-switching signals: a statistical jump model approach. *Journal of Asset Management*, 25:493–507, 2024.
- [36] E. W. Stacy. A generalization of the gamma distribution. *The Annals of Mathematical Statistics*, 33(3):1187 – 1192, 1962. See also https://en.wikipedia.org/wiki/Generalized_gamma_distribution.
- [37] Student. The probable error of a mean. *Biometrika*, 6:1–25, 1908. See also https://en.wikipedia.org/wiki/Student%27s_t-distribution.
- [38] Y.L. Tong. *The Multivariate Normal Distribution*. Research in Criminology. Springer New York, 1990.
- [39] Pauli Virtanen, Ralf Gommers, Travis E. Oliphant, Matt Haberland, Tyler Reddy, David Cournapeau, Evgeni Burovski, Pearu Peterson, Warren Weckesser, Jonathan Bright, Stéfan J. van der Walt, Matthew Brett, Joshua Wilson, K. Jarrod Millman, Nikolay Mayorov, Andrew R. J. Nelson, Eric Jones, Robert Kern, Eric Larson, C J Carey, İlhan Polat, Yu Feng, Eric W. Moore, Jake VanderPlas, Denis Laxalde, Josef Perktold, Robert Cimrman, Ian Henriksen, E. A. Quintero, Charles R. Harris, Anne M. Archibald, Antônio H. Ribeiro, Fabian Pedregosa, Paul van Mulbregt, and SciPy 1.0 Contributors. SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. *Nature Methods*, 17:261–272, 2020.
- [40] E. M. Wright. On the coefficients of power series having exponential singularities. *Journal London Math. Soc.*, s1-8, Issue 1 (1933):71–79, 1933.
- [41] E. M. Wright. The asymptotic expansion of the generalized bessel function. *Proc. London Math. Soc.*, s2-38, Issue 1 (1935):257–270, 1935.