

Topics in Statistical Theory — Example Sheet 1

Lucas Riedstra

8 November 2020

Question 4. Let $X \sim \text{Bin}(n, p)$. Compare the Hoeffding, Bennett, and Bernstein upper bounds on $\mathbb{P}(X/n \geq \frac{1}{2})$ as $p \rightarrow 0$.

Solution. Note that X/n is the average of n i.i.d. random variables $Y_i \sim \text{Bern}(p)$, where $Y_i \in [0, 1]$ for all i .

1. We start with Hoeffding's inequality. In this case, we have

$$\mathbb{P}\left(X/n - p \geq \frac{1}{2}\right) \leq \exp\left(-\frac{2n^2(1/2)^2}{n}\right) = \exp\left(-\frac{n}{2}\right),$$

and this bound also holds as $p \rightarrow 0$.

2. We continue with Bennett's inequality. We consider the mean-zero random variables $Y_i - p$, which are bounded from above by $b = 1 - p$. Now Bennett's inequality tells us that

$$\mathbb{P}\left(X/n \geq \frac{1}{2}\right) \leq \exp\left(-\frac{np(1-p)}{(1-p)^2} h\left(\frac{1-p}{2p(1-p)}\right)\right) = \exp\left(-\frac{np}{1-p} \cdot h\left(\frac{1}{2p}\right)\right).$$

We compute

$$h\left(\frac{1}{2p}\right) = \left(1 + \frac{1}{2p}\right) \log\left(1 + \frac{1}{2p}\right) - \frac{1}{2p} = \log\left(1 + \frac{1}{2p}\right) + \frac{1}{2p} \left(\log\left(1 + \frac{1}{2p}\right) - 1\right),$$

and therefore

$$\frac{np}{1-p} h\left(\frac{1}{2p}\right) \geq \frac{n}{2(1-p)} \left(\log\left(1 + \frac{1}{2p}\right) - 1\right) \xrightarrow{p \rightarrow 0} \infty.$$

It follows that the Bennett upper bound converges to 0 as $p \rightarrow 0$.

3. We finish with Bernstein's inequality. We have for $q \geq 3$ that

$$\frac{2(1-p)}{q!} \leq \frac{2}{q!} \leq 3^{2-q},$$

and therefore we have

$$\begin{aligned} \mathbb{E}[(Y_i - p)_+^q] &= p(1-p)^q = \sigma_p^2(1-p)^{q-1} = (q! \sigma_p^2(1-p)^{q-2}/2) \cdot (2(1-p)/q!) \\ &\leq q! \sigma_p^2((1-p)/3)^{q-2}/2, \end{aligned}$$

so $Y_i - p$ satisfies Bernstein's condition with $c = \frac{1-p}{3}$. Now Bernstein's inequality tells us that

$$\mathbb{P}(X/n \geq \frac{1}{2}) \leq \exp\left(-\frac{n(1/2)^2}{2(\sigma_p^2 + (1-p)/6)}\right) = \exp\left(-\frac{n}{8\sigma_p^2 + 4(1-p)/3}\right) \xrightarrow{p \rightarrow 0} \exp\left(-\frac{3n}{4}\right).$$

Of course, the true limit is 0 for any n , which is only given by Bennett's inequality. We also see that Hoeffding's inequality gives the most loose bound.

Question 10. (a) *Verify the algebraic identity*

$$\varphi_\sigma(x - \mu)\varphi_{\sigma'}(x - \mu') = \varphi_{\sigma\sigma'/(\sigma^2 + \sigma'^2)^{1/2}}(x - \mu^*)\varphi_{(\sigma^2 + \sigma'^2)^{1/2}}(\mu - \mu')$$

where $\mu^* := (\sigma'^2\mu + \sigma^2\mu')/(\sigma^2 + \sigma'^2)$, and φ_σ is the $N(0, \sigma^2)$ density.

(b) Let $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} N(0, \sigma^2)$. Taking K to be the $N(0, 1)$ density, show that the MISE of the kernel density estimate \hat{f}_n with kernel K and bandwidth h can be expressed exactly as

$$\text{MISE}(\hat{f}_n) = \frac{1}{2\pi^{1/2}} \left\{ \frac{1}{nh} + \left(1 - \frac{1}{n}\right) \frac{1}{(h^2 + \sigma^2)^{1/2}} - \frac{2^{3/2}}{(h^2 + 2\sigma^2)^{1/2}} + \frac{1}{\sigma} \right\}.$$

Proof. (a) We have

$$\begin{aligned} & \frac{(x - \mu)^2}{\sigma^2} + \frac{(x - \mu')^2}{\sigma'^2} \\ &= \frac{\sigma'^2(x - \mu)^2 + \sigma^2(x - \mu')^2}{\sigma^2\sigma'^2} \\ &= \frac{(\sigma^2 + \sigma'^2)x^2 - 2(\sigma'^2\mu + \sigma^2\mu')x + \sigma'^2\mu^2 + \sigma^2\mu'^2}{\sigma^2\sigma'^2} \\ &= \frac{(\sigma^2 + \sigma'^2)(x^2 - 2\mu^*x) + \sigma'^2\mu^2 + \sigma^2\mu'^2}{\sigma^2\sigma'^2} \\ &= \frac{(\sigma^2 + \sigma'^2)(x^2 - 2\mu^* + \mu^{*2}) - (\sigma^2 + \sigma'^2)\mu^{*2} + \sigma'^2\mu^2 + \sigma^2\mu'^2}{\sigma^2\sigma'^2} \\ &= \frac{(x - \mu^*)^2}{(\sigma\sigma'/(\sigma^2 + \sigma'^2)^{1/2})^2} + \frac{\sigma'^2\mu + \sigma^2\mu'^2 - (\sigma'^2\mu + \sigma^2\mu')^2/(\sigma^2 + \sigma'^2)}{\sigma^2\sigma'^2} \\ &= \frac{(x - \mu^*)^2}{(\sigma\sigma'/(\sigma^2 + \sigma'^2)^{1/2})^2} + \frac{(\sigma^2 + \sigma'^2)(\sigma'^2\mu + \sigma^2\mu'^2) - (\sigma'^2\mu + \sigma^2\mu')^2}{\sigma^2\sigma'^2(\sigma^2 + \sigma'^2)} \\ &= \frac{(x - \mu^*)^2}{(\sigma\sigma'/(\sigma^2 + \sigma'^2)^{1/2})^2} + \frac{(\mu - \mu')^2}{\sigma^2 + \sigma'^2} \\ &= \frac{(x - \mu^*)^2}{(\sigma\sigma'/(\sigma^2 + \sigma'^2)^{1/2})^2} + \frac{(\mu - \mu')^2}{((\sigma^2 + \sigma'^2)^{1/2})^2}, \end{aligned}$$

which proves the claim.

(b) Let $K = \varphi_1$ and define $K_h(x) := h^{-1}K(x/h)$ so $K_h = \varphi_h$. Then recall from the lectures that

$$\text{MISE}(\hat{f}_n) = \frac{1}{n} \int_{\mathbb{R}} [(\varphi_h^2 * \varphi_\sigma)(x) - (\varphi_h * \varphi_\sigma)^2(x)] dx + \int_{-\infty}^{\infty} [(\varphi_h * \varphi_\sigma)(x) - \varphi_\sigma(x)]^2 dx$$

We will use the previous exercise to compute all these terms. We have in general

$$\begin{aligned} (\varphi_h * \varphi_\sigma)(x) &= \int_{\mathbb{R}} \varphi_h(x - z)\varphi_\sigma(z) dz \\ &= \int_{\mathbb{R}} \varphi_\sigma(z)\varphi_h(z - x) dz \\ &= \varphi_{(\sigma^2 + h^2)^{1/2}}(x) \int_{\mathbb{R}} \varphi_\xi(z - \mu^*) dz \\ &= \varphi_{(\sigma^2 + h^2)^{1/2}}(x). \end{aligned} \tag{1}$$

We also have

$$\varphi_\sigma^2(x - \mu) = \varphi_{\sigma/\sqrt{2}}(x - \mu) \varphi_{\sqrt{2}\sigma}(0) = \frac{1}{2\sigma\sqrt{\pi}} \varphi_{\sigma/\sqrt{2}}(x - \mu). \quad (2)$$

Combining eqs. (1) and (2) we get

$$\begin{aligned} (\varphi_h^2 * \varphi_\sigma)(x) &= \int_{\mathbb{R}} \varphi_h^2(x - z) \varphi_\sigma(z) dz \\ &= \frac{1}{2h\sqrt{\pi}} \int_{\mathbb{R}} \varphi_{h/\sqrt{2}}(x - z) \varphi_\sigma(z) dz \\ &= \frac{1}{2h\sqrt{\pi}} (\varphi_{h/\sqrt{2}} * \varphi_\sigma)(x) \\ &= \frac{1}{2h\sqrt{\pi}} \varphi_{(\sigma^2 + h^2/2)^{1/2}}(x) \end{aligned}$$

We also get

$$(\varphi_h * \varphi_\sigma)^2(x) = \varphi_{(\sigma^2 + h^2)^{1/2}}^2(x) = \frac{1}{2(\sigma^2 + h^2)^{1/2}\sqrt{\pi}} \varphi_{(\sigma^2 + h^2)^{1/2}/\sqrt{2}}(x).$$

Finally, we have

$$((\varphi_h * \varphi_\sigma)(x) - \varphi_\sigma(x))^2 = (\varphi_h * \varphi_\sigma)^2(x) - 2(\varphi_h * \varphi_\sigma)(x) \varphi_\sigma(x) + \varphi_\sigma^2(x).$$

The first term we already computed, the third term is $\frac{1}{2\sigma\sqrt{\pi}} \varphi_{\sigma/\sqrt{2}}(x)$, so we only need to compute

$$(\varphi_h * \varphi_\sigma)(x) \varphi_\sigma(x) = \varphi_{(\sigma^2 + h^2)^{1/2}}(x) \varphi_\sigma(x) = \varphi_\xi(x) \varphi_{(2\sigma^2 + h^2)^{1/2}}(0) = \frac{1}{\sqrt{2\pi}(2\sigma^2 + h^2)^{1/2}} \varphi_\xi(x),$$

where ξ is an irrelevant constant.

Combining all these terms and using that $\varphi_\sigma(x - \mu)$ integrates to 1 for any μ, σ , we get

$$\begin{aligned} \text{MISE } \hat{f}_n &= \frac{1}{n} \left(\frac{1}{2h\sqrt{\pi}} - \frac{1}{2(\sigma^2 + h^2)^{1/2}\sqrt{\pi}} \right) + \frac{1}{2(\sigma^2 + h^2)^{1/2}\sqrt{\pi}} - \frac{\sqrt{2}}{\sqrt{\pi}(2\sigma^2 + h^2)^{1/2}} + \frac{1}{2\sigma\sqrt{\pi}} \\ &= \frac{1}{2\sqrt{\pi}} \left(\frac{1}{nh} + \left(1 - \frac{1}{n}\right) \frac{1}{(\sigma^2 + h^2)^{1/2}} - \frac{2^{3/2}}{(2\sigma^2 + h^2)^{1/2}} + \frac{1}{\sigma} \right). \end{aligned}$$

□

Question 11. Use the expression from 10(b) to derive an upper bound of the form $\text{MISE } \hat{f}_n \leq C_1/(nh) + C_2^2 h^4$ (where you should specify C_1, C_2). Show that $\varphi_\sigma \in \mathcal{F}_{\mathcal{N}}(2, L)$ with $L^2 = 3/(8\pi^{1/2}\sigma^5)$, and hence compare the bound from the first part of this question with that obtained from the general theory from lectures.

Solution. We have

$$\left(1 - \frac{1}{n}\right) \frac{1}{(\sigma^2 + h^2)^{1/2}} - \frac{2^{3/2}}{(2\sigma^2 + h^2)^{1/2}} \leq \frac{1}{(\sigma^2 + h^2)^{1/2}} - \frac{2}{(\sigma^2 + h^2/2)^{1/2}} < 0,$$

I do not know how to obtain an upper bound of the form $C_1/(nh) + C_2^2 h^4$ from this expression.

To show that $\varphi_\sigma \in \mathcal{F}_{\mathcal{N}}(2, L)$, we must show that $\varphi_\sigma \in \mathcal{F}_{\mathcal{N}}(2, L)$. By question 9, it suffices to show that $\|\varphi_\sigma''\|_{L^2}^2 \leq L^2$. A simple computation gives

$$\varphi_\sigma''(x) = \frac{1}{\sqrt{2\pi}\sigma^5} (x^2 - \sigma^2) \exp\left(-\frac{x^2}{2\sigma^2}\right) \leq \frac{1}{\sqrt{2\pi}\sigma^5} x^2 \exp\left(-\frac{x^2}{2\sigma^2}\right)$$

so

$$\|\varphi''_\sigma\|_{L^2}^2 \leq \frac{1}{2\pi\sigma^{10}} \int_{\mathbb{R}} x^4 \exp\left(-\frac{x^2}{\sigma^2}\right) dx \stackrel{*}{=} \frac{1}{2\pi\sigma^{10}} \cdot \frac{3}{4}\sqrt{\pi}\sigma^5 = \frac{3}{8\sqrt{\pi}\sigma^5} = L^2,$$

where \star can be computed using the fact that the integral is, up to scaling, the fourth moment of $N(0, \sqrt{2}\sigma)$ distribution.

Note that for $K = \varphi_1$, we have

$$R(K) = \int_{-\infty}^{\infty} \varphi_1^2(x) dx = \frac{1}{2\sqrt{\pi}},$$

while

$$\mu_2^2(K) = \int_{-\infty}^{\infty} x^2 \varphi_1(x) dx = 1.$$

Plugging all the above into theorem 27 shows that

$$\text{MISE}(\hat{f}_n) \leq \frac{1}{2\sqrt{\pi}} \frac{1}{nh} + \frac{3}{8\sqrt{\pi}\sigma^5} h^4.$$