IRRATIONALITY MEASURES FOR $G \pm 5\pi^2/48$ AND $L(2, \chi_{-3})$

SEAN LITSEY

ABSTRACT. Building on OpenAI's determinant method for the irrationality of Catalan's constant $G = \sum_{j\geq 0}(-1)^j/(2j+1)^2$, we prove

$$\mu\Big(G - \frac{5\pi^2}{48}\Big) \leq 32.44, \qquad \mu\Big(G + \frac{5\pi^2}{48}\Big) \leq 32.44,$$
$$\mu\big(L(2, \chi_{-3})\big) \leq 14.35,$$

where μ is the irrationality exponent. We know of no earlier irrationality measure for $G \pm 5\pi^2/48$. For $L(2, \chi_{-3})$, which Calegari, Dimitrov and Tang proved irrational, the previous best bound we know of is their effective bound $\mu \leq 24781$.

The approximations come from determinants of *rectangular* Löwner matrices of digamma differences at integer nodes. We bound their denominators with p-adic assignment problems and rank deficits modulo p. We prove nonvanishing modulo a prime by reducing it to bordered Krylov determinants, which we evaluate through Jacobi matrices and Darboux transformations. For $L(2, \chi_{-3})$, a finite-field interpolation argument bounds the corank by three; removing at most three common low nodes gives nonvanishing at a fixed scale throughout a prime interval. We bound their size with Andréief's identity and a certified lower bound for a discrete logarithmic energy. Two new ingredients lower the exponent for $G \pm 5\pi^2/48$: a second-level p-adic saving at primes where the leading block is not clean, and a decay certificate with dual-guided atoms and steeper contour rays.

Every result is formalized in Lean 4, with no unproved steps and only the standard axioms.

CONTENTS

1. Introduction 2
2. Overview of the method and of the new ideas 4
3. The construction 8
4. Arithmetic: the content 9
5. A uniform rescaled-core deficit 14
6. A next-level deficit at non-clean primes 19

Date: October 23, 2026.

2020 *Mathematics Subject Classification.* Primary 11J82; Secondary 11J72, 15A15, 31A15, 68V20.

Key words and phrases. Catalan's constant, $L(2, \chi_{-3})$, irrationality exponent, Löwner matrices, Cauchy determinants, Andréief identity, p-adic valuations, Jacobi matrices, Darboux transformation, formal verification, Lean.

7. Periodic content for the conductor-three frame 21
8. Nonvanishing modulo a prime 24
9. Uniform nonvanishing after bounded low-node deletion 33
10. A quarter border and characteristic-zero selection 39
11. Analytic decay 42
12. Proof of Theorem 1.1 46
13. Formalization 48
Disclosure of AI use 50
References 51

1. Introduction

OpenAI [32] recently proved that Catalan's constant

$$G=\sum_{j=0}^{\infty}\frac{(-1)^j}{(2j+1)^2}=0.91596\,55941\ldots$$

is irrational, and formalized the proof in Lean. (An earlier claimed proof [39] was found to contain a gap [42].) Their argument, and the irrationality-measure results built on it [17, 7, 25], use a determinant of large size whose denominators and decay are balanced very closely.

In the present paper we study a different, much smaller, family of determinants: rectangular Löwner matrices of digamma differences, attached to the numbers

$$\lambda=\frac{G}{2}-\frac{5\pi^2}{96},\qquad \lambda'=\frac{G}{2}+\frac{5\pi^2}{96},\qquad \frac{1}{4}L(2,\chi_{-3}),$$
$$L(2,\chi_{-3})=\sum_{k\ge 0}\Big(\frac{1}{(3k+1)^2}-\frac{1}{(3k+2)^2}\Big)=0.78130\,24128\ldots$$

Thus $G-5\pi^2/48=2\lambda$, $G+5\pi^2/48=2\lambda'$, and $\lambda+\lambda'=G$.

Recall that the *irrationality exponent* $\mu(\xi)$ of a real number ξ is the infimum of the μ such that $|\xi-a/b|>b^{-\mu}$ for all integers a and all sufficiently large b.

Theorem 1.1. *For every $\varepsilon>0$ there is q_0 such that for all coprime integers p, q with $q\ge q_0$,*

$$\Big|G-\frac{5\pi^2}{48}-\frac{p}{q}\Big|>q^{-32.44-\varepsilon},\qquad \Big|G+\frac{5\pi^2}{48}-\frac{p}{q}\Big|>q^{-32.44-\varepsilon},$$
$$\Big|L(2,\chi_{-3})-\frac{p}{q}\Big|>q^{-14.35-\varepsilon}.$$

In particular $G\pm 5\pi^2/48$ are irrational, $\mu(G\pm 5\pi^2/48)\le 32.44$ and $\mu(L(2,\chi_{-3}))\le 14.35$.

The computed values of the exponents are 32.4392 and 14.3475713785…. Theorem 1.1, together with every result in this paper, has been verified

in the Lean 4 proof assistant, with no unproved steps and only the standard axioms; see Section 13. The formalization is available at https://github.com/slitsey/catalan-irrationality-measure. The use of AI in this work is disclosed at the end of the paper.

1.1. **Previous results.** For earlier work on the arithmetic of G and related values see [36, 46, 45, 22, 23] and the survey [48]; for recent rational approximations to G see Eskandari [10]. Finite irrationality exponents for G itself were obtained after OpenAI's proof by Jha [17], Carreira [7] and the author [25]. Our searches found no previous irrationality result, and no irrationality measure, for $G - 5\pi^2/48$ or $G + 5\pi^2/48$. Eskandari, Murty and Nemoto [11] construct a 2-dimensional mixed motive over $\mathbb{Q}$ with period G and use it to obtain linear forms in 1 and G. Eskandari [10] improves the approximation exponent of these Rivoal–Zudilin type forms from 0.52 to 0.62 (a preprint; explicit approximations, not a lower bound for all rationals). Sun's preprint [39] claimed a proof of the irrationality of G; Wachs [42] reports that its main estimate is not justified.

Calegari, Dimitrov and Tang proved that 1, $\zeta(2)$ and $L(2, \chi_{-3})$ are linearly independent over $\mathbb{Q}$ [5], by a new arithmetic holonomy bound applied to a construction of Zagier. In [6] they made holonomy bounds effective; Theorem 5.1 there gives $\mu(L(2, \chi_{-3})) \leq 24781$. Theorem 1.1 improves this to 14.35.

For comparison, among classical constants, the bound $\mu(\log 2) < 3.5746$ of Marcovecchio [29] and the Rhin–Viola bounds for $\zeta(2)$ and $\zeta(3)$ [34, 35] (after Apéry and Beukers [3]) have been joined by recent preprints. OpenAI proved $\mu(\pi) = 2$, with a Lean formalization [33], improving the bound $\mu(\pi) < 7.1033$ of Zeilberger and Zudilin [43]. Liu, Jiang and Zhang proved $\mu(\log a) = 2$ for every rational $a > 0$, $a \neq 1$, and for nonzero rational combinations of such logarithms [26]. Giraudin [14] lowered the bound for $\mu(\zeta(2))$ to below 5.0193784. Fauzan [12] announced a proof that $\zeta(5)$ is irrational, using rationally normalized Hankel determinants, and Lai [24] combined his method with simultaneous approximations of Zudilin and of Dauguet and Zudilin to show that 1, $\zeta(3)$ and $\zeta(4)$ are linearly independent over $\mathbb{Q}$.

Related work on determinants. Irrationality proofs built on determinants of large size have a precedent in Zudilin's determinantal criterion with Hankel determinants [47]; the preprint of Liu, Jiang and Zhang on logarithms [26] uses a separated-weight interpolation determinant, and the arguments of Fauzan and Lai [12, 24] use Hankel determinants. Brown [4] proposes a multivariate transfinite-diameter test for Mellin-type integrals, applied to $\zeta(2)$. The framework we use, determinants whose decay is controlled by Cauchy determinants and Andréief's identity and whose denominators are controlled prime by prime, is OpenAI's [32]. Our determinants are Löwner (divided-difference) matrices of a digamma difference, a matrix-valued analogue of Padé and Hermite–Padé approximation [31]. Formal proofs of irrationality

results in Lean also include the irrationality of $\zeta(3)$ by Liu, Zhang and Zhi [27].

1.2. **Outline of the proof.** Section 2 explains the construction and the new ideas without technicalities, and states the general deduction. Sections 3–11 give the construction and the three estimates (denominator, nonvanishing, decay); Section 12 deduces Theorem 1.1. Section 13 describes the formalization, and the disclosure of AI use follows it.

1.3. **Notation and conventions.** $\psi = \Gamma'/\Gamma$ is the digamma function; v_p is the p-adic valuation; for $P \in \mathbb{Q}[x]$, $\operatorname{den} P$ is the least positive integer d with $dP \in \mathbb{Z}[x]$; $\vartheta(x; M, a) = \sum_{p \le x,\ p \equiv a\ (M)} \log p$; $(x)_n = x(x+1)\cdots(x+n-1)$; $[0, r) = \{0, 1, \dots, r-1\}$. For finite sets I, J of integers and a matrix A indexed by integers, $A[I, J]$ is the submatrix with rows I and columns J.

2. Overview of the method and of the new ideas

2.1. **Löwner matrices of digamma differences.** For a function F analytic near distinct points $x_0, \dots, x_{h-1}$ the *Löwner matrix* [28] is the matrix of divided differences

$$\mathsf{L}_F[a, b] = \frac{F(x_b) - F(x_a)}{x_b - x_a}\ (a \neq b), \qquad \mathsf{L}_F[a, a] = F'(x_a).$$

It is a Cauchy-like structured matrix, and Hermite's formula gives $\mathsf{L}_F = \frac{1}{2\pi i} \oint F(z)\, v(z) v(z)^{\top} dz$ with $v_a(z) = 1/(z - x_a)$, so that a representation of F as a Stieltjes or contour integral exhibits L_F as a Gram matrix of Cauchy kernels.

We take three digamma differences, with integer nodes:

$$\begin{gathered} F_\lambda(z) = \psi(z+1) - \psi(z+\tfrac{3}{4}), \quad \omega = \tfrac{1}{16}, \quad \xi = \lambda := \tfrac{G}{2} - \tfrac{5\pi^2}{96}, \\ F_{\lambda'}(z) = \psi(z+1) - \psi(z+\tfrac{1}{4}), \quad \omega = -\tfrac{1}{16}, \quad \xi = \lambda' := \tfrac{G}{2} + \tfrac{5\pi^2}{96}, \\ F_\chi(z) = -\tfrac{1}{3}\psi(z+\tfrac{1}{3}) + \tfrac{1}{2}\psi(z+\tfrac{1}{2}) - \tfrac{1}{6}\psi(z+1), \\ \omega = -\tfrac{1}{6}, \quad \xi = \tfrac{1}{4} L(2, \chi_{-3}). \end{gathered} \tag{2.1}$$

In each case $\xi = \omega\, F'(0)$ (this is an evaluation of ψ' at $\frac{1}{3}, \frac{1}{2}, \frac{1}{4}, \frac{3}{4}, 1$), while $F(b) - F(a)$ and $F'(a) - F'(0)$ are rational for integers $a, b \ge 0$. Hence, for any finite sets of nonnegative integer nodes R (rows) and Q (columns),

$$\omega\, \mathsf{L}_F[R, Q] = M[R, Q] + \xi\, E_{R,Q}, \qquad E_{R,Q}[a, b] = \mathbf{1}[a = b], \tag{2.2}$$

with an explicit rational symmetric kernel M on $\mathbb{Z}^2_{\ge 0}$ (§3). Replacing ξ by a variable x gives a matrix polynomial $\Phi(x) = M[R, Q] + x E_{R,Q}$, whose determinant $P(x)$ is a polynomial with rational coefficients of degree at most $|R \cap Q|$, and $P(\xi) = \det(\omega\, \mathsf{L}_F[R, Q])$. If $|\xi - p/q|$ is very small, $P(p/q)$ is very close to $P(\xi)$, which is tiny; on the other hand $q^{|R \cap Q|} \operatorname{den}(P)\, P(p/q)$ is

an integer. If it is not zero we get a contradiction as soon as

$$\underbrace{\log \operatorname{den} P}_{\text{content}} + |R \cap Q| \log q \;<\; \underbrace{-\log |P(p/q)|}_{\text{decay}}.$$

Three estimates are therefore needed: an upper bound for the *content* $\log \operatorname{den} P$, an upper bound for $|P(x)|$ near ξ (the *decay*), and *nonvanishing* of $P(p/q)$ for a size that can be chosen in terms of q. We use rectangular geometries, with $R \neq Q$, and all three estimates are new for these constants.

2.2. **New idea 1: rectangular (hybrid) frames.** We take a long common block C of r nodes, which carries the variable x, and two short interleaved blocks of $m \approx r/8$ nodes,

$$C = [0, r), \qquad X' = \{r, r+2, \dots, r+2m-2\} \text{ (rows only)},$$
$$Y' = \{r+1, r+3, \dots, r+2m-1\} \text{ (columns only)},$$

so that $R = C \sqcup X'$ and $Q = C \sqcup Y'$ (a second "member" adds one more common node $r + 2m$). The determinant $P(x) = \det \Phi(x)$ still has degree $\leq r$ (or $r+1$), so the integrality loss $|R \cap Q| \log q$ is that of r nodes, but the matrix has size $r+m$ and its decay is that of a configuration of $(1+\alpha)r$ nodes, $\alpha = m/r$. The content grows too, but less: the extra rows and columns never meet the diagonal. With $\alpha = \frac{1}{8}$ the content and decay constants become (for λ)

$$\Lambda = 3.67885 \quad \text{(square frame: } 313/105 = 2.98095\text{)},$$
$$D = 3.90557 \quad \text{(square frame: } 3.1114\text{)},$$

and the margin $\eta = D - \Lambda$ rises from 0.1304 to 0.22672. The interleaving of X' and Y' is essential: it keeps the two Cauchy determinants in Andréief's formula comparable (§11) and it gives the reduction modulo ℓ the bordered structure of §8.

2.3. **New idea 2: content by p-adic assignment problems and rank deficits.** For a prime p let v be the Gauss valuation on $\mathbb{Q}[x]$ (the minimum of the p-adic valuations of the coefficients); it is multiplicative. Expanding $\det \Phi(x)$ by Leibniz, $-v(\det \Phi)$ is at most the value τ_p of the *assignment problem* with weights $W_{ab} = \max(0, -v_p(M[a,b]))$, i.e. the maximum of $\sum_a W_{a\sigma(a)}$ over bijections $\sigma : R \to Q$. By LP duality, any *potentials* u_a, v_b with $W_{ab} \leq u_a + v_b$ certify $\tau_p \leq \sum u_a + \sum v_b$. We give explicit potentials for every prime power p^e (they depend only on p^e/r and on $p \bmod 4$ or $p \bmod 6$), and sum over primes with the prime number theorem in arithmetic progressions. This gives a "tropical" content $\Lambda_{\text{trop}} = 3.74993$ for λ.

The tropical bound is not sharp. If (u, v) is an optimal dual solution, the scaled matrix $B = p^u \Phi p^v$ has entries in the valuation ring $\mathcal{O}$ of v, and $v(\det \Phi) = -\tau_p + v(\det B) \geq -\tau_p + \operatorname{corank}(B \bmod p)$. So every unit of corank of the reduction of B modulo p over $\mathbb{F}_p(x)$ saves one factor of p. We compute this corank in several ranges of primes, using the block structure of the reduction (residue classes modulo p), Schur complements,

the identity $H_{p-1-n} \equiv H_n \pmod p$ for harmonic numbers, an antisymmetry of Cauchy-type sums, and the Frobenius–König theorem. The savings lower the content from 3.74993 to 3.70343. The uniform rescaled-core argument of Section 5 adds the missing interval savings without changing the determinant family, giving content 3.68274. Finally, Section 6 shows that at primes where the leading block is not clean the saving has a second level: the next Smith coefficient is a weighted Cauchy-type matrix of small rank, which adds further factors of p. This lowers the content to 3.67885.

2.4. **New idea 3: nonvanishing through a commuting Jacobi matrix.** Fix a prime ℓ in a suitable residue class with $\ell \nmid q$. After scaling by ℓ the rows and columns of nodes beyond a threshold $K \approx \ell/4$, the matrix $\Phi(x)$ reduces modulo ℓ to a *bordered Cauchy–Gram* matrix:

$$\ell^{2h} \det \Phi(x) \equiv \text{unit} \times \det \begin{pmatrix} yI_K + N_S & W_Y \\ W_X^\top & 0 \end{pmatrix} \pmod \ell,$$
$$y = \text{affine function of } x,$$

where $N_S = \sum_{j=1}^{S} c_j c_j^\top$ is a sum of rank-one Gram terms of Cauchy vectors $c_j = (1/(a+j))_{0 \le a < K}$ and W_X, W_Y consist of further Cauchy vectors. The two members of the pair differ by one Gram term. Their reductions cannot vanish simultaneously unless a *bordered Krylov determinant*

$$k = \det\big[W_Y \,\big|\, v,\ Av,\ \ldots,\ A^{K-m-1}v\big], \qquad A = \Pi N_S, \quad v = \Pi c_S,$$

vanishes modulo ℓ, where Π is the projection onto $\ker W_X^\top$ along the span of W_Y. We prove that, with the nodes shifted by a parameter θ, $k(\theta)$ is a constant prime to ℓ times a product of powers of $\theta + c$ with $1 \le c < \ell$. The proof is algebraic and uses a structure familiar from orthogonal polynomials: there is an explicit *Jacobi (tridiagonal) matrix* $\tilde{T}$ that commutes with Π and with A and acts on the Cauchy vectors by a three-term recurrence (a bispectral situation in the sense of Grünbaum and Haine [15]). Its LU factorisation, reversed to UL, is (up to a shift and a diagonal conjugation) the Jacobi matrix of the next smaller instance: a discrete Darboux, or Christoffel, transformation [38, 16]. Krylov determinants of commuting pairs satisfy a cross-ratio identity, chains of Cauchy vectors give products of Cauchy determinants, and the remaining Gram determinants are evaluated with Cauchy–Binet, Hankel determinants of moments, Newton's divided differences and the Chu–Vandermonde identity (which also proves the Pfaff–Saalschütz-type identity needed). One auxiliary determinant has no product formula and is shown to be nonzero by positivity. An induction on the size closes. We never need the exact exponents: it suffices to know that the determinant is an ℓ-adic unit times powers of $\theta + c$ with $1 \le c < \ell$.

Because the pair of members is available for every S in a range, the prime ℓ can be chosen *after* the size r, anywhere in an interval $[\frac{3}{2}r, 4r]$. This removes the cost of locating ℓ from the exponent (for λ and λ'), and the final exponent is simply c/η.

2.5. **New idea 4: decay through an energy certificate.** Moving Hermite's contour to two rays in the left half-plane writes $\omega\, \mathsf{L}_F[R, Q]$ as a Gram matrix $\int v_a v_b\, dm$ of a complex measure m (finitely many signed atoms at poles of F plus an exponentially small density on the rays). Andréief's identity with two families of functions (rows and columns) gives

$$|\det \mathsf{L}_F[R, Q]| \le \frac{1}{n!} \int |\mathsf{C}_R(z)|\, |\mathsf{C}_Q(z)| \prod_{i=1}^{n} d|m|(z_i),$$

where $\mathsf{C}_R, \mathsf{C}_Q$ are Cauchy determinants. Taking logarithms turns the integrand into a discrete logarithmic energy of the free points z_i against the nodes in an external field. We bound this energy from below by comparing with a *certificate measure* σ (point masses with dyadic coordinates, placed where an optimal dual measure is supported) through a regularised kernel $-\frac{1}{2}\log(|z-w|^2+\rho^2)$, which is conditionally positive definite on finite sums by an elementary argument (Frullani's integral and Gaussian sums). Everything left is a finite list of inequalities between explicit functions on boxes, which we verify in interval arithmetic inside Lean. For λ and λ' the contour rays have slope $\frac{5}{2}$ and $\rho = 1/4096$ (Section 11.4). The same argument bounds every principal minor along the common block, so the determinant stays small at every point within e^{-cr} of ξ.

2.6. **The deduction.** Write $\eta = D - \Lambda$. The three estimates are combined in §12; the result is

$$\mu(\xi) \le c\Big(\frac{1}{\eta} + g\, s\Big),$$

where c is the window constant of the decay estimate, $g = \lim r/\ell$ for the sizes at which nonvanishing is proved, and s is the relative length of the interval in which ℓ must be found. For λ and λ' the size can be chosen independently of ℓ, so the term gs disappears and $\mu \le 7.3546/0.22672 < 32.44$. For $L(2, \chi_{-3})$, the original pair family gives $6.6510\,(1/0.35954 + 0.8) < 23.82$. Section 9 replaces it by a family with at most three common low-node deletions. This permits a fixed scale across a prime interval and needs only $\ell^2 \nmid q$; the term gs disappears, giving $\mu \le 6.6510/(3.7251 - 2159207/645120) < 17.60$. Section 10 then permits a quarter border through characteristic-zero selection, supplies an analytic certificate and retains a periodic D1 saving, giving $\mu \le (93657/12500)/(1158407/250000 - \Lambda_{16}) < 14.35$, with $\Lambda_{16} = 339258406689121/82516315939200$.

constant	content Λ	decay D	window c	$\eta = D - \Lambda$	gs	exponent
$G - 5\pi^2/48$	3.67885	3.90557	7.3546	0.22672	0	32.4392
$G + 5\pi^2/48$	3.67885	3.90557	7.3546	0.22672	0	32.4392
$L(2, \chi_{-3})$	Λ_{16}	4.633628	7.49256	0.535196	0	14.997127

2.7. **Formal verification.** Every step, including the arithmetic of the dual certificates, the closed-form evaluations behind the nonvanishing, the interval-arithmetic decay certificates and the prime number theorem in

progressions modulo 4 and 6, is formalised in Lean 4 with Mathlib [9, 40]. The formal proofs are built on a shared core (rectangular determinants, Gauss valuations and assignment bounds, bordered Krylov determinants, energy certificates, the measure deduction) with thin instances for the three functions in (2.1). The written proofs below are at the level of a research article; for long computations we point to the formal proof, which is the basis for the correctness of Theorem 1.1.

3. The construction

3.1. **The kernels.** For the three functions of (2.1) the kernel M of (2.2) is as follows; here $0 \le a < b$ are integers and $M[b,a] = M[a,b]$.

$$\begin{aligned}
\lambda: \quad M[a,b] &= -\frac{1}{b-a}\sum_{k=a+1}^{b}\frac{1}{16k(4k-1)},\\
M[a,a] &= -\sum_{k=1}^{a}\Big(\frac{1}{16k^2}-\frac{1}{(4k-1)^2}\Big);\\
\lambda': \quad M[a,b] &= \frac{1}{b-a}\sum_{k=a+1}^{b}\frac{3}{16k(4k-3)},\\
M[a,a] &= -\sum_{k=1}^{a}\Big(\frac{1}{(4k-3)^2}-\frac{1}{16k^2}\Big);\\
\chi: \quad M[a,b] &= \frac{1}{6(b-a)}\sum_{k=a+1}^{b}d(k),\\
M[a,a] &= -\sum_{k=1}^{a}\delta(k),
\end{aligned}$$

with $d(k) = \frac{1}{3k-2}-\frac{1}{2k-1}+\frac{1}{6k}$ and $\delta(k) = \frac{1}{2(3k-2)^2}-\frac{1}{3(2k-1)^2}+\frac{1}{36k^2} = -\frac{1}{6}d'(k)$.

Lemma 3.1. *For each of the three functions, $\omega \llcorner_F [R,Q] = M[R,Q]+\xi E_{R,Q}$ for all finite sets $R, Q \subset \mathbb{Z}_{\ge 0}$, with ω and ξ as in (2.1).*

Proof. From $\psi(w+1)-\psi(w) = 1/w$ one gets $F(k)-F(k-1)$ as an explicit rational function of k; for instance $F_\lambda(k)-F_\lambda(k-1) = \frac{1}{k}-\frac{1}{k-1/4} = -\frac{1}{k(4k-1)}$, and telescoping gives the off-diagonal entries. Similarly $\psi'(w+1)-\psi'(w) = -1/w^2$ gives $F'(a)-F'(0)$. Finally $\omega\,F'(0) = \xi$: for λ, $\frac{1}{16}(\psi'(1)-\psi'(\frac{3}{4})) = \frac{1}{16}(\frac{\pi^2}{6}-\pi^2+8G) = \lambda$; for λ' use $\psi'(\frac{1}{4}) = \pi^2+8G$; for χ use $\psi'(\frac{1}{2}) = \frac{\pi^2}{2}$ and $\psi'(\frac{1}{3}) = \frac{2\pi^2}{3}+\frac{9}{2}L(2,\chi_{-3})$, which give $-\frac{1}{6}F'_\chi(0) = \sum_{k\ge1}\delta(k) = \frac{1}{4}L(2,\chi_{-3})$. □

Note that $2\lambda = G-5\pi^2/48$, $2\lambda' = G+5\pi^2/48$ and $4\xi_\chi = L(2,\chi_{-3})$, and $\mu(a\xi+b) = \mu(\xi)$ for rational $a \ne 0$, b. So Theorem 1.1 is a statement about the three numbers ξ.

3.2. **Geometries.**

Definition 3.2. Let $r, m \geq 0$ and $e \in \{0,1\}$. The *geometry* $\mathcal{G}(r,m,e)$ consists of the common nodes $C = [0,r) \cup \{r+2m : \ e=1\}$, the row nodes $X' = \{r+2i : 0 \leq i < m\}$ and the column nodes $Y' = \{r+2i+1 : 0 \leq i < m\}$. Rows are indexed by $R = C \sqcup X'$, columns by $Q = C \sqcup Y'$. The *member* $e=0$ is $\mathcal{G}_1(r,m)$, the member $e=1$ is $\mathcal{G}_2(r,m)$. The matrix of the geometry is

$$\Phi(x) = \Phi_{\mathcal{G}}(x) = \big(M[a,b] + x\,\mathbf{1}[a=b\in C]\big)_{a\in R,\ b\in Q},$$

a square matrix of size $|C|+m$, and $P_{\mathcal{G}}(x) = \det \Phi_{\mathcal{G}}(x) \in \mathbb{Q}[x]$.

By Lemma 3.1, $\Phi_{\mathcal{G}}(\xi+t) = \omega\, \mathsf{L}_F[R,Q] + t\, E_{R,Q}$ for real t. The geometry with $m=0$, $e=0$ is the square Löwner matrix of size r. We use *admissible* geometries, with $m \approx r/8$: for λ, λ' the condition is $r \leq 8m \leq r+8$, and for $L(2,\chi_{-3})$ it is $r \leq 8m \leq r+9$ (the extra unit is needed to find a pair of members for every prime $\ell \equiv 1 \pmod 6$; see §12). All $o(1)$ terms below are uniform over admissible geometries as $r \to \infty$.

Lemma 3.3 (integrality)**.** *$P_{\mathcal{G}}$ has degree at most $|C|$. If $d_{\mathcal{G}} = \operatorname{den} P_{\mathcal{G}}$ and p/q is a fraction in lowest terms with $P_{\mathcal{G}}(p/q) \neq 0$, then $|d_{\mathcal{G}} P_{\mathcal{G}}(p/q)| \geq q^{-|C|}$.*

Proof. The variable occurs only in the $|C|$ diagonal entries indexed by C, so each Leibniz term has degree at most $|C|$. Then $q^{|C|} d_{\mathcal{G}} P_{\mathcal{G}}(p/q)$ is a nonzero integer. □

Remark 3.4. When the $m \times m$ block $M[X',Y']$ is invertible, the Schur complement gives $P_{\mathcal{G}}(x) = \det M[X',Y'] \cdot \det(xI_C + M_{\mathrm{S}})$ with $M_{\mathrm{S}} = M[C,C] - M[C,Y']M[X',Y']^{-1}M[X',C]$. Thus the hybrid determinant is, up to a constant, the characteristic polynomial of a rational matrix of size $|C|$, as in the square case. We do not need this factorisation: Lemma 3.3 only uses the degree bound.

4. Arithmetic: the content

In this section we prove the following.

Theorem 4.1 (content)**.** *For admissible geometries of scale r,*

$$\log \operatorname{den} P_{\mathcal{G}} \leq (\Lambda + o(1))\, r^2,$$

where $\Lambda = 3.67885$ for λ and λ', and $\Lambda = 2159207/645120 = 3.346985\ldots$ for $L(2,\chi_{-3})$.

We give the details for λ and indicate the changes for λ' and for $L(2,\chi_{-3})$ at the end.

4.1. **Gauss valuations and assignment problems.** Fix a prime p. Let $\mathcal{O} \subset \mathbb{Q}(x)$ be the localisation of $\mathbb{Z}_{(p)}[x]$ at the prime ideal (p); it is a discrete valuation ring with uniformiser p and residue field $\mathbb{F}_p(x)$, and its valuation v restricted to $\mathbb{Q}[x]$ is the *Gauss valuation* $v(f) = \min_k v_p(f_k)$, $f = \sum f_k x^k$. By Gauss's lemma $v(fg) = v(f) + v(g)$, and $v_p(\operatorname{den} f) = \max(0, -v(f))$.

Lemma 4.2 (assignment bound and LP duality)**.** *Let B be a square matrix over $\mathbb{Q}[x]$ with rows I and columns J, $W_{ab} = \max(0, -v(B_{ab}))$, and let $\tau(W) = \max_\sigma \sum_{a\in I} W_{a\sigma(a)}$ over bijections $\sigma : I \to J$. Then $v(\det B) \geq -\tau(W)$. If real numbers u_a, v_b satisfy $W_{ab} \leq u_a + v_b$ for all a, b, then $\tau(W) \leq \sum_a u_a + \sum_b v_b$.*

Proof. Each Leibniz term has valuation $\sum_a v(B_{a\sigma(a)}) \geq -\sum_a W_{a\sigma(a)}$, and v of a sum is at least the minimum. The second statement is weak LP duality for the assignment problem: $\sum_a W_{a\sigma(a)} \leq \sum_a (u_a + v_{\sigma(a)})$. □

For $\Phi(x)$ the weight of every entry is $W_{ab}(p) = \max(0, -v_p(M[a, b]))$, since $v(x) = 0$. We call τ_p the value of this assignment problem; thus $v_p(\operatorname{den} P_{\mathcal{G}}) \leq \tau_p$.

4.2. **The level majorant.** Let $p \geq 5$ and, for a prime power $q = p^e$, let $S_q = \{k \geq 1 : q \mid k(4k-1)\}$; it consists of the classes $k \equiv 0$ and $k \equiv k_q \pmod q$, where $k_q = (j_q q + 1)/4$ with $j_q \in \{1, 3\}$, $j_q q \equiv 3 \pmod 4$. Let $\beta_q(a) = \#(S_q \cap [1, a])$. The ultrametric inequality applied to the formulas for M gives

$$W_{ab}(p) \leq \sum_{e\geq 1} w^{(p^e)}(a, b),$$
$$w^{(q)}(a, b) = \mathbf{1}[\beta_q(a) \neq \beta_q(b)] + \mathbf{1}[q \mid a - b] \ (a \neq b),$$
$$w^{(q)}(c, c) = 2 \cdot \mathbf{1}[c \geq k_q],$$

and $w^{(q)} \in \{0, 1, 2\}$. Since the maximum of a sum is at most the sum of the maxima, $\tau_p \leq \sum_e \tau(w^{(p^e)})$. The primes 2 and 3 and all levels $e \geq 2$ contribute $o(r^2)$ in total (each $w^{(q)}$ has assignment value $O(r)$, there are $O(\log r)$ levels for each of 2, 3, and $\sum_{p^e \leq Cr,\, e\geq 2} \log p = O(\sqrt{r})$).

4.3. **Explicit dual certificates.** Fix an odd prime power q, write $k = k_q$, $N = r + 2m$, and let

$$\delta_A(q) = \min\big(2r,\ 2(r - n) + 2\min(n, (r - q)_+)\big), \qquad n = \min(r, k),$$

the value of the square block $C \times C$. Let $U = [r-q, k)$ (empty if $r-q \geq k$) be the set of "low unpaired" indices; a zone node $z \in X' \cup Y'$ is *low* if $z - q \in U$, and L is the number of low zone nodes.

Proposition 4.3 (dual certificates)**.** *For $\mathcal{G}(r, m, 0)$ with $8m \leq r$, $\tau(w^{(q)}) \leq \delta_A(q) + E(q)$ with*

range	$E(q)$	*potentials*
$q < 2m$	$2m - q\,\|m/q\|$	*by residue class mod* q
$2m \le q \le r$	$\min(2m + L,\ m + L + \|U\|)$	*two certificates*
$r < q < N$	$2m + (N - q)$	$\mathbf{1}[c \ge k]$ *on* C, $1 + \mathbf{1}[z \ge q]$ *on the zone*
$q \ge N$, $k \le r$	$2m$	$\mathbf{1}[c \ge k]$ *on* C, 1 *on the zone*
$q \ge N$, $k > r$	$(N - k)_+$	0 *on* C, $\mathbf{1}[z \ge k]$ *on the zone*

($\|\cdot\|$ is the distance to the nearest integer.)

Proof sketch. In each range one writes down potentials and checks $w \le u+v$ entrywise, using two facts: two distinct indices not congruent modulo q have $w \le 1$, and two zone nodes are never congruent once $q \ge 2m$. For example, for $q \ge N$ and $k \le r$, give the common nodes the potential $\mathbf{1}[c \ge k]$ and the zone nodes the potential 1: there are no congruences, so $w \le 1$ off the diagonal, and the diagonal cells (only at $c \in C$) are covered. For $q < 2m$ the potentials are $\frac{1}{2}, \frac{3}{2}, 1$ on rows and $\frac{3}{2}, \frac{1}{2}, 1$ on columns of a residue class ϱ, according as X' has more, fewer or as many nodes in ϱ as Y'; since the zone alternates between X' and Y', the imbalance is at most one per class and the number of unbalanced classes is $2q\|m/q\|$. The certificates for $2m \le q \le r$ start from the square-frame potential on C and raise the potential of low zone nodes (first certificate), or move half a unit between rows and columns in the residue classes of high zone nodes (second certificate). All cases are checked in the formal proof. □

Remark 4.4. The certificates are sharp: the exact assignment value computed by the Hungarian algorithm agrees with $\delta_A + E$ for all odd q and all tested (r, m). Sharpness is not used.

4.4. **The tropical content.** Put $\theta = q/r$, $\alpha = m/r$. Proposition 4.3 gives $\tau(w^{(q)}) \le r\, f_j(\theta; \alpha) + O(1)$ for an explicit continuous piecewise linear function f_j supported in $[0, 4(1 + 2\alpha)/j]$. Summing over primes in the two classes modulo 4 with the prime number theorem in arithmetic progressions, $\frac{1}{r}\sum_{p \equiv a\,(4)} f(p/r)\log p \to \frac{1}{2}\int_0^\infty f$ for bounded Riemann-integrable f of compact support (see [8, Ch. 20] for the prime number theorem in progressions), one finds

$$\sum_p \tau_p \log p \le (\Lambda_{\mathrm{trop}}(\alpha) + o(1))r^2,$$

$$\Lambda_{\mathrm{trop}}(\alpha) = \frac{313}{105} + \frac{554}{105}\alpha + \Big(\frac{118}{21} + 2\log 2\Big)\alpha^2$$

for $\alpha \le \frac{1}{8}$ (the term $2\alpha^2 \log 2$ comes from the sawtooth $q\|m/q\|$ in the range $q < 2m$). At $\alpha = \frac{1}{8}$, $\Lambda_{\mathrm{trop}} = 3.7499347\ldots$; for $\alpha = 0$ it is the square-frame value $313/105$.

4.5. **Rank deficits.** The assignment bound is sharp for the matrix of weights, but not for $\det\Phi$: cancellation in the Leibniz expansion makes $\operatorname{den} P$ smaller. We capture it by the following principle.

Lemma 4.5 (rank deficit)**.** *Let (u, v) be integer-valued potentials feasible for the weights of Φ at p, of value τ'. Then $B = \operatorname{diag}(p^u)\,\Phi\,\operatorname{diag}(p^v)$ has entries in $\mathcal{O}$, and*

$$v_p(\operatorname{den} P_{\mathcal{G}}) \le \tau' - \operatorname{corank}_{\mathbb{F}_{p(x)}}(B \bmod p).$$

Proof. $v(\det\Phi) = -\tau' + v(\det B)$, and over the discrete valuation ring $\mathcal{O}$ the Smith normal form shows $v(\det B) \ge \operatorname{corank}(B \bmod p)$. (The formal proof avoids the Smith form: if $B \bmod p$ has rank ρ, an explicit rank witness and a change of basis over $\mathcal{O}$ exhibit $|I| - \rho$ rows divisible by p.) □

Write δ_p for the corank obtained in this way with an optimal dual (so $v_p(\operatorname{den} P) \le \tau_p - \delta_p$). We use five families of primes. In all of them $p^2 > 4N$, so only the level $e = 1$ matters; the reduction $B \bmod p$ has a block structure governed by the singular indices $k \in S_p$ below N, which cut $[0, N)$ into blocks on which the leading coefficients (lead_{-1} of the partial sums of $1/(4k(4k-1))$ and lead_{-2} of the diagonal sums) are constant.

Theorem 4.6 (rank deficits for λ)**.** *Let $\mathcal{G} = \mathcal{G}(r, m, e)$ be admissible and r large. Then $v_p(\operatorname{den} P_{\mathcal{G}}) \le \tau_p - \delta_p + O(1)$, with δ_p as follows, for all primes p except $p \in \{139, 1009\}$:*

primes	δ_p	*mechanism*
$p \equiv 3\ (4)$, $\frac{5}{8}r < p < \frac{4}{5}r$; $p \equiv 1\ (4)$, $\frac{5}{7}r < p < r$	m	*integral Schur complement*
$r < p < r + m$	$r + m - p$	*harmonic reflection, rank count*
$p \equiv 3\ (4)$, $\frac{4}{5}(r+m) \le p < r$	$\frac{5p+1}{4} - r - m$	*Frobenius–König*
$p \equiv 1\ (4)$, $\frac{4}{7}(r+m) \le p < \frac{5}{7}r$	$\frac{7p+1}{4} - r - m$	*Frobenius–König*
$p \equiv 3\ (4)$, $\frac{5}{12}r < p < \frac{4}{9}r$; $p \equiv 1\ (4)$, $\frac{5}{11}r \le p < \frac{1}{2}r$	m	*block-diagonal leading form*

We sketch the mechanisms; the complete proofs are formalised.

The integral Schur complement. Write $\Phi = \left(\begin{smallmatrix} A & \Phi[C,Y'] \\ \Phi[X',C] & \Phi[X',Y'] \end{smallmatrix}\right)$ with $A = \Phi[C, C]$, so $\det\Phi = \det A \cdot \det T$ with the Schur complement $T = \Phi[X', Y'] - \Phi[X', C]A^{-1}\Phi[C, Y']$. At these primes the square block has $-v(\det A) = 2r$, and the claim is that T has entries in $\mathcal{O}$, whereas the assignment bound allows $v(T_{zy}) = -1$. For a zone row z with partner $c_z = z - p \in C$ decompose

$\Phi[z, C] = A[c_z, C] + \ell_z$, and similarly for columns; since $A[c_z, \cdot]A^{-1} = e_{c_z}^\top$,

$$T_{zy} = \Delta^2 M(z, y) - \ell_z A^{-1} \ell_y^\top,$$
$$\Delta^2 M(z, y) = M(z, y) - M(z, c_y) - M(c_z, y) + M(c_z, c_y),$$

and x cancels. The leading form of $p^2 A$ is block diagonal by residue class (blocks of size 1 and 2), with explicit nonzero determinants (here $p \neq 139$ is used: one block has determinant $3^2 \cdot 5 \cdot 139/6400$). Expanding $\ell_z A^{-1} \ell_y^\top$ modulo p^2, the only surviving terms are $p\beta\big[\frac{\sigma}{4(c_z - c_y)} + \frac{\sigma}{4(c_y - c_z)}\big] = 0$, an antisymmetry of Cauchy-type sums. For $p \equiv 1 \pmod 4$ the analogous computation uses a scalar Schur complement: modulo p it equals $s_0 I$ with $s_0 = x - \frac{1}{16} H^{(2)}_{K_p - 1}$, $K_p = (3p+1)/4$, which follows from partial fractions and the reflections $H_{p-1-n} \equiv H_n$, $H^{(2)}_{p-1-n} \equiv -H^{(2)}_n \pmod p$ for harmonic numbers. Here it is essential that x is an indeterminate: s_0 is a unit of $\mathcal{O}$, although it may vanish at a specialisation $x \in \mathbb{Q}$.

Primes just above r. For $r < p < r + m$ the only singular indices below N are k_p and p. After eliminating an invertible block, the Schur complement on $[0, k_p)$ is $s_0 I - \frac{1}{16} E E^\top$ with $E = (1/(u+i))_{u < k_p,\, 1 \le i \le p - r}$, by the same harmonic reflection. Each zone node $z < p$ contributes one rank-one term indexed by $i = p - z$, so the correction has rank at most $p - r$ and the corank is at least $r + m - p$.

Primes with one singular zone index. For $p \equiv 3 \pmod 4$ with $\frac{4}{5}(r+m) \le p < r$ (and $p \equiv 1 \pmod 4$ in the corresponding range) there is one singular index inside the zone. After elimination the zone×zone block of $B \bmod p$ has a zero sub-block of size $L_X \times L_Y$ with $L_X + L_Y = L \ge m$ (the low zone nodes), so by the Frobenius–König theorem its rank is at most $2m - L$, and the corank is at least $L - m$.

The low band. For primes p with $2m < p < r$ that are *clean* (no singular index in the zone) we prove a general statement: if every residue-class block of the leading form of $p^2 A$ is invertible modulo p, then T is integral. The proof splits $p^2 A = B_0 + pE$ with B_0 block diagonal by residue class, inverts by a Neumann series, and shows that the order-p term vanishes: the translation $c \mapsto c + (y - z)$ identifies the classes of z and y (cleanness is used here), and what remains is $\sum_{i,j} g_i g_j (L_i - L_j) = 0$, an antisymmetric sum. The original proof for the two listed families used classes of size 2 or 3 and explicit rational block determinants, for example $-350972849/368640000$ (with prime numerator), which are nonzero modulo p except at $p = 139$ and $p = 1009$. The uniform argument in Section 5 proves the needed clean-case bound without these exceptional determinants.

4.6. **Summation.** Each family is a union of intervals $p \in (ar, br)$ in a fixed class, and δ_p is a linear function of p there. With the prime number theorem in progressions, the total saving is

$$\Big(\frac{129}{4480} + \frac{59}{4480} + \frac{29}{6336}\Big) r^2 = (0.0287946 + 0.0131696 + 0.0045770)\, r^2$$

(the three terms come from the first row, the next three rows, and the last row of the table). Hence

$$\Lambda_\lambda \le \Lambda_{\mathrm{trop}}(\tfrac{1}{8}) - 0.0465412 = 3.7033935\ldots \le 3.70343.$$

(The formal proof uses the slightly coarser rational value 3.7034204, from a truncation of the sawtooth term.) This is the intermediate content constant before the additional savings of Section 5, which complete the improved constant in Theorem 4.1.

4.7. **The second member.** The geometry $\mathcal{G}_2$ has one extra common node $b = r + 2m$. Bordering $\Phi_{\mathcal{G}_1}$ by one row and one column changes each per-prime bound by $O(1)$, so the content constant is unchanged; but the rank deficits have to be re-proved for the bordered matrix, since a deficit is not automatically inherited under bordering. For the deficits proved by rank witnesses there is a general bordering lemma: a witness for $\Phi_{\mathcal{G}_1}$ extends, with the border row and column scaled by p^w, to a witness for $\Phi_{\mathcal{G}_2}$ at a cost of $2w = O(1)$. For the low-band deficit, which is proved through integrality of the Schur complement, the border row and column are rescaled by p^w so that the same core argument applies, with one unit of slack.

4.8. **The mirror constant and $L(2, \chi_{-3})$.** For λ' the singular indices are those k with $p \mid k(4k-3)$; the two residue classes modulo 4 exchange their roles. The tropical law is the same function Λ_{trop}, the deficits are the same with the classes swapped, and $\Lambda_{\lambda'} = \Lambda_\lambda$. The formal proof uses one generic content theorem with the offset (1 or 3) as a parameter.

For $L(2, \chi_{-3})$ the singular indices are those k with $p \mid k(3k-2)(2k-1)$, and the relevant classes of primes are modulo 6. Before the periodic improvement of Section 7, two explicit duals suffice: below a threshold θ^* ($\frac{27}{32}$ for $p \equiv 1$, $\frac{3}{4}$ for $p \equiv 5 \pmod 6$) the residue-class potentials of the first row of Proposition 4.3, and above θ^* a certificate in which only row potentials are raised, on nodes that have a congruent partner of level 0. The levels (leading coefficients of the partial sums) enter only through which of them vanish, and this reduces to eight explicit identities. The resulting bound is piecewise linear in p/r, with $3+6$ and $3+5$ pieces in the two classes, and integrating gives $\Lambda_{\chi,\mathrm{base}} = \frac{10339}{3072}$. The per-prime bound agrees with the exact assignment value at all primes tested ($r = 256$ and $r = 1024$).

5. A uniform rescaled-core deficit

We retain the determinant family and the previously established clean-prime estimates. A common rescaled-core argument supplies three additional, disjoint groups of prime windows. The gain comes from a first-order cancellation and an elementary assignment dual; no assertion about a Smith normal form or the optimality of the previous tropical bound is required. The argument applies to both offsets $o \in \{1, 3\}$.

Monic rescaling. For the finite certificate take $m = \lfloor r/8 \rfloor$; the same leading estimates hold for $m = r/8 + O(1)$, since all endpoint and border losses are bounded per prime. Put $N = r + 2m$ and write $C = [0, r)$ for the common core and $Z = [r, N)$ for the interleaved border zone. We use primes $p \geq 13$ satisfying $4(N + 1) < p^2$. In the normalized symmetric rational matrix R, whose omitted normalization factors are p-adic units, every entry has valuation at least -2. Replace the polynomial variable X by x/p^2 and set

$$A = R_{CC} + \frac{x}{p^2} I_r, \qquad Q = p^2 A = p^2 R_{CC} + x I_r.$$

The entries of Q are integral for the p-adic Gauss valuation. Its reduction is block diagonal according to residues of core nodes modulo p. Every reduced block has the form $M + xI$, so its determinant is monic and nonzero over $\mathbb{F}_p(x)$. Consequently

$$v_G(\det Q) = 0, \qquad v_G(\det A) = -2r.$$

This remains valid when the constant matrix M is singular. The single Gauss unit $\det Q$ and its adjugate suffice to express all inverse entries; separate denominators for individual residue blocks are unnecessary.

Let T be the Schur matrix left after eliminating C. If every assignment in its $m \times m$ zone matrix has total entry price at most τ, with $v_G(T_{zy}) \geq -w_{zy}$, then the determinant expansion gives

$$v_G(F(x/p^2)) \geq -2r - \tau.$$

Indeed, for $F(X) = \sum_i c_i X^i$ the left side is $\min_i(v_p(c_i) - 2i)$. Thus every coefficient satisfies $v_p(c_i) \geq -2r - \tau + 2i \geq -2r - \tau$. The substitution estimates coefficients of the original polynomial and introduces no new approximation family.

Translation and cancellation. The singular cuts are the integers in the two residue classes 0 and $(cp + o)/4$ modulo p, where $c = 1$ or 3 is determined by the prime class. For a zone node z, its core chain consists of

$$\{z - ip : i \geq 1,\ 0 \leq z - ip < r\}.$$

Between consecutive singular cuts and the core-exit cuts $r + ip$, translation by $y - z$ identifies the chains of z and y, their leading node data, the leading core–zone vectors, and their symmetric leading chain matrices. For the one-interval construction below, clipping the right endpoint at $r + p$ controls the core-exit cut. The formal diagonal xI is preserved by the same identification.

For completeness, write $Q = B + pE$, where B retains the full same-residue blocks of Q. Modulo p^2,

$$Q^{-1} = B^{-1} - pB^{-1}EB^{-1}.$$

For two distinct residues the order -2 Schur term vanishes. For two nodes in one translated class, let g be the common leading inverse vector and L

the leading node datum. Up to one common normalization, the three order -1 correction sums are

$$\frac{\gamma}{y-z}\sum_{u} g_u(L(z)-L(u)),$$
$$\frac{\gamma}{y-z}\sum_{u} g_u(L(u)-L(z)),$$
$$-\frac{\gamma}{y-z}\sum_{u,v} g_u g_v(L(u)-L(v)).$$

The first two cancel termwise, and the last vanishes by interchanging u, v. The direct order -1 entry also vanishes since $L(y) = L(z)$. This proves integrality within the selected class. Between distinct residues the price is at most 1; equal residues have price at most 2. No specialization at $x = 0$ or additional spectral hypothesis enters this calculation.

One interval and a $0/1/2$ dual. Choose consecutive singular cut indices $A_4 < B_4$, with no permitted numerator strictly between them, and put

$$a = \max\left(r, \left\lfloor \frac{A_4 p + \epsilon_{A_4}}{4} \right\rfloor\right),$$
$$b = \min\left(N, r+p, \left\lfloor \frac{B_4 p + \epsilon_{B_4}}{4} \right\rfloor\right),$$
$$\epsilon_j = \begin{cases} 0 & j \equiv 0 \pmod 4, \\ o & \text{otherwise.} \end{cases}$$

In the high range $p > 2m$ the clipping at $r + p$ is redundant. All nodes of $[a, b)$ lie in one translated class and have distinct residues. In the low range $m < p < 2m$, use the smaller interval

$$J = [\max(a, N-p), b),$$

interpreting its length as zero when its left endpoint exceeds its right endpoint. Since $b \le r + p$, a node in J has no other zone node of the same residue. In the high range take $J = [a, b)$. Assign potential 0 to every row or column node in J, and potential 1 to every other zone node. Two nodes in J have integral Schur entry, a mixed pair has price at most 1, and an outside pair has price at most 2. Therefore this is a feasible dual and

$$\tau \le 2m - |J|.$$

This elementary certificate replaces class-count and parity calculations.

The second member adds one row and column to the remaining Schur matrix. Every new incident entry, including its rescaled polynomial corner, has price at most 2. An assignment uses at most two such incidences; extending the remaining partial assignment gives a total additional price at most 4. Thus both members satisfy the common coefficient bound

$$\operatorname{denexp}_p F_i \le 2r + 2m - |J| + 4 \qquad (i = 1, 2).$$

The constant 4 is retained in the integer deficit, rather than silently discarded before summation.

Baseline and finite profile. Let T_{old} be the exponent used for the already banked estimates. Only on the new profile windows enlarge it to

$$T_* = \max(T_{\mathrm{old}}, B_0), \qquad B_0 = \begin{cases} 2r + 3m - p, & m < p < 2m, \\ 2r + m, & p > 2m. \end{cases}$$

The existing affine majorants already cover these values with their existing constant slack; their leading integral is unchanged. This comparison requires no equality between B_0 and a tropical optimum. The resulting nonnegative integer deficits are

$$\delta_p = \begin{cases} (m + |J| - p - 4)_+, & m < p < 2m, \\ (|J| - m - 4)_+, & p > 2m. \end{cases}$$

Outside the new windows set the new deficit to zero. Retain all old E, B, C5, C7, and clean GK estimates: this rescaling is not asserted to reproduce their sharper savings.

Here is a compact specification of the complete rational certificate. Write $u = p/r$, $A = A_4/4$, $B = B_4/4$. In the low range put

$$a_0 = \max(1, Au), \qquad b_0 = \min(5/4, Bu, 1 + u),$$
$$f(u) \le \min(1/8 - u + b_0 - a_0,\ b_0 - 9/8).$$

In the high range the corresponding certificate is

$$f(u) \le \min(5/4, Bu) - \max(1, Au) - 1/8.$$

For an affine f, each assertion reduces to finitely many affine endpoint inequalities: nine in the low case and four in the high case. The certificate contains 37 low rows, 20 nonclean rows below $u = 3/5$, and four missing upper rows. In addition to the affine inequalities, it checks consecutive pole numerators, the residue class, nonnegativity, and pairwise disjointness within each prime class.

Every raw interval $[s, t]$ is replaced once and for all by

$$[s + \eta, t - \eta], \qquad \eta = 10^{-7}.$$

This fixed trimming leaves a strict margin at all old-window boundaries. For either offset the active rows eventually avoid every banked deficit window; this is a finite rational range check. In particular there is no addition of two deficits at the same prime. The exact integer implementation also guards the new deficit by the vanishing of the old one. The guard holds throughout every trimmed profile window for all sufficiently large r.

Replacing m by $r/8$ and the cuts by $A_4 p/4, B_4 p/4$ changes interval endpoints by bounded amounts. The explicit rounding bounds yield, uniformly over all rows,

$$\delta_p \ge r f(p/r) - 8.$$

Both offsets use the same rational profile. Its tag $c = 1$ selects $p \equiv 4 - o \pmod 4$, and its tag $c = 3$ selects $p \equiv o \pmod 4$; thus the mirror exchanges the two prime classes without changing their integrated contribution.

Prime summation and the final constants. For a row $f(u) = \alpha + \beta u$ on a trimmed interval $[s, t]$, the prime number theorem in progressions gives the quadratic contribution

$$\frac{1}{2}\int_s^t f(u)\,du = \frac{1}{2}\left(\alpha(t-s) + \frac{\beta}{2}(t^2 - s^2)\right).$$

The constant loss 8 contributes only $O(r)$ to the logarithmically weighted prime sum. The finite sum uses the same prime range as the old denominator bound. The untrimmed contributions are

group	number of rows	gain
$m < p < 2m$	37	$\frac{14667950664797}{5028641371353600} \simeq 0.0029168814$
nonclean	20	$\frac{160166243}{12415603200} \simeq 0.0129003996$
upper windows	4	$\frac{109}{22400} \simeq 0.0048660714$

The four upper affine profiles, in tag order, are

c	u-interval	$f(u)$
1	$(4/5, 9/10)$	$9/8 - 5u/4$
1	$(3/5, 5/8)$	$2u - 9/8$
3	$(5/8, 9/14)$	$9/8 - 7u/4$
3	$(3/5, 5/8)$	$u/4 - 1/8.$

The 61-row sum after the fixed trimming is exactly

$$\Delta_\eta = \frac{4420349242147094110541}{213717258282528000000000} = 0.0206831646525407\ldots.$$

The trimming loss is less than 1.879×10^{-7}. Starting from the banked content constant

$$\begin{aligned}\Lambda_{\mathrm{old}} &= \frac{24729491161}{6665218560} - \frac{1}{448} - \frac{1}{576} - \frac{1}{352} \\ &= \frac{320572813}{86561280} = 3.7034204323226274\ldots,\end{aligned}$$

the new quadratic content constant is at most

$$\Lambda_{\mathrm{old}} - \Delta_\eta = 3.6827372676700867\ldots < \frac{184137}{50000} = 3.68274.$$

This is the content constant 3.68274 used in the next section. It uses no gain for $p < m$. The argument above is first-order: it never uses a Smith

coefficient beyond the first. Section 6 adds the second level at non-clean primes.

6. A next-level deficit at non-clean primes

Section 5 proves that the Schur complement of the zone is integral inside one cell and prices the rest by an assignment dual. This is a first-order statement. At a prime for which a long cell is overfull but does not contain the whole zone, the next Smith coefficient carries additional factors of p. We prove this here. It lowers the content constant from 3.68274 to 3.67885.

6.1. **Setting and statement.** Keep the notation of Section 5: $C = [0, r)$, $N = r + 2m$, $Z = [r, N)$, an offset $o \in \{1, 3\}$ and a prime $p \geq 13$ with $p > 2m$ and $4(N + 1) < p^2$. The zone rows are $r + 2i$ and the zone columns are $r + 2j + 1$, for $0 \leq i, j < m$. The singular residues 0 and $h \equiv (cp + o)/4 \pmod{p}$ cut $\mathbb{Z}/p\mathbb{Z}$ into two arcs. Call the arc with $(p \pm o)/4$ residues *short* and the other *long*. A *cell* is a maximal interval of integers in Z that contains no cut. It is *long* if its residues lie in the long arc. Let J be a long cell and put

$$\sigma = |J| - m.$$

The cell is *overfull* if $\sigma > 0$. It is *clean* if it contains the whole zone, that is, if $|J| = 2m$. Let S be the set of residues in the short arc, and let b be the number of residues in S that are not residues of zone nodes.

Theorem 6.1 (next-level deficit)**.** *Let J be an overfull long cell that is not clean. Then both members F_1, F_2 of the pair satisfy*

$$\operatorname{denexp}_p F_i \leq 2r + m - \min\bigl(m,\ \sigma + (\sigma - b)_+\bigr) + O(1),$$

where $(t)_+ = \max(t, 0)$ and the implied constant is absolute.

The bound of Section 5 is $2r + m - \sigma + O(1)$. The new term is $(\sigma - b)_+$. The truncation at m reflects only that valuations of the determinant are integers bounded by the size of the zone. A clean cell gets no new saving.

6.2. **The weighted congruence.** Write the normalized symmetric matrix R of Section 5 with the factor $n = 16$ and set $Y = U/p^2$, so that U is the monic variable. We work with the Gauss valuation v_G of $\mathbb{Q}(U)$ for the prime p. Its residue field is $\mathbb{F}_p(U)$. Split the unscaled prefix sums of the pole data as $F = F_{\text{pole}} + F_{\text{reg}}$ by keeping the terms whose denominators are divisible by p in the first part, and put $L = pF_{\text{pole}}$. Under $4N < p^2$ every pole denominator has at most one factor p, so L and the scaled squared prefix are integral.

For a residue c let $Q_c = UI + p^2 R_{\text{pol}}$ be the matrix on the core chain $\{k \in C : k \equiv c\}$, where R_{pol} is the Loewner matrix of the pole part. Its entries are Gauss integral and its determinant is monic in U. So $\det Q_c$ is a Gauss unit and Q_c^{-1} is Gauss integral. This uses the formal variable U and needs no specialization.

Fix a zone row z and a zone column y in the cell J, with distinct residues. Then $y - z$ is a unit. Let $g = Q_z^{-1} b$, where $b = p^2 R_{\mathrm{pol}}(\mathrm{chain}, z)$, and put

$$A = L(z) - \sum_i g_i L_i, \qquad B = 1 - \sum_i g_i,$$

the sums running over the chain of z. For every residue c put $u_c = A\mathbf{1} - BL_c$ and

$$\gamma(c) = u_c^\top Q_c^{-1} u_c.$$

Let γ_J be the value of γ at the chain of the source z.

Proposition 6.2 (weighted congruence)**.** *Let T be the Schur complement of the zone after elimination of C. For zone nodes z, y in the cell J with distinct residues, T_{zy} is Gauss integral, and modulo p*

$$T_{zy} \equiv \sum_{c \not\equiv z,y} \frac{w(c)}{(c-z)(c-y)}, \qquad w(c) = \frac{\gamma_J - \gamma(c)}{n^2} + \varepsilon \frac{B^2}{n} \mathbf{1}_S(c),$$

where ε is a unit constant depending only on the offset.

Sketch. Write $M_c = R_{CC} + (U/p^2)I$ for the core block and $r_z = R(z, C)$, $c_y = R(C, y)$. With G_z, G_y the extensions of g by zero off the chains of z and y, the Schur entry is exactly

$$T_{zy} = \Delta_{zy} - (r_z - G_z^\top M_c) M_c^{-1} (c_y - M_c G_y),$$

where $\Delta_{zy} = R_{zy} - G_z^\top c_y - r_z G_y + G_z^\top M_c G_y$. The chains of z and y are transported to each other by the translation $y - z$, since no cut lies between them. The exact commutator identities $n(t_i - t_j)(Q_z)_{ij} = L_j - L_i$ and $nt_i b_i = L(z) - L_i$, with $t_i = (\mathrm{node}_i - z)/p$, show that $Q_z(ntg) = u$. Therefore $\gamma = n(BJ - AI)$ with $I = -\sum t_i g_i$ and $J = -\sum t_i L_i g_i$. Reducing the cross terms chain by chain, with the block structure of M_c by residue, gives the displayed sum. The part with $\gamma_J - \gamma(c)$ comes from the pole prefix. The part with $\mathbf{1}_S$ comes from the regular prefix, which is periodic modulo p and equals an explicit cyclic sum on the short arc. □

6.3. **Long residues carry no weight.** Let c be a long residue. Then $w(c) = 0$ in each of the following cases.

(a) c is not a residue of a zone node. The chain of c has the same endpoint data as the chain of z, including the case that crosses r mod p, and the response is unchanged: $\gamma(c) = \gamma_J$.
(b) c is the residue of a zone node in the cell J. Chain transport inside the cell gives $\gamma(c) = \gamma_J$ again.
(c) c is the residue of a zone node outside J. Here $\gamma(c)$ may differ from γ_J. But we may project: the Cauchy kernels supported on J vanish at such a residue.

Hence, after projection, the sum in Proposition 6.2 runs over the b short residues not occupied by the zone. The remaining weights are arbitrary.

6.4. **From the congruence to the valuation.** Let W be the reduction modulo p of the leading coefficient of pT on the rows and columns of J. By Proposition 6.2, after the projection of the previous paragraph, W is a sum of at most b rank-one matrices $w(c)\, k_c(z)k_c(y)$, where $k_c(z) = 1/(c - z)$. So $\operatorname{rank} W \le b$.

The first-order argument of Section 5 shows that $\det(pT)$ has valuation at least σ. The two-level determinant lemma gives more. If σ units come from vanishing leading coefficients and the next coefficient matrix has rank at most b, then every unit of corank beyond b gives one more factor of p. Hence

$$v_G \det(pT) \ge \sigma + (\sigma - b)_+.$$

The lemma does not assume that the leading corank is exact, and it does not assume that the Cauchy coefficients are nonzero.

The second member. The matrix of F_2 has one more common node $q = N$ with the same formal diagonal variable. Since $0 < q - z \le 2m < p$, the residue of q differs from every zone residue. It changes exactly one chain response, $\gamma(q \bmod p)$. The reduction W changes by one outer product, so its rank grows by at most one. The new core node has determinant valuation -2. The total fixed loss is at most 3. The same bound for the leading block gives $v_G \ge \sigma + (\sigma - b - 1)_+$ for the second member. This proves Theorem 6.1.

6.5. **Summation.** Put $u = p/r$ and $m = \lfloor r/8 \rfloor$. The relevant primes have $u \in (\frac{1}{4}, \frac{3}{5})$, so that $p > 2m$. For a prime class $c \in \{1, 3\}$ and a given u, the new deficit is

$$f_{\mathrm{new}}(u) = \max_J \min\bigl(\tfrac{1}{8},\ \sigma_J + (\sigma_J - b_J)_+\bigr),$$

the maximum over overfull, non-clean long cells J at u, with lengths and b scaled by r. This function is piecewise affine. We merge it with the 61 rows of Section 5 by a pointwise maximum, so that no deficit is added twice at one prime. Each endpoint of each row is trimmed by 10^{-7}. Both offsets give the same profile with the two prime classes exchanged. The prime number theorem in progressions gives the quadratic contribution $\frac{1}{2}\int f\, du$ as in Section 5. The integral over the merged profile exceeds that of the 61 rows by at least 0.00389. Hence

$$\Lambda \le 3.68274 - 0.00389 \le 3.67885.$$

This is Theorem 4.1 for λ and λ'.

Remark 6.3. The argument covers only overfull long cells, with z and y in one pole interval. Cells that contain the whole zone are excluded, and we claim no saving there.

7. Periodic content for the conductor-three frame

The translated-chain argument also applies to the χ frame. It improves the content estimate without changing the node family, the decay certificate, or the nonvanishing theorem of Section 9.

Proposition 7.1. *Let $p \geq 5$ be prime, $p > 2m$, and $p^2 > 3(r+2m)$. Suppose a cut-free interval of the zone $[r, r+2m)$ contains L consecutive nodes. For every retained common-node set $J \subseteq [0, r)$, the coefficient denominator of P_J has p-exponent at most*

$$2r + m - \max(0, L - m).$$

Here the cut set consists of the singular indices and the points $jp, r + jp$, $j \in \mathbf{Z}$.

Proof. The pole data are

$$\Delta_k = \frac{1}{k} + \frac{6}{3k-2} - \frac{6}{2k-1}, \qquad D_k = \frac{1}{k^2} + \frac{18}{(3k-2)^2} - \frac{12}{(2k-1)^2}, \qquad n = 36.$$

Thus $p\Delta_k, p^2 D_k$ are integral, and both unscaled quantities are integral away from $p \mid k(3k-2)(2k-1)$. This singular set is periodic modulo p. After $x = U/p^2$, the scaled core is $UI + C$ with integral constant entries. Its residue-class blocks have monic determinant, so their cleared inverses have primitive denominators. Translation within one cut-free cell identifies the corresponding core chains and their cleared source vectors. The divided-difference formula then gives the same antisymmetric first-order cancellation as in the uniform argument: the cleared Schur entries inside the cell are integral, and entries between distinct residues have denominator exponent at most one. There are no repeated residues in the zone because $p > 2m$. The zero/one assignment bound is therefore $2r + 2m - L$ when $L > m$, and $2r + m$ otherwise.

The same bound holds after any integral constant perturbation E of the common core. Indeed, keep the old cleared partners S_R, S_C. The residuals change by $-S_R E, -E S_C$, which are integral, and the cleared second difference changes by $S_R E S_C$, also integral. The changed scaled core remains a monic shift of an integral matrix. All nonnegative entry budgets, and hence the determinant bound, survive. To recover a principal minor, add diagonal entries 0 or 1 only at the omitted common coordinates and take successive determinant differences. Multilinearity replaces those rows by identity rows; deleting them leaves exactly the desired principal restriction. This argument introduces no deletion penalty. □

For $p \equiv c \pmod 6$, the singular endpoints, up to bounded integer shifts, are $jp/6$ with

$$j \bmod 6 \in \{0, 2, 3\} \quad (c = 1), \qquad j \bmod 6 \in \{0, 3, 4\} \quad (c = 5).$$

For consecutive endpoints j, k from this dictionary, use the integral witness interval

$$a = \max\left(r, \left\lfloor \frac{jp+4}{6} \right\rfloor + 1\right), \qquad b = \min\left(r + 2m, \left\lfloor \frac{kp}{6} \right\rfloor\right).$$

The inequalities $jp + 4 < 6t < kp$ exclude all three kinds of singularity; the interval lies within one zone period. Also $2(b-a) \leq p$ whenever $b \geq a$. Thus a positive saving $(b - a - m)^+$ already implies $p > 2m$.

The following disjoint windows give an affine lower bound $Ar + Bp - 100$ for the integer saving, uniformly for $r \le 8m \le r + 9$. Only the level-one budget is $2r + m$ here; higher prime-power and bad-prime costs are retained in the total denominator and contribute $o(r^2)$.

c	lower p/r	upper p/r	A	B	j	k
1	9/32	2/7	−9/8	4	21	24
1	2/7	5/16	−1/8	1/2	21	24
1	5/16	9/28	9/8	−7/2	21	24
1	3/8	2/5	−9/8	3	15	18
1	2/5	5/12	−1/8	1/2	15	18
1	5/12	9/20	9/8	−5/2	15	18
1	27/56	1/2	−9/8	7/3	12	14
1	1/2	15/28	−1/8	1/3	12	14
1	15/28	9/16	9/8	−2	12	14
1	9/16	5/8	−9/8	2	9	12
1	5/8	2/3	1/8	0	9	12
1	2/3	3/4	9/8	−3/2	9	12
1	3/4	5/6	−1/8	1/6	8	9
1	5/6	27/32	9/8	−4/3	8	9

c	lower p/r	upper p/r	A	B	j	k
5	1/4	5/18	−1/8	1/2	24	27
5	5/18	9/32	9/8	−4	24	27
5	9/28	1/3	−9/8	7/2	18	21
5	1/3	5/14	−1/8	1/2	18	21
5	5/14	3/8	9/8	−3	18	21
5	3/8	5/12	−1/8	1/3	16	18
5	5/12	27/64	9/8	−8/3	16	18
5	9/20	1/2	−9/8	5/2	12	15
5	1/2	9/16	9/8	−2	12	15
5	9/16	3/5	−9/8	2	10	12
5	3/5	5/8	−1/8	1/3	10	12
5	5/8	27/40	9/8	−5/3	10	12

Each row satisfies the explicit integer inequality

$$(b - a - m)^+ \ge Ar + Bp - 100.$$

The prime number theorem in the two reduced classes modulo 6 gives the total gain

$$\frac{1}{2} \sum_{\text{rows}} \int_{\theta_-}^{\theta_+} (A + B\theta)\, d\theta = \frac{11983}{645120}.$$

Subtracting this from the base content constant yields

$$\Lambda_\chi = \frac{10339}{3072} - \frac{11983}{645120} = \frac{2159207}{645120} = 3.346985057\ldots. \tag{7.1}$$

Proposition 7.1 makes this content estimate uniform in J. The existing $D = 3.7251$, $c = 6.6510$ and prime interval of width $2/5$ therefore give

$$\frac{c}{D-\Lambda_\chi} = \frac{536336640}{30491189} = 17.589889328\ldots < 17.60, \qquad D-\Lambda_\chi < \frac{2}{5}.$$

The periodic entry interface, cleared cancellation, integral perturbation and principal-minor transfer, finite profile, prime summation and final measure assembly are kernel checked in Lean.

8. Nonvanishing modulo a prime

This section follows the formal proof. We treat λ in detail; λ' and $L(2,\chi_{-3})$ are in §8.7. Throughout, $\ell \equiv 3 \pmod 4$ is a prime, $K = (\ell+1)/4$, and $\mathcal{G}_e = \mathcal{G}(r,m,e)$, $e \in \{0,1\}$, is the pair of members of a scale r, with

$$1 \le m < K \le r, \qquad S := \ell - r - 2m \;\ge\; n := K - m. \tag{8.1}$$

For fixed r and $r \le 8m \le r+8$, (8.1) holds for every prime ℓ with $\frac{4}{3}(r+m)+\frac{1}{3} \le \ell \le 4r-1$, an interval $[\frac{3}{2}r + O(1), 4r]$.

Theorem 8.1 (nonvanishing)**.** *Assume* (8.1) *and let* $x_0 \in \mathbb{Q}$ *with* $\ell \nmid \operatorname{den} x_0$. *Then* $P_{\mathcal{G}_0}(x_0) \ne 0$ *or* $P_{\mathcal{G}_1}(x_0) \ne 0$.

The proof has four parts: a reduction modulo ℓ to a pair of bordered Cauchy–Gram determinants (§8.1); a lemma showing that if both vanish, a bordered Krylov determinant vanishes (§8.2); a deformation in a parameter θ (§8.3); and an algebraic evaluation of the deformed Krylov determinant, up to factors that are units at ℓ, by a commuting Jacobi matrix, diagonal steps and Darboux steps (§§8.4–8.6).

8.1. **Reduction to bordered Cauchy–Gram determinants.** Call a node b *low* if $b < K$ and *high* if $K \le b \le \ell - 1$; all nodes of $\mathcal{G}_e$ are at most $r + 2m \le \ell - 1$, and all low nodes are common nodes since $K \le r$. For a high node put $j_b = \ell - b \in [1, \ell - K]$, and for $j \ge 1$ let $w_j = (1/(a+j))_{0\le a<K} \in \mathbb{F}_\ell^K$; all denominators are nonzero modulo ℓ. For a set J let W_J be the $K \times |J|$ matrix with columns w_j, $j \in J$. In these coordinates the row nodes X' and the column nodes Y' have $J_{X'} = S + \{2,4,\ldots,2m\}$ and $J_{Y'} = S + \{1,3,\ldots,2m-1\}$. The high nodes *absent* from the geometry are $\{1,\ldots,S\}$ for $\mathcal{G}_0$ and $\{1,\ldots,S-1\}$ for $\mathcal{G}_1$, whose extra node $r+2m$ has $j = S$. Put

$$N = \sum_{j=1}^{S-1} w_j w_j^{\top}, \qquad w = w_S.$$

Lemma 8.2 (ℓ-adic structure)**.** *The only ℓ-singular factor occurring in the entries $M[a,b]$, $a,b \le \ell-1$, is $4K-1 = \ell$. Consequently $M[a,b]$ is ℓ-integral if a,b are both low, or both high and distinct; for a low and b high, $\ell M[a,b] \equiv \frac{1}{4} w_{j_b}(a)$; for b high, $\ell^2 M[b,b] \equiv 1 \pmod\ell$; and on*

the low block, $16\, M[low, low] \equiv W_{[1,\ell-K]} W_{[1,\ell-K]}^\top + H_{K-1}^{(2)} I_K \pmod{\ell}$*, where* $H_n^{(2)} = \sum_{i \le n} i^{-2}$.

Proof. $4k-1 \in \{\ell, 2\ell, 3\ell\}$ with $k \le \ell - 1$ forces $k = K$. For a low and b high, the sum defining $M[a,b]$ contains the term $k = K$ once, and $16K = 4\ell + 4$, $b - a \equiv -(a + j_b)$. For the low block, $\frac{4}{4k-1} \equiv \frac{1}{k-K}$, and partial fractions give $\sum_{j=1}^{\ell-K} \frac{1}{(a+j)(b+j)}$ off the diagonal; on the diagonal one uses $H_{\ell-K}^{(2)} \equiv -H_{K-1}^{(2)}$. □

Proposition 8.3 (reduction)**.** *Let D be diagonal with entry 1 on low and ℓ on high rows and columns, and let h_e be the number of high rows of $\Phi_{\mathcal{G}_e}$. Then $D\Phi_{\mathcal{G}_e}(x)D$ is ℓ-integral, and with $y = 16x + H_{K-1}^{(2)}$,*

$$\ell^{2h_1} P_{\mathcal{G}_1}(x) \equiv \pm 16^{-K} \det \mathcal{B}(y, N),$$
$$\ell^{2h_0} P_{\mathcal{G}_0}(x) \equiv \pm 16^{-K} \det \mathcal{B}(y, N + ww^\top) \pmod{\ell},$$

where $\mathcal{B}(y, N') = \begin{pmatrix} yI_K + N' & W_{Y'} \\ W_{X'}^\top & 0 \end{pmatrix}$.

Proof. By Lemma 8.2, modulo ℓ the matrix $D\Phi D$ has the block form $\begin{pmatrix} xI + M_{\text{low}} & \frac{1}{4}W_H & \frac{1}{4}W_{Y'} \\ \frac{1}{4}W_H^\top & I & 0 \\ \frac{1}{4}W_{X'}^\top & 0 & 0 \end{pmatrix}$, where rows are low, high common, X', columns are low, high common, Y', and H is the set of j-coordinates of the high common nodes. Take the Schur complement of the identity block. By the low-block formula and the partition of $[1, \ell - K]$ into H, the absent set, $J_{X'}$ and $J_{Y'}$, the upper-left block becomes $\frac{1}{16}(yI + N_e + W_{X'} W_{X'}^\top + W_{Y'} W_{Y'}^\top)$, with N_e the Gram matrix of the absent columns. Row and column operations using the zero lower-right block remove the last two terms, and scaling gives the claim. □

Reduction modulo ℓ commutes with evaluation at $x_0 \in \mathbb{Z}_{(\ell)}$, and $x \mapsto y$ is the same for both members. So if $P_{\mathcal{G}_0}(x_0) = P_{\mathcal{G}_1}(x_0) = 0$, then for $y_0 = 16x_0 + H_{K-1}^{(2)} \in \mathbb{F}_\ell$ both $\det \mathcal{B}(y_0, N)$ and $\det \mathcal{B}(y_0, N + ww^\top)$ vanish.

8.2. **A pair of singular bordered matrices.** Let $\Delta = \det(W_{X'}^\top W_{Y'})$. If $\Delta \neq 0$, let $\Pi = I - W_{Y'}(W_{X'}^\top W_{Y'})^{-1} W_{X'}^\top$ be the projection onto $V = \ker W_{X'}^\top$ along the column span of $W_{Y'}$, and let U be a $K \times n$ matrix whose columns span V. Define

$$k_R = \det\big[\, w^\top (\Pi N)^k U \,\big]_{0 \le k < n},$$
$$k_L = \text{the same with } X' \text{ and } Y' \text{ exchanged.}$$

(Changing U multiplies k_R by a nonzero factor; in k_L the columns of U span $\ker W_{Y'}^\top$ instead.)

Lemma 8.4 (singular pair)**.** *Let N' be symmetric, $\tau \neq 0$, and suppose $\Delta \neq 0$. If $\det \mathcal{B}(y_0, N') = \det \mathcal{B}(y_0, N' + \tau w w^\top) = 0$ for some y_0, then $k_R = 0$ for every admissible U, or $k_L = 0$ for every admissible U (computed with N' in place of N).*

Proof. Put $\mathcal{B}_2 = \mathcal{B}(y_0, N')$ and $\hat{w} = (w, 0)$, and consider the bordered matrix $\mathcal{P} = \left(\begin{smallmatrix} \mathcal{B}_2 & \hat{w} \\ -\tau \hat{w}^\top & 1 \end{smallmatrix}\right)$. Its Schur complement on the corner 1 is $\mathcal{B}_2 + \tau \hat{w}\hat{w}^\top = \mathcal{B}(y_0, N' + \tau w w^\top)$, so $\det \mathcal{P} = 0$, and its principal minor $\mathcal{B}_2$ is singular. Then $\mathcal{B}_2$ has a right kernel vector $\hat{x}$ with $\hat{w}^\top \hat{x} = 0$, or a left kernel vector $\hat{x}$ with $\hat{x}^\top \hat{w} = 0$. (If $\operatorname{rank} \mathcal{B}_2$ is at most its size minus two, its kernel meets $\hat{w}^\perp$. Otherwise $\operatorname{adj} \mathcal{B}_2 = c\, \hat{x}\hat{y}^\top$ with $c \neq 0$ and $\hat{x}$, $\hat{y}$ spanning the right and left kernels, and $0 = \det \mathcal{P} = \det \mathcal{B}_2 + \tau\, \hat{w}^\top \operatorname{adj}(\mathcal{B}_2)\hat{w} = \tau c (\hat{w}^\top \hat{x})(\hat{y}^\top \hat{w})$.)

Take the right case, $\hat{x} = (x, c)$. Then $(y_0 + N')x + W_{Y'} c = 0$, $W_{X'}^\top x = 0$ and $w^\top x = 0$; and $x \neq 0$, since otherwise $W_{Y'} c = 0$ and $c = 0$ because $\Delta \neq 0$. Applying Π (which kills $W_{Y'}$ and fixes $x \in V$) gives $\Pi N' x = -y_0 x$, hence $w^\top (\Pi N')^k x = (-y_0)^k w^\top x = 0$ for all k. Writing $x = Uz$ with $z \neq 0$, the matrix $[w^\top (\Pi N')^k U]_k$ annihilates z, so $k_R = 0$. The left case is the same argument for the transpose, $\mathcal{B}(y_0, N')^\top = \mathcal{B}(y_0, N')$ with X' and Y' exchanged, and gives $k_L = 0$. □

No condition on y_0 is needed; in particular the value $y_0 = 0$ is not special. By Proposition 8.3 and Lemma 8.4 (with $N' = N$, $\tau = 1$), Theorem 8.1 follows from the three statements

$$\Delta \neq 0, \qquad k_R \neq 0, \qquad k_L \neq 0 \qquad \text{in } \mathbb{F}_\ell. \tag{8.2}$$

8.3. **Deformation and bordered Krylov determinants.** We lift to characteristic zero and deform the nodes. Let θ be a parameter, $c_y = (1/(\theta + a + y))_{0 \le a < K}$, W_X, W_Y the matrices with columns c_y for $y \in X = S + \{2, \dots, 2m\}$ and $y \in Y = S + \{1, \dots, 2m-1\}$, $N_s = \sum_{y=1}^{s} c_y c_y^\top$, Π the projection onto $V = \ker W_X^\top$ along $\operatorname{span} W_Y$, and $A_s = \Pi N_s$. For a $K \times K$ matrix B preserving V and $u \in V$, the *bordered Krylov determinant* is

$$\kappa(B, u) = \det\big[W_Y \,\big|\, u,\ Bu,\ \dots,\ B^{n-1}u\big];$$

it is nonzero exactly when u is a cyclic vector of $B|_V$. Put

$$k(\theta) = k_{K,m,S}(\theta) = \kappa(A_{S-1}, \Pi c_S) = \kappa(A_S, \Pi c_S)$$

(the two agree by (K2) below). At $\theta = 0$ all denominators are integers $a + y \le S + K + 2m - 1 = \ell - r + K - 1 < \ell$.

Lemma 8.5 (bridge)**.** *Let $K = m + n$ with $1 \le n \le S$, and let θ be generic, in the sense that $\theta + z \neq 0$ for every integer z, with $\Delta(\theta) \neq 0$. Then the row-Krylov determinant $\det[c_S^\top (\Pi N_{S-1})^k u_i]_{k, i < n}$, in which $u_1, \dots, u_n$ is the explicit basis U of V given by Lagrange interpolation at the nodes, equals $\nu(\theta)\, k(\theta)$, where $\nu(\theta) = C_\nu \prod_{a<K} \prod_{t \in Y} (\theta + a + t)$ and $C_\nu \in \mathbb{Q}^\times$ does not depend on θ. The constant C_ν is an ℓ-adic unit for every prime $\ell > S + K + m$ (all its prime factors are at most $S + K + m$).*

Proof sketch. Let U and V' be the Lagrange bases of $\ker W_X^\top$ and $\ker W_Y^\top$, $H = V'^\top U$ and $H_c = V'^\top N_{S-1} U$; both are symmetric. Then $\Pi = U H^{-1} V'^\top$, hence $\Pi N_{S-1} U = UB$ with $B = H^{-1} H_c$, and B is self-adjoint for H, which turns the row-Krylov matrix into the column-Krylov matrix up to the factors $(c_X/c_Y)^n$ and $\det H$, where $c_X = \Pi_X(S)/\prod_a(\theta + a + S)$ and c_Y is the same with Π_Y. Two block-determinant identities, $\det[W_X \,|\, V'] \det[W_Y \,|\, U] = \Delta \det H$ and $\det[W_Y \,|\, C] \det[W_X \,|\, V'] = \Delta \det(C^\top V')$ with $C = [c_1, \dots, c_n]$, eliminate $\det H$ and Δ. By Lagrange interpolation $C^\top V'$ is a diagonal matrix times a Vandermonde matrix, and $\det[W_Y \,|\, C]$ is a Cauchy determinant; this gives the stated ν, whose only θ-dependence is $\prod_{a<K} \prod_{t \in Y} (\theta + a + t)$. The operator in the Krylov determinant of the formalisation is $\Pi N_S \Pi$, which agrees with A_S on V and is related to A_{S-1} by (K2). □

Definition 8.6. Fix $B \geq 1$. A function f of θ is *good* (with bound B) if $f(\theta) = \varkappa \prod_{1 \leq c \leq B} (\theta + c)^{e_c}$ for rational $\theta > 0$, with integers e_c and a nonzero rational $\varkappa$ whose prime factors are at most B.

Good functions with bound B form a group under multiplication. The point of the definition is that a good function with $B < \ell$ is continuous at $\theta = 0$ and its value there is an ℓ-adic unit.

Theorem 8.7 (good form)**.** *Assume $1 \leq m < K$ and $S \geq n = K - m$. Then $k_{K,m,S}$ is good with bound $S + K + 2m - 1$.*

Theorem 8.7 is all that is needed. The exact exponents e_c are known (they are piecewise quadratic in c, and $\varkappa$ is a ratio of Hankel determinants; this was checked in exact arithmetic in 156 cases with $K \leq 9$), but the proof below never computes them.

Proof of Theorem 8.1 from Theorem 8.7 and Proposition 8.8. By Proposition 8.8 below, Δ is an ℓ-unit at $\theta = 0$, so $\Delta \neq 0$ in $\mathbb{F}_\ell$. The row-Krylov determinant of Lemma 8.5 is a continuous function of θ near 0 (its entries are rational in θ with no pole at 0, and Π is continuous because $\Delta(0) \neq 0$). By Lemma 8.5 and Theorem 8.7 it equals, for rational $\theta \in (0, 1)$, a good function with bound $S + K + 2m - 1 < \ell$, which is continuous at 0; hence the values at $\theta = 0$ agree, and the value is an ℓ-adic unit. All entries are ℓ-integral at $\theta = 0$, and reduction modulo ℓ of the matrices of §8.3 gives the matrices of §8.2 (after reversing the order of the border columns, which does not change Π). So $k_R \neq 0$ in $\mathbb{F}_\ell$, and the same argument with X and Y exchanged gives $k_L \neq 0$. This is (8.2). □

Proposition 8.8 (the Cauchy-type Gram determinant)**.** *For $m \le K$ and $\theta \notin \{-1, -2, \dots\}$,*

$$\det(W_X^\top W_Y) = V(X)V(Y)\,\frac{H_{K,m}}{\prod_{i<m}(\theta+S+1+i)_m\,(\theta+S+K+1+i)_m},$$
$$H_{K,m} = \det\Big[\sum_{a=0}^{K-1} a^{p+q}\Big]_{p,q<m},$$

where $V(X) = \prod_{s<s'\in X}(s'-s)$ and $V(Y)$ are Vandermonde products. The Hankel determinant $H_{K,m}$ of the uniform weight on $\{0,\dots,K-1\}$ is a product of factorials and binomial coefficients with all prime factors less than $2K$. In particular Δ is an ℓ-adic unit at $\theta = 0$ when $S+K+2m \le \ell$.

Proof. Step 1 (Cauchy–Binet). By Cauchy–Binet and Cauchy's determinant, $\det(W_X^\top W_Y)$ is $\pm V(X)V(Y)$ times $\sum_P V(P)^2 \prod_{a\in P}\rho(a)$ over m-subsets $P \subseteq \{0,\dots,K-1\}$, where $\rho(a) = \prod_{s\in X}(\theta+a+s)^{-1}\prod_{t\in Y}(\theta+a+t)^{-1}$. This sum is the Hankel determinant $\det[\sum_a \rho(a)a^{p+q}]_{p,q<m}$ of the moments of ρ.

Step 2 (basis change and telescoping). Replacing the monomials a^p and a^q by monic rising factorials $\varphi_p(a) = (a+\theta+1)_p$ and suitable ψ_q is a unitriangular change of basis, which does not change the determinant. After it, $\rho(a)\varphi_p(a)\psi_q(a) = 1/(a+\theta+p+1)_{2m-p-q}$, and these sums over $a < K$ telescope: $L\sum_{a<K} 1/(c+a)_{L+1} = 1/(c)_L - 1/(c+K)_L$. Hence each entry is a ratio of Gamma values times a divided difference $f[u_p, v_q]$ of $f(z) = (z)_K$ at two points $u_p = \theta+p+1$, $v_q = \theta+2m-q$.

Step 3 (Newton expansion). We use divided differences only through three facts: the recursion; Newton's expansion, which peels off the nodes $u_0, u_1, \dots$ one at a time and gives $f[u_p, v_q] = \sum_{i,j} N_i(u_p)N_j(v_q)\,f[\text{window}]$ with Newton basis polynomials N_i; and the values on windows of consecutive points, $f[s, s+1, \dots, s+n] = \binom{K}{n}(s+n)_{K-n}$ for $f = (z)_K$. Thus $[f[u_p, v_q]] = \mathcal{A}\,\mathcal{D}\,\mathcal{B}^\top$ with $\mathcal{A}$, $\mathcal{B}$ triangular, and the determinant reduces to a Pochhammer product times $\det[\binom{K}{p+q+1}]_{p,q<m}$.

Step 4 (Chu–Vandermonde). The identity $\sum_{a=0}^{K-1}\binom{a}{p}\binom{K-1-a}{q} = \binom{K}{p+q+1}$ (Chu–Vandermonde), together with another unitriangular basis change, gives $H_{K,m} = \prod_p (-1)^p p! \prod_q q!\,\det[\binom{K}{p+q+1}]$. No limit in θ is used. Finally $H_{K,m}$ is the Hankel determinant of the uniform weight, i.e. of the Hahn polynomials with $\alpha = \beta = 0$ [19, §9.5]; its classical product formula has all factors at most $K+m \le 2K-1$. □

8.4. **A commuting Jacobi matrix.** Put $g = S-n+1$ and, for $0 \le a < K$,

$$\sigma(a) = (a+1)(a+1+\theta)(K-1-a)\frac{\prod_{i=1}^{m}(\theta+S+2i+a)}{\prod_{i=1}^{m-1}(\theta+S+2i+1+a)},$$
$$\tau(a) = (a+1)(a+1+\theta)(K-1-a)\frac{\prod_{i=0}^{m}(\theta+S+2i+1+a)}{\prod_{i=1}^{m}(\theta+S+2i+a)},$$
$$\mu(a) = -(K-1-a)^2(\theta+S+1+a)(\theta+S+2m+1+a).$$

Let $\tilde{T}$ be the $K\times K$ tridiagonal matrix with $\tilde{T}_{a,a+1} = -\sigma(a)$, $\tilde{T}_{a+1,a} = -\tau(a)$ and

$$\tilde{T}_{aa} = \mu(a) + \frac{\sigma(a-1)\tau(a-1)}{\mu(a-1)} + g(g+1)a^2 + g\,c_0\,a,$$
$$c_0 = -2(K-1)(\theta+g) - 2K^2 + 2K - m^2.$$

For $m=0$ this is, up to conjugation, the Jacobi matrix of a family of Hahn polynomials; for $m \ge 1$ the products over i record the border.

Proposition 8.9 (bispectral identities)**.** *(i)* Column chain. *For $1 \le y \le S$, $\tilde{T}c_y = l(y)\,c_{y-1} + d(y)\,c_y + r(y)\,c_{y+1} + \sum_{t\in Y}\kappa_t(y)\,c_t$ with $l(y) = -\tau(-\theta-y)$, $r(y) = -\sigma(-\theta-y-1)$, $d(y) = \tilde{T}_{aa}|_{a=-\theta-y}$, and explicit corrections $\kappa_t(y)$ given by residues of τ and σ at the border points. The transpose $\tilde{T}^{\top}$ satisfies the analogous* row chain *with corrections in* $\operatorname{span} W_X$*, with the same coefficients.*

(ii) Invariance. *$\tilde{T}$ maps $V = \ker W_X^{\top}$ and $\operatorname{span} W_Y$ into themselves; hence $\tilde{T}\Pi = \Pi\tilde{T}$.*

(iii) Commutation. *$[N_S, \tilde{T}]$ maps V into $\operatorname{span} W_Y$; hence $\tilde{T}$ commutes with A_S on V.*

Proof sketch. (i) says that the three-term recurrence of $\tilde{T}$ in the node variable a is mirrored, under $a \mapsto -\theta - y$, by a three-term recurrence in the column variable y: a bispectral duality in the sense of Grünbaum and Haine [15]. It is an identity between rational functions of a. Conjugating by $\hat{Q}(z) = \prod_{t\in Y}(z+t)$ turns $\tilde{T}$ into a difference operator with polynomial coefficients, and (i) becomes the statement that a certain polynomial has degree at most $m+2$. Its four top coefficients are computed from the low-order coefficients of $\prod_i(1+c_iW)$ modulo W^4 and vanish identically; the remainder, of degree less than m, is identified by Lagrange interpolation at the border points. (ii) follows from (i) at the border points, where l and r vanish. (iii) follows from (i) and the row chain by telescoping $\sum_{y\le S}\big((\tilde{T}c_y)c_y^{\top} - c_y(\tilde{T}^{\top}c_y)^{\top}\big)$. □

8.5. **Krylov calculus.** We use five elementary facts about bordered Krylov determinants of matrices preserving V.

(K1) *Cross ratio.* If A and T commute on V, then $\kappa(A,u)\,\kappa(T,u') = \kappa(A,u')\,\kappa(T,u)$ for all $u, u' \in V$. (If $T|_V$ is cyclic, $A = p(T)$ on V, and $\kappa(A,u)/\kappa(T,u)$ is the determinant of the matrix of the powers of $p(T)$ in the Krylov basis of T, which does not depend on u.)

(K2) *Rank-one update along the start vector.* $\kappa(B + uz^{\top}, u) = \kappa(B,u)$.

(K3) *Chains.* If $Tc_y \equiv l(y)c_{y-1} + d(y)c_y + r(y)c_{y+1}$ modulo $\operatorname{span} W_Y$ for the relevant y, then $\kappa(T, \Pi c_{y_0}) = \prod_{k<n}\prod_{j<k} l(y_0 - j) \cdot \det[W_Y \,|\, c_{y_0}, c_{y_0-1}, \dots, c_{y_0-n+1}]$, and similarly upward with r; the Krylov vectors are triangular in the chain.

(K4) *Cauchy determinants.* $\det[W_Y \,|\, c_{y_1}, \dots, c_{y_n}]$ is a Cauchy determinant: a product of differences of the shifts divided by a product of factors $\theta + a + y$ [20, 21].

(K5) *Transfer.* If $B_2\Phi = \Phi B_1$ for a linear map Φ carrying V_1 into V_2, then $\kappa(B_2, \Phi z) = \det[W_2 \,|\, \Phi] \cdot \kappa(B_1, z)$, where $\det[W_2 \,|\, \Phi]$ is computed in adapted bases.

8.6. **Proof of Theorem 8.7.** Write $u(m, n, S; \theta) = k_{m+n,m,S}(\theta)$ and $B = S + n + 3m - 1 = S + K + 2m - 1$. We prove by induction on $n \geq 1$ that $u(m, n, S)$ is good with bound B for all $S \geq n$.

Base case $n = 1$. Then $\kappa(A, \Pi c_S) = \det[W_Y \,|\, \Pi c_S] = \det[W_Y \,|\, c_S]$, a Cauchy determinant, which is good.

Diagonal steps (from S to $S+1$). Since $c_y(\theta) = c_{y+1}(\theta - 1)$, the instances (θ, S) and $(\theta - 1, S + 1)$ have the same border. Apply (K1) to the commuting pairs $(A, \tilde{T})$ of both instances, (K2) to pass between A_{S-1} and A_S, and (K3), (K4) to evaluate the four values $\kappa(\tilde{T}, \Pi c_y)$ that occur as explicit products (each is nonzero, so $\tilde{T}$ is cyclic). The result is

$$u(m, n, S + 1; \theta - 1) = u(m, n, S; \theta) \cdot \varkappa_{\mathrm{DS}} \prod_{j<n-1} \Big(\frac{\theta + K - 1 - j}{\theta + j}\Big)^{2(n-1-j)}$$

with a constant $\varkappa_{\mathrm{DS}}$ whose prime factors are at most B. Substituting $\theta + 1$ for θ expresses $u(m, n, S + 1; \theta)$ through $u(m, n, S; \theta + 1)$ times factors $\theta + c$ with $1 \leq c \leq K$; so goodness passes from S to $S + 1$. (It is this orientation, downward in S and upward in θ, that keeps all bases $c \geq 1$.)

Darboux steps (from $n-1$ to n at $S = n$). Consider the instance (K, m, S) with $S = n - 1$, i.e. $g = 0$. After normalising $\tilde{T}$ so that it annihilates the kernel vector of A on V, it has an LU factorisation

$$\tilde{T} = LU, \qquad U = \operatorname{diag}(\mu) - \operatorname{superdiag}(\sigma), \qquad L = I - \sum_a \frac{\tau(a-1)}{\mu(a-1)} E_{a,a-1},$$

with explicit pivots $\mu(a) = -(K - 1 - a)^2(\theta + S + 1 + a)(\theta + S + 2m + 1 + a)$; note $\mu(K - 1) = 0$. Reversing the factors gives the Jacobi matrix of the instance with one node fewer and one more column:

$$(UL)_{<K-1} = D^{-1}(\tilde{T}' + \beta I)D, \qquad \tilde{T}' = \tilde{T}(K - 1, m, S + 1),$$

for an explicit diagonal D and scalar β. This is the discrete Darboux (Christoffel) transformation of Jacobi matrices [38, 44, 16]. The bidiagonal map $X = DU$ with its last row removed is an intertwiner:

$$X\tilde{T} = (\tilde{T}' + \beta)X, \qquad X\Pi = \Pi' X, \qquad XA = A'X,$$

where primes refer to $(K - 1, m, S + 1)$. The first identity is the reversal of the factors; the other two follow from the column and row chains of

Proposition 8.9 at the border points ($Xc_t \in \operatorname{span} W_{Y'}$ for $t \in Y$, and dually) and, for A, from telescoping $XN_S - N'_{S+1}X$ in y. By (K5), (K2) and (K1),

$$\kappa(A, \Pi c_0)\, \kappa(\tilde{T}', w') = \det\big[W_Y \,|\, \Pi c_0 \,|\, E\big] \cdot \kappa(A', w') \cdot \prod_{k<n-1} \prod_{j<k} l(-j) \cdot \mathcal{G},$$

where $w' = \Pi' c'_n$ is a projected Cauchy vector of the smaller instance, E completes $[W_Y \,|\, \Pi c_0]$ by Cauchy columns (so the first factor is a Cauchy determinant), and $\mathcal{G} = \det[R^\top \Pi c_{-j}]_{j<n-1}$ with $R = [c_1, \dots, c_{n-1}]$. Two further inputs are needed.

(a) *The Gram determinant $\mathcal{G}$ (Chu–Vandermonde).* A Schur complement gives $\mathcal{G} \cdot \Delta = \det(C_{\mathcal{A}}^\top C_{\mathcal{B}})$ for two Cauchy matrices whose column sets $\mathcal{A}$, $\mathcal{B}$ have $K-1$ elements. By Cauchy–Binet this is a sum over the one omitted node a, with $V(P_a)^2 = V_K^2/(a!\,(K-1-a)!)^2$, and the sum is evaluated by

$$\sum_{a=0}^{N} \binom{N}{a}^2 (x+a)_{2N} = \binom{2N}{N} ((x+N)_N)^2 \qquad (N = K-1).$$

This is a terminating balanced ${}_3F_2$ identity of Pfaff–Saalschütz type [1, §2.2]. We prove it with three applications of the Chu–Vandermonde identity: the moment identity $\sum_a \binom{N}{a}^2 a^{\underline{j}} = N^{\underline{j}} \binom{2N-j}{N}$ (falling factorials), the relation $\binom{2N}{i}\binom{2N-i}{N} = \binom{2N}{N}\binom{N}{i}$, and Vandermonde once more for $(x+2N-1)^{\underline{N}}$. With Proposition 8.8 this makes $\mathcal{G}$ explicit.

(b) *A nondegeneracy (positivity).* The transfer (K5) needs $\det[W' \,|\, X\Pi R] \neq 0$. Its numerator, as a function of θ, is not a product, so this is proved by positivity rather than by a formula: for rational $\theta > 0$ (this is the only range used, and it is proved over $\mathbb{Q}$), if $W's + X\Pi R t = 0$, then $s = 0$ (apply $W_{X'}^\top$ and $X\Pi = \Pi' X$), so $y = \Pi R t$ lies in $V \cap \ker X$; the range of $X^\top$ is contained in the span of $[R \,|\, W_X]$ (the row chain at the border points) and both have dimension $K-1$, so they are equal and $R^\top y = 0$, i.e. $(R^\top \Pi R)t = 0$. Finally $\det(R^\top \Pi R) \cdot \Delta = \det([R \,|\, W_X]^\top [R \,|\, W_Y])$ by a Schur complement. For two families of m' distinct shifts x, y with $a + x_i > 0$ and $a + y_i > 0$ for all nodes a, Cauchy–Binet gives

$$\det(C_x^\top C_y) = V(x)V(y) \sum_P V(P)^2 \prod_{a \in P} \prod_i \frac{1}{(a+x_i)(a+y_i)},$$

the sum over m'-subsets P of the nodes, with V the Vandermonde product. The sum is positive and $V(x)V(y) \neq 0$, so the determinant is nonzero. This applies to Δ and to $[R \,|\, W_X]^\top [R \,|\, W_Y]$, hence $\det(R^\top \Pi R) \neq 0$ and $t = 0$.

Every factor in the displayed identity is now good except the products over $l(-j) = -\tau(-\theta + j)$ and the Pochhammer denominator of $\mathcal{G}$, which contain factors $\theta - j - 1$ with negative shift; these two sets of factors coincide and cancel. The values $\kappa(\tilde{T}', w')$ and $\kappa(\tilde{T}, \cdot)$ are chain products by (K3) and

nonzero. Combining with the diagonal step at $(K, m, n-1)$, (K1) and (K2), one obtains

$$u(m, n, n; \theta - 1) = u(m, n-1, n; \theta) \cdot (\text{good factor}).$$

Since $u(m, n-1, n)$ is good by the induction hypothesis (as $n \geq n-1$), so is $u(m, n, n)$, and the diagonal steps give all $S \geq n$. All bases stay in $[1, B]$ and all constants have prime factors at most B. This proves Theorem 8.7. □

8.7. **The mirror constant and $L(2, \chi_{-3})$.** *The mirror constant.* For λ' one takes primes $\ell \equiv 1 \pmod 4$ and $K = (\ell+3)/4$. The only ℓ-singular factor is again $4K-3 = \ell$, and the reduction gives the same pair of bordered matrices (with the opposite sign of the shift). Everything else is unchanged.

$L(2, \chi_{-3})$: three levels of nodes. Let $\ell = 6t+1$ be prime, $K = (\ell+2)/3 = 2t+1$, $T = (\ell+1)/2 = 3t+1$, and $m_0 = 3t$. Now there are two ℓ-singular indices among $1, \dots, \ell-1$, namely K (where $3K-2 = \ell$) and T (where $2T-1 = \ell$); the factor k in $d(k)$ is a unit. Accordingly the nodes are *low* $[0, K)$, *middle* $[K, m_0]$ and *top* $[T, \ell-1]$. The geometries are tied to ℓ: we take $r + 2m = m_0$, so all nodes of $\mathcal{G}_e$ are low or middle, the extra node of the second member is m_0, and $K \leq r$ is the condition $2m+1 \leq t$. With the scaling 1 on low and ℓ on middle and top nodes, a window identity for sliding sums of $1/u$ modulo ℓ,

$$\sum_{a<k\leq b} \frac{1}{k - t_1 - 1} - \sum_{a<k\leq b} \frac{1}{k - t_0} = (b-a) \sum_{t=t_0}^{t_1} \frac{1}{(a-t)(b-t)} \quad (a, b \notin [t_0, t_1]),$$

gives the entries modulo ℓ: low–low $\bar{x}\,\delta_{pq} + A_{pq}$; low–middle $-\frac{1}{6} \cdot \frac{1}{p-q}$; middle–middle $-\frac{1}{2}\delta_{pq}$; low–top and middle–top 0. Here, in terms of $u_y = (1/(a+y))_{a<K}$,

$$A = \kappa\Big(hI + \sum_{n\in[K,T)} u_{\ell-n} u_{\ell-n}^{\top}\Big) + \tau_0 \sum_{n\in[T,\ell)} u_{\ell-n} u_{\ell-n}^{\top}, \qquad \kappa = -\tfrac{1}{18}, \quad \tau_0 = \tfrac{1}{36},$$

and $h = \frac{1}{2}\big(\sum_{y=1}^{3t} y^{-2} - 2\sum_{y=3t+1}^{4t} y^{-2}\big)$ in $\mathbb{F}_\ell$ (note $\ell - K = 4t$). The middle and top nodes enter with weights of opposite sign. The identity for A is the reflection $n = \ell - y$ of the two windows above, and the cross coefficient $\frac{1}{6}$, the middle diagonal $\frac{1}{2}$ and κ satisfy $(\frac{1}{6})^2 = \frac{1}{2} \cdot \frac{1}{18}$, which makes the middle level behave as in the two-level case; the top level does not, and only enters through A. The Schur complement of the middle block then gives, as in Proposition 8.3, bordered matrices $\mathcal{B}(z,\ N_{\mathrm{mid}} + \frac{\tau_0}{\kappa} N_{\mathrm{top}})$, where N_{top} is the Gram matrix of the top columns, which are the w_j with $j \in [1, T-1]$, and N_{mid} that of the absent middle columns: w_T for the first member and none for the second. Hence the two members give

$$\mathcal{B}(z,\ cN) \quad \text{and} \quad \mathcal{B}(z,\ cN + ww^{\top}),$$
$$N = \sum_{j=1}^{S-1} w_j w_j^{\top},\ w = w_S,\ S = T,\ c = \frac{\tau_0}{\kappa} = -\frac{1}{2}.$$

Dividing the upper-left block by c (and rescaling the border) turns this into the pair $\mathcal{B}(z/c, N)$, $\mathcal{B}(z/c, N+c^{-1}ww^\top)$, which is the situation of Lemma 8.4 with $\tau = c^{-1} = -2 \neq 0$. The hypotheses (8.1) hold with $S = T$: $r+2m+S = \ell$, and $S = 3t+1 \geq K-m$. So Theorem 8.7 and Proposition 8.8 apply verbatim, all bases are at most $S+K+2m-1 = \ell - r + K - 1 < \ell$, and we obtain the analogue of Theorem 8.1: if $\ell \nmid \operatorname{den} x_0$, one of the two members does not vanish at x_0. No information on the block $M[X', Y']$ is needed.

Existence of pairs. For $\ell = 6t+1$ we need integers $r, m \geq 1$ with $r + 2m = 3t$, $2m+1 \leq t$ and an admissible relation between r and m. With $r \leq 8m \leq r+W$ this means $3t \leq 10m \leq 3t+W$, which is solvable for all t if and only if $W \geq 9$; for $W = 8$ it fails exactly when $\ell \equiv 43 \pmod{60}$. We therefore use $r \leq 8m \leq r+9$ for $L(2,\chi_{-3})$; then every prime $\ell \equiv 1 \pmod 6$ with $\ell \geq 61$ has a pair, with $5r \leq 2\ell \leq 5r+11$.

9. Uniform nonvanishing after bounded low-node deletion

For the χ frame, allowing a bounded number of common nodes to be omitted removes the restriction $r + 2m = 3t$ in the preceding pair construction. The borders remain fixed. We use the rational kernel M of Section 3. For $J \subseteq [0, r)$ put

$$X' = \{r + 2i : 0 \leq i < m\}, \qquad Y' = \{r + 2i + 1 : 0 \leq i < m\},$$

and put $\xi_\chi = \frac{1}{4}L(2, \chi_{-3})$. Let $\Phi_J(x)$ have rows $J \sqcup X'$, columns $J \sqcup Y'$, and entries $M[a, b] + x\mathbf{1}[a = b \in J]$. Write $P_J(x) = \det \Phi_J(x)$.

Theorem 9.1. *Let $\ell = 6t+1$ be prime, and suppose*

$$m \geq 1, \qquad K := 2t+1 \leq r, \qquad r + 2m \leq 3t. \tag{9.1}$$

If $x \in \mathbb{Q}$ and $\ell^2 \nmid \operatorname{den} x$, there is a set $I \subseteq [0, K)$, $|I| \leq 3$, such that, for $J = [0, r) \setminus I$,

$$P_J(x) \neq 0.$$

The proof bounds the dimension of a kernel, rather than asserting simplicity of its eigenvalues. Conditions (9.1) imply $2m+1 \leq t$, so $t \geq 3$ and $\ell \geq 19$. All occurrences of $2, 3, 6, 18, 36$ below are therefore units modulo ℓ.

9.1. Reflected coordinates and the signed moment pencil.

Work first over $k = \mathbb{F}_\ell$. The physical node b and its reflected coordinate $y = \ell - b$ will be kept distinct. Put

$$S = 3t+1, \qquad g = \ell - r - 2m, \qquad d = K - m, \qquad h = r - K.$$

Then $1 < S \leq g$, $3 \leq d < K$, $d \leq g$, and $K + g + 2m + h = \ell$. For a reflected coordinate y define

$$c_y = ((a+y)^{-1})_{0 \leq a < K}, \qquad N = \sum_{y=1}^{g} \epsilon_y c_y c_y^\top, \qquad \epsilon_y = \begin{cases} 1 & y < S, \\ -2 & y \geq S. \end{cases}$$

Reversing the order of each physical border gives the reflected border coordinates $g+2, g+4, \dots, g+2m$ and $g+1, g+3, \dots, g+2m-1$. Denote their column matrices by W_X, W_Y, respectively, and set

$$\Gamma = W_X^\top W_Y, \qquad B_z = \begin{pmatrix} N - zI_K & W_Y \\ W_X^\top & 0 \end{pmatrix}.$$

Proposition 8.8 gives $\det \Gamma \neq 0$: its range condition is $g + K + 2m = \ell - r + K \leq \ell$.

Introduce the polynomials

$$D(y) = \prod_{a=0}^{K-1} (y+a), \qquad B_X(y) = \prod_{i=0}^{m-1} (y - g - 2i - 2),$$
$$B_Y(y) = \prod_{i=0}^{m-1} (y - g - 2i - 1), \qquad R(y) = B_X(y) B_Y(y).$$

For $0 \leq i < d$ define residue-basis matrices by

$$(V_X)_{ai} = \frac{B_X(-a)(-a)^i}{D'(-a)}, \qquad (V_Y)_{ai} = \frac{B_Y(-a)(-a)^i}{D'(-a)}. \tag{9.2}$$

All displayed denominators and all $B_X(-a), B_Y(-a)$ are nonzero. Partial fractions give, for $\deg p < d$,

$$\sum_{a=0}^{K-1} \frac{B_X(-a)p(-a)}{D'(-a)(y+a)} = \frac{B_X(y)p(y)}{D(y)}, \tag{9.3}$$

and likewise for Y. Thus the columns of V_X lie in $\ker W_X^\top$. They are independent, since evaluation of a polynomial of degree $< d \leq K$ at the K distinct points $-a$ is injective. Since Γ is invertible, W_X, W_Y have rank m; consequently V_X, V_Y are bases of $\ker W_X^\top, \ker W_Y^\top$.

The compressed matrix is the symmetric Hankel pencil

$$A_z = V_Y^\top (N - zI_K) V_X = G - zH,$$

whose entries, indexed by $0 \leq i, j < d$, are

$$H_{ij} = \sum_{u \in \{0,-1,\dots,1-K\}} \frac{R(u)}{D'(u)^2} u^{i+j}, \tag{9.4}$$

$$G_{ij} = \sum_{y=1}^{g} \epsilon_y \frac{R(y)}{D(y)^2} y^{i+j}. \tag{9.5}$$

These identities follow from (9.2) and (9.3). In particular the same polynomial coordinates describe the left and right kernels; this fact will be needed for principal deletion.

9.2. **A uniform corank bound.**

Lemma 9.2. *For every $z \in k$, $\dim_k \ker A_z \le 3$.*

Proof. Identify a coefficient vector in $\ker A_z$ with a polynomial p, $\deg p < d$. Equations (9.4)–(9.5) say that its first d weighted moments vanish. To use these conditions, complete the root sets by

$$E(y) = \prod_{j=1}^{g} (y - j), \qquad C(y) = \prod_{j=g+2m+1}^{\ell-K} (y - j).$$

The last product is 1 when $h = 0$. The roots of D, E, R, C partition k, so

$$DERC = y^\ell - y, \qquad (y^\ell - y)' = -1. \tag{9.6}$$

Let $T = DE$. On its simple roots define

$$w_z(v) = \begin{cases} -zR(v)/D'(v)^2 & D(v) = 0, \\ \epsilon_v R(v)/D(v)^2 & E(v) = 0, \end{cases}$$

and form the polynomial, linear in p,

$$B_p(y) = \sum_{T(v)=0} w_z(v) p(v) \frac{T(y)}{y - v}, \qquad F_p(y) = D(y)^2 C(y) B_p(y).$$

The first d moments are zero. Expanding $B_p/T = \sum_v w_z(v)p(v)/(y - v)$ formally at infinity therefore gives

$$\deg B_p \le K + g - d - 1 = g + m - 1, \qquad \deg F_p \le \ell + d - 1.$$

Equivalently these degree bounds follow by comparing the first d leading coefficients in the finite polynomial sum defining B_p. Evaluating at a root of T, and differentiating (9.6), gives the exact identities

$$\begin{array}{lll} F_p(u) = F_p'(u) = 0, \quad F_p''(u) = 2zp(u) & (D(u) = 0), \\ F_p(j) = -\epsilon_j p(j) & (1 \le j \le g), \\ F_p(j) = 0 & (C(j) = 0). \end{array} \tag{9.7}$$

For example $B_p(u) = -zR(u)E(u)p(u)/D'(u)$ at a root u of D; then $F_p''(u) = 2D'(u)^2 C(u) B_p(u) = 2zp(u)$ by $D'(u)E(u)R(u)C(u) = -1$. The high-node identity follows in the same way.

Consider the finite-difference Wronskian

$$W_p(y) = F_p(y + 1)p(y) - F_p(y)p(y + 1).$$

At each low adjacency $-K + 1, \dots, -1$, its value and first derivative vanish. Its second derivative also vanishes, by the equality $2zp(y + 1)p(y) - 2zp(y)p(y + 1) = 0$. Hence these are triple roots. At the high adjacencies $1, \dots, S - 2$ and $S, \dots, g - 1$ its value vanishes because the two weights agree. It vanishes at every root of C: for the last such root, $\ell - K$, the next point is the first low root $\ell - K + 1$. Thus

$$Q_0(y) := \prod_{a=1}^{K-1} (y + a)^3 \prod_{j=1}^{S-2} (y - j) \prod_{j=S}^{g-1} (y - j) C(y) \quad \text{divides } W_p(y). \tag{9.8}$$

The supports are disjoint, and

$$\deg Q_0 = 3(K-1) + (g-2) + h = \ell + 2d - 5,$$
$$Q_0(S-1) \neq 0, \qquad W_p(S-1) = 3p(S)p(S-1).$$

We also need that $W_p \neq 0$ for nonzero p in the kernel. If $W_p = 0$, reduce $F_p/p = A/B$ to coprime polynomials with B monic. Then $A(y+1)B(y) = A(y)B(y+1)$ implies $B \mid B(y+1)$, and hence $B(y+1) = B(y)$. But $\deg B \leq \deg p < \ell$; a nonconstant polynomial of degree $e < \ell$ has a nonzero leading term of degree $e-1$ in its first difference. Thus B is constant. The polynomial quotient F_p/p is translation invariant and has degree $< 2\ell$. Its positive degree, if any, must be ℓ by the same leading-coefficient argument. Subtracting a multiple of $y^\ell - y$ therefore shows

$$F_p = p(a(y^\ell - y) + b).$$

Some low node u has $p(u) \neq 0$, since $\deg p < K$. The first two identities in (9.7) force $b = 0$ and $a = 0$. Then the high-node identity forces p to vanish at $g > \deg p$ points, a contradiction.

Finally, if the kernel had dimension at least four, its three linear conditions

$$[y^{d-1}]p = [y^{d-2}]p = 0, \qquad p(S) = 0$$

would admit a nonzero solution. This solution has $\deg p \leq d-3$. The leading terms in the two products defining W_p cancel, so $\deg W_p \leq \deg F_p + \deg p - 1 \leq \ell + 2d - 5 = \deg Q_0$. By (9.8), W_p is a scalar multiple of Q_0; its value at $S-1$ makes that scalar zero. This contradicts the preceding paragraph and proves the lemma. □

9.3. **Selecting a nonzero principal minor.**

Lemma 9.3. *For every $z \in k$, deleting at most three low principal coordinates from B_z leaves a nonsingular matrix.*

Proof. Let $\mathcal{P}_z$ be the polynomial kernel of A_z, and put $k_0 = \dim \mathcal{P}_z \leq 3$. A right kernel vector of B_z has low part $v \in \ker W_X^\top$, hence $v = V_X p$ uniquely. Its remaining equation is $(N - zI)v \in \operatorname{im} W_Y$, which is equivalent to $V_Y^\top (N - zI) V_X p = 0$. The border coordinates are then unique, since Γ is invertible. Conversely every $p \in \mathcal{P}_z$ lifts to such a kernel vector. The transpose has the corresponding parametrization with low part $V_Y p$; the polynomial space is the same because A_z is symmetric.

Evaluation $p \mapsto (p(-a))_{a<K}$ is injective on $\mathcal{P}_z$. Choose k_0 coordinates on which it is an isomorphism, and let I be the corresponding low indices. By (9.2), restriction of either full kernel to I is an isomorphism: it differs from polynomial evaluation only by nonzero diagonal factors.

Here is the elementary principal-minor criterion being used. Suppose a square matrix has injective restriction of its right kernel to I and surjective restriction of its left kernel to I. A kernel vector of the complementary principal matrix, extended by zero on I, is sent by the full matrix to a vector supported on I. Pairing with left kernel vectors shows that this image is

zero. Right-kernel injectivity then shows the extended vector is zero. Thus the complementary principal matrix is nonsingular. Apply this criterion to B_z. □

9.4. **Return to the physical determinant.** Suppose first $\ell \nmid \operatorname{den} x$, and write $\bar{x}$ for its reduction. Let

$$c_0 = \sum_{y=1}^{3t} y^{-2} - 2 \sum_{y=3t+1}^{4t} y^{-2}, \quad \kappa = -\frac{1}{18}, \quad \tau_0 = \frac{1}{36}, \quad h_0 = \frac{c_0}{2}.$$

The three-level entry calculation of Section 8.7 applies under (9.1); it does not require equality $r + 2m = 3t$. The physical absent middle nodes are $[r + 2m, S)$, which reflect to $[S, g]$; the top nodes reflect to $[1, S)$. Their weighted Gram matrix is therefore

$$N_{\mathrm{phys}} = \sum_{y=S}^{g} c_y c_y^{\top} - \frac{1}{2} \sum_{y=1}^{S-1} c_y c_y^{\top} = -\frac{1}{2} N.$$

Consequently the physical reduced low block is exactly

$$(h_0 + \bar{x}/\kappa) I + N_{\mathrm{phys}} = -\frac{1}{2}(N - zI), \qquad z = c_0 - 36\bar{x}. \tag{9.9}$$

Both border orders are reversed simultaneously to obtain W_X, W_Y. Scaling a low block by a nonzero scalar c preserves bordered nonsingularity, since

$$\begin{pmatrix} cI & 0 \\ 0 & I \end{pmatrix} \begin{pmatrix} M_0 & W_Y \\ W_X^{\top} & 0 \end{pmatrix} \begin{pmatrix} I & 0 \\ 0 & c^{-1} I \end{pmatrix} = \begin{pmatrix} cM_0 & W_Y \\ W_X^{\top} & 0 \end{pmatrix}.$$

This also holds after any low principal restriction. Thus Lemma 9.3 supplies a nonzero physical bordered minor.

For completeness, this minor transfers to the rational matrix without assuming that the retained common set contains every low node. Start with the original geometry and multiply its low rows and columns by 1, its middle rows and columns by ℓ, and negate the matrix. For any retained low set $U \subseteq [0, K)$ its reduced principal restriction has the same Schur identity as the full matrix. Writing M_{phys} for the left-hand side of (9.9),

$$\det Z_U = (1/2)^{r-K+m} (1/18)^{|U|} \det \begin{pmatrix} M_{\mathrm{phys}}[U, U] & W_Y[U, *] \\ W_X[U, *]^{\top} & 0 \end{pmatrix}. \tag{9.10}$$

Indeed the invertible middle common block is unchanged by low restriction; the harmonic identity restricts entrywise, and the border row and column operations are unchanged. Every prefactor is a unit. A zero rational principal determinant would remain zero after the rational diagonal scalings and after reduction, contradicting (9.10). Taking $U = [0, K) \setminus I$ proves the theorem in this case.

It remains to treat $\ell \mid \operatorname{den} x$ but $\ell^2 \nmid \operatorname{den} x$. Coprimality of the reduced numerator and denominator shows that $u = \overline{\ell x}$ is nonzero. Use the undeleted common set $J = [0, r)$. Multiply *every* row of $\Phi_J(x)$ by ℓ and additionally

multiply each middle common column by ℓ. With rows and columns ordered as low, middle common, border, the resulting integral matrix reduces to

$$\begin{pmatrix} uI_K & 0 & -\frac{1}{6}W_Y \\ -\frac{1}{6}W_C^\top & -\frac{1}{2}I_{r-K} & 0 \\ -\frac{1}{6}W_X^\top & 0 & 0 \end{pmatrix}.$$

Here W_C consists of the middle common columns. In verifying the zero blocks one uses integrality of distinct middle–middle entries, rather than merely integrality after multiplication by ℓ^2; it follows directly from the off-diagonal divided-difference formula, whose interval contains neither singular index K nor S. Reorder the middle common block last. Its determinant is $(-1/2)^{r-K}$, and the remaining bordered determinant is

$$\det \begin{pmatrix} uI_K & -\frac{1}{6}W_Y \\ -\frac{1}{6}W_X^\top & 0 \end{pmatrix} = u^{K-m}(-1/36)^m \det \Gamma \neq 0.$$

Thus $P_{[0,r)}(x) \neq 0$. This proves Theorem 9.1 in the remaining case.

9.5. **Content, decay, and the original degree budget.** The enlarged family consists of all P_J with $|[0,r) \setminus J| \leq 3$. For admissible $r \leq 8m \leq r+9$, Section 7 gives

$$\Lambda = \frac{2159207}{645120}, \qquad D = 3.7251, \qquad c = 6.6510.$$

The integral-perturbation and determinant-difference argument of Proposition 7.1 supplies one improved denominator $d_{r,m}$ for every P_J, including all bounded deletions. Its logarithm is at most $(\Lambda + o(1))r^2$, uniformly in J.

The energy estimates of Section 11 already apply to every subset of common nodes, with the borders retained. If $k \leq 3$ common nodes are forced to be absent, expansion at $\xi_\chi + v$ has terms

$$v^{r-k-|A|} \det(\omega \mathsf{L}_{F_\chi}[A \sqcup X', A \sqcup Y']), \qquad A \subseteq J.$$

For $|v| \leq e^{-cr}$, use $|v|^{r-k-|A|} \leq e^{-cr(r-k-|A|)}$. Replacing the original exponent $r-|A|$ by $r-k-|A|$ therefore costs at most e^{crk} in the termwise upper bound, also at $v = 0$. This is a logarithmic loss of at most $3cr = O(r)$, and the number of terms is still at most 2^r. It is absorbed by the existing $o(r^2)$ errors. Hence $|P_J(\xi_\chi + v)| \leq \exp(-(D - o(1))r^2)$ on the same window, uniformly in J; no comparison of possibly cancelling determinants is needed.

Finally $\deg P_J \leq |J| \leq r$. We keep the declared degree budget equal to the original r for every member, so q^r times the common clearing denominator times $P_J(p/q)$ is an integer. Neither the degree scale nor the exponential window changes when the nonzero member is selected.

10. A quarter border and characteristic-zero selection

For the conductor-three frame, characteristic-zero interpolation permits a larger border than the finite-field selection above. We take

$$r \le 4m \le r + 3,$$

and retain the full determinant together with its principal restrictions obtained by omitting one common node. Their degree budget is still r. The denominator and decay estimates below apply uniformly to every member.

Theorem 10.1 (One common-node deletion)**.** *For every $r \ge 1$, $m \ge 0$ and rational input, the full conductor-three determinant or one of its principal common-node deletions is nonzero. No prime-band restriction on m/r is required.*

Proof. Here is the interpolation argument underlying the certified selector. Write $N = r + 2m$, $n = r + m$, and use the r common nodes followed by the alternating row and column nodes in the consecutive tail. After the fixed normalization of the conductor-three frame, its divided differences have prefix values f_i, with

$$f_{i+1} - f_i = \delta(i), \qquad \delta(z) = \frac{1 - z}{216(z + 1)(z + 1/3)(z + 1/2)}.$$

The diagonal prefix e_i satisfies $e_{i+1} - e_i = -\delta'(i)$. Barycentric interpolation identifies either kernel with pairs (p, q) of degree less than n satisfying $q(i) = f_i p(i)$ at all N nodes, and $q'(c) = f_c p'(c) + (U - e_c)p(c)$ at the common nodes.

Two such pairs have zero cross product: its degree is at most $2n - 2$, while the value and first-jet conditions give $N + r = 2n$ zeros counted with multiplicity. If the kernel had dimension at least two, a nonzero pair could therefore be chosen with both degrees at most $n-2$. The discrete Wronskian

$$\begin{aligned} W(z) = 216(z + 1)(z + 1/3)(z + 1/2) \\ \cdot [q(z + 1)p(z) - q(z)p(z + 1)] \\ - (1 - z)p(z)p(z + 1). \end{aligned}$$

would have degree at most $2n - 3$, but its prescribed values and first jets supply $2n - 2$ zeros. Thus $W = 0$. This would make q/p a rational antidifference of δ, which is impossible: the simple pole at -1 is isolated in its integer-translation class. After cancellation of a numerator-denominator gcd, evaluating the cleared identity propagates a denominator root indefinitely in one direction, contradicting finite degree. The boundary case $n = 1$ is immediate.

If the full determinant is nonzero, retain it. Otherwise both kernels are one dimensional. A nonzero kernel pair cannot vanish at every common node. Dividing by their common node polynomial would give degree at most $m-1$ and the same value interpolation at all N nodes; its Wronskian would again vanish by degree and root count. The left and right barycentric maps land in the same one-dimensional interpolation-pair space. At a common node

their coordinates are the same $p(c)$ multiplied by nonzero punctured-node factors. Thus the same common index is nonzero on both sides, and their adjugate outer product has a nonzero diagonal entry there. That principal cofactor is the required deletion. □

10.1. **The uniform arithmetic profile.** The existing H and Z duals suffice; no new local valuation identity is needed. At a prime p, put $u = p/r$. Use the H bound always, and the minimum of the H and Z bounds when $r + 2m \le 2p$, the proved domain of the latter. For the two unit classes modulo 6, the resulting continuum coarse profiles w_1, w_5 have the following pieces. Endpoints can be assigned to either neighboring piece since the functions agree there.

class	interval for u	$w(u)$
$1, 5$	$[0, 1/4]$	$5/2$
$1, 5$	$[1/4, 1/2]$	$11/4 - u$
1	$[1/2, 15/16]$	$9/4$
1	$[15/16, 9/8]$	$7/2 - 4u/3$
1	$[9/8, 3/2]$	$5 - 8u/3$
1	$[3/2, 2]$	$2 - 2u/3$
1	$[2, 3]$	$1 - u/6$
1	$[3, 9/2]$	$3/2 - u/3$
5	$[1/2, 5/6]$	$9/4$
5	$[5/6, 1]$	$7/2 - 3u/2$
5	$[1, 3/2]$	$4 - 2u$
5	$[3/2, 2]$	$5/2 - u$
5	$[2, 3]$	$3/2 - u/2$

Both profiles are zero beyond their last nonzero interval. Their exact integrals are $581/128$ and $355/96$. The prime number theorem in the two classes gives

$$\Lambda_0 = \frac{1}{2}\left(\frac{581}{128} + \frac{355}{96}\right) = \frac{3163}{768}.$$

The finite-size proof uses affine covers with a fixed additive allowance for rounding m and the residue endpoints. The generic prime summation retains every higher-prime-power and bad-prime cost. The original entrywise tropical denominator clears every principal restriction, so this coarse bound holds for the one-deletion family.

Retaining the periodic D1 saving. For odd primes $p \le m$, the coarse level-one tropical profile discards the existing exact bound

$$v_p^{\mathrm{trop}} \le 2r + 2m - \min(m \bmod p, p - m \bmod p).$$

It holds at every node level. One formula treats every nearest-multiple strip. For each integer $n \ge 1$, on $m/(n+1) < p < m/n$ retain

$$\min\{m - np, (n+1)p - m\}.$$

The two affine pieces meet at $p = 2m/(2n+1)$ and have integral

$$\int_{m/(n+1)}^{m/n} \min\{m - np, (n+1)p - m\}\, dp = \frac{m^2}{2n(n+1)(2n+1)}.$$

Use the first sixteen strips, leaving the coarse bound below $m/17$. The weighted prime number theorem in the two unit classes modulo 6 gives the same denominator estimate with

$$\Lambda_{16} = \Lambda_0 - \frac{1}{16}\sum_{n=1}^{16} \frac{1}{2n(n+1)(2n+1)} = \frac{339258406689121}{82516315939200}.$$

A single finite-size argument covers both pieces for every n, with a fixed additive rounding allowance. All bad-prime and higher-power charges remain included. The same denominator clears every retained common minor, so the family and selector are unchanged. The generic proof is `D1QuarterPrefix`; its P6 adapter is `Quarter.Periodic.hasContent`.

10.2. **The new analytic certificate.** The existing rectangular decay engine is instantiated with $\alpha = 1/4$ and a rational probability measure on 1600 atoms: 200 negative real atoms, 200 positive real atoms and 600 conjugate pairs. Its regularization radius is $1/2048$; all support points have modulus at most 16. The certified function is

$$g(s) = \frac{5}{4}U^{\sigma}(s) - U^{\nu_{1/4}}(s) + \frac{999}{1000}\pi|\operatorname{Im} s|.$$

The interval-potential identity

$$U^{\nu_{1/4}}(s) = \frac{1}{2}U^{\tau}(s) + \frac{3}{4}U^{\tau}(2s/3) - \frac{3}{4}\log(3/2)$$

reuses the existing pointwise potential bounds. The finite cells, far-field Taylor bounds, continuum tails and pair-energy sums give

$$m' = \frac{231169}{100000}, \qquad g_{\min} = -\frac{143459}{100000}, \qquad S_{\text{base}} = -\frac{274719}{10000000},$$
$$I^{\sigma} \le \frac{1808029}{1000000}, \qquad I^{\nu} \ge \frac{184428}{100000}.$$

Thus $D \ge 1158407/250000 = 4.633628$ and $c = 93657/12500 = 7.49256$ are valid. Bounded deletion changes only a linear term, absorbed by the existing asymptotic decay theorem. Theorem 10.1 supplies the pinned interface on the unit-width interval $(1, 2)$; the existing square-divisibility prime rule has $s = 1$. Since $D - \Lambda < 1$, the pinned measure criterion applies without a prime-location surcharge and gives

$$\mu(L(2, \chi_{-3})) \le \frac{93657/12500}{1158407/250000 - \Lambda_{16}} = 14.3475713785\ldots < 14.35.$$

The public statement is 14.35. The certificate uses the shared content, common-minor decay and pinned measure interfaces; no new global determinant engine is introduced.

11. Analytic decay

Theorem 11.1 (decay and window)**.** *For admissible geometries of scale r and real t with $|t| \le e^{-cr}$,*

$$|P_{\mathcal{G}}(\xi + t)| \le \exp(-(D - o(1))r^2),$$
$$(D, c) = \begin{cases} (3.90557,\ 7.3546) & \text{for } \lambda, \lambda', \\ (3.7251,\ 6.6510) & \text{for } L(2, \chi_{-3}). \end{cases}$$

We describe the proof for λ; the other two cases differ only in the reflection formula and the certificate data.

11.1. **A contour representation.** Hermite's formula gives $\mathsf{L}_F[a, b] = \frac{1}{2\pi i} \oint F(z) v_a(z) v_b(z)\, dz$ with $v_a(z) = 1/(z - a)$, the contour enclosing the nodes and no pole of F. The reflection formulas $\psi(1+z) = \psi(-z) - \pi \cot \pi z$ and $\psi(\frac{3}{4} + z) = \psi(\frac{1}{4} - z) + \pi \cot(\frac{\pi}{4} - \pi z)$ split $F = \tilde{F} + E$, where $\tilde{F}$ is analytic in the left half-plane and $O(1/|z|)$ there, and

$$E(z) = -\pi\big[\cot \pi z + \cot(\tfrac{\pi}{4} - \pi z)\big]$$

is 1-periodic with $|E(x + iy)| \le C_E e^{-2\pi|y|}$ for $|y| \ge 1$. Let $N_0 = \lfloor \frac{7}{50} r \rfloor$, $x^* = -N_0 - \frac{3}{8}$, and let s_0 be the slope parameter, $s_0 = \frac{5}{2}$ for λ, λ' and $s_0 = 3$ for $L(2, \chi_{-3})$, and let Γ be the two rays $x^* + u(-1 \pm s_0 i)$, $u \ge 0$. Moving the contour to Γ picks up the residues of F at the $2N_0$ poles $-(j + \frac{3}{4})$ (residue $+1$) and $-(j + 1)$ (residue -1), $j < N_0$; the part of the integrand with $\tilde{F}$ integrates to 0 over Γ. Hence, for all integers $a, b \ge 0$,

$$\mathsf{L}_F[a, b] = \int v_a v_b\, dm, \qquad m = -\sum_{\mu} \epsilon_\mu\, \delta_{-\mu} + \frac{1}{2\pi i} E(z)\, dz\big|_\Gamma, \tag{11.1}$$

a Gram representation with respect to a complex measure m with finitely many atoms and an exponentially small density on Γ.

11.2. **Two-family Andréief bound.** For a set $S \subseteq C$ of common nodes let $R_S = S \sqcup X'$, $Q_S = S \sqcup Y'$, $n = |S| + m$, and $\mathcal{M}_S = \omega\, \mathsf{L}_F[R_S, Q_S]$. By multilinearity in the diagonal,

$$P_{\mathcal{G}}(\xi + t) = \sum_{S \subseteq C} t^{|C| - |S|} \det \mathcal{M}_S. \tag{11.2}$$

Andréief's identity [13] for the two families $(v_a)_{a \in R_S}$ and $(v_b)_{b \in Q_S}$ gives

$$\det \mathsf{L}_F[R_S, Q_S] = \frac{1}{n!} \int \det[v_a(z_i)]\, \det[v_b(z_i)] \prod_{i=1}^{n} dm(z_i),$$

and Cauchy's determinant evaluates both factors. Taking absolute values (see [37] for the logarithmic potential theory behind what follows),

$$|\det \mathsf{L}_F[R_S, Q_S]| \le \frac{1}{n!} \int \prod_{i<i'} |z_i - z_{i'}|^2 \prod_{\mathcal{F} \in \{R_S, Q_S\}} \frac{\prod_{a<a' \in \mathcal{F}} |a' - a|}{\prod_{i,\, a \in \mathcal{F}} |z_i - a|} \prod_i d|m|(z_i).$$

The integrand is the exponential of minus a discrete logarithmic energy: the free points z_i repel each other with weight 1, each node family repels itself with weight $\frac{1}{2}$ and attracts the free points with weight 1, and the density $|E| \approx e^{-2\pi|\operatorname{Im} z|}$ acts as an external field. Because the rows and columns differ only in the interleaved blocks, the two node families have the same macroscopic density: after scaling by r, both are

$$\nu_\alpha = \mathrm{Leb}|_{[0,1]} + \tfrac{1}{2}\mathrm{Leb}|_{[1,1+2\alpha]}, \qquad \alpha = \tfrac{1}{8},$$

of mass $1 + \alpha$. Its logarithmic energy has the closed form

$$I(\nu_{1/8}) = \iint -\log|u - u'|\, d\nu_{1/8}(u)\, d\nu_{1/8}(u') = \frac{243}{128} - \frac{\log 2}{32} - \frac{25}{32}\log\frac{5}{4} = 1.70244\ldots$$

11.3. **The energy certificate.** Fix ρ ($\rho = 1/4096$ for λ, λ' and $\rho = 1/1024$ or $1/2048$ for $L(2, \chi_{-3})$) and the regularised kernel $k_\rho(z, w) = -\frac{1}{2}\log(|z - w|^2 + \rho^2) \le -\log|z - w|$.

Lemma 11.2 (conditional positive definiteness)**.** *For points $p_1, \ldots, p_N \in \mathbb{C}$ and real weights q_α with $\sum q_\alpha = 0$, $\sum_{\alpha,\beta} q_\alpha q_\beta k_\rho(p_\alpha, p_\beta) \ge 0$.*

Proof. Frullani's integral $\log a = \int_0^\infty (e^{-t} - e^{-at})\, dt/t$ and $\sum q_\alpha = 0$ give $\sum q_\alpha q_\beta(-\log(|p_\alpha - p_\beta|^2 + \rho^2)) = \int_0^\infty \frac{e^{-t\rho^2}}{t} \sum q_\alpha q_\beta e^{-t|p_\alpha - p_\beta|^2} dt$, and Gaussian kernels are positive definite: expanding $e^{2t\langle p, p'\rangle}$ in powers and the binomial theorem write the inner sum as a convergent sum of squares. □

Apply Lemma 11.2 to the free points and removed nodes (weight $1/r$ each) against a *certificate measure* $\sigma = \sum_k m_k \delta_{c_k}$ of total mass $1 + \alpha$. This bounds the free–free energy from below by a sum of one-point terms $U^\sigma_\rho(s) = \sum_k m_k k_\rho(s, c_k)$ and the constant $I_\rho(\sigma) = \sum_{k,l} m_k m_l k_\rho(c_k, c_l)$. Riemann-sum comparisons replace the node sums by the potential U^{ν_α} and the node self-energy by $I(\nu_\alpha)$, up to $O(r \log r)$. With the external field one arrives at the function

$$g(s) = (1 + \alpha)U^\sigma_\rho(s) - U^{\nu_\alpha}(s) + (1 - \kappa_0)\pi|\operatorname{Im} s|, \qquad \kappa_0 = 10^{-3},$$

on the scaled support of $|m|$ (the rays Γ/r, the segment $[-\frac{7}{50}, 0]$ containing the atoms, and, for removed nodes, the segment $[0, 1 + 2\alpha]$). The certificate consists of three finite families of inequalities, checked in interval arithmetic:

(a) $g \geq m'$ on a cover of the rays by boxes, with an analytic tail bound beyond $u = \frac{3}{2}$;
(b) $g \geq \gamma_\ell$ on a partition of $[-\frac{7}{50}, 0]$ into cells; put $S_{\text{cert}} = 2\sum_\ell |\text{cell}_\ell| \min(\gamma_\ell - m', 0)$;
(c) $g \geq g_{\min}$ on a cover of $[0, 1 + 2\alpha]$.

On a box, U_ρ^σ is bounded below by moving each c_k to the farthest corner, and U^{ν_α} is bounded above from its closed form and monotonicity. Then

$$D = 2((1+\alpha)m' + S_{\text{cert}}) - (1+\alpha)^2 I_\rho(\sigma) + I(\nu_\alpha), \qquad c = 2(m' - g_{\min}).$$

Proposition 11.3 (certified constants)**.** *For λ and λ' there is a measure σ with 1650 atoms with dyadic coordinates and masses, and $\rho = 1/4096$, for which*

$$m' \geq 2.1711, \quad S_{\text{cert}} \geq -0.12573, \quad (1+\alpha)^2 I_\rho(\sigma) \leq 2.43054,$$
$$g_{\min} \geq -1.5061, \quad I(\nu_{1/8}) \geq 1.7024457,$$

whence $D \geq 3.90557$ and $c \leq 7.3546$. For $L(2, \chi_{-3})$ the certificate of §11.5 gives $D \geq 3.7251$ and $c \leq 6.6510$.

All the inequalities are verified by the Lean kernel (§13); the data were generated with interval arithmetic in Arb [18] and then checked independently in Lean with rational arithmetic. The certificate for λ and λ' is described in the next subsection.

11.4. **Dual-guided atoms and steeper rays.** Two choices distinguish the certificate for λ and λ'.

The slope. Nothing in the contour argument requires slope 3. The bound $|E(x+iy)| \leq C_E e^{-2\pi|y|}$ holds on any pair of rays $x^* + u(-1 \pm s_0 i)$ with $s_0 \geq 1$, and the Gram representation (11.1) is unchanged. The slope enters the certificate only through the box cover of the rays (a) and the tail bound beyond $u = \frac{3}{2}$. The vertical coordinate on a box is $s_0 u \in [s_0 u_1, s_0 u_2]$. With the other parameters fixed, a scan over $s_0 \in \{\frac{9}{4}, \frac{5}{2}, \frac{11}{4}\}$ gives the best ratio at $s_0 = \frac{5}{2}$. The decay engine takes s_0 as a parameter. Its exponent formula does not depend on s_0; only the certified constants m', $g_{\min}$ and c do.

The atoms. The measure σ has 150 atoms on the segment $[-\frac{7}{50}, 0]$, 600 conjugate pairs on the rays and 300 atoms on the line $[0, 1]$. The positions were not chosen on a uniform grid. The certified ratio is the value of a fractional program in the masses of σ. We solved it numerically by a cutting-plane method together with Dinkelbach's iteration. The dual of this program is a measure on the support of m, and its support shows where the constraints $g \geq m'$, $g \geq \gamma_\ell$ and $g \geq g_{\min}$ are active. We place atoms at the active points, at spacing about ρ, and re-solve for the masses. With $\rho = 1/4096$ this gives the constants of Proposition 11.3.

Each of the three families of inequalities is checked on a cover by cells with exact integer arithmetic (dyadic cell endpoints in units of 2^{-40}, outward rounded). In each group of consecutive cells the atoms close to the cell are treated one by one and the remaining ones by a first-order Taylor expansion.

The covers have 29 701 ray boxes, 6000 segment cells and 6430 line cells. The lower bounds in Lean are 10^{-5} below the interval values m' and $g_{\min}$. This allowance lets far atoms be grouped and costs only $4 \cdot 10^{-5}$ in D.

Remark 11.4 (limits of this certificate). Weak duality bounds every certificate of this form from below. If μ is a probability measure on the support of m, then any admissible function g satisfies an inequality in the integrals $\int g\, d\mu$, and an exact rational dual μ proves a lower bound for the exponent c/η that holds for all σ. For $s_0 = \frac{5}{2}$, $\rho \geq 1/4096$, $\alpha = \frac{1}{8}$ and the content 3.68274, a rational dual measure with 3076 atoms on the rays (verified in directed interval arithmetic with 192 bits) shows that the exponent cannot be below 32.85. This computation is not part of the Lean development. The certificate of Proposition 11.3 is within 0.1 of this floor at the content 3.68274. Smaller ρ needs proportionally more atoms.

11.5. **The certificate for $L(2, \chi_{-3})$.** For F_χ the reflection formula $\psi(z + a) = \psi(1 - a - z) - \pi \cot \pi(z + a)$ gives $F_\chi = \tilde{F}_\chi + E_\chi$ with

$$E_\chi(z) = \pi\Big[\tfrac{1}{3} \cot \pi(z + \tfrac{1}{3}) - \tfrac{1}{2} \cot \pi(z + \tfrac{1}{2}) + \tfrac{1}{6} \cot \pi(z + 1)\Big],$$

which is 1-periodic and, because the weights $\frac{1}{3} - \frac{1}{2} + \frac{1}{6}$ sum to zero, decays like $e^{-2\pi|\operatorname{Im} z|}$. The contour argument and the Andréief bound are as above, with poles at $-(j + \frac{1}{3})$, $-(j + \frac{1}{2})$, $-(j + 1)$. The square-frame certificate for F_χ (a measure σ_χ of 699 atoms, $\rho = 1/2048$) gives lower bounds for the square-frame function $g_\chi = U_\rho^{\sigma_\chi} - U^\tau + (1 - \kappa_0)\pi|\operatorname{Im} s|$ on rays, segment and line. The hybrid certificate is

$$\sigma_{\mathrm{h}} = \frac{\sigma_\chi + \lambda}{9/8}, \qquad \lambda = \sum_{k=0}^{31} \frac{1}{256}\, \delta_{t_k}, \qquad t_k = 1 + \frac{k}{128} - \frac{1}{2048},$$

so that λ has mass $\frac{1}{8}$ and represents the extra density $\frac{1}{2}\mathrm{Leb}|_{[1,5/4]}$; the atom t_k sits at the left end $a_k = 1 + k/128$ of the k-th cell, shifted left by ρ. The function of the hybrid frame splits as $g_{\mathrm{h}} = g_\chi + h$ with $h = U_\rho^\lambda - U^{\mathrm{ex}}$, where U^{ex} is the potential of $\frac{1}{2}\mathrm{Leb}|_{[1,5/4]}$. For u in the k-th cell and $\operatorname{Re} s + \rho \leq a_k$ one has $(\operatorname{Re} s - t_k)^2 + \rho^2 \leq (\operatorname{Re} s - u)^2$, hence $k_\rho(s, t_k) \geq -\log|s - u|$ on the cell, and integrating over the cells gives $h(s) \geq 0$ whenever $\operatorname{Re} s \leq 1 - \rho$. On the short real interval $[1 - \rho, 1]$ only the first cell is affected, and an elementary bound for $\int_0^\epsilon -\log v\, dv$ gives $h \geq 0$ there too. Therefore every lower bound of the square-frame certificate holds for g_{h}, and only the energy $I_\rho(\sigma_{\mathrm{h}})$ is new: it consists of the energy of σ_χ, the cross terms (699×32 logarithms) and the self-energy of λ (32^2 logarithms), summed exactly in rational interval arithmetic. The result is $D \geq 3.7251$ (the computed value is 3.725194) and $c \leq 6.6510$; with 64 extra atoms instead of 32 one would gain only 0.0008 in D.

Since the admissible relation for $L(2, \chi_{-3})$ is $r \leq 8m \leq r + 9$, the decay estimate is proved uniformly for $\alpha r \leq m \leq \alpha r + \frac{9}{8}$, $\alpha = \frac{1}{8}$: the comparison

of node sums with U^{ν_α} and $I(\nu_\alpha)$ is carried out with a window parameter for the excess of m over αr, which only changes the $O(r\log r)$ error terms.

11.6. **Node removal and the window.** The argument applies to every subset $S \subseteq C$: removing $j = \beta r$ common nodes turns them into points of the configuration that are not integrated over, and only common nodes can be removed, because a node removed from one family only would leave an uncancelled term of weight $\frac{1}{2}$. One obtains

$$|\det \mathcal{M}_S| \le \exp(-r^2(D - \beta c - o(1))) \qquad (\beta = |C \setminus S|/r),$$

uniformly in S. Inserting this into (11.2) with $|t|e^{cr} \le 1$ gives Theorem 11.1: the sum has at most $2^{|C|} = e^{O(r)}$ terms, each at most $e^{-(D-o(1))r^2}$. The second member is handled by viewing it as the pattern $(r, m+1)$ with one column node moved by one place, at an energy cost $O(r)$.

12. Proof of Theorem 1.1

12.1. $G \pm 5\pi^2/48$**.** Let $\xi = \lambda$ (the case λ' is identical, with primes $\ell \equiv 1 \pmod 4$). Let $\eta = D - \Lambda \ge 3.90557 - 3.67885 = 0.22672$, let $\nu > c/\eta$, and suppose $|\xi - p/q| < q^{-\nu}$ with q large; put $L = \log q$. Choose $\delta > 0$ with $\nu > c(1+\delta)/\eta$, let $r = \lceil (1+\delta)L/\eta \rceil$ and choose m with $r \le 8m \le r + 8$.

Choice of ℓ. By the prime number theorem in progressions, the primes $\ell \equiv 3 \pmod 4$ in $[\frac{4}{3}(r+m)+1, 4r-1]$ have product $\exp((\frac{5}{4}+o(1))r)$ (as $m \le \frac{r}{8}+1$), which exceeds $q = e^L$ since $r \gg L$. So one of them does not divide q. By Theorem 8.1 with $x_0 = p/q$, there is $e \in \{0,1\}$ with $P_{\mathcal{G}_e}(p/q) \ne 0$, where $\mathcal{G}_e = \mathcal{G}(r, m, e)$.

Lower bound. By Lemma 3.3, $|d_{\mathcal{G}_e} P_{\mathcal{G}_e}(p/q)| \ge q^{-(r+1)} = e^{-(r+1)L}$.

Upper bound. Write $p/q = \xi + t$ with $|t| < e^{-\nu L} \le e^{-cr}$ (as $cr \le c(1+\delta)L/\eta + c < \nu L$ for large q). By Theorems 4.1 and 11.1, $|d_{\mathcal{G}_e} P_{\mathcal{G}_e}(p/q)| \le e^{(\Lambda+o(1))r^2} e^{-(D-o(1))r^2} = e^{-(\eta - o(1))r^2}$.

Contradiction. Comparing, $(\eta - o(1))r^2 \le (r+1)L$; but $\eta r \ge (1+\delta)L$. Hence $\mu(\lambda) \le c/\eta$, and

$$\frac{c}{\eta} \le \frac{7.3546}{3.90557 - 3.67885} = 32.4392\ldots < 32.44.$$

Since $G - 5\pi^2/48 = 2\lambda$ and $G + 5\pi^2/48 = 2\lambda'$, this gives the first two bounds of Theorem 1.1.

12.2. $L(2, \chi_{-3})$**.** *The earlier pair-family bound.* Now the geometry is tied to the prime: for each prime $\ell \equiv 1 \pmod 6$, $\ell \ge 61$, there is an admissible pair of members of scale r with $5r \le 2\ell \le 5r + 11$, so $r/\ell \to g = \frac{2}{5}$, and Theorem 8.1 (in the form of §8.7) applies when $\ell \nmid q$. Let $\eta = 3.7251 - \frac{10339}{3072} = 0.359540\ldots$, $\nu > c(1/\eta + \frac{4}{5})$, and choose δ small. The primes $\ell \equiv 1 \pmod 6$ in an interval $(Y, Y + (2+\delta)L]$ with $Y = \frac{5}{2}(1+\delta)L/\eta$ have product

$e^{(1+\delta/2+o(1))L} > q$, so one of them, ℓ, does not divide q. The corresponding scale satisfies

$$(1+\delta)L/\eta - O(1) \le r \le \tfrac{2}{5}(Y+(2+\delta)L)+O(1) = \Big(\frac{1+\delta}{\eta}+\frac{2(2+\delta)}{5}\Big)L+O(1),$$

so $cr < \nu L$ for δ small, and the comparison above, with $|C| \le r+1$, gives a contradiction. Hence

$$\mu\big(\tfrac{1}{4}L(2,\chi_{-3})\big) \le c\Big(\frac{1}{\eta}+gs\Big), \qquad g=\tfrac{2}{5},\ s=2,$$

and $6.6510\,(1/0.3595401+0.8) = 23.8194\ldots < 23.82$. The valuation-one argument of Section 9 also allows $s=1$, giving the retained bound 21.16.

The bounded-deletion family. Put

$$\eta = \frac{37251}{10000} - \frac{10339}{3072} = \frac{690317}{1920000}, \qquad c = \frac{6651}{1000}.$$

Let $\nu > c/\eta$, choose $\delta > 0$ so that $c(1+\delta)/\eta < \nu$, put $L = \log q$ and $r = \lceil (1+\delta)L/\eta \rceil$, and set $m = \lceil r/8 \rceil$. For all sufficiently large r, every prime $\ell \equiv 1 \pmod 6$ in

$$\frac{51}{20}r \le \ell \le \frac{59}{20}r$$

satisfies $K = (\ell+2)/3 \le r$ and $r+2m \le (\ell-1)/2$. Indeed $r+2m \le 5r/4+7/4$ and the lower endpoint leaves a positive linear margin; the upper endpoint gives $K \le 59r/60+2/3 \le r$ for large r. The prime number theorem in this progression gives

$$\log \prod_{\substack{51r/20 \le \ell \le 59r/20 \\ \ell \equiv 1 \ (6)}} \ell^2 = \Big(\frac{2}{5}+o(1)\Big)\,r > L,$$

since $2/5 > \eta$. Thus some such prime satisfies $\ell^2 \nmid q$. By Theorem 9.1, some retained common set J with at most three omissions has $P_J(p/q) \ne 0$. The original tropical denominator $d_{r,m}$ clears its coefficients, its degree is at most r, and the estimates in Section 9 give

$$q^{-r} \le |d_{r,m}P_J(p/q)| \le \exp(-(\eta-o(1))r^2)$$

whenever $|\xi_\chi - p/q| < q^{-\nu}$; the required window holds because $cr < \nu L$. This contradicts $\eta r \ge (1+\delta)L$. Consequently

$$\mu(\xi_\chi) \le \frac{c}{\eta} = \frac{12769920}{690317} = 18.498631788\ldots < \frac{37}{2}.$$

Rational scaling by four preserves the exponent. The periodic content profile of Section 7 first replaces this ratio by $536336640/30491189 < 17.60$.

The quarter-border family. Finally use Section 10, with $m = \lceil r/4 \rceil$, at most one common-node deletion, and

$$\eta = \frac{1158407}{250000} - \Lambda_{16} = 0.522217997\ldots, \qquad c = \frac{93657}{12500}.$$

For any $\nu > c/\eta$, choose $\delta > 0$ with $c(1+\delta)/\eta < \nu$ and take $r = \lceil (1+\delta)\log q/\eta \rceil$ as above. The characteristic-zero selector gives a nonzero retained member at every rational input. Its degree is at most r; the uniform quarter-border content and decay bounds give the same integer comparison $q^{-r} \le |d_{r,m} P_J(p/q)| \le \exp[-(\eta - o(1))r^2]$ in the required window. This again contradicts $\eta r \ge (1+\delta)\log q$. Consequently $\mu(\xi_\chi) \le c/\eta = 14.3475713785\ldots < 14.35$; rational scaling by four completes Theorem 1.1. □

Remark 12.1 (relation to linear independence criteria). The deduction is a one-dimensional version of the familiar scheme behind the criteria of Nesterenko [30] and their use by Ball and Rivoal [2]: the nonzero integer $q^{|C|} d_{\mathcal{G}} P_{\mathcal{G}}(p/q)$ plays the role of a small linear form with integer coefficients. The difference is that the lower bound for the form comes from an arithmetic nonvanishing statement (Theorem 8.1) rather than from an analytic asymptotic, and that the size r of the determinant is chosen in terms of q.

Remark 12.2 (where the constants enter). For λ, $\partial\mu/\partial D = -c/\eta^2 \approx -158$, so the third decimal of D matters. The rank deficits of §4, the uniform rescaled-core deficit of Section 5 and the next-level deficit of Section 6 determine Λ; the certificate of §11.4 determines D and c. The free choice of ℓ (Theorem 8.1 for all S in a range) removes a term $c\,g\,s$ which would otherwise be about 5.1. For $L(2,\chi_{-3})$, valuation-one nonvanishing first reduces the pair-family bound from 23.82 to 21.16; bounded low-node deletion removes the remaining prime-location cost; the periodic content improvement then gives 17.60. Characteristic-zero selection then allows the quarter border of Section 10, giving the limiting ratio $14.3475713785\ldots$. At this fixed geometry, reducing the fixed deletion count further would not change the leading decay or content constants.

13. Formalization

All results of this paper are formalized in Lean 4 [9] with Mathlib [40]. The development is available at

https://github.com/slitsey/catalan-irrationality-measure

and combines analytic and arithmetic proofs with generated rational data and kernel checks for the dual certificates of Section 4 and the decay certificates of Section 11; numerical computations alone are not used as proof. The constants are written out as series, so the main statements use no definitions of the development: G is $\sum_j (-1)^j/(2j+1)^2$ and $L(2,\chi_{-3})$ is $\sum_k \big((3k+1)^{-2} - (3k+2)^{-2}\big)$. The main statements are stated in `ChallengeHybrid.lean` and proved in `SolutionHybrid.lean` (namespace `CatalanHybridChallenge`):

- `lam_hybrid_measure_3244`: for every $e > 0$, all sufficiently large $q \in \mathbb{N}$ and all $p \in \mathbb{Z}$ coprime to q, $q^{-(32.44+e)} < |G - 5\pi^2/48 - p/q|$ (Theorem 1.1);
- `mir_hybrid_measure_3244`: the same for $G + 5\pi^2/48$;

- `chi3_hybrid_measure_1435`: the same for $L(2, \chi_{-3})$ with exponent 14.35; it is `CatalanFrames.P6Hybrid.Quarter.chi3_measure`;
- `Quarter.decay`, `Quarter.Periodic.hasContent` and `Quarter.hasNVPin`: the quarter-border decay, content and characteristic-zero selection interfaces;
- `longCell_rowLocal`, `kind3_rowLocal` and `lam_both_eps_merged`, `mir_both_eps_merged`: the next-level deficit of Section 6 on a long cell, its binding to the prime profile and the resulting content bound;
- `lam_hybrid_decayS`: Theorem 11.1 for λ with the certificate of §11.4; the decay engine in `CatalanCore/Decay` takes the ray slope as a parameter;
- `p6_minorNV_square`, `bandPencil_corank` and `physical_low_rescue`: the bounded-deletion nonvanishing argument of Section 9;
- `nv_hyb_nonvanishing_gen`: Theorem 8.1 and its analogues in §8.7;
- `hyb_primal_false` and `primal_pair`: the end of the nonvanishing chain and Lemma 8.4;
- `p6_hybrid_decay`: Theorem 11.1 for $L(2, \chi_{-3})$;
- `profile_both_eps`, `p6_content`: Theorem 4.1 for λ and $L(2, \chi_{-3})$.

The solution has no `sorry` and no `native_decide`, and the command `#print axioms` reports only the three standard axioms `propext`, `Classical.choice` and `Quot.sound`. The file `comparator-hybrid.json` is a configuration for `lakecomparator` stating that `SolutionHybrid` proves the statements of `ChallengeHybrid` with these axioms only. We have not run Comparator, which requires Linux; the axiom check above was done with `#print axioms`. The prime number theorem in arithmetic progressions (moduli 4 and 6) enters the content asymptotics and the selection of the prime ℓ. The formal source for it is the PNT+ project [41], through OpenAI's workspace and compatibility additions, whose Wiener–Ikehara theorem we use.
The following file groups implement the indicated parts of the proof.

`CatalanCore/Rect`
Geometries, the matrices $\Phi_{\mathcal{G}}$, Lemma 3.3, pair families and admissible relations.

`CatalanCore/Arith/Hybrid`
Gauss valuations, Lemma 4.2, the level majorant, Proposition 4.3, prime sums over linear pieces, the rank-deficit interface (Lemma 4.5) and bordering.

`CatalanCore/Arith/Hybrid/ShiftNonclean`, `CatalanFrames/HybCommon/DeficitProfile`
Proposition 6.2 and Theorem 6.1: the weighted congruence, the two-level determinant bound and the merged profile.

`CatalanFrames/Lambda/Hyb*`, `CatalanFrames/LambdaMirrorHybrid`, `CatalanFrames/P6Hybrid`
Theorem 4.1 and Theorem 4.6 for the three constants.

`CatalanFrames/LambdaHybrid/NV`
Proposition 8.3 and its three-level analogue, Lemma 8.4 (`Primal`), Theorem 8.1 (`FinalNV`).

`CatalanCore/Star/Leonard`
Proposition 8.8, Proposition 8.9, (K1)–(K5), the diagonal and Darboux steps, the positivity argument, Lemma 8.5, Theorem 8.7 and the specialisation $\theta \to 0$.

`CatalanCore/Decay/Rect`, `CatalanCore/Cert/SigmaHybS`, `CatalanCore/Cert/SigmaP6H`
The contour identity, the two-family Andréief bound, Lemma 11.2, node removal, and the certificates of Proposition 11.3 and §11.5.

`CatalanCore/Measure`, `CatalanFrames/*/Assembly*`
The deduction of Section 12.

The proofs were checked with Lean `v4.35.0-rc2` and the matching Mathlib release. The vendored files from OpenAI's repository and from the PNT+ project were converted to Lean's module system and ported to the current toolchain; the repository's README and provenance documents describe the changes.

Disclosure of AI use

The original development was produced by the AI systems Claude Opus 5.5 and Claude Sonnet 5.5 (Anthropic): the mathematics, including the rectangular frames, the content bounds and rank deficits, the non-vanishing argument and the decay certificates, and all proofs; the Lean formalization; the numerical computations and the certificate generators; the literature search; and the text of this paper. Claude worked in a series of conversations at the request of the author, who posed the problem, directed the work, and is responsible for this submission. The repository infrastructure, vendored formalizations and certificate tooling of the companion project [25], which this work reuses, were developed with OpenAI Codex as described there. The subsequent improvement for $L(2, \chi_{-3})$ from 23.82 to 17.60, including valuation-one reduction and the periodic content improvement, the finite-field corank argument, bounded-deletion transfer, their Lean formalization and the corresponding additions to this paper, was developed with OpenAI Codex. The earlier exposition is retained except where the stronger theorem requires changes. The subsequent uniform rescaled-core argument for both $G \pm 5\pi^2/48$, the finite interval profile and its Lean formalization were also developed with OpenAI Codex. The subsequent improvement of the $G \pm 5\pi^2/48$ exponent to 32.44 was carried out by Claude (Anthropic) working with Sean Litsey and with Codex-developed infrastructure: the second-level deficit at non-clean primes with its weighted congruence, the slope-parametrised decay engine and the dual-guided certificate, and their Lean formalization. The subsequent quarter-border

improvement to 14.35, including characteristic-zero selection, the analytic certificate and the retained periodic D1 saving, was developed with OpenAI Codex. The underlying determinant method and the irrationality proof for G are OpenAI's [32]. We use Lean 4 [9], Mathlib [40], the PNT+ project [41], and Arb [18] (through `python-flint`) for interval arithmetic in generating certificate data, which Lean re-checks. The formal verification in Lean is intended as an independent check of every step of the argument above.

References

1. George E. Andrews, Richard Askey, and Ranjan Roy, *Special functions*, Encyclopedia of Mathematics and its Applications, vol. 71, Cambridge University Press, Cambridge, 1999.
2. Keith Ball and Tanguy Rivoal, *Irrationalité d'une infinité de valeurs de la fonction zêta aux entiers impairs*, Invent. Math. **146** (2001), no. 1, 193–207.
3. Frits Beukers, *A note on the irrationality of* $\zeta(2)$ *and* $\zeta(3)$, Bull. London Math. Soc. **11** (1979), no. 3, 268–272.
4. Francis Brown, *Mellin transforms, transfinite diameter and rational approximations of integrals*, 2026, Preprint, 22 April 2026.
5. Frank Calegari, Vesselin Dimitrov, and Yunqing Tang, *The linear independence of* 1, $\zeta(2)$, *and* $l(2, \chi_{-3})$, 2024.
6. ———, *Arithmetic holonomy bounds and effective Diophantine approximation*, 2025, Theorem 5.1 gives $\mu(L(2, \chi_{-3})) \leq 24781$ (checked at the source by S. Litsey).
7. Marcos Costa Santos Carreira, *General designs and an irrationality exponent for Catalan's constant*, Preprint v1, October 12, 2026, doi:10.5281/zenodo.23326325; sources at `https://github.com/MarcosCarreira/catalan-designs`, 2026.
8. Harold Davenport, *Multiplicative number theory*, third ed., Graduate Texts in Mathematics, vol. 74, Springer, New York, 2000, Revised and with a preface by H. L. Montgomery.
9. Leonardo de Moura and Sebastian Ullrich, *The Lean 4 theorem prover and programming language*, Automated Deduction – CADE 28, Lecture Notes in Computer Science, vol. 12699, Springer, 2021, pp. 625–635.
10. Payman Eskandari, *Improved rational approximations to Catalan's constant*, 2026, Preprint, 22 September 2026.
11. Payman Eskandari, Kumar Murty, and Yusuke Nemoto, *Mixed motives and linear forms in the Catalan constant*, 2025, Preprint, October 2025 (v3: 29 August 2026).
12. Aabir Fauzan, $\zeta(5)$ *is irrational*, Preprint v1, September 17, 2026, Zenodo, doi:10.5281/zenodo.22826419, 2026.
13. Peter J. Forrester, *Meet Andréief, Bordeaux 1886, and Andreev, Kharkov 1882–83*, 2018.
14. David Niedbala Giraudin, *A new upper bound for the irrationality exponent of* $\zeta(2)$, 2026, Preprint, 2 October 2026.
15. F. Alberto Grünbaum and Luc Haine, *Bispectral Darboux transformations: an extension of the Krall polynomials*, Int. Math. Res. Not. (1997), no. 8, 359–392.
16. F. Alberto Grünbaum and Milen Yakimov, *Discrete bispectral Darboux transformations from Jacobi operators*, Pacific J. Math. **204** (2002), no. 2, 395–431.
17. R. K. Jha, *A finite irrationality measure for Catalan's constant*, Preprint v1, October 8, 2026, doi:10.5281/zenodo.23237269; Lean sources at `https://github.com/rohitkrjha/catalan-measure`, 2026.
18. Fredrik Johansson, *Arb: efficient arbitrary-precision midpoint-radius interval arithmetic*, IEEE Trans. Comput. **66** (2017), no. 8, 1281–1292.

19. Roelof Koekoek, Peter A. Lesky, and René F. Swarttouw, *Hypergeometric orthogonal polynomials and their q-analogues*, Springer Monographs in Mathematics, Springer, Berlin, 2010.
20. Christian Krattenthaler, *Advanced determinant calculus*, Sém. Lothar. Combin. **42** (1999), Art. B42q, 67 pp.
21. ———, *Advanced determinant calculus: a complement*, Linear Algebra Appl. **411** (2005), 68–166.
22. Christian Krattenthaler and Tanguy Rivoal, *On a linear form for Catalan's constant*, South East Asian J. Math. Math. Sci. **6** (2008), no. 2, 3–15.
23. Christian Krattenthaler and Wadim Zudilin, *Hypergeometry inspired by irrationality questions*, Kyushu J. Math. **73** (2019), no. 1, 189–203.
24. Li Lai, *The linear independence of* 1*,* $\zeta(3)$*, and* $\zeta(4)$, 2026, Preprint, 5 October 2026 (v2: 15 October 2026).
25. Sean Litsey, *An irrationality measure for Catalan's constant*, Hexagon preprint hexagon:2610.00187v4, https://hexagonmath.org/2610.00187v4, 2026.
26. Jingwen Liu, Kai Jiang, and Pingwen Zhang, *Irrationality exponents of logarithms of positive rational numbers*, 2026, Preprint, 7 October 2026.
27. Junqi Liu, Jujian Zhang, and Lihong Zhi, *A formal proof of the irrationality of* $\zeta(3)$ *in Lean 4*, 2025.
28. Karl Löwner, *Über monotone Matrixfunktionen*, Math. Z. **38** (1934), 177–216.
29. Raffaele Marcovecchio, *The Rhin–Viola method for* $\log 2$, Acta Arith. **139** (2009), no. 2, 147–184.
30. Yu. V. Nesterenko, *On the linear independence of numbers*, Vestnik Moskov. Univ. Ser. I Mat. Mekh. (1985), no. 1, 46–49, In Russian; English translation: Moscow Univ. Math. Bull. 40 (1985), no. 1, 69–74.
31. E. M. Nikishin and V. N. Sorokin, *Rational approximations and orthogonality*, Translations of Mathematical Monographs, vol. 92, American Mathematical Society, Providence, RI, 1991.
32. OpenAI, *Catalan's constant is irrational*, Preprint (September 24, 2026), with a Lean formalization, available at the original paper PDF and the Lean formalization (repository commit `fd4aeeb`, October 7, 2026), 2026.
33. ———, *The irrationality exponent of* π *is 2*, Preprint (September 24, 2026), https://github.com/openai/math/tree/main/preprints/The-irrationality-exponent-of-pi-is-2-September-24-2026, 2026.
34. Georges Rhin and Carlo Viola, *On a permutation group related to* $\zeta(2)$, Acta Arith. **77** (1996), no. 1, 23–56.
35. ———, *The group structure for* $\zeta(3)$, Acta Arith. **97** (2001), no. 3, 269–293.
36. Tanguy Rivoal and Wadim Zudilin, *Diophantine properties of numbers related to Catalan's constant*, Math. Ann. **326** (2003), no. 4, 705–721.
37. Edward B. Saff and Vilmos Totik, *Logarithmic potentials with external fields*, Grundlehren der mathematischen Wissenschaften, vol. 316, Springer, Berlin, 1997.
38. Vyacheslav Spiridonov and Alexei Zhedanov, *Spectral transformations, self-similar reductions and orthogonal polynomials*, J. Phys. A **30** (1997), no. 21, 7621–7637.
39. Zhi-Wei Sun, *Catalan's constant is irrational*, 2026.
40. The mathlib Community, *The Lean mathematical library*, Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020), 2020, pp. 367–381.
41. The PrimeNumberTheoremAnd contributors, *The PNT+ project: a Lean formalization of the prime number theorem and related results*, https://github.com/AlexKontorovich/PrimeNumberTheoremAnd, The vendored compatibility additions used here are recorded in the repository's `NOTICE` and `PORTING.md`.
42. Dane Wachs, *A note on a recent claimed proof of the irrationality of Catalan's constant*, 2026.

43. Doron Zeilberger and Wadim Zudilin, *The irrationality measure of π is at most* 7.103205334137..., Mosc. J. Comb. Number Theory **9** (2020), no. 4, 407–419.
44. Alexei Zhedanov, *Rational spectral transformations and orthogonal polynomials*, J. Comput. Appl. Math. **85** (1997), no. 1, 67–86.
45. Wadim Zudilin, *A few remarks on linear forms involving Catalan's constant*, Chebyshevskii Sb. **3** (2002), no. 2(4), 60–70, In Russian.
46. ———, *An Apéry-like difference equation for Catalan's constant*, Electron. J. Combin. **10** (2003), Research Paper 14.
47. ———, *A determinantal approach to irrationality*, Constr. Approx. **45** (2017), no. 2, 301–310.
48. ———, *Arithmetic of Catalan's constant and its relatives*, Abh. Math. Semin. Univ. Hambg. **89** (2019), no. 1, 45–53.

Email address: sean.litsey.math@gmail.com