

# Small RNA dynamics in cholinergic systems

DISSERTATION  
ZUR ERLANGUNG DES DOKTORGRADES  
DER NATURWISSENSCHAFTEN

VORGELEGT BEIM FACHBEREICH 14  
DER JOHANN WOLFGANG VON GOETHE-UNIVERSITÄT  
IN FRANKFURT AM MAIN

VON  
SEBASTIAN LOBENTANZER  
AUS SCHLÜCHTERN

FRANKFURT 2019  
(D30)

©2019 – SEBASTIAN LOBENTANZER  
ALL RIGHTS RESERVED.

## Small RNA dynamics in cholinergic systems

### ABSTRACT

Science still is very much in the discovery stage when it comes to transcriptional interactions, be it the long known workings of transcription factors or the recently discovered subtle fine-tuning of expression by small RNA, including microRNAs and transfer RNA fragments.

Quisque facilisis erat a dui. Nam malesuada ornare dolor. Cras gravida, diam sit amet rhoncus ornare, erat elit consectetur erat, id egestas pede nibh eget odio. Proin tincidunt, velit vel porta elementum, magna diam molestie sapien, non aliquet massa pede eu diam. Aliquam iaculis. Fusce et ipsum et nulla tristique facilisis. Donec eget sem sit amet ligula viverra gravida. Etiam vehicula urna vel turpis. Suspendisse sagittis ante a urna. Morbi a est quis orci consequat rutrum. Nullam egestas feugiat felis. Integer adipiscing semper ligula. Nunc molestie, nisl sit amet cursus convallis, sapien lectus pretium metus, vitae pretium enim wisi id lectus. Donec vestibulum. Etiam vel nibh. Nulla facilisi. Mauris pharetra. Donec augue. Fusce ultrices, neque id dignissim ultrices, tellus mauris dictum elit, vel lacinia enim metus eu nunc.

# Contents

<b>1</b>	<b>INTRODUCTION</b>	<b>1</b>
1.1	Cholinergic Systems . . . . .	1
1.2	Transcriptional Connectomics . . . . .	4
<b>2</b>	<b>MIRNET: CREATION OF A COMPREHENSIVE CONNECTOMICS DATABASE</b>	<b>11</b>
2.1	Implementation . . . . .	12
2.2	Materials . . . . .	14
2.3	Usage . . . . .	18
2.4	Statistical Approach to Transcriptional Connectomics . . . . .	21
<b>3</b>	<b>MICRORNA DYNAMICS IN CHOLINERGIC DIFFERENTIATION OF HUMAN NEURONAL CELLS</b>	<b>22</b>
3.1	Neuronal Transcriptomes - Background . . . . .	23
3.2	Cortical Single-Cell RNA Sequencing . . . . .	24
3.3	The Cellular Model . . . . .	26
3.4	Small RNA Sequencing and Differential Expression Analysis . . . . .	29
3.5	Network Generation . . . . .	29
3.6	The Cholinergic/Neurokinin Interface . . . . .	29
3.7	Application to Schizophrenia and Bipolar Disorder . . . . .	29
<b>4</b>	<b>DYNAMICS BETWEEN SMALL AND LARGE RNA IN THE BLOOD OF STROKE VICTIMS</b>	<b>30</b>
4.1	Background . . . . .	31
4.2	Cohort . . . . .	31
4.3	RNA Sequencing and Differential Expression Analysis . . . . .	31
4.4	tRF Homology . . . . .	31
4.5	WGCNA . . . . .	31
4.6	Co-correlation . . . . .	31
4.7	Networks . . . . .	31
4.8	Direct Interaction . . . . .	31
4.9	Feedforward Loops . . . . .	31
<b>5</b>	<b>DISCUSSION</b>	<b>32</b>
5.1	Methods . . . . .	32
<b>6</b>	<b>CONCLUSION</b>	<b>35</b>
	<b>REFERENCES</b>	<b>45</b>
<b>A</b>	<b>TRANSCRIPTION FACTOR REGULATORY CIRCUITS - TISSUE TYPES</b>	<b>46</b>

THIS IS THE DEDICATION.

*»Ever tried. Ever failed. No matter.  
Try again. Fail again.  
Fail better.«*

Simon Beckett

## Acknowledgments

THANKS ARE DUE, for every scientist is not only standing on the shoulders of giants, but also on those of very real persons, without whom this dissertation would not have been possible. consectetur adipiscing elit. Morbi commodo, ipsum sed pharetra gravida, orci magna rhoncus neque, id pulvinar odio lorem non turpis. Nullam sit amet enim. Suspendisse id velit vitae ligula volutpat condimentum. Aliquam erat volutpat. Sed quis velit. Nulla facilisi. Nulla libero. Vivamus pharetra posuere sapien. Nam consectetur. Sed aliquam, nunc eget euismod ullamcorper, lectus nunc ullamcorper orci, fermentum bibendum enim nibh eget ipsum. Donec porttitor ligula eu dolor. Maecenas vitae nulla consequat libero cursus venenatis. Nam magna enim, accumsan eu, blandit sed, blandit a, eros.

# Abbreviations

<b>ACh</b>	acetylcholine
<b>Ago</b>	argonaute (protein)
<b>API</b>	application programming interface
<b>CAGE</b>	5' cap analysis of gene expression
<b>CNS</b>	central nervous system
<b>DMEM</b>	Dulbecco's modified eagle medium
<b>FCS</b>	fetal calf serum
<b>gp130</b>	see IL6ST (gene)
<b>LA-N-2</b>	human neuroblastoma cell line (female)
<b>LA-N-5</b>	human neuroblastoma cell line (male)
<b>miRNA</b>	microRNA
<b>PBS</b>	phosphate buffered saline
<b>RT-qPCR</b>	real-time quantitative polymerase chain reaction
<b>RIN</b>	RNA integrity number (RNA quality measure)
<b>RISC</b>	RNA-induced silencing complex
<b>RPMI1640</b>	Roswell Park Memorial Institute medium
<b>RNA-seq</b>	RNA sequencing
<b>SQL</b>	structured query language
<b>TF</b>	transcription factor
<b>tiRNA</b>	transfer RNA half
<b>tRF</b>	transfer RNA fragment
<b>tRNA</b>	transfer RNA
<b>UTR</b>	untranslated region
<b>vAChT</b>	vesicular acetylcholine transporter (from SLC18A3 gene)

## GENES

**ACHE** acetylcholinesterase

**AIF1** allograft inflammatory factor 1 (microglia marker protein)

**CHAT** choline acetyltransferase

**CHRNA7** nicotinic acetylcholine receptor subunit  $\alpha 7$

**CNTF** ciliary neurotrophic factor

**CNTFR** ciliary neurotrophic factor receptor (soluble)

**GFAP** glial fibrillary acidic protein (central astrocyte marker)

**IL-6** interleukin 6

**IL6R** interleukin 6 receptor (soluble)

**IL6ST** interleukin 6 signal transducer (membrane bound; also known as gp130)

**JAK** janus kinase

**LIF** leukaemia inhibiting factor

**LIFR** leukaemia inhibiting factor receptor (soluble)

**NGF** nerve growth factor

**OLIG1** oligodendrocyte transcription factor 1

**RBFOX3** RNA-binding Fox-1 homolog 3 (neuronal marker gene; also known as NeuN)

**SLC18A3** vesicular acetylcholine transporter (official gene symbol)

**SST** somatostatin

**STAT** signal transducer and activator of transcription

**TYK** tyrosine kinase

**VIP** vasoactive intestinal peptide



*I know words. I have the best words.*

Donald Trump


# 1

## Introduction

### 1.1 CHOLINERGIC SYSTEMS

NARY A PROCESS IN THE MAMMALIAN BODY CAN COMMENCE WITHOUT PARTICIPATION OF CHOLINERGIC SYSTEMS. Acetylcholine (ACh) was chemically and pharmacologically described by Henry Dale more than 100 years ago<sup>1</sup>. A short time later, Otto Loewi published the first proof of signal transmission by small molecules: he transferred physiological solutions from electrically stimulated frog hearts to naive hearts and observed their reactions; the solution that provoked a parasympathetic response he proposed to contain a »vagus substance«<sup>2</sup>. Finally, in 1929, Henry Dale completed the picture by isolating acetylcholine from mammalian tissue and identifying it as the molecule responsible for the parasympathetic response<sup>3</sup>. Dale and Loewi's joint effort in »Discoveries in Chemical Transmission of Nerve Impulses« was rewarded with the Nobel Prize in Physiology or Medicine in 1936.

Although we have learned much about cholinergic systems in these past 100 years, our understanding of the mammalian nervous system still is fairly limited. Even when disregarding peripheral nervous systems, the complexity of cholinergic transmission is immense, and a myriad of functions have been attributed to cholinergic circuits in the central nervous system (CNS). Central nervous projections of cholinergic fibres were extensively mapped by M. Marsel Mesulam and others in the 1980s<sup>4</sup>, with a majority of long projection neurons originating in one of the eight cholinergic nuclei, Ch1-Ch8. While many of these anatomical structures have been filled with meaning by associations with both rudimentary as well as higher brain functions, there are still as many cholinergic pathways whose function is entirely unclear (Figure 1.1, from my first manuscript<sup>5</sup>).



**Figure 1.1:** This is a figure that floats inline and here is its caption.

This holds particularly true for the only recently discovered cortical cholinergic interneurons, which, in comparison to their projecting counterparts, are very small and numerically vastly inferior to other neuron types in the cortex. Thus, their detection and analysis with current methods is challenging.

### 1.1.1 CHOLINERGIC ASPECTS OF DISEASE

SINCE CHOLINERGIC SYSTEMS ARE INTEGRAL FOR A MYRIAD PHYSIOLOGICAL FUNCTIONS, they are naturally involved in aetiologies and phenotypes of a number of central and peripheral diseases. Of interest to this dissertation are the cholinergic aspects of degenerative and non-degenerative central nervous diseases (such as Alzheimer's Disease, Bipolar Disorder, Schizophrenia), ischemic conditions in stroke, and peripheral modulation of immune responses, particularly in the context of the aforementioned diseases.

## ALZHEIMER'S DISEASE

## SCHIZOPHRENIA AND BIPOLAR DISORDER

## STROKE

## IMMUNITY

### 1.1.2 NEUROKINES

In comparison to the widely studied cholinergic projection neurons originating in the basal forebrain (Ch1-Ch4) that are known to depend on a retrograde survival signal by means of nerve growth factor (NGF), trophic influences on other cholinergic populations such as the cortical interneurons are unclear. NGF was described by Rita Levi-Montalcini in the 1950s as the first known instance of trophic peptides required for the survival of sympathetic ganglia<sup>6</sup>, and the dependence of basal forebrain cholinergic neurons on retrograde NGF signalling was discovered in the 1980s<sup>7</sup>.

A second group of trophic peptides with cholinergic implications are the so-called »neurokines«; the name results from the fact that this particular subgroup of cytokines has been associated with neuronal function in the central and peripheral nervous systems. Most prominently they include the ciliary neurotrophic factor (CNTF), leukaemia inhibiting factor (LIF), and interleukin 6 (IL-6), all of which coincidentally have been known under the acronym CDF. In the end of the 1980s, two groups of scientists (McManaman<sup>8</sup> and Rao<sup>9</sup>) independently identified proteins in extracts of muscle fibre that induced a differentiation of neurons towards a cholinergic type, and thus termed these proteins »choline acetyltransferase development factor« or »cholinergic differentiation factor« (both abbreviated CDF). Only later, through sequencing of the peptides, it became known that they had in fact discovered two distinct neurokines, LIF (Rao) and CNTF (McManaman, personal communication). IL-6, on the other hand, is abbreviated CDF for an entirely different reason: in this case it is short for »CTL (cytolytic T lymphocyte) differentiation factor«.

CNTF, LIF, and IL-6 convey their impact on neuronal activity through a partly redundant neurokine receptor pathway. There are two basic types of neurokine receptors: soluble and transmembrane. The primary receptors for CNTF (CNTFR) and IL-6 (IL6R) are soluble proteins that are secreted into the extracellular space and, upon binding of a neurokine, bind to transmembrane receptor dimers on the cell surface. These transmembrane receptors are the LIF receptor (LIFR) and the »interleukin 6 signal transducer« (IL6ST, also known as gp130). Every neurokine has its preferred constellation of soluble and transmembrane receptors: CNTF binds to the soluble CNTF receptor and a dimer consisting of one gp130 and one LIFR protein; IL-6 binds to the soluble IL6R and a dimer of two units of gp130; LIF does not usually bind a soluble receptor but rather binds immediately to a dimer comprising one of each gp130 and LIFR; however, there is significant redundancy and crosstalk between those systems<sup>10,11</sup>.

All receptor constellations result in a main effect of activation of the JAK/STAT cascade. More

specifically, neurokinins can activate janus kinases (JAKs) 1 and 2 or the homologous tyrosine kinase (TYK) 2, and, successively, STAT («signal transducer and activator of transcription») isoforms 1, 3, 5A, and 5B, which then convey a multitude of cellular effects (e.g. in immunity or differentiation) through transcriptional activation. The STAT cascade is inherently self-limiting in that it usually leads to expression of transcription factors that serve as repressors of the STAT genes (XXX).

Neurokinins, particularly IL-6(?), might serve as a link between the immunological and cholinergic aspects of physiological or disease processes.

find a place

FIGURE  
STAT

## 1.2 TRANSCRIPTIONAL CONNECTOMICS

NO MATTER THEIR LOCATION, CHOLINERGIC NEURONS ARE DEFINED BY THEIR ABILITY TO SYNTHESISE ACh AND RELEASE IT TO NEIGHBOURING CELLS TO A CERTAIN EFFECT. To fulfil this task, two particular proteins are essential: the choline acetyltransferase (CHAT) to synthesise ACh from choline and acetyl-Coenzyme A, and the vesicular acetylcholine transporter (vAChT, official gene symbol *SLC18A3*), which concentrates ACh in vesicles for later release. A notable genetic feature connects these two proteins beyond their functional association: the small *SLC18A3* gene (2420 nucleobases) sits inside the first intron of the *CHAT* gene and thus is already included in its primary transcript, and is subject to the *CHAT* promoter. However, oftentimes the (mature) transcript levels of *CHAT* and *SLC18A3* mRNA seem to be independently regulated; from the perspective of the organism, the possibility of differential regulation between these two genes makes sense. Since *SLC18A3* does not possess its own promoter, this differential regulation has to be conveyed epigenetically.

This dissertation deals in large parts with approaches aiming to decipher these interactions; and while its primary topic revolves around cholinergic systems, the methods described in the following are designed to be applicable to the entirety of the genome/epigenome. Four particular types of cellular actors are subject of these methods and therefore will be briefly introduced: genes in the classical sense as the conveyors of cellular function by encoding for proteins; transcription factors (TFs), a subclass of protein coding genes that are able to regulate the expression of other genes; microRNAs (miRNAs), a class of small non-coding RNA that has been known for approximately two decades and is reasonably well described functionally and mechanistically; and transfer RNA fragments (tRFs), a second class of small non-coding RNA that has only recently been rediscovered and is significantly less well described regarding its functionality.

Where to put:

where goes  
this?

Distinguish neuronal connectomics from transcriptional connectomics

For the sake of simplicity, all descriptions of genomics and transcriptomics matters, genes, miRNAs, and tRFs in this dissertation are to be seen in the context of *Homo sapiens*, unless explicitly stated oth-

erwise.

### 1.2.1 TRANSCRIPTION FACTORS

TFs were among the first intracellular regulatory mechanisms to be discovered (the earliest article referencing the term »transcription factor« in its title on PubMed was published in 1972). TFs commonly translocate from the cytosol into the nucleus upon activation (often by phosphorylation), where they bind specific DNA sequences that usually range in size from 6 to 12 nucleobases. The regions containing these binding sites (about 100 - 1000 bases in size) determine the effect upon binding, which can be one of two main modes: either a promoter, leading to an increased activity of transcription in the downstream vicinity of the binding site, or a repressor, having the opposite effect.

There exists a vast body of knowledge on TF interactions with genes, mostly due to the long period of time since their discovery and the multitude of scientific publications, most often studying single TFs and their interactions with few genes, but cumulatively curated by several organisations. One of the currently largest curations of TF data, TRANSFAC, saw its original release in 1988. While these curation efforts can be extensive, they may present with serious bias towards particular TFs that might hold more scientific interest and thus are published far more frequently than others. Recently, comprehensive efforts have extended the available data significantly. Driven by the advent of RNA sequencing (RNA-seq), computational approaches have become able to not only comprehensively predict TF-gene interactions, but to do so in a highly tissue-specific manner (see 2.2.3). The human body is estimated to express up to 2600 distinct DNA-binding proteins, most of them presumed TFs<sup>12</sup>, although other studies give lower estimates.

### 1.2.2 MICRORNAS

THE FIRST ENDOGENOUS »SMALL RNA WITH ANTISENSE COMPLEMENTARITY« was described in 1993<sup>13</sup>, but miRNAs were only recognised as a distinct regulatory class of molecules in the early 2000s. They are typically between 18 and 22 nucleobase-long, single stranded RNA fragments, and their function is now largely undisputed: miRNAs serve as targeting molecules for a protein complex whose primary purpose is to repress translation of mRNA, and, in some cases, lead to mRNA degradation. The complex, therefore, is called RNA-induced silencing complex (RISC); central to its function is the family of argonaute (Ago) proteins, which can bind the mature miRNA and orient it for interaction with its targets. Guidance of RISC to the target mRNA is generally mediated via sequence complementarity between miRNA and the targeted mRNA. Specifically, a »seed« region, usually bases 2-8 on the miRNA, is mainly responsible for the interaction; in case of perfect complementarity of this seed to the mRNA sequence, the interaction is considered »canonical«.

In early miRNA research, the 3' untranslated region (UTR) of the mRNA was believed to contain most miRNA binding sites due to its greater accessibility (i.e., the lack of active ribosomes); however, cumulative recent reports suggest that binding inside the coding region of the mRNA is a regular occurrence. The rules governing miRNA binding to target sequences show considerable flexibility; a recent study shows about 30% of analysed relationships to be of »non-canonical« nature. In those cases, seed pairing with the mRNA is often imperfect. To ameliorate this loss of stability, compensation occurs typically by a secondary complementary structure after a small gap of non-complementary bases, leading to a »bridge«-type constellation. This flexibility has implications in applications involving targeting algorithms; those that consider only the seed region are more prone to false negatives than models that consider, for instance, the free energy of the whole molecule (see 2.2.4).

cite

cite

FIGURE?

miRNAs, similar to coding genes, are transcribed from loci on the genome, many inside introns or even exons of coding genes<sup>14</sup>. The primary transcript (primary miRNA or pri-miRNA) typically contains a hairpin-like structure that usually results in a double-stranded molecule because of internal complementarity, and can contain up to six mature miRNAs. This hairpin structure is recognised by the DGCR8 protein (DiGeorge Syndrome Critical Region 8, in invertebrates called »Pasha«); the complex then associates with the RNA-cleaving protein »Drosha«, which removes bases on the opposite side of the hairpin, creating a miRNA precursor (or pre-miRNA), which is subsequently exported from the nucleus by the shuttle protein Exportin-5. In a final step in the cytosol, the ribonuclease »Dicer« removes the loop joining the 3' and 5' arms of the pre-miRNA, resulting in a duplex of mature miRNA, about 20 nucleotides long. Initially, it was thought to contain only one active miRNA, resulting in a designation of »miRNA\*« for the complementary strand (commonly, the strand with lower expression). However, this notion has been disproven, and to reflect the possibility of both strands performing miRNA functions, nomenclature has changed to specify the arm of the pre-miRNA from which the mature form originates (suffix »-3p« for the 3' arm, and »-5p« for the 5' arm).

miRNAs are organised and curated by means of a periodically updated web-based platform, miRBase<sup>15</sup>. For *Homo sapiens*, miRBase v2.1 contains 2588 mature miRNAs from XXX precursors. Evolutionarily, the miRNA repertoire has grown from rodents to primates, resulting in a number of primate-specific miRNAs that may convey additional function. miRNA nomenclature is organised<sup>16</sup> in a way that assigns evolutionarily conserved miRNAs the same designation (number) in all species in which they are expressed. In their full names, a prefix stating the organism of origin is added; for example, hsa-miR-125b-5p (for *Homo sapiens*) and mmu-miR-125b-5p (for *Mus musculus*) share the same sequence and most of their functionalities.

Disease?

miRNA genes, in the same way as protein coding genes, can also be subject to promoters and repressors, adding another layer of expression control by TFs. However, these TF-miRNA relationships are far less well described than common coding gene interactions, because miRNAs due to their shortness are not amenable to many standard gene expression assay forms. Estimation of the number

of distinct targets of any one miRNA varies widely; however, it is generally accepted to not be less than several dozen targets per miRNA, and up to thousands of genes per miRNA (although that estimate might be overenthusiastic).

Prediction?

### 1.2.3 TRANSFER RNA FRAGMENTS

TRANSFER RNA (tRNA) BREAKDOWN PRODUCTS HAVE BEEN KNOWN FOR DECADES, with first descriptions in the 1970s; back then, they were associated with a higher turnover of tRNA in cancer cells<sup>17</sup>, and proposed as urine-based biomarkers for certain malignancies<sup>18</sup>. However, their genesis was attributed to random processes, and due to lacking molecular biology characterisation techniques, interest in those fragments quickly faded. It was not until recently that studies have shown tRNA to be a major source of stable expression of small noncoding RNA<sup>19,20</sup> in most mammalian tissues. Indeed, replicating the reports from the 1970s, tRNA breakdown products are the dominant form of small RNA in secreted fluids, such as urine and bile, and make up large parts of other bodily fluids as well<sup>21</sup>. They exist in two major forms: transfer RNA halves (tiRNAs), and the smaller transfer RNA fragments (tRFs). *from stroke paper* tiRNAs derive from either end of the tRNA, and are created by angiogenin cleavage at the anticodon loop<sup>22,23</sup>. Smaller fragments are derived from the 3' and 5' ends of the tRNA (3'-tRF/5'-tRF) or internal tRNA parts (i-tRF), respectively, and may incorporate into Ago protein complexes and act like miRNAs to suppress their targets<sup>24,25</sup>.

However, there is considerable controversy about the generalisation of tRF functions, as distinct publications discover very different and sometimes opposing mechanisms of action for their respective fragments. An obvious assumption is the miRNA-like functionality, at least for those tRFs that are in the length range of miRNAs. There have been several instances of tRFs proven to act as miRNA-like suppressors of translation in a RISC-associated manner<sup>25</sup>, and of Dicer playing a large part in their biogenesis<sup>19</sup>. There are even instances of small RNA molecules previously mislabeled miRNAs that have been discovered to actually be tRNA-derived, such as miR-1280<sup>26</sup>.

On the other hand, multiple groups have identified tRFs to function not in an antisense-complementary manner, but by homology aspects. A valine-derived tRF was found to regulate translation by competing with mRNA directly at the binding site at the initiation complex and thereby displacing the original mRNA, leading to its translational repression<sup>27</sup>. Others have found multiple classes of tRFs derived from glutamine, aspartate, glycine, and tyrosine tRNAs, that displace multiple oncogenic transcripts from an RNA-binding protein (YBX1), conveying tumor-suppressive activity<sup>28</sup>. Most counterintuitive is the recent finding of a tRF proven to bind to several ribosomal protein mRNAs and enhancing their translation, and, when specifically inhibited, leading to apoptosis in rapidly dividing cells<sup>29</sup>.

There is no consistent nomenclature yet to describe and organise tRFs, which are by nature more heterogeneous than miRNAs; considering their biogenesis, one tRNA molecule can be the origin



of several hundred distinct tRF molecules. Multiple approaches are common in current literature, most prominently tRFs are tied to the parent tRNA and the amino acid coded for by this tRNA. For example, the 22-nucleotide-long LeuCAG<sub>3'</sub> tRF (meaning: a fragment of 22 bases starting at the 3' end of the leucine-carrying tRNA with anticodon »CAG«) was shown to play an important role in regulating ribosome biogenesis<sup>29</sup>. Since there is no repository of the likes of miRBase yet, this approach can be cumbersome for replication purposes, and explicit statement of the exact sequence of each fragment is a must in publication. In fact, since the aforementioned paper does not mention the sequence explicitly, there exist 6 distinct possibilities of fragments fitting this description. While manageable on this small scale, this system prohibits efficient analysis of larger sets of tRFs that cannot be individually controlled. For this reason, the approach of Loher and colleagues<sup>30</sup> might be preferable: they propose the generation of a "license plate" based on the sequence of the fragment directly, composed of the prefix »tRF«, the length of the fragment, and a custom oligonucleotide string encoding (e.g., »B3« stands for »AAAGT«). This way, tRF names are unique and unmistakably linked to the sequence, nomenclature is species-independent, and tRNA origin can be quickly determined by sequence lookup.

Disease?

? Levels of tRFs may be modulated even more rapidly than levels of miRNAs, since tRNA molecules are very abundant in the cell and generation of mature tRFs requires only enzymatic degradation of tRNA but no de-novo transcription of the molecule in the nucleus (citation).

#### 1.2.4 NESTED MULTIMODAL TRANSCRIPTIONAL INTERACTIONS

##### - THE NEED FOR CONNECTOMICS

..., multiple levels of obstacles have to be overcome. The ultimate aim of any such approach is the generation of a robust model for the studied phenomenon; the theoretical and practical hurdles to be surmounted to reach this goal are many. The more we know about the functioning of these intertwined systems, the more we understand how much there is still to learn.

For example, only recently it has become clear how complex transcriptional regulation by means of TFs really is, and, incidentally, the two systems studied foremost in this dissertation (nerve and immune cells) are the two most transcriptionally complex systems in any mammal. Through study of comprehensive genomic information of 394 tissue types in approximately 1000 human primary cell, tissue, and culture samples (from the FANTOM5 consortium) it was estimated that the mean number of active TFs towards any given gene is highest in immune (12 TFs per gene) and nervous cells (10), and that any one TF in nervous and immune cells controls expression of a mean of 175 and 160 genes, respectively<sup>31</sup> (see also Section 2.2.3).

Similarly, it has been found that miRNAs, particularly in the nervous system, possess a much higher tissue specificity than coding genes, resulting in an expression landscape that varies widely between individual neuron types that are in close proximity in the brain. With the exception of single cell RNA-seq, no modern analysis method is capable of a resolution appropriate for accurate charac-



terisation of these expression patterns, resulting in extinction of the signal of miRNAs that are not expressed consistently across cell types (similar to »housekeeping« genes) because of statistical interference. Very recent studies show that miRNA-gene co-expression networks are tightly linked to cell types in the nervous system, and that groups of miRs as functional modules associate with particular phenotypes in developmental and mature states<sup>32</sup>. This functional association with cell phenotype was found in quality comparable to the expression patterns of TFs, yet in quantity conveys smaller impact and thus is thought to be a fine-tuning mechanism, subtle and precise in purpose.

Another aspect of the tissue specificity of CNS-associated miRNAs is the high likelihood of underrepresentation for those very specifically expressed miRNAs. Adding to the problem is the experimental bias towards rodent models when it comes to thorough studies of the CNS, where human or other primate samples are a rarity compared to rats or mice. Assessments of the numbers of yet unknown novel primate- and tissue specific miRNAs estimate their magnitude in the thousands<sup>33</sup>, resulting in an effective doubling of currently known miRNAs.

These high numbers of potentially interacting players present computational challenges: If estimating the number of expressed genes in a human cell at 20 000 (and the number of TFs at a low 1000), this makes for an estimated minimum of 200 000 »real« interactions in the possible  $C = \frac{1000!}{10!(1000-10)!} \cdot 20\,000$ , which practically equals infinity; this is without accounting for different tissue types or cell states (e.g., differentiation or disease). Similarly, the amount of mature miRNAs (2588 in miRBase v21) and their ability to target even more distinct transcripts than TFs with one single molecule present immense computational requirements for even listing all possible or actual relationships. An interaction table describing targeting of genes by miRNAs in one type of tissue has  $2588 \cdot 20\,000 \approx 50\,000\,000$  individual fields.

Combining all aspects of transcriptional interaction presents additional challenges. A simple model system to visualise (in only one type of cell) the interaction of TFs targeting genes, and of miRNAs targeting genes as well as TFs, contains about 20 000 genes (a subset of which of the size of about 2000 are TFs), 2588 mature miRNAs, and a total of  $2588 \cdot 20\,000 + 2000 \cdot 20\,000 \approx 90\,000\,000$  potential interactions. In standard application scenarios, such as the generation of an interaction network around a group of genes (e.g., the cholinergic genes), the processing requirements grow linearly with each added interaction partner, and exponentially with every regulatory layer that is added.

Practically, this information has to be provided, gathered, and integrated, which further multiplies the amount of storage and processing power required. miRWalk 2.0, a collection of miRNA interaction data, has collected 12 of the most popular miRNA-targeting prediction datasets, each of which has their strengths and weaknesses (see 2.2.4). Experimentally validated interactions (e.g. as collected in DIANA TarBase or miRTarBase) are gold standard, but far from comprehensive and strictly speaking only relevant for the cellular context in which the experiment was originally performed; there are also different evidence qualities to be accounted for, depending on the type of experiment performed. Ideally, all of these data are still accessible when performing the analysis, so a database created for this

example of  
standard  
interaction  
gene x miR  
x TF

purpose should be able to incorporate all this information without any data loss while still remaining feasible in terms of computation time as well as space and working memory requirements.

This dissertation will first describe the creation of such a database and what has been learned during its various stages, and then go on to apply the database to different biological problems from real world experiments, such as the cholinergic differentiation of male and female cultured cells, or the blood of stroke victims.

»Wir sehen in der Natur nie etwas als Einzelheit, sondern wir sehen alles in Verbindung mit etwas anderem, das vor ihm, neben ihm, hinter ihm, unter ihm und über ihm sich befindet.«

Johann Wolfgang von Goethe

# 2

## *miRNet*: Creation of a Comprehensive Connectomics Database

THE NEED FOR BIOINFORMATICAL SUPPORT IN CONNECTOMICS is immediately obvious from the sheer multitude of possible interactions between the participating factors. However, when I began working on this project (October 2015), there was no integrative database available for this purpose. Earlier that year, miRWalk 2.0 had been published, for the first time providing a relatively comprehensive source of predicted as well as experimentally validated miRNA targeting data<sup>34</sup> (see 1.2.2). One year later, Marbach's »regulatory circuits« were published<sup>31</sup>, enabling analysis of comprehensive TF-gene relationships in 394 human tissues (see Section 1.2.1). These collections (as well as the data they were derived from) are the basis of the database further called *miRNet*, the development of which will be described in the following chapter.

Since a large part of the scientific progress of this dissertation deals with practical problems of multimodal connectomics, I will begin by describing the infrastructure that makes effective computation of these problems possible. After this technical description of database structure and creation, I will explain the types and organisation of its content. The remainder of the chapter will then deal with the application of this infrastructure to real-world problems in transcriptional connectomics, and the statistical approaches suited to this special case.

## 2.1 IMPLEMENTATION

For any biological question to be asked in a bioinformatics setting, the effectiveness of the computational query determines the practicality of the approach. Because resources (i.e., processing power, storage, and working memory) are limited, the database that is queried should be organised in a way that facilitates retrieval of the desired information without excess processing of useless information. In the simplified case of only miRNAs interacting with genes in one direction (miRNA  $\rightarrow$  gene), this means retrieval of only those interactions relevant for the queried genes or miRNAs.

Traditional table-based approaches (also known as relational databases) such as SQL («Structured Query Language») cannot provide such an implementation, since individual entries for genes and miRNAs (rows and columns) have to be accessed in their entirety, whether there is a connection between gene and miRNA (1) or not (0). Additionally, adding layers to these interactions (e.g., distinct prediction algorithms, tissues, or the interaction between TFs and genes) require the addition of entire tables the same size as the database, which is detrimental to effective use of space; and more complex queries also necessitate the transfer of information between those distinct tables (in SQL typically via a JOIN command), which claims additional working memory and processing time. Overall, the so-called «many-to-many» organisation of data does not lend itself to representation in a relational database.

Figure to explain tables?

The actual performance is determined by the processing power of the machine it is running on and several structural properties, such as organisation, indexing, monotony, and of course the size of the database; therefore, an estimation of processing time for queries is bound to be inaccurate. However, processing times typically do not vary on the scale of orders of magnitude, and thus general estimations can be made. Well optimised SQL databases with a size of 5 to 10 GB on disk usually require tens of minutes if not hours to complete one single complex query<sup>35</sup>; *miRNet* in its current form takes up approximately 15 GB of storage. Since one analysis typically consists of several hundreds (and, in the case of permutation analyses, several hundreds of thousands) of these queries, processing times in SQL implementation are too long to be practically useful. (It seems important to note that, as of 2018, SQL also offers a graph-based organisation in addition to the traditional, relational layout. These two are separate systems, and not to be confused. The advantages of Neo4j as explained in the following should be seen from the perspective of 2015, when the database was established, and when there was no graph-based SQL implementation.)

### 2.1.1 NEO4J: A GRAPH-BASED INFRASTRUCTURE

To query and display biological data that are organised in a network-like structure (many-to-many), a database that lends itself to the efficient processing and storage of network data is optimal. «Neo4j» utilises a database structure that is built on the save and recall of data points in «nodes» and «edges», which represent entities (nodes) and relationships between those entities (edges); both nodes and

edges can have any number of attributes and a unique property called »type«, typically used to describe the class of the entry (such as »gene« or »miRNA«). This database organisation replicates the network-like structure of the biological data studied. Neo4j combines the network-like data structure with an efficient indexing system for quickly finding the entries queried for, and then »walks« along the edges of the nodes that have been found, thus only searching and returning the data that is relevant to the current query. Theoretically, this makes the database more likely to be efficient in the setting of transcriptional interactions, an estimation that turned out to be true.

Fig DB  
structure

Depending on the input, these queries can also be rather large; however, the main pitfall of tabular databases such as SQL is circumvented: there is no need to process entire rows or columns of the table to make sure that the query is satisfied in its entirety. This is particularly useful in a setting of sparse information. For example: only 30 of the 2588 miRNAs target a specific gene, which is common; a relational database, after finding the index of the queried gene, would have to search 2588 fields for 1/0; the graph database, on the other hand, has to execute only 30 searches (or, more accurately, 30 »walks« along the edges). In practice, even in the very first prototype implementations, this accelerated standard-case computations approximately thousand-fold, and was even able to accommodate advanced approaches in situations that had been inaccessible in the tabular implementation.

#### 2.1.2 HIGH-THROUGHPUT DATABASE GENERATION

Neo4j provides several API (»application programming interface«) possibilities in implementation. For the purpose of entering large amounts of data into the database at once, the Java implementation is superior to the other forms in that it provides a batch processing mode via its `BatchInserter` class. I thus wrote a custom Java program for the purpose of creating an initial state of the database from the largest set of data, the complete miRWalk 2.0 content with 12 algorithms and validated interactions. The downloaded data was organised in a plain text based file format, with one text file for each miRNA, totalling in size about 6 GB (for *H. sapiens*). The database was set up in a way that allows only one node for each individual miRNA and gene entered to avoid duplications, using the commands

Java and R  
description?  
Where?

- `createDeferredConstraint()`
- `assertPropertyIsUnique()`
- `createDeferredSchemaIndex()`

of the Neo4j Java package. This approach made sure to create only one node for each miRNA (type: MIR) and gene (type: GENE) in the data, which is essential for proper functioning of the database. Each of these nodes received several properties to store individual data, such as the various gene/miRNA identifiers, origin of data, and species.

Between those basic nodes, the batch insertion process created edges for each relationship that was found in the original data, assigning a type identifier to each edge detailing the origin of this interaction (type: name of the prediction algorithm or »VALIDATED« for experimental data). Thus, while the nodes for genes and miRNAs themselves are unique, an arbitrary number of relationships can exist between any two nodes, depending on how many interactions they share.

aggregation  
here?

### 2.1.3 MAINTENANCE AND QUALITY CONTROL

All additional datasets, such as the TF regulatory circuits or tRF targeting predictions, were entered into *miRNet* using the regular operation mode. Testing was also performed in regular operation, with manual as well as automated tests to assert the correct transfer of information from raw data to the graph database, and to avoid unpredictable behaviour. At times, conflicts had to be resolved manually, for instance when miRNA names conflicted between old »miRNA\*« and new »3p/5p« notation; all manual edits are documented in the code, which was published alongside my first manuscript<sup>5</sup>.

Except for the rapid import of large amounts of data in creation of a database, the Java implementation of Neo4j does not offer many advantages over the native R implementation, »RNeo4j«. Thus, after creation and a short period of experimentation with graphical user interfaces, I abandoned the Java program in favour of the more flexible R programming.

## 2.2 MATERIALS

All materials used in the creation of *miRNet* have been acquired from resources that are non-commercial, web-available, and open-source (in the case of code). All properties and relationships derived from this data were entered into *miRNet* as either nodes, properties of nodes, edges, or properties of edges.

### 2.2.1 GENE ANNOTATION

Even though »regular« protein coding genes have been known for a long time, there is no consensus yet about their nomenclature and organisation. Complicated by newly discovered functions and properties of phylogenetic nature, the scientific representation of the human genome is in constant flux. Several large organisations strive to provide a robust annotation of the human gene catalog, but also in many cases contradict one another. There are three nomenclature systems that are of high importance in modern genomics:

- The traditional naming system of acronyms and fantasy-names (e.g. CHAT), also occasionally called »gene symbol«, is still widely popular because of its accessibility to humans, but is also not particularly robust because of a high amount of synonyms with high confusion potential and instances of genes without names having to carry unwieldy systematic names.
- The American Center for Biotechnology Information (NCBI), a branch of the National Institute of Health (NIH), curates and hosts a multitude of biological and medical data, and for the

organisation of gene information uses its own systematic nomenclature termed »Entrez« ID. Entrez is a molecular biology database that integrates many aspects of biology and medicine in a gene-centered manner, and therefore Entrez IDs are useful to quickly connect a gene to its function, nucleotide sequence, or associated diseases. Entrez IDs are regular integers without additional characters.

- Akin to the NCBI effort, ENSEMBL is a project of the European Bioinformatics Institute (EBI) as part of the European Molecular Biology Laboratory (EMBL). Compared to the Entrez database, it is more focused on study and maintenance of the genome itself, and therefore has a more intricate nomenclature that allows for differentiation of, for example, genes and their various transcript isoforms (ENSEMBL IDs carry character prefixes for class identification, e.g., ENSG for genes, ENST for transcripts).

All of these are being used on a regular basis in many publications, and, often, they are used exclusively. As a result, the end user of the published data has to have access to all possible annotation forms, or, at least, a means to translate one into the other; often, this also introduces conflicts. For this reason, all ID types were entered into *miRNet* upon creation or during maintenance, for convenience and to minimise analysis prolongations due to conflict resolution.

### 2.2.2 MICRORNA ANNOTATION

miRBase provides a consistent annotation for miRNAs. Due to their relatively recent discovery, there still are major changes from version to version; the syntax, however, is stable. In addition to the miRNA »names« that are composed of species, the string »miR«, pre-miRNA designation number, and strand origin (not in all cases!), such as »hsa-miR-125b-5p«, miRBase provides IDs for pre-miRNA molecules (also called ancestors) termed »MIID«, and IDs for mature miRNA molecules termed »MIMAT«. However, in practice, these are rarely used.

### 2.2.3 TRANSCRIPTION FACTOR TARGETING

The FANTOM<sub>5</sub> project has applied 5' cap analysis of gene expression (CAGE) to a large number of human samples from diverse tissues to determine the accurate 5' ends of each transcript<sup>36</sup>. Knowledge of this fact enables accurate prediction of promoters likely to control a transcript's expression. Marbach and colleagues used this information in combination with detailed human gene expression data to derive a complex interaction network of TFs and genes (»regulatory circuits«), and in doing so aggregated samples with similar expression patterns and origins into 394 fictional tissues<sup>31</sup>. For every tissue, each TF was assigned transcriptional activities towards all genes that it supposedly targets (with the sum of all activities in any given tissue being 1); and the cumulative transcriptional activities towards any given gene correlate well with the actual gene expression in corresponding samples from an independent repository.

Even in its fifth iteration, FANTOM data is not entirely comprehensive, which came to my attention due to a cholinergic anomaly: the 5' CAGE peaks of the *CHAT* and *CHRNA7* (the nicotinic  $\alpha 7$  receptor subunit) genes in raw FANTOM<sub>5</sub> data do not pass the expression threshold, and therefore are not included in, e.g., Marbach's »regulatory circuits«. Both are critically important not only for neuronal cholinergic systems, but also for the non-neuronal aspect of immune processes. For instance, macrophages have been shown to produce ACh via CHAT(cite), and the  $\alpha 7$  homomeric ACh receptor conveys direct immune suppression by its expression on monocytes(cite). Paradoxically, the CAGE peak of *SLC18A3*, which lies in the first intron of *CHAT*, crosses the threshold and therefore is included in the data. Unfortunately, I was not able to remedy these circumstances even upon personal communication with Daniel Marbach (author of »regulatory circuits«) and Hideya Kawaji of the FANTOM<sub>5</sub> consortium, although the latter acknowledged the possibility of a gene annotation deficit leading to misattribution of the *CHAT* signal to *SLC18A3* due to the closeness of their 5' ends.

The entire collection of transcriptional activities in all tissues was downloaded from the project's web page<sup>31</sup>, and neuronal and immune tissues were entered into *miRNet*. The collected data comprises XX neuronal tissues and XX immune cell tissues (Appendix A), and XX TF-gene relationships in total.

#### 2.2.4 MICRORNA INTERACTIONS

The content of miRWalk 2.0 is freely available online<sup>37</sup>; however, there is no option of downloading the complete set. The targeting data thus was downloaded per miRNA with standard options for all 12 prediction algorithms (miRWalk, miRDB, PITA, MicroT4, miRMap, RNA22, miRanda, miRNAMap, RNAhybrid, miRBridge, PICTAR2, and TargetScan) in plain text format. For experimentally validated interactions, the main sources were DIANA TarBase<sup>38</sup> and miRTarBase<sup>39</sup>, both of which offer complete download options. As of 2019, the 3.0 version of miRWalk allows complete species downloads; however, the developers have abandoned their third party algorithm plurality reducing the number of available alternatives from 12 to 4, which can be considered a significant disadvantage:

mention  
custom  
crawler?

While sequence complementarity, particularly of the »seed«-region, is the primary paradigm of miRNA-mRNA interaction, prediction algorithms vary widely in their implementation, general purpose, and approach to interaction prediction (for a comprehensive review of approaches and rules, see<sup>40</sup>). A large group of available options utilise sequence conservation aspects to increase candidate viability (such as miRanda, PicTar, TargetScan, and microT4). Others, such as RNA22 and PITA, utilise biophysical aspects such as free energy of binding or the accessibility of target sites due to secondary RNA structures as prediction arguments. All of these approaches have their up- and down-sides, e.g. considering their general precision and sensitivity, or their adequate prediction of particular cases, such as multiple site targeting. Thus, it has been proposed to use a combination of complemen-



algorithm	hit frequency
RNAHYBRID	71.62%
MIRMAP	19.90%
MIRWALK	19.74%
TARGETSCAN	16.33%
RNA <sub>22</sub>	12.34%
MICROT <sub>4</sub>	11.81%
MIRANDA	10.65%
PITA	4.90%
MIRDB	1.17%
MIRNAMAP	0.75%
PICTAR <sub>2</sub>	0.62%
MIRBRIDGE	0.15%

**Table 2.1:** Prediction algorithms ordered by the fraction of all possible interactions they predict as being real (positive rate). Different algorithms display a wide variation of hit rates in the entirety of predicted interactions between any miRNA and gene. Red: excluded from analysis.

tary approaches instead of only one algorithm per analysis<sup>41</sup>. For this reason, I might have preferred the 2.0 version of miRWalk, even if 3.0 had been available at the time.

One advantage of the collection of all data in a quickly accessible database is the opportunity to compare the different approaches to target prediction. A statistical evaluation of the collected interaction data from miRWalk 2.0 showed vast differences in general prediction quantity (Table 2.1) as well as prediction accuracy and sensitivity when compared to the validated subset of data (Table 2.2). Since the ground truth is not known, this is an additional argument for the combination of multiple algorithms instead of the use of a single set. Apart from RNAhybrid and miRBridge, all algorithms presented reasonable base hit frequencies and increases in the validated test set. Therefore, the remaining 10 algorithms were included in *miRNet* targeting data. For ease of use, an additional relationship type was created from the aggregated single algorithm hits of any miRNA→gene relationship, with the sum of algorithms predicting the interaction as a score variable. This yields a theoretical score range from 1 to 10. To account for experimentally validated interactions, each miRNA→gene relationship that was supported by strong evidence of interaction was modified by addition of 10.5 score points (a half point for quick identification of a validated relationship). The resulting optimised graph contains XX miRNA→gene targeting relationships with a distinct score distribution (Figure XX).

### 2.2.5 DE-NOVO PREDICTION OF TRF TARGETING

Due to the recency of their (re-)discovery, no comprehensive interaction sources exist for transfer RNA fragments. There have been documented cases of miRNA-like behaviours of distinct RNA fragments<sup>19,25</sup>, justifying an attempt to predict interactions in a comprehensive manner. Of the available options for nucleotide interaction prediction algorithms, TargetScan<sup>42</sup> seems particularly suited

FIGURE:  
Histogram  
of score dis-  
tributions?

TFs not the  
only CHAT  
anomaly

algorithm	validated hit frequency	hit rate increase
PICTAR <sub>2</sub>	6.98%	1129.40%
MIRDB	9.80%	838.43%
MIRANDA	51.73%	485.94%
TARGETSCAN	70.63%	432.51%
MIRNAMAP	3.10%	410.95%
PITA	15.57%	317.20%
MICROT <sub>4</sub>	32.60%	276.10%
MIRMAP	53.86%	270.65%
MIRWALK	50.95%	258.15%
RNA <sub>22</sub>	22.51%	182.38%
RNAHYBRID	90.47%	126.32%
MIRBRIDGE	0.01%	0.00%

**Table 2.2:** Prediction algorithms ordered by their increase in true positive rate when considering only validated interactions. The hit rate increase when comparing experimentally validated interactions with the entire predicted data (Table 2.1) is also subject to strong variation. Hit rate increase is the increase of hit rate if only considering validated data as opposed to all predicted interactions. None of the studied algorithms unite a good precision (hit rate increase) and coverage (validated hit frequency).

for this task because it provides the option of evaluating the evolutionary conservation of target sites in the putatively targeted genes, thereby providing an additional layer of security: The sequence of 3' UTRs is evolutionarily less stable than the coding part of genes; thus, high conservation of the binding site might indicate evolutionary pressure to keep up the interaction with the fragment, making an actual function of the interaction more likely. TargetScan also presents with reasonable sensitivity and specificity as confirmed by an independent group<sup>43</sup>, and through an additional algorithm allows the attribution of a score based on the branch length (on the species tree) of conserved targeting<sup>44</sup>.

miRNA-like behaviour implies the existence of a region on the tRF similar to a miRNA »seed«, and TargetScan also expects a seed as input to its targeting algorithm. Since there has been no definitive answer to the question as to where the seed region in tRFs might be, it is safest to assume nothing and explore all possibilities, i.e., simulate every possible seed position for interaction discovery. For this purpose, all discovered sequences of tRFs were chopped into 7-base pieces (7mers), which is the length of miRNA seeds, and statistically improbable enough to appear in the genome at random; the average length of a human 3' UTR is 800 bases, so the probability of finding any 7mer randomly in any one 3' UTR is  $p = \frac{800}{4^7} = 0.049$ .

Describe  
Targetscan  
process

## 2.3 USAGE

Neo4j uses a language (called »Cypher«) akin to SQL, which utilises keyphrases to issue commands, but combines it with a semi-graphical syntax to account for the graph-based layout of the data. In the following, I will describe its basic usage and the advantages it provides in the matter of transcriptional connectomics. The basic »finder« function (similar to **SELECT** in SQL) is called **MATCH**

in Cypher, and, when combined with the semi-graphical syntax, can be used to identify nodes or more complex patterns in the database. The graphical syntax consists of two main building blocks that represent the basic types of data inside the database: nodes as regular brackets »( )« and edges between nodes as a construct of hyphens and box brackets, that can also have a direction indicated by the greater sign »( )-[ ]->( )«. To specify the elements to be found, attributes of nodes and/or edges can be filtered by using curly brackets in the node definition, or the **WHERE** clause. To be returned, elements need to be assigned arbitrary variable names:

---

**Listing 2.1: MATCH**

---

```
1 MATCH (gene:GENE {species: 'HSA'})
2 WHERE gene.name = 'CHAT'
3 RETURN gene
```

---

Query 2.1 identifies a node (arbitrarily designated »gene«) with type GENE (indicated by the colon), with attributes »species« (HSA, i.e. *H. sapiens*) and »name« (CHAT), and returns the node with all its attributes. Since the nodes of type GENE are restrained, there can only be one gene of species *H. sapiens* with this name in the database, and thus, only one data point will be returned. The graphical syntax further allows for pattern matching of, for instance, miRNA→gene relationships:

---

**Listing 2.2: Patterns**

---

```
1 MATCH (mir:MIR)-[rel:TARGETS]->(gene:GENE {species: 'HSA'})
2 WHERE gene.name = 'CHAT'
3 RETURN mir, rel, gene
```

---

Query 2.2, similar to query 2.1, starts by identifying the node of species HSA with the name CHAT, and proceeds to look for miRNA→gene relationship edges arriving at this node; the relationships have to be of the type TARGETS (the pre-aggregated score-based accumulation of targeting). As soon as no further edges are found, the process terminates and returns all found miRNAs (»mir«), relationships (»rel«), and genes (»gene«) in discrete form, including all their attributes, such as the ENSG and Entrez IDs, the MIMAT IDs for all found miRNAs, or the score value of their targeting relationship. In this query, since there is a constraint on genes, the only gene returned is *CHAT*. However, Cypher is not limited to filtering on unique attributes; it allows for query and return of as many data points as are needed. For example, if one is interested in all miRNA→gene interactions in the cholinergic system, the query might look as follows:

---

**Listing 2.3: Filtering**

---

```
1 MATCH (mir:MIR)-[rel:TARGETS]->(gene:GENE {species: 'HSA'})
2 WHERE gene.name IN {cholinergic_genes}
```

3 **RETURN** mir, rel, gene

---

The effectiveness of graph-based databases becomes clear in this approach: Query 2.3 is processed starting at a user-defined filter, the list of cholinergic genes as an input (containing *CHAT*, *SLC18A3*, cholinergic receptor genes, acetylcholinesterase, etc). In a first step, all nodes are found that fulfil the criteria: type GENE, from species *H. sapiens*, that are in the list of names given. Since the gene nodes are indexed, this only requires milliseconds. Then, through the connection of edges to these nodes, it finds all miRNA nodes that have a miRNA→gene relationship towards any of the cholinergic genes. By using the gene nodes as starting point, the query can end as soon as no other edges fulfilling these criteria are found on any of the nodes. In comparison, to satisfy this query in a relational database, the rows representing these cholinergic genes would have to be assessed in their entirety, not only in those columns that represent an extant relationship, thus prolonging execution.

The database then returns all miRNA→gene relationships in this set, representing the network of cholinergic miRNA regulators, including all of their attributes. The advantages of graph-based data do not end there; say one wants to return only »master« regulators of cholinergic systems, defined as miRNAs that target at least 4 of the genes in the cholinergic set. In a relational database, this would have to be done post-hoc, by aggregation of relationships and removal of any results that do not exceed this threshold. This requires storage of the entire result in memory, and additional computational steps that can be very taxing depending on the size of the result table. In Cypher, this can be done during the query (code comments indicated by »//« explain single steps):

**Listing 2.4:** Two-stage Filtering

---

```
1 MATCH (gene:GENE {species: 'HSA'})
2 WHERE gene.name IN {cholinergic_genes}
3 WITH gene //the found genes are used as input for the second query
4 MATCH (mir:MIR)-[rel:TARGETS]->(gene)
5 WHERE count(rel) >= 4
6 RETURN mir, rel, gene
```

---

Query 2.4 essentially proceeds in the same way as query 2.3 in that it identifies the gene nodes filtered for and looks for the miRNAs connected to those nodes by TARGETS-type relationships; however, in the second step (which is performed per gene node as returned by the **WITH** clause), it returns only those patterns that have at least 4 incoming miRNA→gene relationships. Query 2.4 only requires little additional processing compared to query 2.3, and thus does not require nearly as much time as the post-hoc filtering required in a relational database query. This filtering can be applied in many stages, and in many forms, such as sums, averages, maximum and minimum, or other combinations of arithmetic and logical classifiers. Additionally, the patterns can be extended to represent complex relationships inside the graph. For instance, the following query 2.5 was used

to find miRNAs that regulate any given gene in the database, and, simultaneously, affect TFs that are involved in regulation of this same gene (this type of interaction is called feedforward loop, see also Section 4.9).

**Listing 2.5:** Feedforward Loop Identification

---

```

1 MATCH (gene:GENE) //find gene
2 WHERE gene.id = ID //by identifier (Entrez)
3 WITH gene //use as input for next step
4 MATCH (tf:GENE {species: 'HSA', tf:TRUE})-[rel]->(gene)
5 //find TFs targetting that gene
6 WHERE type(rel) IN {tissue_types} //TFs only from specific tissues
7 //for instance, CNS cell types (Appendix A)
8 WITH gene, rel, tf //use as input for next step
9 MATCH (gene)<-[rel_m1:TARGETS]-(mir:MIR {species:
    'HSA'})-[rel_m2:TARGETS]->(tf)
10 //find miRNAs that target both gene and TF
11 WHERE rel_m1.score > 5 AND rel_m2.score > 5
12 //filter by minimum cumulative score
13 RETURN gene, tf, rel, type(r) AS tissue, mir, rel_m1, rel_m2

```

---

This analysis can be done in real time on the whole genome and miRnome and only takes seconds for one iteration, a performance unimaginable in a relational database approach.

## 2.4 STATISTICAL APPROACH TO TRANSCRIPTIONAL CONNECTOMICS

### Permutation

*There is no scientific study more vital to man  
than the study of his own brain. Our entire view  
of the universe depends on it.*


Francis Crick

# 3

## microRNA Dynamics in Cholinergic Differentiation of Human Neuronal Cells

EVEN THOUGH MUCH HAS BEEN ACHIEVED in the integrative study of miRNA control of gene expression, computational analysis of transcriptional interactions has not yet reached the level of sophistication needed for the accurate prediction of events inside mammalian cells. For this reason, combination of a bioinformatical assay with modern molecular biology methods can strengthen the message and reproducibility of any approach. The spectrum of processes worthy of study is as wide as modern biomedicine. Similarly, experimental models can span the entire repertoire available to a modern laboratory. The selection of a model adequate to the research question therefore is as important as diligent analysis and careful interpretation of results.

This chapter will discuss the current state of knowledge on brain transcriptomics, generally and in the specific case of cholinergic neurons in the CNS, and then go on to explain the steps I undertook to elucidate the relevance of small RNA processes in central cholinergic systems. First, my aim was to clarify co-expression patterns of central cholinergic neurons, which required analysis of transcriptome data in single-cell resolution. Based on this information, I selected two human models of cholinergic neuronal differentiation and established a differentiation protocol amenable to RNA extraction and successive molecular biology assays, most importantly, RNA-seq. The expression patterns so obtained were then used to perform bioinformatics analyses using the database introduced in Chapter 2, *miRNet*.



**Figure 3.1: Schematic display of the blood-brain-barrier.** The blood-brain-barrier surrounds virtually every capillary in the CNS. A: Astrocyte, B: Basal Membrane, E: Endothelial Cell, T: Tight Junction. Modified from Lobentanzer & Klein, 2019<sup>46</sup>.

### 3.1 NEURONAL TRANSCRIPTOMES - BACKGROUND

The mammalian brain requires a constant supply of oxygen and nutrients, because it does not provide storage for either. Though it only makes up approximately 2% of the entire human body mass, its energy expenditure is around 20% of the whole<sup>45</sup>. For this reason, there is no cubic micrometre of brain tissue (except for the ventricles) that is not infiltrated by multiple capillaries. Since the blood-brain-barrier is essentially provided by supporting glia cells surrounding all capillaries from the »inside« (see Fig. 3.1, adapted from my second publication<sup>46</sup>), neurons constitute only a minority of brain tissues (but burn two thirds of its energy).

Before the advent of single-cell RNA-seq, studies that endeavoured to clarify the transcriptional profiles of neurons used microarray technology, which was only recently succeeded by RNA-seq (also known as deep sequencing or next generation sequencing). For these methods, several cubic millimetres of brain tissue are required at the least; often, cubic centimetres are used. In contrast, the diameter of neuronal somata is usually in the micrometre range. Thus, the resolution of the method and the actual cellular resolution differ by a factor of approximately 10 000. Additionally, even among the neuronal population, there is considerable heterogeneity and transcriptomic plurality; single brain regions rarely consist of less than 30 different neuron types, tightly packed next to each other, each with their own transcriptional identity<sup>47,48,49,50</sup>. Newest studies, deciphering the murine nervous system by sequencing of 500 000 individual cells, show that neuron diversity is very similar regardless of brain region<sup>51</sup>. These circumstances hold true for any mammal, and most of our knowledge stems from the analysis of our favourite research animal, the mouse. In humans, the complexity is only exacerbated; in fact, the elevation in CNS transcriptional complexity might be the reason for our superior cognitive abilities(cite).


Cholinergic neurons always constitute a minority in any neuronal population, sometimes to extremes. Most tissues are dominated by few neuron types, such as pyramidal cells in the cortex; the most common neurotransmitter types are GABAergic (inhibitory) and glutamatergic (excitatory), each with several subtypes. It is estimated that more than 80% of cortical neurons are excitatory, and more than 90% of synapses release glutamate<sup>45</sup>. The striatum is fairly well-populated with rather large cholinergic interneurons, and the basal forebrain holds a large amount of (smaller) cholinergic projection neurons (compare Fig. 1.1). However, in transcriptomic analyses, these tissues are seldom used, for lack of scientific interest, or because they are notoriously hard to access (the basal forebrain is small and deeply imbedded in the midbrain). The cortex, particularly the neocortex, is most often the tissue of choice in these studies, due to its scientific interest and accessibility. Though it contains only a minuscule amount of cholinergic interneurons whose identity still is a matter of debate, several of the recent single-cell RNA-seq approaches have independently identified cholinergic interneurons in cortical regions (see Fig. 3.2).

All of the above taken into consideration, several limitations apply when it comes to the selection of a cellular model for the cholinergic processes we aim to understand. A multicellular model is prohibited by the novelty of the subject; possibilities include *in vivo* or *ex vivo* experimentation on rodents or human (3D-)cell culture with multiple cell types. While the former certainly is closest to reality, the diseases of interest display a noticeable lack of transferability from lower mammals to human(cite). The latter, on the other hand, introduces a complexity not well suited to the level of knowledge we possess about the studied processes; in addition, these models are very new, and thus, too many variables would be unknown. For these reasons, I selected two mono-cultures of human neuronal cells for my experiments. I chose to introduce a second cellular model during the experimental phase, because some of the studied diseases display a clear sexual dimorphism; the first experiments were performed on the female-originated LA-N-2 cell line, and later, to explore sex-related differences, the male-originated LA-N-5 cells were added.

### 3.2 CORTICAL SINGLE-CELL RNA SEQUENCING

To estimate the potential impact of the transcriptomes studied in our model on the diseases of interest, co-expression of the relevant genes has to be asserted. Similarly, if neurokines are to possess any relevance for cholinergic properties of central nervous cells, the cells in question would have to express molecular machinery required to receive neurokine signals. The advent of single-cell RNA-seq for the first time enables the resolution of gene expression on a cellular basis, and thus the disentangling of spatially close individual neuron types (and other, non-neuronal CNS cells); most of this information would be lost in RNA-seq performed on brain homogenate, even of a small biopsy. Differences in genes would be reduced to the universally expressed »housekeeping« genes, and in miRNAs, the situation would be worse, in parallel to their even more tissue-specific expression.





**Figure 3.2:** This is a figure that floats inline and here is its caption.

For this reason, I analysed all publicly available single-cell RNA-seq datasets relevant to our questions, which provide a detailed tally of transcriptional subtypes in the CNS. More specifically, all studies that were available at the time focused on some subsection of the cortex (visual or somatosensory) or the hippocampus. Additionally, the data provided by those studies was in some cases pre-aggregated to represent groups of single neurons with similar transcriptomes<sup>48,49</sup>; in other cases, every single neuron was represented<sup>47,50</sup>.

An important quality-related parameter of a single-cell RNA-seq experiment is the sequencing depth achieved per individual sequenced cell. Some datasets I screened do not provide sufficient depth to resolve genes with medium expression, which includes our primary cholinergic markers *CHAT* and *SLC18A3*. The datasets which did provide adequate sequencing depth were filtered for their expression of these markers, and additionally characterised by their expression of common markers for cell types to be expected in the CNS. This resulted in a collection of transcriptomes for potentially cholinergic cells in the sampled brain regions, and allowed an assessment of the functional type and gene co-expression patterns in central cholinergic cells (Fig. 3.2).

Most cells identified as cholinergic by this definition expressed the general neuronal marker *RBFOX3*, also known by its trivial name NeuN, but not the microglial marker *AIF1*. Few cells (or clusters of cells) expressed non-neuronal markers such as *GFAP* (astrocytes) or *OLIG1* (oligoden-

drocytes), hinting at sparse non-neuronal cholinergic functions. In agreement with my findings, cells or clusters identified as cholinergic by the authors of the respective studies had been classified as interneurons and co-expressed a number of known phenotypic neuronal markers, such as *somatostatin* (*SST*) and *vasoactive intestinal peptide* (*VIP*).

The identified cholinergic cells also revealed a constant co-expression with neurokinine-related genes, particularly the transmembrane neurokinine receptors LIFR and IL6ST, demonstrating a capacity to receive and process neurokinine signals. In contrast, the high affinity receptor for NGF, *NTRK1*, is not expressed by many of the cholinergic neurons, fundamentally distinguishing these cells from the basal forebrain cholinergic projection neurons.

Clusters?

### 3.3 THE CELLULAR MODEL

A prominent example of human neuronal cell culture used in the identification and elucidation of cholinergic processes is the immortalised neuroblastoma cell line SH-SY5Y<sup>52</sup>. Derived from its parent line SK-N-SH, an adrenergic neuroblastoma<sup>53</sup>, it expresses ample amounts of *ACHE*, and thus had become a work horse in many cholinergic fields, such as Alzheimer's Disease (which is treated with *ACHE* antagonists), pesticide development, and warfare(cite). However, in spite of its usefulness for processes involving *ACHE*, it turned out a less than optimal choice for the study of molecular events surrounding *CHAT* and *SLC18A3*, as it barely expresses both genes(cite), and cannot be coerced to elevate *CHAT* expression by the usual differentiation techniques (own experimentation, data not shown). Thus, for the questions asked in due course of this dissertation, SH-SY5Y does not qualify as adequate representation of a »cholinergic neuron«.

Following the elimination of SH-SY5Y as a suitable subject, I scoured the literature for candidates representing a cholinergic neuronal transcriptome, and found, among others, representatives of the LA-N neuroblastoma cell lines developed by R.C. Seeger around 1980<sup>54,55</sup>. Neuroblastoma is a form of neuronal cancer often affecting small children, and, consequentially, the two cell lines used in my experiments are immortalised biopsies of a 3 year old girl (LA-N-2<sup>54</sup>) and of a 4 month old boy (LA-N-5<sup>55</sup>). The decision to use LA-N-2 as my initial cellular model was influenced by three factors: it is well described in literature, although most studies had been published in the 1980s and 90s; it expresses a substantial amount of *CHAT* and *SLC18A3*; and it responds to neurokinine-mediated differentiation by assuming a neuronal morphology accompanied by further elevation of *CHAT* and *SLC18A3* expression. LA-N-5 was not nearly as well described as LA-N-2, but later added to the experimental roster because of the complementary sex and hints towards cholinergic differentiation under retinoic acid<sup>56</sup>.

### 3.3.1 CULTURE

LA-N cell culture is not as straightforward as many »go-to« human cell lines used in today's laboratories. Because LA-N-2 and LA-N-5 are very similar in this regard, they were treated similarly and I will describe the procedure for both. They have comparatively high duplication times, which can be lowered by using certain conditions that affect medium composition, nutrition, and CO<sub>2</sub> content. The cells were acquired at DSMZ (Braunschweig, Germany), which recommends keeping them in a 50:50 mixture of Dulbecco's modified eagle medium (DMEM) and Roswell Park Memorial Institute medium (RPMI1640), with 20% fetal calf serum (FCS) added. Sometimes, recommendations also suggest Leibovitz's L-15 medium, which is specifically designed for low CO<sub>2</sub> conditions, and others have suggested increased CO<sub>2</sub> levels inside the incubator. I found a combination of the DSMZ-recommended medium with 8% CO<sub>2</sub> atmosphere inside a 37°C incubator to accelerate growth to a degree that the cells could be split 1:3 to 1:4 in a weekly cycle. This protocol was used for all further experiments, which were performed between splits 2 to 8 after thawing of a batch from -80°C. All handling during maintenance and experimentation was performed under a laminar flow hood.

### 3.3.2 DIFFERENTIATION

Neuronal differentiation of neuroblastoma cell lines has been performed in many instances, utilising a wide variety of differentiation agents such as the very general retinoic acid or 5-bromo-uracil, or very specific reagents, such as the neurokines IL-6 and CNTF(cite). LA-N cells have also been described to react to a selection of these substances; however, due to the elevated interest in neurokine mechanisms, I chose to go ahead with a reagent from this group. Somewhat arbitrarily, I used CNTF, because upon my inquiry, James McManaman revealed in personal communication that the »*CHAT* development factor« that he had discovered<sup>8</sup> was, in fact, CNTF, which had never been published. Additionally, of the neurokines used for differentiation purposes, CNTF is best described in literature and easily acquired in dried form from Merck (Darmstadt, formerly SigmaAldrich, Germany).

LA-N cells are very sensitive to repeated temperature changes, which resulted in increased amounts of apoptotic cells following repeated removal from the incubator after seeding or medium changes during the experiment. For this reason, I chose to only add the differentiation reagent once, 24h after initial seeding of the cells into the wells of a 12-well-plate, and avoid further disturbances until the time of lysis. For the maximum duration of my experiments, 120h from seeding until lysis, the initially supplied medium was sufficient for survival.

Differentiation was performed in regular growth medium without changes in FCS content, and CNTF was added to the medium after an initial growth period of 24h. Cells were seeded into 12-well plates at approximately 200 000 cells/well, with 1 ml of growth medium. To determine the optimal amount of CNTF for differentiation, time-dose-curves were determined for both cell lines in a range from 1 ng ml<sup>-1</sup> to 100 ng ml<sup>-1</sup>. Here, I discovered the first pharmacological difference between LA-

N-2 and LA-N-5: the maximum of their cholinergic response to neurokinine stimulation (i.e., an elevation in *CHAT* and *SLC18A3* transcription) occurs at different concentrations of CNTF. While LA-N-2 cells respond most strongly to 100 ng ml<sup>-1</sup>, LA-N-5 cells show an »inverted u«-type dose response with a maximum around 10 ng ml<sup>-1</sup> CNTF (Fig. ??). James McManaman, who studied LA-N differentiation thoroughly in the 1990s<sup>57</sup>, believes both lines to respond in an »inverted u«-type manner (personal communication); thus, I assume that the LA-N-2 response would also diminish at CNTF concentrations significantly higher than 100 ng ml<sup>-1</sup>.

To study the small RNA dynamics following CNTF exposure of LA-N-2 and LA-N-5, I selected 4 time points at which the experiment was stopped and the cells were quickly lysed *in situ* to preserve total RNA in that state: for the quick, immediate-early-like phase, at 30 and 60 minutes after the addition of CNTF, and, for the long-term effects of differentiation, at 48 and 96 hours after the addition of CNTF. Each time point was controlled by a pseudo-treated culture (using pure water) from the same batch that had been seeded at the same time as the experimental group. In the final series used for the parallel sequencing of LA-N-2 and LA-N-5, all experiments were carried out in quadruplicates.

### 3.3.3 RNA ISOLATION

Total RNA was isolated using TRIzol (ThermoFisher Scientific), essentially as suggested by the manufacturer, with slight changes to the protocol to enrich small RNA species. The cells, growing in a monolayer in 12-well-plates, were cleared of medium, washed two times with 500 µl of cell culture grade phosphate buffered saline (PBS) (Gibco), and immediately suspended in 1 ml of TRIzol, pipetting up and down until visibly dissolved. After incubation for 5 minutes at room temperature, the samples were stored in -20°C for short periods of time until RNA isolation.

TRIzol-suspended lysates (1 ml) were added to RNA-separation centrifuge tubes (PhaseMaker Tubes, ThermoFisher Scientific), adding 200 µl of pure chloroform and mixing vigorously for 15 seconds. After two minutes, the mixture was centrifuged at 12 000 g for 15 minutes, and the upper, watery phase containing the RNA was extracted. This was mixed with approximately 2 parts of pure ethanol and incubated for 10 minutes at room temperature to precipitate the RNA. The precipitate was spun at 12 000 g for another 10 minutes, and the supernatant discarded. The pellet was washed with 85% ethanol (vortexed briefly) and centrifuged again for 5 minutes at 7500 g.

After the final centrifugation step, the samples were transferred to the laminar flow hood, and air dried after removal of most of the supernatant via micropipettors. The pellet was allowed to dry almost until completion and resuspended in 30 µl to 50 µl pure RNase-free water. RNA concentration was measured at a Nanodrop 2000 instrument (ThermoFisher Scientific) and samples were diluted to a uniform concentration of 100 ng µl<sup>-1</sup>. Finally, RNA samples were aliquoted according to later purpose and stored at -80°C.

RNA quality was determined by analysis on a 2100 Bioanalyzer instrument (Agilent) using a nano chip and 1  $\mu$ l of sample; RNA integrity number (RIN) was near optimal for all samples ( $> 9$ ).

### 3.4 SMALL RNA SEQUENCING AND DIFFERENTIAL EXPRESSION ANALYSIS

For the detection and analysis of small RNA species, RNA-seq is the current gold standard method. It allows the mapping of a comprehensive transcriptome and thus is vastly superior to small scale and consecutive methods such as real-time quantitative polymerase chain reaction (RT-qPCR). Assuming an adequate sequencing depth ( $\gtrsim 1$  M reads/sample), it allows a comparison of all expressed small RNA species at once, which is immensely helpful when dealing with processes on the combinatorial scale of miRNA regulation.

#### 3.4.1 SEQUENCING

For small RNA sequencing, the aliquoted samples were shipped on dry ice to the cooperating institute at the Hebrew University of Jerusalem, the Silberman Institute of Molecular Biology, the laboratory of Prof. Hermona Soreq. The RNA was prepared for sequencing using the NEBNext small RNA preparation kit for Illumina. Briefly, this includes ligation of sequencing adapters to both ends of all (single-stranded) RNA fragments in the sample, followed by a size selection gel electrophoresis. The band representing small RNA species on the gel was excised and prepared for loading onto the sequencing chip. After loading, the chip was sequenced in a NextSeq 550 series instrument (Illumina) with a read length of 80 bases, single-ended.

#### 3.4.2 SEQUENCE ALIGNMENT

#### 3.4.3 DIFFERENTIAL EXPRESSION ANALYSIS - R/DESeq2

#### 3.4.4 MICRORNA FAMILY ENRICHMENT

### 3.5 NETWORK GENERATION

### 3.6 THE CHOLINERGIC/NEUROKINE INTERFACE

### 3.7 APPLICATION TO SCHIZOPHRENIA AND BIPOLAR DISORDER

*I realized, "Oh my gosh! I'm having a stroke!" And the next thing my brain says to me is, Wow! This is so cool! How many brain scientists have the opportunity to study their own brain from the inside out?"*

Jill Bolte Taylor

# 4

## Dynamics Between Small and Large RNA in the Blood of Stroke Victims

LOREM IPSUM DOLOR SIT AMET, consectetur adipiscing elit. Morbi commodo, ipsum sed pharetra gravida, orci magna rhoncus neque, id pulvinar odio lorem non turpis. Nullam sit amet enim. Suspendisse id velit vitae ligula volutpat condimentum. Aliquam erat volutpat. Sed quis velit. Nulla facilisi. Nulla libero. Vivamus pharetra posuere sapien. Nam consectetur. Sed aliquam, nunc eget euismod ullamcorper, lectus nunc ullamcorper orci, fermentum bibendum enim nibh eget ipsum. Donec porttitor ligula eu dolor. Maecenas vitae nulla consequat libero cursus venenatis. Nam magna enim, accumsan eu, blandit sed, blandit a, eros.

- 4.1 BACKGROUND
- 4.2 COHORT
- 4.3 RNA SEQUENCING AND DIFFERENTIAL EXPRESSION ANALYSIS
- 4.4 TRF HOMOLOGY
- 4.5 WGCNA
- 4.6 CO-CORRELATION
- 4.7 NETWORKS
- 4.8 DIRECT INTERACTION
- 4.9 FEEDFORWARD LOOPS

*This is some random quote to start off the chapter.*

Firstname lastname

# 5

## Discussion

LOREM IPSUM DOLOR SIT AMET, consectetur adipiscing elit. Morbi commodo, ipsum sed pharetra gravida, orci magna rhoncus neque, id pulvinar odio lorem non turpis. Nullam sit amet enim. Suspendisse id velit vitae ligula volutpat condimentum. Aliquam erat volutpat. Sed quis velit. Nulla facilisi. Nulla libero. Vivamus pharetra posuere sapien. Nam consectetur. Sed aliquam, nunc eget euismod ullamcorper, lectus nunc ullamcorper orci, fermentum bibendum enim nibh eget ipsum. Donec porttitor ligula eu dolor. Maecenas vitae nulla consequat libero cursus venenatis. Nam magna enim, accumsan eu, blandit sed, blandit a, eros.

### 5.1 METHODS

Quisque facilisis erat a dui. Nam malesuada ornare dolor. Cras gravida, diam sit amet rhoncus ornare, erat elit consectetur erat, id egestas pede nibh eget odio. Proin tincidunt, velit vel porta elementum, magna diam molestie sapien, non aliquet massa pede eu diam. Aliquam iaculis. Fusce et ipsum et nulla tristique facilisis. Donec eget sem sit amet ligula viverra gravida. Etiam vehicula urna vel turpis. Suspendisse sagittis ante a urna. Morbi a est quis orci consequat rutrum. Nullam egestas feugiat felis. Integer adipiscing semper ligula. Nunc molestie, nisl sit amet cursus convallis, sapien lectus pretium metus, vitae pretium enim wisi id lectus. Donec vestibulum. Etiam vel nibh. Nulla facilisi. Mauris pharetra. Donec augue. Fusce ultrices, neque id dignissim ultrices, tellus mauris dictum elit, vel lacinia enim metus eu nunc.

Proin at eros non eros adipiscing mollis. Donec semper turpis sed diam. Sed consequat ligula nec tortor. Integer eget sem. Ut vitae enim eu est vehicula gravida. Morbi ipsum ipsum, porta nec,



tempor id, auctor vitae, purus. Pellentesque neque. Nulla luctus erat vitae libero. Integer nec enim. Phasellus aliquam enim et tortor. Quisque aliquet, quam elementum condimentum feugiat, tellus odio consectetur wisi, vel nonummy sem neque in elit. Curabitur eleifend wisi iaculis ipsum. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. In non velit non ligula laoreet ultrices. Praesent ultricies facilisis nisl. Vivamus luctus elit sit amet mi. Phasellus pellentesque, erat eget elementum volutpat, dolor nisl porta neque, vitae sodales ipsum nibh in ligula. Maecenas mattis pulvinar diam. Curabitur sed leo.

Nulla facilisi. In vel sem. Morbi id urna in diam dignissim feugiat. Proin molestie tortor eu velit. Aliquam erat volutpat. Nullam ultrices, diam tempus vulputate egestas, eros pede varius leo, sed imperdiet lectus est ornare odio. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Proin consectetur velit in dui. Phasellus wisi purus, interdum vitae, rutrum accumsan, viverra in, velit. Sed enim risus, congue non, tristique in, commodo eu, metus. Aenean tortor mi, imperdiet id, gravida eu, posuere eu, felis. Mauris sollicitudin, turpis in hendrerit sodales, lectus ipsum pellentesque ligula, sit amet scelerisque urna nibh ut arcu. Aliquam in lacus. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Nulla placerat aliquam wisi. Mauris viverra odio. Quisque fermentum pulvinar odio. Proin posuere est vitae ligula. Etiam euismod. Cras a eros.

Nunc auctor bibendum eros. Maecenas porta accumsan mauris. Etiam enim enim, elementum sed, bibendum quis, rhoncus non, metus. Fusce neque dolor, adipiscing sed, consectetur et, lacinia sit amet, quam. Suspendisse wisi quam, consectetur in, blandit sed, suscipit eu, eros. Etiam ligula enim, tempor ut, blandit nec, mollis eu, lectus. Nam cursus. Vivamus iaculis. Aenean risus purus, pharetra in, blandit quis, gravida a, turpis. Donec nisl. Aenean eget mi. Fusce mattis est id diam. Phasellus faucibus interdum sapien. Duis quis nunc. Sed enim.

Pellentesque vel dui sed orci faucibus iaculis. Suspendisse dictum magna id purus tincidunt rutrum. Nulla congue. Vivamus sit amet lorem posuere dui vulputate ornare. Phasellus mattis sollicitudin ligula. Duis dignissim felis et urna. Integer adipiscing congue metus. Nam pede. Etiam non wisi. Sed accumsan dolor ac augue. Pellentesque eget lectus. Aliquam nec dolor nec tellus ornare venenatis. Nullam blandit placerat sem. Curabitur quis ipsum. Mauris nisl tellus, aliquet eu, suscipit eu, ullamcorper quis, magna. Mauris elementum, pede at sodales vestibulum, nulla tortor congue massa, quis pellentesque odio dui id est. Cras faucibus augue.

Suspendisse vestibulum dignissim quam. Integer vel augue. Phasellus nulla purus, interdum ac, venenatis non, varius rutrum, leo. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Duis a eros. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos hymenaeos. Fusce magna mi, porttitor quis, convallis eget, sodales ac, urna. Phasellus luctus venenatis magna. Vivamus eget lacus. Nunc tincidunt convallis tortor. Duis eros mi, dictum vel, fringilla sit amet, fermentum id, sem. Phasellus nunc enim, faucibus ut, laoreet in, consequat id, metus. Vivamus dignissim. Cras lobortis tempor velit. Phasellus nec diam ac nisl lacinia tristique. Nullam nec metus id mi dictum dignissim. Nullam quis wisi non sem lobortis condimentum. Phasel-

lus pulvinar, nulla non aliquam eleifend, tortor wisi scelerisque felis, in sollicitudin arcu ante lacinia leo.

Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vestibulum tortor quam, feugiat vitae, ultricies eget, tempor sit amet, ante. Donec eu libero sit amet quam egestas semper. Aenean ultricies mi vitae est. Mauris placerat eleifend leo. Quisque sit amet est et sapien ullamcorper pharetra. Vestibulum erat wisi, condimentum sed, commodo vitae, ornare sit amet, wisi. Aenean fermentum, elit eget tincidunt condimentum, eros ipsum rutrum orci, sagittis tempus lacus enim ac dui. Donec non enim in turpis pulvinar facilisis. Ut felis.

Cras sed ante. Phasellus in massa. Curabitur dolor eros, gravida et, hendrerit ac, cursus non, massa. Aliquam lorem. In hac habitasse platea dictumst. Cras eu mauris. Quisque lacus. Donec ipsum. Nullam vitae sem at nunc pharetra ultricies. Vivamus elit eros, ullamcorper a, adipiscing sit amet, porttitor ut, nibh. Maecenas adipiscing mollis massa. Nunc ut dui eget nulla venenatis aliquet. Sed luctus posuere justo. Cras vehicula varius turpis. Vivamus eros metus, tristique sit amet, molestie dignissim, malesuada et, urna.

Cras dictum. Maecenas ut turpis. In vitae erat ac orci dignissim eleifend. Nunc quis justo. Sed vel ipsum in purus tincidunt pharetra. Sed pulvinar, felis id consectetur malesuada, enim nisl mattis elit, a facilisis tortor nibh quis leo. Sed augue lacus, pretium vitae, molestie eget, rhoncus quis, elit. Donec in augue. Fusce orci wisi, ornare id, mollis vel, lacinia vel, massa.

# 6

## Conclusion

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Morbi commodo, ipsum sed pharetra gravida, orci magna rhoncus neque, id pulvinar odio lorem non turpis. Nullam sit amet enim. Suspendisse id velit vitae ligula volutpat condimentum. Aliquam erat volutpat. Sed quis velit. Nulla facilisi. Nulla libero. Vivamus pharetra posuere sapien. Nam consectetur. Sed aliquam, nunc eget euismod ullamcorper, lectus nunc ullamcorper orci, fermentum bibendum enim nibh eget ipsum. Donec porttitor ligula eu dolor. Maecenas vitae nulla consequat libero cursus venenatis. Nam magna enim, accumsan eu, blandit sed, blandit a, eros.

Quisque facilisis erat a dui. Nam malesuada ornare dolor. Cras gravida, diam sit amet rhoncus ornare, erat elit consectetur erat, id egestas pede nibh eget odio. Proin tincidunt, velit vel porta elementum, magna diam molestie sapien, non aliquet massa pede eu diam. Aliquam iaculis. Fusce et ipsum et nulla tristique facilisis. Donec eget sem sit amet ligula viverra gravida. Etiam vehicula urna vel turpis. Suspendisse sagittis ante a urna. Morbi a est quis orci consequat rutrum. Nullam egestas feugiat felis. Integer adipiscing semper ligula. Nunc molestie, nisl sit amet cursus convallis, sapien lectus pretium metus, vitae pretium enim wisi id lectus. Donec vestibulum. Etiam vel nibh. Nulla facilisi. Mauris pharetra. Donec augue. Fusce ultrices, neque id dignissim ultrices, tellus mauris dictum elit, vel lacinia enim metus eu nunc.

Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vestibulum tortor quam, feugiat vitae, ultricies eget, tempor sit amet, ante. Donec eu libero sit amet quam egestas semper. Aenean ultricies mi vitae est. Mauris placerat eleifend leo. Quisque sit amet est et sapien ullamcorper pharetra. Vestibulum erat wisi, condimentum sed, commodo vitae, ornare sit amet, wisi. Aenean fermentum, elit eget tincidunt condimentum, eros ipsum rutrum orci, sagittis tempus lacus enim ac dui. Donec non enim in turpis pulvinar facilisis. Ut felis.

Cras sed ante. Phasellus in massa. Curabitur dolor eros, gravida et, hendrerit ac, cursus non, massa. Aliquam lorem. In hac habitasse platea dictumst. Cras eu mauris. Quisque lacus. Donec ipsum. Nullam vitae sem at nunc pharetra ultricies. Vivamus elit eros, ullamcorper a, adipiscing sit amet, porttitor ut, nibh. Maecenas adipiscing mollis massa. Nunc ut dui eget nulla venenatis aliquet. Sed luctus posuere justo. Cras vehicula varius turpis. Vivamus eros metus, tristique sit amet, molestie dignissim, malesuada et, urna.

Cras dictum. Maecenas ut turpis. In vitae erat ac orci dignissim eleifend. Nunc quis justo. Sed vel ipsum in purus tincidunt pharetra. Sed pulvinar, felis id consectetur malesuada, enim nisl mattis elit, a facilis tortor nibh quis leo. Sed augue lacus, pretium vitae, molestie eget, rhoncus quis, elit. Donec in augue. Fusce orci wisi, ornare id, mollis vel, lacinia vel, massa.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Morbi commodo, ipsum sed pharetra gravida, orci magna rhoncus neque, id pulvinar odio lorem non turpis. Nullam sit amet enim. Suspendisse id velit vitae ligula volutpat condimentum. Aliquam erat volutpat. Sed quis velit. Nulla facilisi. Nulla libero. Vivamus pharetra posuere sapien. Nam consectetur. Sed aliquam, nunc eget euismod ullamcorper, lectus nunc ullamcorper orci, fermentum bibendum enim nibh eget ipsum. Donec porttitor ligula eu dolor. Maecenas vitae nulla consequat libero cursus venenatis. Nam magna enim, accumsan eu, blandit sed, blandit a, eros.

Quisque facilis erat a dui. Nam malesuada ornare dolor. Cras gravida, diam sit amet rhoncus ornare, erat elit consectetur erat, id egestas pede nibh eget odio. Proin tincidunt, velit vel porta elementum, magna diam molestie sapien, non aliquet massa pede eu diam. Aliquam iaculis. Fusce et ipsum et nulla tristique facilis. Donec eget sem sit amet ligula viverra gravida. Etiam vehicula urna vel turpis. Suspendisse sagittis ante a urna. Morbi a est quis orci consequat rutrum. Nullam egestas feugiat felis. Integer adipiscing semper ligula. Nunc molestie, nisl sit amet cursus convallis, sapien lectus pretium metus, vitae pretium enim wisi id lectus. Donec vestibulum. Etiam vel nibh. Nulla facilisi. Mauris pharetra. Donec augue. Fusce ultrices, neque id dignissim ultrices, tellus mauris dictum elit, vel lacinia enim metus eu nunc.

# References

- [1] H. H. Dale. THE ACTION OF CERTAIN ESTERS AND ETHERS OF CHOLINE, AND THEIR RELATION TO MUSCARINE. *Journal of Pharmacology and Experimental Therapeutics*, 6(2), 1914.
- [2] O. Loewi. Über humorale Übertragbarkeit der Herznervenwirkung. *Pflügers Arch. Ges. Physiol.*, 189:239–242, 1921.
- [3] H H Dale and H W Dudley. THE PRESENCE OF HISTAMINE AND ACETYLCHOLINE IN THE SPLEEN OF THE OX AND THE HORSE. *J. Physiol.*, 68: 97, 1929. URL <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1402860/pdf/jphysiol01676-0019.pdf>.
- [4] M. M. Mesulam, E. J. Mufson, A. I. Levey, and B. H. Wainer. Atlas of cholinergic neurons in the forebrain and upper brainstem of the macaque based on monoclonal choline acetyltransferase immunohistochemistry and acetylcholinesterase histochemistry. *Neuroscience*, 12(3): 669–686, 1984. ISSN 03064522. doi: 10.1016/0306-4522(84)90163-5.
- [5] Sebastian Lobentanzer, Geula Hanin, Jochen Klein, and Hermona Soreq. Integrative Transcriptomics Reveals Sexually Dimorphic Control of the Cholinergic/Neurokinin Interface in Schizophrenia and Bipolar Disorder. *CellReports*, pages 1–19, 2019. ISSN 2211-1247. doi: 10.1016/j.celrep.2019.09.017. URL <https://doi.org/10.1016/j.celrep.2019.09.017>.
- [6] R. Levi-Montalcini and B. Booker. Destruction of the sympathetic ganglia in mammals by an antiserum to a nerve-growth protein. *Proceedings of the National Academy of Sciences*, 46(3):384–391, mar 1960. ISSN 0027-8424. doi: 10.1073/pnas.46.3.384. URL <http://www.ncbi.nlm.nih.gov/pubmed/16578497><http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC222845><http://www.pnas.org/cgi/doi/10.1073/pnas.46.3.384>.
- [7] F. Hefti. Nerve growth factor promotes survival of septal cholinergic neurons after fimbrial transections. *Journal of Neuroscience*, 6(8):2155–2162, 1986. ISSN 02706474.
- [8] James L. McManaman, Frances G. Crawford, S. Scott Stewart, and Stanley H. Appel. Purification of a Skeletal Muscle Polypeptide Which Stimulates Choline Acetyltransferase Activity in Cultured Spinal Cord Neurons. *Journal of Biological Chemistry*, 263(12):5890–5897, 1988.

- [9] M S Rao, P H Patterson, and S C Landis. Multiple cholinergic differentiation factors are present in footpad extracts: comparison with known cholinergic factors. *Development (Cambridge, England)*, 116(3):731–44, nov 1992. ISSN 0950-1991. URL <http://www.ncbi.nlm.nih.gov/pubmed/1289063>.
- [10] J. S. Rawlings. The JAK/STAT signaling pathway. *Journal of Cell Science*, 117(8):1281–1283, 2004. ISSN 0021-9533. doi: 10.1242/jcs.00963. URL <http://jcs.biologists.org/cgi/doi/10.1242/jcs.00963>.
- [11] Neil M. Nathanson. Regulation of neurokinin receptor signaling and trafficking. *Neurochemistry International*, 61(6):874–878, nov 2012. ISSN 01970186. doi: 10.1016/j.neuint.2012.01.018. URL <https://linkinghub.elsevier.com/retrieve/pii/S0197018612000307>.
- [12] M Madan Babu, Nicholas M Luscombe, L Aravind, Mark Gerstein, and Sarah A Teichmann. Structure and evolution of transcriptional regulatory networks. *Current Opinion in Structural Biology*, 14(3):283–291, jun 2004. ISSN 0959440X. doi: 10.1016/j.sbi.2004.05.004. URL <https://linkinghub.elsevier.com/retrieve/pii/S0959440X04000788>.
- [13] Rosalind C. Lee, Rhonda L. Feinbaum, and Victor Ambros. The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell*, 75(5):843–854, dec 1993. ISSN 00928674. doi: 10.1016/0092-8674(93)90529-Y. URL <https://linkinghub.elsevier.com/retrieve/pii/009286749390529Y>.
- [14] A. Rodriguez. Identification of Mammalian microRNA Host Genes and Transcription Units. *Genome Research*, 14(10a):1902–1910, sep 2004. ISSN 1088-9051. doi: 10.1101/gr.2722704. URL <http://www.genome.org/cgi/doi/10.1101/gr.2722704>.
- [15] Ana Kozomara, Maria Birgaoanu, and Sam Griffiths-Jones. miRBase: from microRNA sequences to function. *Nucleic Acids Research*, 47(D1):D155–D162, jan 2019. ISSN 0305-1048. doi: 10.1093/nar/gky1141. URL <https://academic.oup.com/nar/article/47/D1/D155/5179337>.
- [16] Victor Ambros, Bonnie Bartel, David P Bartel, Christopher B Burge, James C Carrington, Xuemei Chen, Gideon Dreyfuss, Sean R Eddy, Sam Griffiths-Jones, Mhairi Marshall, Marjori Matzke, Gary Ruvkun, and Thomas Tuschl. A uniform system for microRNA annotation. *RNA (New York, N.Y.)*, 9(3):277–9, mar 2003. ISSN 1355-8382. doi: 10.1261/rna.2183803. URL <http://www.ncbi.nlm.nih.gov/pubmed/12592000http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC1370393>.
- [17] Ernest Borek, B S Baliga, Charles W Gehrke, C W Kuo, Sidney Belman, Walter Troll, and T Phillip Waalkes. High Turnover Rate of Transfer RNA in Tumor Tissue. *CANCER*

- RESEARCH*, 37:3362–3366, 1977. URL <https://cancerres.aacrjournals.org/content/37/9/3362.full-text.pdf>.
- [18] John Speer, Charles W Gehrke, Kenneth C Kuo, T Phillip Waalkes, and Ernest Borek. tRNA breakdown products as markers for cancer. *Cancer*, 44(6): 2120–2123, dec 1979. ISSN 0008-543X. doi: 10.1002/1097-0142(197912)44:6<2120::AID-CNCR2820440623>3.0.CO;2-6. URL <http://www.ncbi.nlm.nih.gov/pubmed/509391>[http://doi.wiley.com/10.1002/1097-0142\(197912\)44:6<2120::AID-CNCR2820440623>3.0.CO;2-6](http://doi.wiley.com/10.1002/1097-0142(197912)44:6<2120::AID-CNCR2820440623>3.0.CO;2-6).
- [19] Christian Cole, Andrew Sobala, Cheng Lu, Shawn R Thatcher, Andrew Bowman, J. W.S. Brown, Pamela J Green, Geoffrey J Barton, and Gyorgy Hutvagner. Filtering of deep sequencing data reveals the existence of abundant Dicer-dependent small RNAs derived from tRNAs. *RNA*, 15(12):2147–2160, dec 2009. ISSN 1355-8382. doi: 10.1261/rna.1738409. URL <http://www.ncbi.nlm.nih.gov/pubmed/19850906><http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC2779667><http://rnajournal.cshlp.org/cgi/doi/10.1261/rna.1738409>.
- [20] Yong Sun Lee, Yoshiyuki Shibata, Ankit Malhotra, and Anindya Dutta. A novel class of small RNAs: tRNA-derived RNA fragments (tRFs). *Genes & development*, 23(22):2639–49, nov 2009. ISSN 1549-5477. doi: 10.1101/gad.1837609. URL <http://www.ncbi.nlm.nih.gov/pubmed/19933153><http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC2779758>.
- [21] Paula M Godoy, Nirav R Bhakta, Andrea J Barczak, Hakan Cakmak, Susan Fisher, Tippi C. MacKenzie, Tushar Patel, Richard W Price, James F Smith, Prescott G Woodruff, and David J Erle. Large Differences in Small RNA Composition Between Human Biofluids. *Cell Reports*, 25(5):1346–1358, oct 2018. ISSN 22111247. doi: 10.1016/j.celrep.2018.10.014. URL <https://doi.org/10.1016/j.celrep.2018.10.014><https://linkinghub.elsevier.com/retrieve/pii/S2211124718315778>.
- [22] Satoshi Yamasaki, Pavel Ivanov, Guo-fu Hu, and Paul Anderson. Angiogenin cleaves tRNA and promotes stress-induced translational repression. *The Journal of Cell Biology*, 185(1):35–42, apr 2009. ISSN 0021-9525. doi: 10.1083/JCB.200811106. URL <http://jcb.rupress.org/content/185/1/35.long>.
- [23] Pavel Ivanov, Mohamed M. Emara, Judit Villen, Steven P. Gygi, and Paul Anderson. Angiogenin-Induced tRNA Fragments Inhibit Translation Initiation. *Molecular Cell*, 43(4):613–623, aug 2011. ISSN 10972765. doi: 10.1016/j.molcel.

- 2011.06.022. URL <https://www.sciencedirect.com/science/article/pii/S1097276511005247?via=ihubhttps://linkinghub.elsevier.com/retrieve/pii/S1097276511005247>.
- [24] Alexander Maxwell Burroughs, Yoshinari Ando, Michiel Laurens de Hoon, Yasuhiro Tomaru, Harukazu Suzuki, Yoshihide Hayashizaki, and Carsten Olivier Daub. Deep-sequencing of human Argonaute-associated small RNAs provides insight into miRNA sorting and reveals Argonaute association with RNA fragments of diverse origin. *RNA Biology*, 8(1):158–177, jan 2011. ISSN 1547-6286. doi: 10.4161/rna.8.1.14300. URL <http://www.tandfonline.com/doi/abs/10.4161/rna.8.1.14300>.
- [25] Pankaj Kumar, Jordan Anaya, Suresh B Mudunuri, and Anindya Dutta. Meta-analysis of tRNA derived RNA fragments reveals that they are evolutionarily conserved and associate with AGO proteins to recognize specific RNA targets. *BMC Biology*, 12(1):78, dec 2014. ISSN 1741-7007. doi: 10.1186/s12915-014-0078-0. URL <http://www.ncbi.nlm.nih.gov/pubmed/25270025http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC4203973http://bmcbiol.biomedcentral.com/articles/10.1186/s12915-014-0078-0>.
- [26] Bingqing Huang, Huipeng Yang, Xixi Cheng, Dan Wang, Shuyu Fu, Wencui Shen, Qi Zhang, Lijuan Zhang, Zhenyi Xue, Yan Li, Yurong Da, Qing Yang, Zesong Li, Li Liu, Liang Qiao, Ying Kong, Zhi Yao, Peng Zhao, Min Li, and Rongxin Zhang. tRF/miR-1280 Suppresses Stem Cell-like Cells and Metastasis in Colorectal Cancer. *Cancer Research*, 77(12):3194–3206, jun 2017. ISSN 0008-5472. doi: 10.1158/0008-5472.CAN-16-3146. URL <http://cancerres.aacrjournals.org/http://cancerres.aacrjournals.org/lookup/doi/10.1158/0008-5472.CAN-16-3146>.
- [27] Jennifer Gebetsberger, Leander Wyss, Anna M Mleczko, Julia Reuther, and Norbert Polacek. A tRNA-derived fragment competes with mRNA for ribosome binding and regulates translation during stress. *RNA Biology*, 14(10):1364–1373, oct 2017. ISSN 1547-6286. doi: 10.1080/15476286.2016.1257470. URL <https://www.tandfonline.com/action/journalInformation?journalCode=krnb20https://www.tandfonline.com/doi/full/10.1080/15476286.2016.1257470>.
- [28] Hani Goodarzi, Xuhang Liu, Hoang C.B. Nguyen, Steven Zhang, Lisa Fish, and Sohail F. Tavazoie. Endogenous tRNA-Derived Fragments Suppress Breast Cancer Progression via YBX1 Displacement. *Cell*, 161(4):790–802, may 2015. ISSN 00928674. doi: 10.1016/j.cell.2015.02.053. URL <https://www.sciencedirect.com/science/article/pii/S0092867415003189?via=ihubhttps://linkinghub.elsevier.com/retrieve/pii/S0092867415003189>.



- [29] Hak Kyun Kim, Gabriele Fuchs, Shengchun Wang, Wei Wei, Yue Zhang, Hyesuk Park, Biswajoy Roy-Chaudhuri, Pan Li, Jianpeng Xu, Kirk Chu, Feijie Zhang, Mei-Sze Chua, Samuel So, Qiangfeng Cliff Zhang, Peter Sarnow, and Mark A. Kay. A transfer-RNA-derived small RNA regulates ribosome biogenesis. *Nature*, 552(7683):57, nov 2017. ISSN 0028-0836. doi: 10.1038/nature25005. URL <http://www.nature.com/doifinder/10.1038/nature25005>.
- [30] Phillipe Loher, Aristeidis G Telonis, and Isidore Rigoutsos. MINTmap: fast and exhaustive profiling of nuclear and mitochondrial tRNA fragments from short RNA-seq data. *Scientific Reports*, 7(1):41184, mar 2017. ISSN 2045-2322. doi: 10.1038/srep41184. URL <http://dx.doi.org/10.1038/srep41184><http://www.nature.com/articles/srep41184>.
- [31] Daniel Marbach, David Lamparter, Gerald Quon, Manolis Kellis, Zoltán Kutalik, and Sven Bergmann. Tissue-specific regulatory circuits reveal variable modular perturbations across complex diseases. *Nature Methods*, 13(4):366–370, apr 2016. ISSN 1548-7091. doi: 10.1038/nmeth.3799. URL <http://www.ncbi.nlm.nih.gov/pubmed/26950747><http://www.nature.com/articles/nmeth.3799><http://regulatorycircuits.org>.
- [32] Tomasz J Nowakowski, Neha Rani, Mahdi Golkaram, Hongjun R Zhou, Beatriz Alvarado, Kylie Huch, Jay A West, Anne Leyrat, Alex A Pollen, Arnold R Kriegstein, Linda R Petzold, and Kenneth S Kosik. Regulation of cell-type-specific transcriptomes by microRNA networks during human brain development. *Nature Neuroscience*, 21(12):1784–1792, dec 2018. ISSN 1097-6256. doi: 10.1038/s41593-018-0265-3. URL <http://www.ncbi.nlm.nih.gov/pubmed/30455455><http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC6312854><http://www.nature.com/articles/s41593-018-0265-3>.
- [33] Eric Londin, Phillipe Loher, Aristeidis G Telonis, Kevin Quann, Peter Clark, Yi Jing, Eleftheria Hatzimichael, Yohei Kirino, Shozo Honda, Michelle Lally, Bharat Ramratnam, Clay E. S. Comstock, Karen E. Knudsen, Leonard Gomella, George L. Spaeth, Lisa Hark, L. Jay Katz, Agnieszka Witkiewicz, Abdolmohamad Rostami, Sergio A. Jimenez, Michael A. Hollingsworth, Jen Jen Yeh, Chad A. Shaw, Steven E. McKenzie, Paul Bray, Peter T Nelson, Simona Zupo, Katrien Van Roosbroeck, Michael J Keating, George A Calin, Charles Yeo, Masaya Jimbo, Joseph Cozzitorto, Jonathan R. Brody, Kathleen Delgrosso, John S. Mattick, Paolo Fortina, and Isidore Rigoutsos. Analysis of 13 cell types reveals evidence for the expression of numerous novel primate- and tissue-specific microRNAs. *Proceedings of the National Academy of Sciences*, 112(10):E1106–E1115, mar 2015. ISSN 0027-8424. doi: 10.1073/pnas.1420955112. URL <http://www.pnas.org/lookup/doi/10.1073/pnas.1420955112>.

- [34] Harsh Dweep and Norbert Gretz. miRWalk2.0: a comprehensive atlas of microRNA-target interactions. *Nature Methods*, 12(8):697–697, 2015. ISSN 1548-7091. doi: 10.1038/nmeth.3485. URL <http://www.nature.com/doifinder/10.1038/nmeth.3485>.
- [35] Surajit Chaudhuri, Vivek Narasayya, and Ravi Ramamurthy. Estimating Progress of Execution for SQL Queries, jun 2004. URL <https://www.microsoft.com/en-us/research/publication/estimating-progress-of-execution-for-sql-queries/?from=https%3A%2F%2Fresearch.microsoft.com%2Fapps%2Fpubs%2F%3Fid%3D76556>.
- [36] Chung-chau Hon, Jordan A Ramilowski, Jayson Harshbarger, Nicolas Bertin, Owen J L Rackham, Julian Gough, Elena Denisenko, Sebastian Schmeier, Thomas M Poulsen, Jessica Severin, Marina Lizio, Hideya Kawaji, Takeya Kasukawa, Masayoshi Itoh, A Maxwell Burroughs, Shohei Noma, Sarah Djebali, Tanvir Alam, Soichi Kojima, Yukio Nakamura, Harukazu Suzuki, Carsten O Daub, Michiel J L De Hoon, Erik Arner, and Long Rnas. An atlas of human long non-coding RNAs with accurate 5' ends. *Nature Publishing Group*, 543(7644):199–204, 2017. ISSN 0028-0836. doi: 10.1038/nature21374. URL <http://dx.doi.org/10.1038/nature21374>.
- [37] H Dweep and N Gretz. miRWalk2 web page.
- [38] Dimitra Karagkouni, Maria D Paraskevopoulou, Serafeim Chatzopoulos, Ioannis S Vlachos, Spyros Tastsoglou, Ilias Kanellos, Dimitris Papadimitriou, Ioannis Kavakiotis, Sofia Maniou, Giorgos Skoufos, Thanasis Vergoulis, Theodore Dalamagas, and Artemis G Hatzigeorgiou. DIANA-TarBase v8: a decade-long collection of experimentally supported miRNA–gene interactions. *Nucleic Acids Research*, 46(D1):D239–D245, jan 2018. ISSN 0305-1048. doi: 10.1093/nar/gkx1141. URL <http://academic.oup.com/nar/article/46/D1/D239/4634010>.
- [39] Chih-Hung Chou, Sirjana Shrestha, Chi-Dung Yang, Nai-Wen Chang, Yu-Ling Lin, Kuang-Wen Liao, Wei-Chi Huang, Ting-Hsuan Sun, Siang-Jyun Tu, Wei-Hsiang Lee, Men-Yee Chiew, Chun-San Tai, Ting-Yen Wei, Tzi-Ren Tsai, Hsin-Tzu Huang, Chung-Yu Wang, Hsin-Yi Wu, Shu-Yi Ho, Pin-Rong Chen, Cheng-Hsun Chuang, Pei-Jung Hsieh, Yi-Shin Wu, Wen-Liang Chen, Meng-Ju Li, Yu-Chun Wu, Xin-Yi Huang, Fung Ling Ng, Waradee Buddhakosai, Pei-Chun Huang, Kuan-Chun Lan, Chia-Yen Huang, Shun-Long Weng, Yeong-Nan Cheng, Chao Liang, Wen-Lian Hsu, and Hsien-Da Huang. miRTarBase update 2018: a resource for experimentally validated microRNA-target interactions. *Nucleic Acids Research*, 46(D1):D296–D302, jan 2018. ISSN 0305-1048. doi: 10.1093/nar/gkx1067. URL <http://www.ncbi.nlm.nih.gov/pubmed/29126174><http://www.mirdb.org/>

<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC5753222>  
<http://academic.oup.com/nar/article/46/D1/D296/4595852>.

- [40] Dong Yue, Hui Liu, and Yufei Huang. Survey of Computational Algorithms for MicroRNA Target Prediction. *Current Genomics*, 10(7):478–492, nov 2009. ISSN 13892029. doi: 10.2174/138920209789208219. URL <http://www.ncbi.nlm.nih.gov/pubmed/20436875><http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC2808675><http://www.eurekaselect.com/openurl/content.php?genre=article&issn=1389-2029&volume=10&issue=7&spage=478>.
- [41] T M Witkos, E Koscińska, and W J Krzyżosiak. Practical Aspects of microRNA Target Prediction. *Current molecular medicine*, 11(2):93–109, 2011. ISSN 15665240. doi: 10.2174/156652411794859250.
- [42] Robin C Friedman, Kyle K.-H. Farh, Christopher B Burge, and David P Bartel. Most mammalian mRNAs are conserved targets of microRNAs. *Genome Research*, 19(1):92–105, oct 2009. ISSN 1088-9051. doi: 10.1101/gr.082701.108. URL <http://www.ncbi.nlm.nih.gov/pubmed/18955434><http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC2612969><http://genome.cshlp.org/cgi/doi/10.1101/gr.082701.108>.
- [43] Panagiotis Alexiou, Manolis Maragkakis, Giorgos L. Papadopoulos, Martin Reczko, and Artemis G. Hatzigeorgiou. Lost in translation: an assessment and perspective for computational microRNA target identification. *Bioinformatics*, 25(23):3049–3055, dec 2009. ISSN 1460-2059. doi: 10.1093/bioinformatics/btp565. URL <http://www.ncbi.nlm.nih.gov/pubmed/19789267><https://academic.oup.com/bioinformatics/article-lookup/doi/10.1093/bioinformatics/btp565>.
- [44] Vikram Agarwal, George W Bell, Jin-Wu Nam, and David P Bartel. Predicting effective microRNA target sites in mammalian mRNAs. *eLife*, 4, aug 2015. ISSN 2050-084X. doi: 10.7554/eLife.05005. URL <https://elifesciences.org/articles/05005>.
- [45] M. E. Raichle and D. A. Gusnard. Appraising the brain’s energy budget. *Proceedings of the National Academy of Sciences*, 99(16):10237–10239, aug 2002. ISSN 0027-8424. doi: 10.1073/pnas.172399499. URL <http://www.pnas.org/cgi/doi/10.1073/pnas.172399499>.
- [46] Sebastian Lobentanzer and Jochen Klein. Zentrales und Peripheres Nervensystem. In Wichmann and Fromme, editors, *Handbuch für Umweltmedizin*, chapter 11. Erg. lfg. edition, 2019.
- [47] Spyros Darmanis, Steven A. Sloan, Ye Zhang, Martin Enge, Christine Caneda, Lawrence M. Shuer, Melanie G. Hayden Gephart, Ben A. Barres, and Stephen R. Quake. A survey of human

- brain transcriptome diversity at the single cell level. *Proceedings of the National Academy of Sciences*, 112(23):201507125, 2015. ISSN 0027-8424. doi: 10.1073/pnas.1507125112. URL <http://www.pnas.org/content/112/23/7285.abstract>.
- [48] a. Zeisel, a. B. M. Machado, S. Codeluppi, P. Lonnerberg, G. La Manno, A. Jureus, S. Marques, H. Munguba, L. He, C. Betsholtz, C. Rolny, G. Castelo-Branco, J. Hjerling-Leffler, and S. Linnarsson. Cell types in the mouse cortex and hippocampus revealed by single-cell RNA-seq. *Science*, 347(6226):1138–42, 2015. ISSN 0036-8075. doi: 10.1126/science.aaa1934. URL <http://science.sciencemag.org.doelec.univ-lyon1.fr/content/347/6226/1138.abstract>.
- [49] Bosiljka Tasic, Vilas Menon, Thuc Nghi TN Nguyen, TK Tae Kyung Kim, Tim Jarsky, Zizhen Yao, Boaz BP Levi, Lucas T Gray, Staci A Sorensen, Tim Dolbeare, Darren Bertagnolli, Jeff Goldy, Nadiya Shapovalova, Sheana Parry, Changkyu CK Lee, Kimberly Smith, Amy Bernard, Linda Madisen, Susan M Sunkin, Michael Hawrylycz, Christof Koch, Hongkui Zeng, Zizhen Yao, Changkyu CK Lee, Nadiya Shapovalova, Sheana Parry, Linda Madisen, Susan M Sunkin, Michael Hawrylycz, Christof Koch, and Hongkui Zeng. Adult mouse cortical cell taxonomy revealed by single cell transcriptomics. *Nature Neuroscience*, advance on(January):1–37, 2016. ISSN 1097-6256. doi: 10.1038/nn.4216. URL <http://dx.doi.org/10.1038/nn.4216>.
- [50] Naomi Habib, Yinqing Li, Matthias Heidenreich, Lukasz Swiech, Inbal Avraham-Davidi, John J. Trombetta, Cynthia Hession, Feng Zhang, and Aviv Regev. Div-Seq: Single-nucleus RNA-Seq reveals dynamics of rare adult newborn neurons. *Science*, 353(6302):925–928, 2016. ISSN 0036-8075. doi: 10.1126/science.aad7038. URL <http://www.sciencemag.org/lookup/doi/10.1126/science.aad7038>.
- [51] Amit Zeisel, Hannah Hochgerner, Peter Lönnerberg, Anna Johnsson, Fatima Memic, Job van der Zwan, Martin Häring, Emelie Braun, Lars E Borm, Gioele La Manno, Simone Codeluppi, Alessandro Furlan, Kawai Lee, Nathan Skene, Kenneth D Harris, Jens Hjerling-Leffler, Ernest Arenas, Patrik Ernfors, Ulrika Marklund, and Sten Linnarsson. Molecular Architecture of the Mouse Nervous System. *Cell*, 174(4):999–1014.e22, aug 2018. ISSN 00928674. doi: 10.1016/j.cell.2018.06.021. URL <http://www.ncbi.nlm.nih.gov/pubmed/30096314><http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC6086934><https://linkinghub.elsevier.com/retrieve/pii/S009286741830789X>.
- [52] June L. Biedler, Suzanne Roffler-Tarlov, Melitta Schachner, and Lewis S. Freedman. Multiple Neurotransmitter Synthesis by Human Neuroblastoma Cell Lines and Clones. *Cancer Res.*, 38(11\_Part\_1):3751–3757, nov 1978. URL <http://cancerres.aacrjournals.org/content/38/11{ }Part{ }1/3751.short>.

- [53] June L. Biedler, Lawrence Helson, and Barbara A. Spengler. Morphology and Growth, Tumorigenicity, and Cytogenetics of Human Neuroblastoma Cells in Continuous Culture. *Cancer Research*, 33(11):2643–2652, 1973. ISSN 15387445.
- [54] Robert C. Seeger, Sylvia A. Rayner, Walter E. Laug, Harry B. Neustein, and William F. Benedict. Morphology, Growth, Chromosomal Pattern, and Fibrinolytic Activity of two New Human Neuroblastoma Cell Lines. *Cancer Research*, 37(5):1364–1371., 1977. ISSN 15387445.
- [55] R C Seeger, Y L Danon, S A Rayner, and F Hoover. Definition of a Thy-1 determinant on human neuroblastoma, glioma, sarcoma, and teratoma cells with a monoclonal antibody. *Journal of immunology (Baltimore, Md. : 1950)*, 128(2):983–9, feb 1982. ISSN 0022-1767. URL <http://www.ncbi.nlm.nih.gov/pubmed/6172518>.
- [56] David P Hill and Kent a Robertson. Characterization of the cholinergic neuronal differentiation of the human neuroblastoma cell line LA-N-5 after treatment with retinoic acid. *Developmental Brain Research*, 102(1):53–67, aug 1997. ISSN 01653806. doi: 10.1016/S0165-3806(97)00076-X. URL <http://www.ncbi.nlm.nih.gov/pubmed/9298234><https://linkinghub.elsevier.com/retrieve/pii/S016538069700076X>.
- [57] James L. McManaman and Frances G. Crawford. Skeletal Muscle Proteins Stimulate Cholinergic Differentiation of Human Neuroblastoma Cells. *Journal of neurochemistry*, pages 258–266, 1991.



# Transcription Factor Regulatory Circuits - Tissue Types