

CATEGORICAL RECONSTRUCTION THEORY

2025-07-04

Tony Zorman

tony.zorman@tu-dresden.de

The classical story

Tannaka duality studies algebraic structures through their categories of representations.

Groups, as people, will be known by their actions

Motto: groups are everywhere.

Groups, as people, will be known by their actions

Motto: G -sets are everywhere.

Groups, as people, will be known by their actions

Motto: G -sets are everywhere.

Problem: Understanding G -sets is difficult.

Groups, as people, will be known by their actions

Motto: G -sets are everywhere.

Problem: Understanding G -sets is difficult.

One solution: Linearise!

Representations of a group

$$\text{Sym}(\triangle)$$

Representations of a group


$$\text{Sym}\left(\triangle\right) = S_3$$

Representations of a group

$$\text{Sym}\left(\triangle\right) = S_3 \hookrightarrow \text{GL}_2(\mathbb{R})$$


Representations of a group

$$\text{Sym}\left(\triangle\right) = S_3 \hookrightarrow \text{GL}_2(\mathbb{R})$$



Representations of a group


$$\text{Sym}\left(\triangle\right) = S_3 \hookrightarrow \text{GL}_2(\mathbb{R})$$



Representations of a group


$$\text{Sym}\left(\triangle\right) = S_3 \hookrightarrow \text{GL}_2(\mathbb{R})$$

$$(1\,2\,3) \rightsquigarrow \begin{pmatrix} \cos \frac{2\pi}{3} & -\sin \frac{2\pi}{3} \\ \sin \frac{2\pi}{3} & \cos \frac{2\pi}{3} \end{pmatrix}$$



Representations of a group


$$\text{Sym}\left(\triangle\right) = S_3 \hookrightarrow \text{GL}_2(\mathbb{R})$$



$$(1\ 3) \rightsquigarrow \begin{pmatrix} \cos \frac{2\pi}{3} & \sin \frac{2\pi}{3} \\ \sin \frac{2\pi}{3} & -\cos \frac{2\pi}{3} \end{pmatrix}$$

Representations of a group

$$\text{Sym}\left(\triangle\right) = S_3 \hookrightarrow \text{GL}_2(\mathbb{R})$$



Representations of a group

$$\text{Sym}\left(\triangle\right) = S_3 \hookrightarrow \text{GL}_2(\mathbb{R})$$

A **representation** of a finite group G consists of a finite-dimensional \mathbb{k} -vector space V and a group homomorphism, called the **action**:

$$\rho: G \longrightarrow \text{GL}(V) := \text{Aut}(V).$$

Representations of a group

$$\text{Sym}\left(\triangle\right) = S_3 \hookrightarrow \text{GL}_2(\mathbb{R})$$

A **representation** of a finite group G consists of a **finite-dimensional** \mathbb{k} -vector space V and a group homomorphism, called the **action**:

$$\rho: G \longrightarrow \text{GL}_{\dim V}(\mathbb{k}).$$

Representations of a group

$$\mathrm{Sym}\left(\triangle\right) = S_3 \hookrightarrow \mathrm{GL}_2(\mathbb{R})$$

A **representation** of a finite group G consists of a finite-dimensional \mathbb{k} -vector space V and a group homomorphism, called the **action**:

$$\rho: G \longrightarrow \mathrm{GL}(V) := \mathrm{Aut}(V).$$

A **morphism of representations** from (V, ρ) to (W, σ) is a linear map $f: V \longrightarrow W$ such that $f\rho_g = \sigma_g f$ for all $g \in G$.

Special building blocks

Representations can be **tensor**ed:

$$(V, \rho) \otimes (W, \sigma) := (V \otimes_{\mathbb{k}} W, \rho \otimes \sigma),$$

Special building blocks

Representations can be tensored

$$V \otimes_{\mathbb{k}} W \cong \mathbb{k}\{v_i \otimes w_j \mid v_i \in B_V, w_j \in B_W\}$$

$$(V, \rho) \otimes (W, \sigma) := (V \otimes_{\mathbb{k}} W, \rho \otimes \sigma),$$

Special building blocks

Representations can be tensored:

$$(n, \rho) \otimes (m, \sigma) := (nm, \rho \otimes \sigma)$$

Special building blocks

Representations can be tensored:

$$(n, \rho) \otimes (m, \sigma) := (nm, \rho \otimes \sigma)$$

where


$$\rho \otimes \sigma: G \longrightarrow \mathrm{GL}_{nm}(\mathbb{k})$$

$$g \longmapsto \rho_g \otimes \sigma_g := \begin{pmatrix} (\rho_g)_{1,1}\sigma_g & \cdots & (\rho_g)_{1,n}\sigma_g \\ \vdots & \ddots & \vdots \\ (\rho_g)_{n,1}\sigma_g & \cdots & (\rho_g)_{n,n}\sigma_g \end{pmatrix}$$

Special building blocks

Representations can be tensored:

where

$$\rho \otimes \sigma: G \longrightarrow \mathrm{GL}_{nm}(\mathbb{K})$$

$$g \longmapsto \rho_g \otimes \sigma_g := \begin{pmatrix} (\rho_g)_{1,1}\sigma_g & \cdots & (\rho_g)_{1,n}\sigma_g \\ \vdots & \ddots & \vdots \\ (\rho_g)_{n,1}\sigma_g & \cdots & (\rho_g)_{n,n}\sigma_g \end{pmatrix}$$

Special building blocks

Representations can be tensored:

$$(n, \rho) \otimes (m, \sigma) := (nm, \rho \otimes \sigma)$$

where

$$\rho \otimes \sigma: G \longrightarrow \mathrm{GL}_{nm}(\mathbb{k})$$

$$g \longmapsto \rho_g \otimes \sigma_g := \begin{pmatrix} (\rho_g)_{1,1}\sigma_g & \cdots & (\rho_g)_{1,n}\sigma_g \\ \vdots & \ddots & \vdots \\ (\rho_g)_{n,1}\sigma_g & \cdots & (\rho_g)_{n,n}\sigma_g \end{pmatrix}$$

Duals of representations

If (V, ρ) is a representation, then (V^*, ρ^*) is one as well.

Duals of representations

If (V, ρ) is a representation, then (V^*, ρ^*) is one as well; with action

$$\begin{aligned}\rho^*: G &\longrightarrow \mathrm{GL}_n(\mathbb{K}) \\ g &\longmapsto (\rho_{g^{-1}})^T.\end{aligned}$$

Putting it all together

There is a distinguished map U :

$$\begin{aligned} U: \operatorname{rep}(G) &\longrightarrow \operatorname{vect} \\ (V, \rho) &\longmapsto V. \end{aligned}$$

Putting it all together

There is a dist

Representations of G

$$U: \text{rep}(G) \longrightarrow \text{vect}$$

$$(V, \rho) \longmapsto V.$$

Putting it all together

There is a dist

Representations of G

Finite-dimensional vector spaces

$$U: \text{rep}(G) \longrightarrow \text{vect}$$

$$(V, \rho) \longmapsto V.$$

Putting it all together

There is a distinguished map U :

$$\begin{aligned} U: \text{rep}(G) &\longrightarrow \text{vect} \\ (V, \rho) &\longmapsto V \\ ((V, \rho) \longrightarrow (W, \sigma)) &\longmapsto (f: V \longrightarrow W). \end{aligned}$$

Putting it all together

There is a distinguished map U :

$$\begin{aligned} U: \operatorname{rep}(G) &\longrightarrow \operatorname{vect} \\ (V, \rho) &\longmapsto V \\ ((V, \rho) \longrightarrow (W, \sigma)) &\longmapsto (f: V \longrightarrow W). \end{aligned}$$

Theorem (Tannaka reconstruction)

The monoidal automorphisms of U are, as a group, isomorphic to G .

Representations can be defined for any monoid M .

Representations can be defined for any monoid M . We can detect whether M is a group based on whether $\text{rep}(M)$ has duals.

Representations can be defined for any monoid M . We can detect whether M is a group based on whether $\text{rep}(M)$ has duals. This is called Tannaka duality.

Categorical generalisations

- [HZ23] with Sebastian Halbig: *Duality in Monoidal Categories*. arXiv: 2301.03545.
- [HZ24a] with Sebastian Halbig: *Diagrammatics for Comodule Monads*. In: Appl. Categ. Struct. 32 (2024).
- [HZ24b] with Sebastian Halbig: *Pivotality, twisted centres, and the anti-double of a Hopf monad*. In: Theory Appl. Categ. 41 (2024).
- [SZ24] with Mateusz Stroiński: *Reconstruction of module categories in the infinite and non-rigid settings*. arXiv: 2409.00793.
- [CSZ25] with Kevin Coulembier and Mateusz Stroiński: *Simple algebras and exact module categories*. arXiv: 2501.06629.
- [Zor25]: *Duoidal R-Matrices*. arXiv: 2503.03445.

- [HZ23] with Sebastian Halbig: *Duality in Monoidal Categories*. arXiv: 2301.03545.
- [HZ24a] with Sebastian Halbig: *Diagrammatics for Comodule Monads*. In: Appl. Categ. Struct. 32 (2024).
- [HZ24b] with Sebastian Halbig: *Pivotality, twisted centres, and the anti-double of a Hopf monad*. In: Theory Appl. Categ. 41 (2024).
- [SZ24] with Mateusz Stroiński: *Reconstruction of module categories in the infinite and non-rigid settings*. arXiv: 2409.00793.
- [CSZ25] with Kevin Coulembier and Mateusz Stroiński: *Simple algebras and exact module categories*. arXiv: 2501.06629.
- [Zor25]: *Duoidal R-Matrices*. arXiv: 2503.03445.

Categories and their ilk


- A **category** is a collection of objects and morphisms between them.

Categories and their ilk

- A **category** is a collection of objects and morphisms between them.
 - Sets, groups, vector spaces, representations, ...

Categories and their ilk

- A **category** is a collection of objects and morphisms between them.
 - Sets, groups, vector spaces, representations, ...
 - A monoid M is a category with a single object:



Categories and their ilk

- A **category** is a collection of objects and morphisms between them.
 - Sets, groups, vector spaces, representations, ...
 - A monoid M is a category with a single object.
 - Homomorphisms of categories are called **functors**.

Categories and their ilk

- A **category** is a collection of objects and morphisms between them.
 - Sets, groups, vector spaces, representations, ...
 - A monoid M is a category with a single object.
 - Homomorphisms of categories are called **functors**.
- A category is **monoidal** if there exists a way to tensor two objects.

Categories and their ilk

- A **category** is a collection of objects and morphisms between them.
 - Sets, groups, vector spaces, representations, ...
 - A monoid M is a category with a single object.
 - Homomorphisms of categories are called **functors**.
- A category is **monoidal** if there exists a way to tensor two objects.
 - Sets with \times , sets with \sqcup , groups with \times , vector spaces with $\otimes_{\mathbb{k}}$, ...

Categories and their ilk

- A **category** is a collection of objects and morphisms between them.
 - Sets, groups, vector spaces, representations, ...
 - A monoid M is a category with a single object.
 - Homomorphisms of categories are called **functors**.
- A category is **monoidal** if there exists a way to tensor two objects.
 - Sets with \times , sets with \sqcup , groups with \times , vector spaces with $\otimes_{\mathbb{k}}$, ...
- A monoidal category is **rigid** if there exists a dual for every object.

Categories and their ilk

- A **category** is a collection of objects and morphisms between them.
 - Sets, groups, vector spaces, representations, ...
 - A monoid M is a category with a single object.
 - Homomorphisms of categories are called **functors**.
- A category is **monoidal** if there exists a way to tensor two objects.
 - Sets with \times , sets with \sqcup , groups with \times , vector spaces with $\otimes_{\mathbb{k}}$, ...
- A monoidal category is **rigid** if there exists a dual for every object.
 - Finite-dimensional vector spaces, representations, finite-dimensional modules over a finite-dimensional Hopf algebra, ...

**Can we generalise Tannaka duality to
other algebraic structures and their
categories of representations?**

Given a (rigid monoidal) category \mathcal{C} , can one tell that this category is of the form $\text{rep}(A)$, for some algebraic object A ?

Duality theory for monads

Any monoidal category \mathcal{C} admits the notion of a **monoid**.

Duality theory for monads

Any monoidal category \mathcal{C} admits the notion of a **monoid**.

- (Set, \times) : ordinary monoids.
- (Group, \times) : abelian groups.
- $(\text{Vect}, \otimes_{\mathbb{k}})$: \mathbb{k} -algebras.

Duality theory for monads

Any monoidal category \mathcal{C} admits the notion of a **monoid**.

- (\mathbf{Set}, \times) : ordinary monoids.
- $(\mathbf{Vect}, \otimes_{\mathbb{k}})$: \mathbb{k} -algebras.
- (\mathbf{Group}, \times) : abelian groups.
- $(\mathbf{End}(\mathcal{C}), \circ)$: monads.

Duality theory for monads

Any monoidal category \mathcal{C} admits the notion of a **monoid**.

- (\mathbf{Set}, \times) : ordinary monoids.
- (\mathbf{Group}, \times) : abelian groups.
- $(\mathbf{Vect}, \otimes_{\mathbb{k}})$: \mathbb{k} -algebras.
- $(\mathbf{End}(\mathcal{C}), \circ)$: monads.

Theorem (Moerdijk, McCrudden, Bruguières–Virelizier)

Given a monad T on a rigid monoidal category \mathcal{C} , there exists a bijection

$$\left\{ \text{Hopf monad structures on } T \right\} \xleftrightarrow{\cong} \left\{ \begin{array}{l} \text{Rigid monoidal structure on } \mathbf{rep}(T) \\ \text{such that } U \text{ is strong monoidal} \end{array} \right\}$$

Duality theory for monads

Any monoidal category \mathcal{C} admits the notion of a **monoid**.

- (\mathbf{Set}, \times) : ordinary monoids.
- (\mathbf{Group}, \times) : abelian groups.
- $(\mathbf{Vect}, \otimes_{\mathbb{k}})$: \mathbb{k} -algebras.
- $(\mathbf{End}(\mathcal{C}), \circ)$: monads.

Theorem (Moerdijk, McCrudden, Bruguières–Virelizier)

Given a monad T on a rigid monoidal category \mathcal{C} , there exists a bijection

$$\left\{ \text{Hopf monad structures on } T \right\} \xleftrightarrow{\cong} \left\{ \begin{array}{l} \text{Rigid monoidal structure on } \mathbf{rep}(T) \\ \text{such that } U \text{ is strong monoidal} \end{array} \right\}$$


$$U: \mathbf{rep}(T) \longrightarrow \mathcal{C}$$

Representation categories

Monoidal categories \mathcal{C} are also monoids.

Representation categories

Monoidal categories \mathcal{C} are also monoids—a \mathcal{C} -**representation** consists of a category \mathcal{M} and a compatible **action**

$$\rho: \mathcal{C} \longrightarrow \text{End}(\mathcal{M}).$$

Representation categories

Monoidal categories \mathcal{C} are also monoids—a \mathcal{C} -**representation** consists of a category \mathcal{M} and a **compatible action**

$$\rho: \mathcal{C} \longrightarrow \text{End}(\mathcal{M}).$$

We, for example, require the existence of coherent isomorphisms

$$\alpha_{x,y,m}: \rho_{x \otimes y}(m) \xrightarrow{\sim} \rho_x(\rho_y(m)), \quad \text{for all } x, y \in \mathcal{C} \text{ and } m \in \mathcal{M}.$$

Representation categories

Monoidal categories \mathcal{C} are also monoids—a \mathcal{C} -**representation** consists of a category \mathcal{M} and a compatible **action**

$$\rho: \mathcal{C} \longrightarrow \text{End}(\mathcal{M}).$$

We, for example, require the existence of coherent isomorphisms

$$\alpha_{x,y}: \rho_{x \otimes y} \xrightarrow{\sim} \rho_x \rho_y, \quad \text{for all } x, y \in \mathcal{C}.$$

Representation categories


Monoidal categories \mathcal{C} are also monoids—a \mathcal{C} -**representation** consists of a category \mathcal{M} and a compatible **action**

$$\rho: \mathcal{C} \longrightarrow \text{End}(\mathcal{M}).$$

We, for example, require the existence of coherent isomorphisms

$$\alpha_{x,y}: \rho_{x \otimes y} \xrightarrow{\sim} \rho_x \rho_y,$$

for all $x, y \in \mathcal{C}$.


$$\rho_{xy} = \rho_x \rho_y$$

Representation categories

Monoidal categories \mathcal{C} are also monoids—a \mathcal{C} -**representation** consists of a category \mathcal{M} and a compatible **action**

$$\rho: \mathcal{C} \longrightarrow \text{End}(\mathcal{M}).$$

We, for example, require the existence of coherent isomorphisms

$$\alpha_{x,y}: \rho_{x \otimes y} \xrightarrow{\sim} \rho_x \rho_y, \quad \text{for all } x, y \in \mathcal{C}.$$

In representation categories, we can define **actions** and **coactions** of functors.

A Kelly-type theorem for representation categories

Theorem (Halbig-Z)

Given

$$\mathcal{C} \begin{array}{c} \xrightarrow{F} \\ \perp \\ \xleftarrow{U} \end{array} \mathcal{D}$$

A Kelly-type theorem for representation categories

Theorem (Halbig-Z)


Given

$$\mathcal{C} \begin{array}{c} \xrightarrow{F} \\ \xleftarrow[\mathcal{U}]{\perp \otimes} \end{array} \mathcal{D}$$

A Kelly-type theorem for representation categories

Theorem (Halbig-Z)

Given



A Kelly-type theorem for representation categories

Theorem (Halbig-Z)


Given

$$\begin{array}{ccc} \mathcal{M} & \begin{array}{c} \xrightarrow{G} \\ \xleftarrow{\perp} \\ \xleftarrow{V} \end{array} & \mathcal{N} \\ \downarrow \text{hook} & \begin{array}{c} \xrightarrow{F} \\ \xleftarrow{\perp \otimes} \\ \xleftarrow{U} \end{array} & \downarrow \text{hook} \\ \mathcal{C} & & \mathcal{D} \end{array}$$

A Kelly-type theorem for representation categories

Theorem (Halbig-Z)

Given



then G -coactions on F are in bijection with V -actions on U .

Tannaka duality for representation categories

Corollary (Halbig-Z)

Let T be a Hopf monad on the rigid monoidal category \mathcal{C} , and K a monad on a \mathcal{C} -representation \mathcal{M} . Then there is a bijection:

$$\left\{ T\text{-Coactions on } K \right\} \xleftrightarrow{\cong} \left\{ \begin{array}{l} \text{rep}(T)\text{-representations on } \text{rep}(K) \\ \text{such that } U^T \text{ acts strongly on } U^K \end{array} \right\}$$

Reconstruction up to Morita equivalence

Theorem (Stroiński–Z)

Let \mathcal{C} be an abelian monoidal category and (\mathcal{M}, ρ) an abelian representation of \mathcal{C} .

Reconstruction up to Morita equivalence

Theorem (Stroiński–Z)

Let \mathcal{C} be an abelian monoidal category and (\mathcal{M}, ρ) an abelian representation of \mathcal{C} . Suppose \mathcal{M} contains a “nice” object ℓ .

Reconstruction up to Morita equivalence

Theorem (Stroiński–Z)

Let \mathcal{C} be an abelian monoidal category and (\mathcal{M}, ρ) an abelian representation of \mathcal{C} . Suppose \mathcal{M} contains a “nice” object ℓ . Then there exists a monad Γ_ℓ such that there is an equivalence of representations

$$\mathcal{M} \simeq \text{rep}(\Gamma_\ell).$$

Reconstruction up to Morita equivalence

Theorem (Stroiński–Z)


Let \mathcal{C} be an abelian monoidal category and (\mathcal{M}, ρ) an abelian representation of \mathcal{C} . Suppose \mathcal{M} contains a “nice” object ℓ . Then there exists a monad Γ_ℓ such that there is an equivalence of representations

$$\mathcal{M} \simeq \text{rep}(\Gamma_\ell).$$

Furthermore, there is a bijection

$$\{(\mathcal{M}, \ell) \text{ as above}\} / \mathcal{M} \simeq \mathcal{N} \xleftrightarrow{\cong} \left\{ \begin{array}{l} \text{Right exact lax } \mathcal{C}\text{-rep} \\ \text{monads on } \mathcal{C} \end{array} \right\} / \text{rep}(T) \simeq \text{rep}(S)$$

Thanks!



Results from other articles

Theorem (Halbig–Z)

There exists a tensor representable category that is not rigid.

Theorem (Halbig–Z)

There are pivotal structures on the centre of a monoidal category that are not induced by the Picard heap of its twisted centre.

Theorem (Z)

The formalism of R-matrices for bialgebras extends to separately oplax monoidal monads in preduoidal categories.

Results from other articles

Theorem (Halbig–Z)

There exists a tensor representable category that is not rigid.

Theorem (Halbig–Z)

There are pivotal structures on the centre of a monoidal category that are not induced by the Picard heap of its twisted centre.

Theorem (Z)

The formalism of R-matrices for bialgebras extends to separately oplax monoidal monads in preduoidal categories.

Results from other articles

Theorem (Halbig–Z)

There exists a tensor representable category that is not rigid.

Theorem (Halbig–Z)

There are pivotal structures on the centre of a monoidal category that are not induced by the Picard heap of its twisted centre.

Theorem (Z)

The formalism of R-matrices for bialgebras extends to separately oplax monoidal monads in preduoidal categories.

References i

- [BV07] Alain Bruguières and Alexis Virelizier. **Hopf monads**. In: *Advances in Mathematics* 215.2 (2007), pp. 679–733. ISSN: 0001-8708. DOI: 10.1016/j.aim.2007.04.011.
- [CSZ25] Kevin Coulembier, Mateusz Stroiński, and Tony Zorman. **Simple algebras and exact module categories**. In: *arXiv e-prints* (2025). arXiv: 2501.06629 [math.RT].
- [HZ23] Sebastian Halbig and Tony Zorman. **Duality in Monoidal Categories**. In: *arXiv e-prints* (2023). arXiv: 2301.03545 [math.CT].
- [HZ24a] Sebastian Halbig and Tony Zorman. **Diagrammatics for Comodule Monads**. In: *Appl. Categ. Struct.* 32 (2024). Id/No 27, p. 17. ISSN: 0927-2852. DOI: 10.1017/CB09781139542333.
- [HZ24b] Sebastian Halbig and Tony Zorman. **Pivotality, twisted centres, and the anti-double of a Hopf monad**. English. In: *Theory Appl. Categ.* 41 (2024), pp. 86–149. ISSN: 1201-561X.
- [Kel74] G. M. Kelly. **Doctrinal adjunction**. English. *Category Sem., Proc., Sydney 1972/1973, Lect. Notes Math.* 420, 257–280 (1974). 1974.
- [McC02] Paddy McCrudden. **Opmonoidal monads**. In: *Theory and Applications of Categories* 10 (2002), No. 19, 469–485.
- [Moe02] Ieke Moerdijk. **Monads on tensor categories**. In: *J. Pure Appl. Algebra* 168.2-3 (2002). *Category theory 1999 (Coimbra)*, pp. 189–208. ISSN: 0022-4049. DOI: 10.1016/S0022-4049(01)00096-2.

References ii

- [SZ24] Mateusz Stroiński and Tony Zorman. **Reconstruction of module categories in the infinite and non-rigid settings**. In: *arXiv e-prints* (2024). arXiv: 2409.00793 [math.QA].
- [Tan38] Tadao Tannaka. **Über den Dualitätssatz der nichtkommutativen topologischen Gruppen**. German. In: *Tôhoku Math. J.* 45 (1938), pp. 1–12. ISSN: 0040-8735.
- [Zor25] Tony Zorman. **Duoidal R-Matrices**. In: *arXiv e-prints* (2025). arXiv: 2503.03445 [math.CT].