

RECONSTRUCTION FOR LAX MODULE MONADS

Foregoing the fibre functor in n easy steps!

Based on joint work with Matti Stroiński: [arXiv:2409.00793](https://arxiv.org/abs/2409.00793)



2025-07-17

Tony Zorman

tony.zorman@tu-dresden.de

Tannaka Reconstruction



Tannaka Reconstruction

$$\begin{array}{c} \mathcal{C}^T \\ u^T \left(\begin{array}{c} \downarrow \quad \uparrow \\ \mathcal{C} \end{array} \right) F^T \\ \mathcal{C} \\ \begin{array}{c} \uparrow \quad \downarrow \\ T \end{array} \end{array}$$

Tannaka Reconstruction

$$\begin{array}{c} \mathcal{C}^T \\ \begin{array}{c} \downarrow U^T \quad \uparrow F^T \end{array} \\ (\mathcal{C}, \otimes, 1) \\ \begin{array}{c} \curvearrowright \\ T \end{array} \end{array}$$

Tannaka Reconstruction

$$\begin{array}{c}
 (\mathcal{C}^T, \otimes, 1) \\
 (U^T, U_2^T, U_0^T) \left(\begin{array}{c} \downarrow \\ \uparrow \end{array} \right)_{F^T} \\
 (\mathcal{C}, \otimes, 1) \\
 \text{\scriptsize T} \curvearrowright
 \end{array}$$

Given a monoidal category \mathcal{C} , are all left \mathcal{C} -module categories equivalent to the modules of an algebra object in \mathcal{C} ?

Theorem ([Ost03; EGNO15])

Let \mathcal{C} be a finite tensor category and let \mathcal{M} be a finite abelian \mathcal{C} -module category, such that the evaluation functor $- \triangleright \ell : \mathcal{C} \longrightarrow \mathcal{M}$ is exact, for all $\ell \in \mathcal{M}$. Then there exists an algebra object $A \in \mathcal{C}$ such that there is an equivalence of \mathcal{C} -module categories $\text{mod}_{\mathcal{C}}(A) \simeq \mathcal{M}$.

Theorem ([Ost03; EGNO15])

Let \mathcal{C} be a *finite* tensor category and let \mathcal{M} be a *finite* abelian \mathcal{C} -module category, such that the evaluation functor $- \triangleright \ell : \mathcal{C} \longrightarrow \mathcal{M}$ is exact, for all $\ell \in \mathcal{M}$. Then there exists an algebra object $A \in \mathcal{C}$ such that there is an equivalence of \mathcal{C} -module categories $\text{mod}_{\mathcal{C}}(A) \simeq \mathcal{M}$.

Finiteness assumptions,

Theorem ([Ost03; EGNO15])

Let \mathcal{C} be a finite *tensor category* and let \mathcal{M} be a finite abelian \mathcal{C} -module category, such that the evaluation functor $- \triangleright \ell : \mathcal{C} \longrightarrow \mathcal{M}$ is *exact*, for all $\ell \in \mathcal{M}$. Then there exists an algebra object $A \in \mathcal{C}$ such that there is an equivalence of \mathcal{C} -module categories $\text{mod}_{\mathcal{C}}(A) \simeq \mathcal{M}$.

Finiteness assumptions, exactness assumptions,

Theorem ([Ost03; EGNO15])

Let \mathcal{C} be a finite *tensor category* and let \mathcal{M} be a finite abelian \mathcal{C} -module category, such that the evaluation functor $- \triangleright \ell : \mathcal{C} \longrightarrow \mathcal{M}$ is exact, for all $\ell \in \mathcal{M}$. Then there exists an algebra object $A \in \mathcal{C}$ such that there is an equivalence of \mathcal{C} -module categories $\text{mod}_{\mathcal{C}}(A) \simeq \mathcal{M}$.

Finiteness assumptions, exactness assumptions, and rigidity assumptions.

Theorem ([Ost03; EGNO15])

Let \mathcal{C} be a finite tensor category and let \mathcal{M} be a finite abelian \mathcal{C} -module category, such that the evaluation functor $- \triangleright \ell : \mathcal{C} \longrightarrow \mathcal{M}$ is exact, for all $\ell \in \mathcal{M}$. Then there exists an algebra object $A \in \mathcal{C}$ such that there is an equivalence of \mathcal{C} -module categories $\text{mod}_{\mathcal{C}}(A) \simeq \mathcal{M}$.

Finiteness assumptions, exactness assumptions, and **rigidity assumptions**.

Theorem ([Ost03; EGNO15])

Let \mathcal{C} be a finite tensor category and let \mathcal{M} be a finite abelian \mathcal{C} -module category, such that the evaluation functor $- \triangleright \ell : \mathcal{C} \longrightarrow \mathcal{M}$ is exact, for all $\ell \in \mathcal{M}$. Then there exists an algebra object $A \in \mathcal{C}$ such that there is an equivalence of \mathcal{C} -module categories $\text{mod}_{\mathcal{C}}(A) \simeq \mathcal{M}$.

Finiteness assumptions, exactness assumptions, and rigidity assumptions.

Proposition ([DSPS19])

In the absence of rigidity, there are finite abelian \mathcal{C} -module categories that cannot be realised as the modules of an algebra object in \mathcal{C} .

Given a monoidal category \mathcal{C} , are all left \mathcal{C} -module categories equivalent to the modules of a **monad** on \mathcal{C} ?

The setup

Let \mathcal{C} be a \mathbb{k} -linear monoidal, and \mathcal{M}, \mathcal{N} \mathbb{k} -linear left \mathcal{C} -module categories.

The setup

Let \mathcal{C} be a \mathbb{k} -linear monoidal, and \mathcal{M}, \mathcal{N} \mathbb{k} -linear left \mathcal{C} -module categories.

$$\otimes : \mathcal{C} \otimes_{\mathbb{k}} \mathcal{C} \longrightarrow \mathcal{C}, \qquad \triangleright : \mathcal{C} \otimes_{\mathbb{k}} \mathcal{M} \longrightarrow \mathcal{M}.$$

The setup

Let \mathcal{C} be a \mathbb{k} -linear monoidal, and \mathcal{M}, \mathcal{N} \mathbb{k} -linear left \mathcal{C} -module categories.

$$\otimes: \mathcal{C} \otimes_{\mathbb{k}} \mathcal{C} \longrightarrow \mathcal{C}, \quad \triangleright: \mathcal{C} \otimes_{\mathbb{k}} \mathcal{M} \longrightarrow \mathcal{M}.$$

such that for all $x, y, z \in \mathcal{C}$ and $m \in \mathcal{M}$, e.g.,

$$(x \otimes y) \otimes z \cong x \otimes (y \otimes z) \quad \text{and} \quad (x \otimes y) \triangleright m \cong x \triangleright (y \triangleright m).$$

The setup

Let \mathcal{C} be a \mathbb{k} -linear monoidal, and \mathcal{M}, \mathcal{N} \mathbb{k} -linear left \mathcal{C} -module categories.

$$\otimes: \mathcal{C} \otimes_{\mathbb{k}} \mathcal{C} \longrightarrow \mathcal{C}, \quad \triangleright: \mathcal{C} \otimes_{\mathbb{k}} \mathcal{M} \longrightarrow \mathcal{M}.$$

such that for all $x, y, z \in \mathcal{C}$ and $m \in \mathcal{M}$, e.g.,

$$(x \otimes y) \otimes z \cong x \otimes (y \otimes z) \quad \text{and} \quad (x \otimes y) \triangleright m \cong x \triangleright (y \triangleright m).$$

A functor $F: \mathcal{M} \longrightarrow \mathcal{N}$ is a *lax \mathcal{C} -module functor* if there exists an associative and unital natural transformation

$$F_2: - \triangleright F(=) \Longrightarrow F(- \triangleright =).$$

The setup

Let \mathcal{C} be a \mathbb{k} -linear monoidal, and \mathcal{M}, \mathcal{N} \mathbb{k} -linear left \mathcal{C} -module categories.

$$\otimes: \mathcal{C} \otimes_{\mathbb{k}} \mathcal{C} \longrightarrow \mathcal{C}, \quad \triangleright: \mathcal{C} \otimes_{\mathbb{k}} \mathcal{M} \longrightarrow \mathcal{M}.$$

such that for all $x, y, z \in \mathcal{C}$ and $m \in \mathcal{M}$, e.g.,

$$(x \otimes y) \otimes z \cong x \otimes (y \otimes z) \quad \text{and} \quad (x \otimes y) \triangleright m \cong x \triangleright (y \triangleright m).$$


A functor $F: \mathcal{M} \longrightarrow \mathcal{N}$ is a *lax \mathcal{C} -module functor* if there exists an associative and unital natural transformation

$$F_2: - \triangleright F(=) \Longrightarrow F(- \triangleright =).$$

The functor F is *oplax* if F_2 goes the other way, and *strong* if it is invertible.


Module categories as deloopings

The *delooping* $\mathbf{B}\mathcal{C}$ of a monoidal category \mathcal{C} is a 2-category with one object.



Module categories as deloopings

The *delooping* $\mathbf{B}\mathcal{M}$ of a **module** category \mathcal{M} is a 2-category with two objects.



The Yoneda lemma™

Proposition

There is an equivalence of \mathcal{C} -module categories

$$\mathcal{M} \simeq \mathrm{Str}\mathcal{C}\mathrm{Mod}(\mathcal{C}, \mathcal{M}),$$

The Yoneda lemmaTM

Proposition

There is an equivalence of \mathcal{C} -module categories

$$\mathcal{M} \simeq \mathrm{Str}\mathcal{C}\mathrm{Mod}(\mathcal{C}, \mathcal{M}), \quad m \longmapsto - \triangleright m,$$

The Yoneda lemma™

Proposition

There is an equivalence of \mathcal{C} -module categories

$$\mathcal{M} \simeq \text{Str}\mathcal{C}\text{Mod}(\mathcal{C}, \mathcal{M}), \quad m \longmapsto - \triangleright m, \quad F1 \longleftarrow F.$$

The Yoneda lemmaTM

Proposition

There is an equivalence of \mathcal{C} -module categories

$$\mathcal{M} \simeq \text{Str}\mathcal{C}\text{Mod}(\mathcal{C}, \mathcal{M}), \quad m \longmapsto - \triangleright m, \quad F1 \longleftarrow F.$$

In particular, $\mathcal{C}^{\text{rev}} \simeq \text{Str}\mathcal{C}\text{Mod}(\mathcal{C}, \mathcal{C})$.

The Yoneda lemma™

Proposition

There is an equivalence of \mathcal{C} -module categories

$$\mathcal{M} \simeq \mathrm{Str}\mathcal{C}\mathrm{Mod}(\mathcal{C}, \mathcal{M}), \quad m \longmapsto - \triangleright m, \quad F1 \longleftarrow F.$$

In particular, $\mathcal{C}^{\mathrm{rev}} \simeq \mathrm{Str}\mathcal{C}\mathrm{Mod}(\mathcal{C}, \mathcal{C})$.

Study cases in which $- \triangleright m$ admits a right adjoint.

Study cases in which $- \triangleright m$ admits a right adjoint. The resulting monad canonically has a lax \mathcal{C} -module structure.

Study cases in which $- \triangleright m$ admits a right adjoint. The resulting monad canonically has a lax \mathcal{C} -module structure. Then apply Beck's monadicity theorem.

To Kelly and Beck

Theorem (Kelly's doctrinal adjunctions, [Kel74])

Given an adjunction $F: \mathcal{C} \rightleftarrows \mathcal{D} : U$ between monoidal categories, oplax monoidal structures on F are in bijective correspondence with lax monoidal structures on U .

To Kelly and Beck

Theorem (Kelly's doctrinal adjunctions, [Kel74])

*Given an adjunction $F: \mathcal{C} \rightleftarrows \mathcal{D} : U$ between **monoidal** categories, oplax **monoidal** structures on F are in bijective correspondence with lax **monoidal** structures on U .*

To Kelly and Beck

Theorem (Kelly's doctrinal adjunctions, [Kel74; HZ24])

*Given an adjunction $F: \mathcal{M} \rightleftarrows \mathcal{N} : U$ between \mathcal{C} -*module* categories, oplax \mathcal{C} -*module* structures on F are in bijective correspondence with lax \mathcal{C} -*module* structures on U .*

To Kelly and Beck

Theorem (Kelly's doctrinal adjunctions, [Kel74; HZ24])

Given an adjunction $F: \mathcal{M} \rightleftarrows \mathcal{N} : U$ between \mathcal{C} -module categories, oplax \mathcal{C} -module structures on F are in bijective correspondence with lax \mathcal{C} -module structures on U .

To Kelly and Beck

Theorem (Kelly's doctrinal adjunctions, [Kel74; HZ24])

Given an adjunction $F: \mathcal{M} \rightleftarrows \mathcal{N} : U$ between \mathcal{C} -module categories, oplax \mathcal{C} -module structures on F are in bijective correspondence with lax \mathcal{C} -module structures on U .

Theorem (Beck's monadicity theorem)

An adjunction $F: \mathcal{C} \rightleftarrows \mathcal{D} : U$ is monadic if and only if U is conservative, \mathcal{D} has coequalisers of U -split pairs, and U preserves them.

To Kelly and Beck

Theorem (Kelly's doctrinal adjunctions, [Kel74; HZ24])

Given an adjunction $F: \mathcal{M} \rightleftarrows \mathcal{N} : U$ between \mathcal{C} -module categories, oplax \mathcal{C} -module structures on F are in bijective correspondence with lax \mathcal{C} -module structures on U .

Theorem (Abelian monadicity)

An adjunction $F: \mathcal{C} \rightleftarrows \mathcal{D} : U$ is monadic if U is exact and reflects zero objects.

Internal projectives

Let \mathcal{C} and \mathcal{M} be abelian.

Internal projectives

Let \mathcal{C} and \mathcal{M} be abelian. An object $\ell \in \mathcal{M}$ is *closed* if there is an adjunction

$$- \triangleright \ell : \mathcal{C} \rightleftarrows \mathcal{M} : [\ell, -].$$

Internal projectives

Let \mathcal{C} and \mathcal{M} be abelian. An object $\ell \in \mathcal{M}$ is *closed* if there is an adjunction

$$-\triangleright \ell: \mathcal{C} \rightleftarrows \mathcal{M} : \lfloor \ell, - \rfloor.$$

A closed object is called *\mathcal{C} -projective* if $\lfloor \ell, - \rfloor$ is (right) exact

Internal projectives

Let \mathcal{C} and \mathcal{M} be abelian. An object $\ell \in \mathcal{M}$ is *closed* if there is an adjunction

$$-\triangleright \ell: \mathcal{C} \rightleftarrows \mathcal{M} : \lfloor \ell, - \rfloor.$$

A closed object is called *\mathcal{C} -projective* if $\lfloor \ell, - \rfloor$ is (right) exact and a *\mathcal{C} -generator* if it is faithful.

Internal projectives

Let \mathcal{C} and \mathcal{M} be abelian. An object $\ell \in \mathcal{M}$ is *closed* if there is an adjunction

$$-\triangleright \ell: \mathcal{C} \rightleftarrows \mathcal{M} : \lfloor \ell, - \rfloor.$$

A closed object is called *\mathcal{C} -projective* if $\lfloor \ell, - \rfloor$ is (right) exact and a *\mathcal{C} -generator* if it is faithful.

Example

- Every object in a rigid monoidal category \mathcal{C} is \mathcal{C} -projective.

Internal projectives

Let \mathcal{C} and \mathcal{M} be abelian. An object $\ell \in \mathcal{M}$ is *closed* if there is an adjunction

$$- \triangleright \ell : \mathcal{C} \rightleftarrows \mathcal{M} : \lfloor \ell, - \rfloor.$$

A closed object is called *\mathcal{C} -projective* if $\lfloor \ell, - \rfloor$ is (right) exact and a *\mathcal{C} -generator* if it is faithful.

Example

- Every object in a rigid monoidal category \mathcal{C} is \mathcal{C} -projective.
- Finite \mathcal{C} -module categories over finite tensor categories always admit \mathcal{C} -projective \mathcal{C} -generators [EGNO15; DSPS19].

Only the Eilenberg–Moore category of an
oplax \mathcal{C} -module monad has a canonical
 \mathcal{C} -module structure.

Linton coequalisers

Definition

The *Linton coequaliser* of $x \in \mathcal{C}$ and $m \in \mathcal{M}^T$ is:

$$T(x \triangleright Tm) \begin{array}{c} \xrightarrow{T(x \triangleright \nabla_m)} \\ \xrightarrow{\mu_{x \triangleright m} \circ TT_{\mathbf{a};x,m}} \end{array} T(x \triangleright m) \longrightarrow\!\!\!\twoheadrightarrow x \blacktriangleright m.$$

Linton coequalisers

Definition

The *Linton coequaliser* of $x \in \mathcal{C}$ and $m \in \mathcal{M}^T$ is:

$$T(x \triangleright Tm) \begin{array}{c} \xrightarrow{T(x \triangleright \nabla_m)} \\ \xrightarrow{\mu_{x \triangleright m} \circ TT_{a;x,m}} \end{array} T(x \triangleright m) \longrightarrow x \triangleright m.$$

Theorem ([AHLF18; SZ24])

The Eilenberg–Moore category of any right exact lax \mathcal{C} -module monad can be equipped with a canonical \mathcal{C} -module structure by mean of Linton coequalisers.

The reconstruction result

Theorem ([SZ24])

Let \mathcal{C} be an abelian monoidal category, \mathcal{M} an abelian \mathcal{C} -module category, and assume that $\ell \in \mathcal{M}$ is a closed \mathcal{C} -projective \mathcal{C} -generator.

The reconstruction result

Theorem ([SZ24])

Let \mathcal{C} be an abelian monoidal category, \mathcal{M} an abelian \mathcal{C} -module category, and assume that $\ell \in \mathcal{M}$ is a closed \mathcal{C} -projective \mathcal{C} -generator. Then

$$\mathcal{M} \simeq_{\triangleright} \mathcal{C}^{[\ell, -\triangleright \ell]}.$$

The reconstruction result

Theorem ([SZ24])

Let \mathcal{C} be an abelian monoidal category, \mathcal{M} an abelian \mathcal{C} -module category, and assume that $\ell \in \mathcal{M}$ is a closed \mathcal{C} -projective \mathcal{C} -generator. Then

$$\mathcal{M} \simeq_{\triangleright} \mathcal{C}^{[\ell, - \triangleright \ell]}.$$

Furthermore, there is a bijection

$$\{(\mathcal{M}, \ell) \text{ as before}\} / \mathcal{M} \simeq \mathcal{N} \xleftrightarrow{\cong} \left\{ \begin{array}{l} \text{Right exact lax } \mathcal{C}\text{-module} \\ \text{monads on } \mathcal{C} \end{array} \right\} / \mathcal{C}^T \simeq \mathcal{C}^S$$

(Op)lax module functors in action

Hopf trimodules

Theorem ([SZ24])

Let B be a bialgebra, and define $\mathcal{V} := {}^B\text{Vect}$.

Hopf trimodules

Theorem ([SZ24])

Let B be a bialgebra, and define $\mathcal{V} := {}^B\mathbf{Vect}$. There is a monoidal equivalence

$$\begin{aligned} {}^B_B\mathbf{Vect} &\longrightarrow \mathbf{LexLax}\mathcal{V}\mathbf{Mod}(\mathcal{V}, \mathcal{V}) \\ X &\longmapsto (X \square_B -, \chi) \end{aligned}$$

between the category of Hopf trimodules, and the category of left exact finitary lax \mathcal{V} -module endofunctors on \mathcal{V} .

The interchanger

For all $M, N \in {}^B\mathbf{Vect}$, the arrow


$$\chi_{M,N}: M \otimes_{\mathbb{k}} (X \square_B N) \longrightarrow X \square_B (M \otimes_{\mathbb{k}} N)$$

The interchanger

For all $M, N \in {}^B\mathbf{Vect}$, the arrow

$$\chi_{M,N}: M \otimes_{\mathbb{k}} (X \square_B N) \longrightarrow X \square_B (M \otimes_{\mathbb{k}} N)$$

is defined by



Deducing a theorem for Hopf trimodules

Proposition

Let \mathcal{C} be a left closed monoidal category such that every lax \mathcal{C} -module endofunctor of \mathcal{C} is strong—in other words, that the monoidal embedding

$$\mathrm{Str}\mathcal{C}\mathrm{Mod}(\mathcal{C}, \mathcal{C}) \hookrightarrow \mathrm{Lax}\mathcal{C}\mathrm{Mod}(\mathcal{C}, \mathcal{C})$$

is an equivalence. Then \mathcal{C} is left rigid.

Deducing a theorem for Hopf trimodules

Proposition

Let \mathcal{C} be a left closed monoidal category such that every lax \mathcal{C} -module endofunctor of \mathcal{C} is strong—in other words, that the monoidal embedding

$$\mathrm{Str}\mathcal{C}\mathrm{Mod}(\mathcal{C}, \mathcal{C}) \hookrightarrow \mathrm{Lax}\mathcal{C}\mathrm{Mod}(\mathcal{C}, \mathcal{C})$$

is an equivalence. Then \mathcal{C} is left rigid.

Corollary ([SZ24])

A bialgebra B admits a twisted antipode if and only if the canonical functor $B \otimes_{\mathbb{k}} - : {}^B\mathrm{Vect} \longrightarrow {}^B_B\mathrm{Vect}^B$ is an equivalence.

Fusion operators for Hopf monads

Proposition ([SZ24])

Let $F: \mathcal{C} \rightleftarrows \mathcal{D} : U$ be an oplax monoidal adjunction.

Fusion operators for Hopf monads

Proposition ([SZ24])

Let $F: \mathcal{C} \rightleftarrows \mathcal{D} : U$ be an oplax monoidal adjunction. The strong monoidal structure of U turns \mathcal{C} into a \mathcal{D} -module category via $- \triangleright = := U(-) \otimes =$.

Fusion operators for Hopf monads

Proposition ([SZ24])

Let $F: \mathcal{C} \rightleftarrows \mathcal{D} : U$ be an oplax monoidal adjunction. The strong monoidal structure of U turns \mathcal{C} into a \mathcal{D} -module category via $- \triangleright = := U(-) \otimes =$. The bimonad $T := UF$ on \mathcal{C} becomes an oplax \mathcal{D} -module monad.

Fusion operators for Hopf monads

Proposition ([SZ24])

Let $F: \mathcal{C} \rightleftarrows \mathcal{D} : U$ be an oplax monoidal adjunction. The strong monoidal structure of U turns \mathcal{C} into a \mathcal{D} -module category via $- \triangleright = := U(-) \otimes =$. The bimonad $T := UF$ on \mathcal{C} becomes an oplax \mathcal{D} -module monad. In particular, the right fusion operator is the “free part” of the coherence morphism:

$$T_{2;F,\text{Id}} = T_{\text{rf}}.$$

Fusion operators for Hopf monads

Proposition ([SZ24])

Let $F: \mathcal{C} \rightleftarrows \mathcal{D} : U$ be an oplax monoidal adjunction. The strong monoidal structure of U turns \mathcal{C} into a \mathcal{D} -module category via $- \triangleright = := U(-) \otimes =$. The bimonad $T := UF$ on \mathcal{C} becomes an oplax \mathcal{D} -module monad. In particular, the right fusion operator is the “free part” of the coherence morphism:

$$T_{2;F,\text{Id}} = T_{\text{rf}}.$$

Further, T_{rf} is an isomorphism if and only if T_2 is.

Thanks!



tony-zorman.com/ct2025

Reconstruction of module categories
in the infinite and non-rigid settings.

[arXiv:2409.00793](https://arxiv.org/abs/2409.00793)

- [AHLF18] Marcelo Aguiar, Mariana Haim, and Ignacio López Franco. **Monads on higher monoidal categories.** In: *Appl. Categ. Struct.* 26.3 (2018), pp. 413–458. ISSN: 0927-2852. DOI: 10.1007/s10485-017-9497-8.
- [BLV11] Alain Bruguières, Steve Lack, and Alexis Virelizier. **Hopf monads on monoidal categories.** In: *Advances in Mathematics* 227.2 (2011), pp. 745–800. ISSN: 0001-8708. DOI: 10.1016/j.aim.2011.02.008.
- [Del02] Pierre Deligne. **Catégories tensorielles. (Tensor categories).** French. In: *Mosc. Math. J.* 2.2 (2002), pp. 227–248. ISSN: 1609-3321.

- [DSPS19] Christopher L. Douglas, Christopher Schommer-Pries, and Noah Snyder. **The balanced tensor product of module categories**. In: *Kyoto J. Math.* 59.1 (2019), pp. 167–179. ISSN: 2156-2261. DOI: 10.1215/21562261-2018-0006.
- [EGNO15] Pavel Etingof, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik. **Tensor categories**. Vol. 205. Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2015, pp. xvi+343. ISBN: 978-1-4704-2024-6. DOI: 10.1090/surv/205.
- [HN99] Frank Hausser and Florian Nill. **Integral Theory for Quasi-Hopf Algebras**. In: *arXiv e-prints* (1999). arXiv: math/9904164 [math.QA].

References iii

- [HZ24] Sebastian Halbig and Tony Zorman. **Diagrammatics for Comodule Monads**. In: *Appl. Categ. Struct.* 32 (2024). Id/No 27, p. 17. ISSN: 0927-2852. DOI: 10.1017/CB09781139542333.
- [Kel74] Gregory M. Kelly. **Doctrinal adjunction**. Proceedings Sydney Category Seminar, 1972/1973, Lect. Notes Math. 420, 257-280 (1974). 1974.
- [Moe02] Ieke Moerdijk. **Monads on tensor categories**. In: *J. Pure Appl. Algebra* 168.2-3 (2002). Category theory 1999 (Coimbra), pp. 189–208. ISSN: 0022-4049. DOI: 10.1016/S0022-4049(01)00096-2.
- [Ost03] Viktor Ostrik. **Module categories, weak Hopf algebras and modular invariants**. In: *Transform. Groups* 8.2 (2003), pp. 177–206. ISSN: 1083-4362. DOI: 10.1007/s00031-003-0515-6.

- [Ost04] Victor Ostrik. **Tensor categories (after P. Deligne)**. In: *arXiv e-prints* (2004). arXiv: math/0401347 [math.CT].
- [Sar17] Paolo Saracco. **On the structure theorem for quasi-Hopf bimodules**. In: *Appl. Categ. Struct.* 25.1 (2017), pp. 3–28. ISSN: 0927-2852. DOI: 10.1007/s10485-015-9408-9.
- [SZ24] Mateusz Stroiński and Tony Zorman. **Reconstruction of module categories in the infinite and non-rigid settings**. In: *arXiv e-prints* (2024). arXiv: 2409.00793 [math.QA].