Planetary Atmospheres Equations and Values

Joseph Glenn Ladd

May 13, 2025

1 Gasses and Equation of State

Mole-based Equation	**Mass-based Equation**
Ideal Gas Constant (R)	Specific Gas Constant (R_s)
$pV = nRT \quad (n = \text{mol})$	$pV = mR_sT \quad (m = \text{mass})$

Table 1: Comparison of Mole-based and Mass-based Ideal Gas Equations

Symbol	**Unit**	**Note**
p	Pa	Pressure
V	m^3	Volume
m	kg	Mass
R_s	$\text{J}\cdot\text{kg}^{-1}\cdot\text{K}^{-1}$	Specific Gas Constant
R	$\text{J}\cdot\text{mol}^{-1}\cdot\text{K}^{-1}$	Ideal Gas Constant
T	K	Temperature
M	$\text{kg}\cdot\text{mol}^{-1}$	Molecular Weight (Molar mass)
n_V	m^{-3}	Number of molecules per unit volume
ρ	$\text{kg}\cdot\text{m}^{-3}$	Density
α	$\text{m}^3\cdot\text{kg}^{-1}$	Specific Volume
n	–	Moles
N	$\text{kg}\cdot\text{m}\cdot\text{s}^{-2}$	Newton (force)
Pa	$\text{kg}\cdot\text{m}^{-1}\cdot\text{s}^{-2}$	Pascal (pressure)
J	$\text{kg}\cdot\text{m}^2\cdot\text{s}^{-2}$	Joule (energy)
N_A	mol^{-1}	Avogadro's Number (6.022×10^{23} particles/mol)

Table 2: Physical symbols, units, and associated meanings

2 Wave Symbols and Quantities

Symbol	**Name**	**Meaning (Wave Context)**
λ	Lambda	Wavelength – distance between wave crests (m)
ν	Nu	Frequency – cycles per second (Hz = 1/s)
$\bar{\nu}$	Nu-bar	Wave number – cycles per meter (1/m)
k	k	Angular (circular) wave number – $k = 2\pi/\lambda$ (rad/m)
ω	Omega	Angular (circular) frequency – $\omega = 2\pi\nu$ (rad/s)
T	T	Period – time per cycle (s)
v_p	v-sub-p	Phase speed – speed at which wave phase propagates (m/s)

Table 3: Wave Symbols and Their Meanings

3 Radiomentric Quantities

Quantity	**Symbol**	**Units**	**Physical Meaning**	**Equation**
Radiant Power (Radiative Flux)	Φ, F	W	Total radiant energy emitted, transferred, or received per second.	$\Phi = \frac{dQ}{dt}$
Radiant Energy (Thermal energy)	Q_e, E, W	J	Total electromagnetic energy accumulated over time.	$Q = \int \Phi(t)\, dt$
Radiant Power per Unit Area (Irradiance, Radiative Flux Density, Exitance)	E, I	$\mathrm{W/m^2}$	Power received per unit surface area (in, through, or out).	$E = \frac{d\Phi}{dA}$
Radiance (Specific Intensity)	L	$\mathrm{W/(m^2\,sr)}$	Radiant power per unit area per solid angle in a specific direction.	$L = \frac{d^2\Phi}{dA \cos\theta \, d\omega}$

Table 4: Radiometric Quantities: Symbols, Units, and Definitions