

SOC 5050: Week 02 Equations Quick Reference

Christopher Prener, Ph.D.

August 29th, 2016

Median (odd n of items)

Let m = the median item's term:

$$me = \left(\frac{n+1}{2} \right)^{th} \quad (1)$$

Median (even n of items)

Let m_a = the lower, middlemost item's term:

$$m_a = \left(\frac{n+1}{2} \right)^{th} \quad (2a)$$

Let m_b = the higher, middlemost item's term:

$$m_b = \left(\frac{n}{2+1} \right)^{th} \quad (2b)$$

Let m = the median, which is the mean of the two middlemost items' terms:

$$m = \frac{m_a + m_b}{2} \quad (2c)$$

Mean

$$\bar{x} = \frac{\sum_{i=1}^n x}{n} \quad (3)$$

Variance

$$s^2 = \frac{\sum_{i=1}^n (x - \bar{x})^2}{n-1} \quad (4)$$

Standard Deviation

$$s = \sqrt{\frac{\sum_{i=1}^n (x - \bar{x})^2}{n-1}} \quad (5)$$

Document Details

Document produced by [Christopher Prener, Ph.D.](#) for the Saint Louis University course SOC 5050 - QUANTITATIVE ANALYSIS: APPLIED INFERENCEAL STATISTICS. See the [course website](#) and the repository README.md file for additional details.



This work is licensed under a [Creative Commons Attribution 4.0 International License](#).