SOC 5050: Week 06 Equations Quick Reference

Christopher Prener, Ph.D.

September 26^{th}, 2016

Binomial Distribution

For the binomial distribution, let:
n = number of independent trials
p = probability of success in each trial

Mean

$$\mu = np \tag{1}$$

Standard Deviation

$$\sigma = \sqrt{np\left(1-p\right)} \tag{2}$$

Poisson Distribution

For the Poisson distribution, let:
n = count of independent events
p = probability of success in each event

$$\lambda = np \tag{3}$$

Standard Normal Distribution

Standardized Scores: Population

$$z = \frac{x-\mu}{\sigma} \tag{4}$$

Standardized Scores: Sample

$$z = \frac{x-\bar{x}}{s} \tag{5}$$

Skew

$$sk = \sqrt{n}\,\frac{\sum_{i=1}^{n}\left(x_i - \bar{x}\right)^3}{\left(\sum_{i=1}^{n}\left(x_i - \bar{x}\right)^2\right)^{\frac{3}{2}}} \tag{6}$$

Note that *sk* is an abbreviation that I use in my classes to differentiate skew from standard deviation. There is no single accepted abbreviation for skew. Similarly, there are a number of equations in use to calculate skew; this is one that I teach because it simplifies some of the required calculations.

Kurtosis

$$k = n\,\frac{\sum_{i=1}^{n}\left(x_i - \bar{x}\right)^4}{\left(\sum_{i=1}^{n}\left(x_i - \bar{x}\right)^2\right)^2} \tag{7}$$

Note that there are a number of accepted abbreviations for kurtosis including *k*. There are also a number of equations in use to calculate kurtosis. As with skew, this is one that I teach because it simplifies some of the required calculations.

Document Details

Document produced by Christopher Prener, Ph.D. for the Saint Louis University course SOC 5050 - QUANTITATIVE ANALYSIS: APPLIED INFERENTIAL STATISTICS. See the course wiki and the repository `README.md` file for additional details.