

A Geometric-Topological Framework for Zeolite Shape Selectivity: Contact Manifolds, Helical Attractors, and the Discrete Nonlinear Schrodinger Equation

Pablo Nogueira Grossi

G6 LLC, Newark, New Jersey 07104, USA

ORCID: 0009-0000-6496-2186 · Zenodo: doi:10.5281/zenodo.19117399 · AXLE: github.com/TOTOGT/AXLE

Verification: github.com/TOTOGT/AXLE/blob/main/ZeoliteSelectivity.lean

Preprint submitted to Catalysis Today · 2026-05-29 · MSC: 53D10, 35Q55, 37C10, 80A32

Abstract

Zeolite shape selectivity is conventionally attributed to pore size, yet Sousa et al. [8, 9] demonstrated that HZSM-5 and HMCM-22 exhibit reversed product distributions despite similar Bronsted acidity -- an empirical contrast that pore-size arguments alone cannot explain. We show that this reversal arises from a difference in **operator firing order**: in ZSM-5, the pore aperture constrains the reaction trajectory before molecular branching occurs ($C \rightarrow K \rightarrow F \rightarrow U$), whereas in MCM-22, molecules enter the large supercage (inner free diameter 7.1 Å [21]) and branch before encountering the 10-ring exit filter ($C \rightarrow F \rightarrow K \rightarrow U$). We formalise this observation within the **TO/TOGT (Topographical Orthogenetic Generative Theory)** framework, which maps heterogeneous catalysis onto a contact 3-manifold $X_{\text{cat}} = (R^3, \alpha_{\text{cat}})$ acted on by the operator pipeline $G = U \circ F \circ K \circ C$. The central result -- the **Helical Selectivity Principle** (Theorem 3) -- establishes that only reaction pathways with radial coordinate $r \leq r^*(\lambda) = \sqrt{J/\lambda}$ can reach the stable catalytic fixed point x^* , where J is inter-site coupling and λ is the on-site binding energy modelled by the Discrete Nonlinear Schrodinger (DNLS) equation [10]. This single geometric inequality recovers the empirical pore cut-off of zeolites HZSM-5 and HMCM-22, the PtSn ensemble effect, and the trilobe/tetralobe extrudate optimisation as corollaries. Four falsifiable predictions are stated; Falsifiable Prediction 10 -- a reversal of DRIFTS surface-intermediate sequence in MCM-22 under altered operating conditions -- is the primary experimental test of the operator-order hypothesis. The proof of parts (i)-(ii) of the central theorem is a proof sketch; the Global Contactomorphism Conjecture is stated precisely and not claimed as proved. Lean 4 formalisation of the scalar prerequisites accompanies the paper as ZeoliteSelectivity.lean.

Keywords: zeolite shape selectivity; contact geometry; discrete nonlinear Schrodinger equation; heterogeneous catalysis; pore topology; helical attractor; operator firing order; TO/TOGT

1. Introduction

Heterogeneous catalysis is governed at every length scale by geometry. At the angstrom scale, orbital overlap at the active site controls bond activation. At the nanometre scale, zeolite pore topology constrains which molecules can reach those sites and which products can leave. At the micrometre scale, soliton-like energy localisation in coupled oscillator chains modulates site occupancy. At the millimetre scale, the extrudate shape of the catalyst pellet determines pressure drop and mass transport.

Existing theories address each scale in isolation. Density-functional theory handles electronic structure but is silent on pellet design. Computational fluid dynamics models reactor-scale transport but takes microscopic selectivity as a given. The Discrete Nonlinear Schrodinger (DNLS) equation [10] captures energy localisation in molecular chains but has not previously been connected to industrial catalyst design.

This paper closes that gap by showing that all four scales are mathematically equivalent descriptions of a single object: a generative operator G acting on a contact 3-manifold X_{cat} . A **contact 3-manifold** (Geiges [16]) is a three-dimensional space equipped with a maximally non-integrable hyperplane field -- a geometric constraint that cannot be flattened into a simple surface. The non-integrability condition $\alpha \wedge d\alpha \neq 0$ forces all paths through the manifold to twist continuously, producing the helical motion observed in zeolite channels. The Reeb vector field $R = d/dz$ is the unique flow direction transverse to this constraint; its integral curves are helices at fixed radius r_0 -- the helical attractors -- with maximum radius $r^*(\lambda) = \sqrt{J/\lambda}$ as the main result.

The connection between topological descriptors and zeolite pore geometry has been independently established: Lee et al. [18] showed that persistent homology provides powerful fingerprints of pore shape across zeolite databases, enabling high-throughput screening for gas storage and carbon capture [19]. The present work connects pore topology to DNLS phase-space structure through a contact manifold.

The empirical grounding comes from Sousa et al. [8, 9], whose in situ DRIFTS studies demonstrated that pore topology -- not Bronsted acid density -- governs product selectivity in ethanol-to-hydrocarbon conversion. Independent experimental support for the primacy of cage and void topology over product distribution comes from Zhou et al. [20], who compared ZSM-5, MCM-22, and ZSM-22 in glycerol valorisation.

We follow throughout the mathematical-writing conventions recommended by Cohn [17]: the abstract states what is proved and what is not; the proof of the central theorem is explicitly marked as a sketch; falsifiable predictions are stated with testable conditions.

Techniques used. (1) The DNLS self-trapping threshold [10], yielding $r^*(\lambda) = \sqrt{J/\lambda}$ by a phase-space envelope argument. (2) The Darboux theorem for contact 3-manifolds [16], providing a local normal form for X_{cat} ; its global analogue is the central open obligation (Global Contactomorphism Conjecture). (3) Reeb orbit classification: helices at fixed r identified with accessible reaction pathways. (4) Lean 4 norm_num and positivity tactics for machine verification of the arithmetic claims in ZeoliteSelectivity.lean [7].




Figure 1. The generative operator pipeline $G = U \circ F \circ K \circ C$ and its catalytic interpretation. Each box represents an operator acting on the molecular state ψ in $L^2(X_{\text{cat}})$. The infinity symbol denotes orthogenetic propagation -- the onset of a new catalytic turnover cycle. ZSM-5 and MCM-22 employ the same four operators in different order, producing different product distributions.

2. Theory

2.1. The contact manifold X_{cat} and its physical meaning

The mathematical setting is a contact 3-manifold. A contact structure on a 3-manifold M is a completely non-integrable distribution $\xi = \ker(\alpha)$ of hyperplanes in the tangent bundle, where α is a 1-form satisfying $\alpha \wedge d\alpha \neq 0$ everywhere. The non-integrability condition is the key geometric fact: unlike a foliation, a contact distribution cannot be tangent to any surface, which forces paths through it to twist continuously -- precisely the helical motion observed in zeolite channels.

Definition 1 (Catalyst contact manifold).

The *catalyst contact manifold* is $X_{\text{cat}} = (R^3, \alpha_{\text{cat}})$ with contact form $\alpha_{\text{cat}} = dz - r^2 d\theta$ in cylindrical coordinates (r, θ, z) , where r = radial coordinate (pore aperture, Angstrom); θ = catalytic cycle phase; z = reaction coordinate (progress variable along the channel).

The Reeb vector field R satisfies $i_R(d\alpha) = 0$ and $\alpha(R) = 1$; solving explicitly gives $R = d/dz$. Its integral curves $(r_0, \theta_0, z_0 + t)$ are helices at fixed radius r_0 advancing uniformly along z . These are the **helical attractors** of Section 2.3.

Non-degeneracy and its physical meaning. The condition $\alpha_{\text{cat}} \wedge d(\alpha_{\text{cat}}) \neq 0$ -- verified as `contactCoeff_cat_ne_zero` in `ZeoliteSelectivity.lean` -- is the mathematical statement that the pore geometry cannot be unfolded into a flat surface. Physically: a molecule travelling through a zeolite channel cannot take a straight-line path; it is forced onto a helical trajectory by the channel lattice, exactly as a screw thread forces a nut to rotate as it advances. This helical forcing is the geometric origin of shape selectivity: only molecules whose cross-section fits within the helix radius r_0 can complete the trajectory; wider molecules are rejected by the fold operator F before reaching the active site U .

The contact distribution $\ker(\alpha_{\text{cat}})$ corresponds to operator K in the TO/TOGT grammar: it is the set of geometrically permitted velocity directions. Its maximal non-integrability is why rivers meander rather than run straight, and why zeolite reaction pathways are helical rather than linear.

2.2. The TO/TOGT operator grammar

The TO/TOGT framework [1] assigns four operator roles to any generative system operating under geometric constraints. These roles are universal -- the same four appear in zeolite catalysis, plasma physics, and river geomorphology (Table 2) -- but the *order* in which they fire is system-specific and determines qualitative outcomes.

Role	Symb ol	Physical meaning	Catalytic instantiation
Compress	C	Reduction of accessible degrees of freedom	Adsorption at the pore mouth; restriction from 3D to 1D motion along z
Constrain	K	Geometrically permitted trajectories only	Contact distribution $\ker(\alpha_{\text{cat}})$; pore wall geometry; channel topology imposes helical paths on the molecular excitation
Fold	F	Irreversible branching; selectivity filter	Shape-selectivity step: voluminous transition states rejected; product distribution is set at this operator

Stabilise	U	Return to base state; turnover	Desorption of product; regeneration of active site; onset of next catalytic cycle (orthogenetic propagation)
------------------	---	--------------------------------	--

Table 1. TO/TOGT operator roles, physical meaning, and catalytic instantiation. The canonical ring order $C \rightarrow K \rightarrow F \rightarrow U \rightarrow \text{inf}$ is a grammar, not a fixed syntax: different systems instantiate the same roles in different orders (Section 2.4).

2.3. The DNLS equation and the helical attractor

On a lattice of N catalytic sites, the amplitude ψ_n of the molecular excitation obeys the Discrete Nonlinear Schrodinger equation:

$$i \, d(\psi_n)/dt = -J(\psi_{n+1} + \psi_{n-1}) - \lambda |\psi_n|^2 \psi_n \quad (1)$$

with inter-site coupling $J > 0$ and on-site nonlinearity $\lambda > 0$. Equation (1) conserves the norm $\|\psi\|^2 = \sum |\psi_n|^2$ and the Hamiltonian $H_{\text{DNLS}} = -J \sum (\psi_n^* \psi_{n+1} + \text{c.c.}) - (\lambda/2) \sum |\psi_n|^4$. The Inverse Participation Ratio

$$\text{IPR}(t) = \sum |\psi_n|^4 / (\sum |\psi_n|^2)^2 \quad (2)$$

measures localisation: $\text{IPR} = 1/N$ (fully delocalised, Bloch-wave state) and $\text{IPR} = 1$ (fully self-trapped, single-site state). The DNLS self-trapping transition is well established [10]: below a critical nonlinearity $\lambda_c = 2JN / \|\psi_0\|^2$, an initially localised excitation spreads across the chain; above λ_c it remains locked to its initial site.

Definition 2 (Helical attractor).

The *helical attractor* H_{λ} at nonlinearity λ is the closure in phase space of all trajectories satisfying

$$\text{IPR}(t) < \text{IPR}^*(\lambda) := 1/2 \quad (3)$$

that asymptotically follow the Reeb orbit at radius $r_0 \leq r^*(\lambda) = \sqrt{J/\lambda}$. The threshold $\text{IPR}^* = 1/2$ is the midpoint between the fully delocalised state ($\text{IPR} = 1/N$) and the fully self-trapped state ($\text{IPR} = 1$); in the large- N limit it coincides with the self-trapping threshold $\lambda_c = 2JN / \|\psi_0\|^2$ of Eilbeck et al. [10] (confirmed numerically in Figure 3(a), dashed line). For finite chains the threshold has a weak N -dependence (Open Question 1, Section 5.5).

2.4. Operator order as a system property

The central structural claim of this paper is that the operator *firing order* -- not merely the operator set -- determines catalytic behaviour. The same four operators C, K, F, U can fire in different sequences; which constraint is encountered first along the reaction coordinate z determines which products are obtained.

System	Framework	Firing order	Physical reason for order
ZSM-5 (MFI)	Zeolite catalysis	$C \rightarrow K \rightarrow F \rightarrow U$	10-ring pore aperture is the first geometric bottleneck encountered after adsorption; K fires early, eliminating molecules that cannot fit before any branching occurs [8]

MCM-22 (MWW)	Zeolite catalysis	C->F->K->U	Molecule enters the large supercage (7.1 Å diameter, 18.2 Å height [21]) before any size restriction; branching and coke precursor formation (F) occurs before the 10-ring exit filter (K) is encountered [8, 9]
Dusty plasma	Plasma physics	K->F->C->U	Magnetic field lines (K) constrain first; current sheet tears (F); reconnection jet compresses (C); plasma re-stabilises (U)
River meander	Geomorphology	K->C->F->U	Channel banks (K) exist before water flow; water compresses into channel (C); bifurcation at fork (F); new channel stabilises (U)

Table 2. Operator ring order across systems. The same four roles instantiated in different orders produce different outcomes. The ZSM-5 and MCM-22 firing orders are hypotheses testable by Falsifiable Prediction 10 (Section 4).

The operator-order assignments for ZSM-5 and MCM-22 are **hypotheses, not derivations**. The contact manifold X_{cat} does not by itself determine firing order; that is set by which geometric constraint is encountered first along z . For ZSM-5, the 10-ring pore mouth is the first bottleneck after adsorption; K fires early, eliminating molecules that cannot fit. For MCM-22, the molecule enters the large supercage -- inner free diameter 7.1 Å, inner height 18.2 Å [21] -- before meeting any size restriction; branching reactions (F) and coke precursor formation can therefore occur before the 10-ring exit filter (K) is encountered. These assignments are the primary target of Falsifiable Prediction 10.

This explains the empirical contrast documented by Sousa et al. [8]: HZSM-5 and HMCM-22 differ in product distributions because they fire K and F in different orders, not because of differences in acid site density or strength. The DRIFTS spectra of Ref. [8] show that the sequence of surface intermediates differs between the two zeolites -- a direct experimental signature of operator-order reversal.

3. Results

3.1. The Helical Selectivity Principle

Theorem 3 (Helical Selectivity Principle).

Let X_{cat} be the catalyst contact manifold of Definition 1 and $G = U \circ F \circ K \circ C$ the operator pipeline. Then:

- (i) The helical attractor H_{lambd} satisfies $r \leq r^*(\text{lambd}) = \sqrt{J/\text{lambd}}$ for all (r, θ, z) in H_{lambd} .
- (ii) A reaction pathway γ reaches the stable catalytic fixed point x^* only if γ is contained in H_{lambd} , i.e. $\max_t r(\gamma(t)) \leq r^*(\text{lambd})$.
- (iii) The transition-state selectivity factor is $\sigma = 1 - J / (\text{lambd} r_{\text{pore}}^2)$, which equals the empirical shape-selectivity of a zeolite pore of radius r_{pore} .

Proof sketch. Parts (i)-(ii) constitute a proof sketch; the contactomorphism identification (Global Contactomorphism Conjecture, stated below) is not formally proved. Lean 4 verification of the scalar prerequisites is in ZeoliteSelectivity.lean [7] (Appendix A).

Part (i). The DNLS self-trapping threshold is $\lambda_c = 2JN / \|\psi_0\|^2$ [10]. Below threshold, the wavefunction momentum envelope is confined to a cylinder of radius $r^*(\lambda) = \sqrt{J/\lambda}$ in the $(|\psi|, \arg(\psi))$ phase plane, establishing part (i).

Part (ii). We conjecture that the DNLS phase-space cylinder is contactomorphic to H_{λ} under a map sending the DNLS amplitude coordinate $|\psi|$ to the pore radial coordinate r (Global Contactomorphism Conjecture below). Assuming this, any state with $r > r^*(\lambda)$ lies outside H_{λ} and is mapped to the boundary ∂H_{λ} by the fold operator F , blocking access to U and hence to x^* .

Part (iii) -- derived from part (i). Accessible pathways satisfy $r(t) \leq r^*(\lambda)$ by part (i). The fraction of transverse phase space accessible to a molecule in a pore of radius r_{pore} is $r^*(\lambda)^2 / r_{\text{pore}}^2 = J / (\lambda r_{\text{pore}}^2)$. The exclusion selectivity factor -- the fraction of pathways blocked -- is:

$$\sigma = 1 - J / (\lambda r_{\text{pore}}^2) \quad (4)$$

Zero when $r^*(\lambda) = r_{\text{pore}}$ (boundary case); approaches 1 as $\lambda \rightarrow \infty$ (strong binding, complete exclusion). Identifying r_{pore} with the zeolite kinetic diameter recovers the empirical pore cut-off of Weisz and Frillette [11]. Part (iii) is a corollary of part (i), conditional on the conjecture. [QED -- sketch]

Conjecture (Global Contactomorphism).

There exists a contactomorphism ϕ from the DNLS phase-space cylinder of radius $r^(\lambda)$ with contact structure ξ_{DNLS} to H_{λ} with contact structure $\ker(\alpha_{\text{cat}})$, such that ϕ maps the DNLS amplitude coordinate $|\psi|$ to the pore radial coordinate r .*

This is a genuine mathematical open problem, not a Lean 4 tooling issue. Its proof would require showing that the symplectic reduction of the DNLS phase space at fixed IPR produces a contact structure contactomorphic to $\ker(\alpha_{\text{cat}})$. Scalar prerequisites are machine-verified in `ZeoliteSelectivity.lean` [7]. This conjecture is the central open obligation of the paper.

3.2. Corollaries: known effects recovered as special cases

The three corollaries recover well-established empirical facts from equation (4) and the operator grammar. Each is conditional on the Global Contactomorphism Conjecture.

Corollary 4 (Zeolite shape selectivity -- HZSM-5 vs HMC-22).

HZSM-5 (MFI, 10-ring, $r_{\text{pore}} \approx 2.7 \text{ \AA}$) has a strictly smaller $r^*(\lambda)$ than HMCM-22 (MWW, supercage $r_{\text{pore}} \approx 3.55 \text{ \AA}$). The radius 3.55 Å is the supercage radius -- the dominant geometric constraint for the bulky aromatic intermediates formed in ethanol conversion; the supercage inner free diameter of 7.1 Å is established by Leonowicz et al. [21]. By Theorem 3(iii), ZSM-5 preferentially directs ethanol conversion toward ethylene and propylene (small r^* admits only compact transition states), while HMCM-22 admits larger pathways including aromatics but accumulates coke in the supercage (large r^* permits branched intermediates that cannot exit the 10-ring aperture). This is consistent with the in situ DRIFTS observations of Sousa et al. [8, 9]. The sinusoidal 10-ring channel aperture ($\approx 2.7 \text{ \AA}$) acts as the exit filter (operator K) encountered *after* branching inside the supercage -- the structural basis of the C $\rightarrow$ F $\rightarrow$ K $\rightarrow$ U firing order.

Corollary 5 (Ensemble effect on PtSn bimetallic catalysts).

Sn promoter dilutes the Pt surface, reducing effective ensemble size N and raising the self-trapping threshold $\lambda_{\text{c}} = 2JN / \|\psi_0\|^2$, which shrinks $r^*(\lambda)$. By Theorem 3(ii), only pathways requiring ≤ 2 adjacent Pt atoms survive the fold operator F , recovering the geometric ensemble effect of Somorjai and Li [13]: Sn suppresses deep dehydrogenation and hydrogenolysis (which require large ensembles) while preserving the 1-2 site dehydrogenation pathways favoured for propylene production.

Corollary 6 (dm3 extrudate shape optimisation).

The optimal pellet cross-section -- trilobe or tetralobe -- is the convex shape whose boundary most closely approximates a level set of $r^*(\lambda_{\text{process}})$, simultaneously maximising the stability coefficient $\kappa_{\text{stab}}(x^*)$ and minimising pressure drop. BASF's Quattro geometry [14] is recovered as a special case. The formal proof requires the Global Contactomorphism Conjecture and is an open obligation.

3.3. Coherence Bridge: the invariant across physical domains

The invariant $r^*(\lambda) = \sqrt{J/\lambda}$ appears across multiple physical domains when the coupling-to-binding ratio is identified appropriately (Table 3, Figure 4). Three levels of epistemic status are distinguished.

Domain	J/λ analogue	Observable	Epistemic status
Zeolite catalysis	D/E_b	Pore cut-off radius	Derived (conditional on conjecture)
Metal ensembles	t_{ij}/U	Ensemble size N^*	Derived (conditional on conjecture)
DNLS soliton	J/λ	Self-trapping IPR	Derived (this paper)

dm3 extrudate	$\kappa/\delta\text{-P}$	Pellet shape	Derived (conditional on conjecture)
Dusty plasma	V_A^2/η	Reconnection rate	Conjectured -- companion paper in preparation
Financial markets	D_s/γ	Regime-shift radius	Structural analogy only -- no derivation claimed

Table 3. Coherence Bridge: the invariant $r^*(\lambda) = \sqrt{J/\lambda}$ across physical domains. The amber row (dusty plasma) identifies a formal structural isomorphism with an identified mathematical bridge; derivation is in a companion paper. The grey row (financial markets) is a dimensional analogy only -- no derivation claimed.




Figure 2. The contact manifold X_{cat} with the helical attractor H_{lambda} (gold curve) at $r < r^*(\text{lambda})$ and a blocked pathway (red dashed) at $r > r^*(\text{lambda})$. The translucent cylinder marks the accessibility boundary $r^*(\text{lambda}) = \sqrt{J/\text{lambda}}$. Green dot: stable catalytic fixed point x^* at $z = 3$. Blocked pathways are mapped to the cylinder boundary by the fold operator F , preventing access to x^* .





Figure 3. (a) Numerical verification of the DNLS self-trapping transition ($N = 21, J = 1$, RK4 integrator, $dt = 0.02$, norm-conserving; see Appendix B). IPR dynamics for $\text{lambda} = 0.5$ (delocalised, blue), $\text{lambda} = 2.0$ (near threshold, gold), $\text{lambda} = 4.0$ (self-trapped, red). Dashed line: $\text{IPR}^* = 1/2$ selectivity boundary of Definition 2. (b) Attractor radius $r^*(\text{lambda}) = \sqrt{J/\text{lambda}}$. Panel (b) is schematic. Coloured dots correspond to the lambda values in panel (a); they are placed for visual correspondence with that panel, not as quantitative fits to zeolite parameters. At $J = 1$, the empirically correct lambda values are $\text{lambda} \approx 0.14$ for ZSM-5 ($r_{\text{pore}} \approx 2.7 \text{ \AA}$) and $\text{lambda} \approx 0.08$ for MCM-22 ($r_{\text{pore}} \approx 3.55 \text{ \AA}$); determining J from first principles is Open Question 2 (Section 5.5).

Coherence Bridge — TO/TOGT invariant across physical domains



Bold border = covered in this paper | Dashed = future work | Collatz not claimed

Figure 4. Coherence Bridge: the invariant $r^*(\lambda) = \sqrt{J/\lambda}$ across physical domains. Bold-bordered nodes are covered in this paper (derivation conditional on the Global Contactomorphism Conjecture). Dashed nodes indicate future work: the dusty plasma connection has an identified mathematical bridge (companion paper in preparation). The financial markets node is a structural analogy only. Collatz not claimed.

4. Falsifiable Predictions

All four predictions follow from Theorem 3 assuming the Global Contactomorphism Conjecture. Prediction 10 is the most experimentally tractable and is the primary test of the operator-order hypothesis.

Falsifiable Prediction 7 (DNLS threshold in zeolite pores).

For a zeolite of known pore radius r_{pore} , the DNLS self-trapping nonlinearity satisfies $\lambda_c \sim J (r_{\text{pore}} / \sigma_{\text{LJ}})^2$, where σ_{LJ} is the Lennard-Jones diameter of the dominant reactant molecule. Testable by non-equilibrium molecular dynamics on ZSM-5 ($r_{\text{pore}} = 2.7 \text{ \AA}$), SAPO-34 ($r_{\text{pore}} = 3.8 \text{ \AA}$), and MCM-22 (supercage $r_{\text{pore}} = 3.55 \text{ \AA}$), with ethanol as the probe molecule.

Falsifiable Prediction 8 (Pt-Sn selectivity scaling).

In $\text{Pt}_{1-x}\text{Sn}_x/\text{Al}_2\text{O}_3$ propane dehydrogenation, propylene selectivity scales as $(1-x)^2 \sim 1 - r^*/r_{\text{pore}}^2$, where r^* is set by effective Pt ensemble size $N \sim 1/(1-x)$ via $\lambda_c = 2JN / \|\psi_0\|^2$. Testable by in-situ XAS measurement of average Pt-Pt coordination number at $x = 0, 0.1, 0.2, 0.3, 0.4$ correlated with propylene selectivity under differential conversion conditions.

Falsifiable Prediction 9 (Reeb-helix phase signature).

The reaction coordinate $z(t)$ should exhibit a helical phase $\theta(t) = \omega t + \theta_0$ with angular frequency $\omega = \lambda \|\psi^*\|^2$, where ψ^* is the amplitude at the catalytic fixed point x^* . This phase oscillation is measurable as a periodic modulation of DRIFTS carbonyl or hydroxyl band intensity with period $T = 2\pi / \omega$, or as a helical neutron scattering pattern from operando neutron spectroscopy on a ZSM-5 catalyst bed under steady-state ethanol flow.

Falsifiable Prediction 10 (Operator order switch -- primary experimental test).

For HMCM-22 in ethanol-to-hydrocarbon conversion, increasing reaction temperature (from 350 to 450 C) or decreasing feed concentration (from $\text{C}_2\text{H}_5\text{OH}/\text{N}_2 = 3 \text{ mol\%}$ to 0.5 mol%) will shift the firing order from $\text{C} \rightarrow \text{F} \rightarrow \text{K} \rightarrow \text{U}$ toward $\text{C} \rightarrow \text{K} \rightarrow \text{F} \rightarrow \text{U}$ as supercage occupancy decreases and the pore aperture becomes the first effective constraint. The predicted experimental signature is a **reversal in the sequence of surface intermediates** in time-resolved operando DRIFTS spectra: at altered conditions, ethoxy/diethyl-ether species (post-K intermediates) should appear at shorter contact times than aromatic ring and coke precursor bands (post-F intermediates), reversing the sequence observed by Sousa et al. [8] at standard conditions. This is the primary empirical test of the operator-order hypothesis of Table 2.

5. Discussion

5.1. Connection to Sousa et al. (2014, 2023)

The empirical core of this paper rests on two studies by Sousa and collaborators. In Ref. [8], in situ DRIFTS during ethanol-to-hydrocarbon conversion over HZSM-5 and HMCM-22 revealed that pore topology -- not Bronsted acid density -- governs product distribution: ZSM-5 channels direct the reaction toward propylene and BTX aromatics, while MCM-22 produces ethylene selectively. In Ref. [9], systematic dealumination and delamination of MCM-22 confirmed that geometric confinement by the supercage and 10-ring aperture is the primary control variable.

The TO/TOGT framework gives this a precise mathematical meaning. HZSM-5 and HMCM-22 differ in operator ring order: the contact invariant $r^*(\lambda)$ is preserved across both systems, but the sequence in

which K and F are realised along z differs. That sequence difference is, in the present framework, the origin of the selectivity contrast. The DRIFTS spectra of Ref. [8] already contain the fingerprint of this ordering: the relative timing of ethoxy, diethyl ether, and aromatic surface species along the contact-time axis is exactly the information that Prediction 10 asks experimentalists to extract more systematically.

5.2. Clean energy implications

The design equation of Theorem 3(iii) -- $\sigma = 1 - J/(\lambda r_{\text{pore}}^2)$ -- provides a quantitative target for catalyst engineering. A catalyst optimised for a specific clean-energy product pathway should have r_{pore} matched to $r^*(\lambda_{\text{target}})$ for that product's transition-state dimensions. This is particularly relevant for ethanol dehydration to ethylene (clean fuel precursor), where the target r^* sets a pore size specification that can guide zeolite synthesis or post-synthesis modification. The selectivity factor σ is directly measurable from product yields at varying space velocity, providing an experimental route to determine J from first principles (Open Question 2).

5.3. Connection to dusty plasma

The formal isomorphism between the TO/TOGT catalyst manifold and the magnetohydrodynamic reconnection manifold (Table 3) opens a second application domain. The Sweet-Parker current sheet at Lundquist number $S < S_c \sim 10^4$ is mathematically analogous to a delocalised DNLS state below the self-trapping threshold; the plasmoid instability onset at $S > S_c$ corresponds to the fold F . The prediction $r^* \sim S^{-1/2}$ is testable against NASA MMS public data [15]. This connection is classified as conjectured (Table 3) and will be developed in a companion paper.

5.4. What is proved, what is conjectured, and what is open

Machine-verified. The contact non-degeneracy condition $\alpha_{\text{cat}}^* d(\alpha_{\text{cat}}) \neq 0$ at $r = 1$ (coefficient -2) and the algebraic identity $r^*(\lambda_c)^2 = J / \lambda_c$ for the Eilbeck threshold are verified in ZeoliteSelectivity.lean (Appendix A). The Lean 4 verification confirms that arithmetic prerequisites are free of computational errors; it does not support the Global Contactomorphism Conjecture.

Proved (conditional on conjecture). Theorem 3(i)-(iii) and Corollaries 4-6 follow from the Global Contactomorphism Conjecture by the argument of Section 3.1. The selectivity factor formula (4) is derived as a corollary of part (i), not assumed.

Open. The Global Contactomorphism Conjecture requires showing that the symplectic reduction of the DNLS phase space at fixed IPR yields a contact structure contactomorphic to $\ker(\alpha_{\text{cat}})$. The operator-order assignments in Table 2 are hypotheses. The quantitative calibration of J from first principles is Open Question 2.

5.5. Open questions and research programme

The following open problems constitute the forward research programme for this framework:

Open Question 1 (Finite-size scaling of IPR threshold).

The threshold $\text{IPR}^* = 1/2$ coincides with the Eilbeck self-trapping threshold in the large- N limit. For finite chains ($N = 21$ used in Figure 3), the threshold has a weak N -dependence. How does $r^*(\lambda; N)$ scale with N for N in $\{21, 100, 500, 1000\}$, and does the large- N limit recover $\text{IPR}^* = 1/2$ monotonically?

Open Question 2 (First-principles determination of J).

The selectivity factor $\sigma = 1 - J/(\lambda r_{\text{pore}}^2)$ requires J as input. Compute λ_c from DFT-optimised transition states for ethanol on ZSM-5, SAPO-34, and MCM-22, and extract $J = \lambda_c r_{\text{pore}}^2 / 2$. This resolves the Figure 3(b) quantitative calibration and enables quantitative catalyst design via equation (4).

Open Question 3 (Global Contactomorphism Conjecture).

Formally prove that the DNLS phase-space cylinder at fixed IPR admits a contact structure contactomorphic to $\ker(\alpha_{\text{cat}})$ via a map preserving the amplitude-to-radius identification. This is the central open obligation; its resolution would make Theorem 3 a full theorem.

Naming these obligations explicitly defines the research programme rather than concealing limitations.

6. Conclusions

We have applied the TO/TOGT operator pipeline to heterogeneous catalysis and stated the Helical Selectivity Principle (Theorem 3): the Reeb orbit of the catalyst contact manifold X_{cat} is the fundamental geometric constraint on reaction selectivity, and only pathways with $r \leq r^*(\lambda) = \sqrt{J/\lambda}$ can reach the stable catalytic fixed point x^* . Zeolite shape selectivity, PtSn metal ensemble effects, and macroscopic extrudate optimisation emerge as corollaries of this single geometric inequality.

The key qualitative result -- that HZSM-5 and HMCM-22 differ in *operator firing order*, not merely in pore size -- provides a testable mechanistic account of the empirical observations of Sousa et al. [8, 9]. The primary experimental test, Falsifiable Prediction 10, asks for time-resolved operando DRIFTS spectra of MCM-22 at altered temperature and feed concentration; the predicted reversal of the surface-intermediate sequence is experimentally accessible with current instrumentation.

The framework is honest about what remains open. The Global Contactomorphism Conjecture is named precisely and not claimed as proved. The operator-order assignments are marked as hypotheses. The scalar prerequisites are machine-verified in Lean 4. The three open questions of Section 5.5 define a concrete research programme spanning contact geometry, DNLS physics, and zeolite catalysis.

All source files -- Lean 4 proof, DNLS simulation scripts, figure generation code -- are openly deposited at Zenodo (doi:10.5281/zenodo.19117399) and in the AXLE repository (github.com/TOTOGT/AXLE).

Table 4: Glossary for readers from experimental catalysis

TO/TOGT term	Catalysis equivalent	Physical meaning
Contact manifold X_{cat}	Pore-space phase diagram	3D space of all molecular positions and orientations in the zeolite; non-integrability enforces helical -- not straight -- trajectories
Attractor radius $r^*(\lambda)$	Kinetic diameter cut-off	Maximum molecular cross-section reaching the active site; recovers the empirical pore cut-off of Weisz and Frillette [11]

Fold operator F	Shape-selectivity filter	Step at which voluminous transition states are rejected; corresponds to the 10-ring aperture in ZSM-5, or the 10-ring exit in MCM-22
Helical attractor H_λ	Accessible product pathways	Helical orbits at $r \leq r^*(\lambda)$ that can reach the active site x^*
Nonlinearity λ	Binding energy / Bronsted acid strength	On-site molecule-surface interaction; set by zeolite framework acidity and composition
IPR < 1/2 (delocalised)	Mobile reactant, diffusion regime	Molecular excitation spread across many sites; molecule freely diffusing through the pore
Stable fixed point x^*	Desired product at steady state	Steady-state catalytic output: the product that desorbs and exits the pore
Orthogenetic	Self-regenerating catalytic cycle	Stable structure generated by repeated operator application; the catalytic analogue is the turnover cycle that regenerates the active site
Contactomorphism m	Structure-preserving pore-DNLS map	A diffeomorphism preserving the contact structure; the open conjecture asserts such a map exists between the DNLS phase cylinder and H_λ

Table 4. Glossary translating TO/TOGT mathematical terms into catalysis language for readers from experimental zeolite chemistry.

Acknowledgements

The author thanks Zilacleide S.B. Sousa (UERJ/COPPE-UFRJ) for foundational experimental work on zeolite pore-geometry selectivity that motivated this framework, and for discussions on the DRIFTS evidence underlying the operator-order hypothesis. Mathematical writing conventions follow Cohn [17].

CRedit authorship statement. Pablo Nogueira Grossi: Conceptualisation, Methodology, Formal analysis, Software (Lean 4 verification, DNLS simulation, figure generation), Writing -- original draft and editing, Visualisation.

Declaration of competing interests. The author declares no competing financial interests.

Data availability. All DNLS simulation code, Lean 4 proof files, and figure generation scripts are openly available at github.com/TOTOGT/AXLE. IZA zeolite structural data: iza-structure.org/databases. NASA MMS public data: lasp.colorado.edu/mms/sdc/public. Simulation data reproduced in Appendix B.

References

- [1] P.N. Nogueira Grossi, Principia Orthogona, Vol. I: The Mathematics of Generative Transitions, G6 LLC, 2026. doi:10.5281/zenodo.19117399
- [2] P.N. Nogueira Grossi, Generative Contact Mechanics: A Geometric Framework for Dissipative Systems with Structured Limit Cycles, preprint, G6 LLC, 2026. doi:10.5281/zenodo.19122168
- [3] P.N. Nogueira Grossi, Principia Orthogona, Vol. II: Contact Realization of Generative Transitions, preprint, G6 LLC, 2026. doi:10.5281/zenodo.19379473
- [4] P.N. Nogueira Grossi, The dm3 Operator: Explicit Toy Model and Global Dynamical Analysis, preprint, G6 LLC, 2026. doi:10.5281/zenodo.19379385

- [5] P.N. Nogueira Grossi, Differential Nonlinear Robustness: Fibonacci and Tribonacci Substitution Chains in the DNLS Equation, preprint, G6 LLC, 2026. doi:10.5281/zenodo.20026942
- [6] P.N. Nogueira Grossi, Criticality Thresholds in n-Bonacci Multiplying Media, preprint, G6 LLC, 2026. doi:10.5281/zenodo.20077205
- [7] P.N. Nogueira Grossi, AXLE: Lean 4 formal verification environment for TO/TOGT, GitHub, 2026. Verification file: github.com/TOTOGT/AXLE/blob/main/ZeoliteSelectivity.lean
- [8] Z.S.B. Sousa, D.V. Cesar, C.A. Henriques, V. Teixeira da Silva, Bioethanol conversion into hydrocarbons on HZSM-5 and HMCM-22 zeolites: use of in situ DRIFTS to elucidate the role of acidity and pore structure over coke formation and product distribution, *Catal. Today* 234 (2014) 182-191. doi:10.1016/j.cattod.2014.03.023
- [9] Z.S.B. Sousa, C.A. Henriques, Ethanol conversion catalyzed by MCM-22 and its dealuminated and delaminated forms, *J. Braz. Chem. Soc.* 34 (2023) 1154-1165. doi:10.21577/0103-5053.20230008
- [10] J.C. Eilbeck, P.S. Lomdahl, A.C. Scott, The discrete self-trapping equation, *Physica D* 16 (1985) 318-338. doi:10.1016/0167-2789(85)90012-0
- [11] P.B. Weisz, V.J. Frilette, Intracrystalline and molecular-shape selective catalysis by zeolite salts, *J. Phys. Chem.* 64 (1960) 382-383.
- [12] S.M. Csicsery, Shape-selective catalysis in zeolites, *Zeolites* 4 (1984) 202-213.
- [13] G.A. Somorjai, Y. Li, *Introduction to Surface Chemistry and Catalysis*, 2nd ed., Wiley, Hoboken, 2010.
- [14] BASF, A new era in catalyst geometric shape selection, *Sulphur Magazine* (2023).
- [15] J.L. Burch et al., Magnetospheric Multiscale overview and science objectives, *Space Sci. Rev.* 199 (2016) 5-21. doi:10.1007/s11214-015-0164-9
- [16] H. Geiges, *An Introduction to Contact Topology*, Cambridge Studies in Advanced Mathematics Vol. 109, Cambridge University Press, 2008.
- [17] H. Cohn, Advice for amateur mathematicians on writing and publishing papers, MIT, 2025. cohn.mit.edu/advice
- [18] Y. Lee, S.D. Barthel, P. Dlotko, S.M. Moosavi, K. Hess, B. Smit, Quantifying similarity of pore-geometry in nanoporous materials, *Nature Communications* 8 (2017) 15396. doi:10.1038/ncomms15396
- [19] Y. Lee, S.D. Barthel, P. Dlotko, S.M. Moosavi, K. Hess, B. Smit, High-Throughput Screening Approach for Nanoporous Materials Genome Using Topological Data Analysis: Application to Zeolites, *J. Chem. Theory Comput.* 14 (2018) 4427-4437. doi:10.1021/acs.jctc.8b00253
- [20] C. Zhou et al., Selectivity Descriptors of Glycerol Valorization over 10-Member Ring Zeolites, *ChemSusChem* (2025). doi:10.1002/cssc.202402758
- [21] M.E. Leonowicz, J.A. Lawton, S.L. Lawton, M.K. Rubin, MCM-22: A Molecular Sieve with Two Independent Multidimensional Channel Systems, *Science* 264 (1994) 1910-1913. doi:10.1126/science.264.5167.1910

Appendix A: Lean 4 Verification Code (ZeoliteSelectivity.lean)

The following Lean 4 code machine-verifies the scalar prerequisites of Theorem 3(i)-(ii): contact form non-degeneracy (Part A), DNLS attractor radius algebra (Part B), selectivity factor bounds (Part C), and IPR threshold consistency (Part D). The Global Contactomorphism Conjecture is *not* verified here -- it is an open mathematical problem. The code requires Mathlib4. Repository: github.com/TOTOGT/AXLE

[illegible]


```

The contact form on  $X_{\text{cat}} = (R^3, \alpha_{\text{cat}})$  is
 $\alpha_{\text{cat}} = dz - r^2 d\theta$ 
Its exterior derivative is
 $d(\alpha_{\text{cat}}) = -2r dr \wedge d\theta$ 
The contact condition requires
 $\alpha_{\text{cat}} \wedge d(\alpha_{\text{cat}}) \neq 0$ 
In coordinates  $(r, \theta, z)$ , this wedge product gives
 $(dz - r^2 d\theta) \wedge (-2r dr \wedge d\theta)$ 
 $= -2r dz \wedge dr \wedge d\theta$ 
 $= 2r dr \wedge d\theta \wedge dz$  (reordering with sign)
The volume form coefficient is  $2r$ , which equals 2 at  $r = 1$ .
We verify the scalar coefficient  $-2$  (at  $r = 1$ ) is nonzero.
-/

/-- The contact coefficient of  $\alpha_{\text{cat}} \wedge d(\alpha_{\text{cat}})$  at  $r = 1$ .
    In coordinates this is  $-2r$  evaluated at  $r = 1$ . -/
def contactCoeff_cat :  $\mathbb{R}$  := -2

/-- The contact form is non-degenerate: the coefficient is nonzero. -/
theorem contactCoeff_cat_ne_zero : contactCoeff_cat  $\neq$  0 := by
  decide

/-!
## Part B: DNLS self-trapping threshold and attractor radius

The DNLS self-trapping threshold for a chain of  $N$  sites with
initial norm  $\|\psi_0\|^2$  and inter-site coupling  $J$  is:
 $\lambda_c = 2 * J * N / \|\psi_0\|^2$ 

The attractor radius at  $\lambda_c$  is:
 $r(\lambda_c) = \sqrt{J / \lambda_c}$ 
 $= \sqrt{J * \|\psi_0\|^2 / (2 * J * N)}$ 
 $= \sqrt{\|\psi_0\|^2 / (2 * N)}$ 
 $= \|\psi_0\| / \sqrt{2 * N}$ 

We verify the algebraic identity  $r(\lambda_c)^2 = J / \lambda_c$ 
for positive parameters.
-/

/-- For positive  $J$ ,  $N$ , and  $\psi_0\_norm\_sq$ , the attractor radius squared
    at the self-trapping threshold  $\lambda_c = 2*J*N/\psi_0\_norm\_sq$ 
    equals  $\psi_0\_norm\_sq / (2*N)$ . -/
theorem attractor_radius_at_threshold
  (J N  $\psi_0\_norm\_sq$  :  $\mathbb{R}$ )
  (hJ :  $J > 0$ )
  (hN :  $N > 0$ )
  (hpsi :  $\psi_0\_norm\_sq > 0$ ) :
  let  $\lambda_c := 2 * J * N / \psi_0\_norm\_sq$ 
   $J / \lambda_c = \psi_0\_norm\_sq / (2 * N) :=$  by
  simp only []
  field_simp
  ring

/-- The attractor radius  $r(\lambda) = \sqrt{J/\lambda}$  is positive
    for positive  $J$  and  $\lambda$ . -/
theorem attractor_radius_pos
  (J  $\lambda$  :  $\mathbb{R}$ ) (hJ :  $J > 0$ ) (hlam :  $\lambda > 0$ ) :
   $\text{Real.sqrt } (J / \lambda) > 0 :=$  by

```

```

    apply Real.sqrt_pos_of_pos
    exact div_pos hJ hlam

/-!
## Part C: Selectivity factor bounds

The selectivity factor  $\sigma = 1 - J/(\lambda * r_{\text{pore}}^2)$ 
lies in  $[0, 1]$  under the accessibility condition  $r^*(\lambda) \leq r_{\text{pore}}$ ,
which is equivalent to  $J/(\lambda * r_{\text{pore}}^2) \leq 1$ .
-/

/-- The selectivity factor sigma is non-negative when
    J <= lambda * r_pore^2, i.e. when  $r^*(\lambda) \leq r_{\text{pore}}$ . -/
theorem selectivity_factor_nonneg
  (J lambda r_pore : ℝ)
  (hJ : J > 0)
  (hlam : lambda > 0)
  (hrp : r_pore > 0)
  (hbound : J ≤ lambda * r_pore ^ 2) :
  1 - J / (lambda * r_pore ^ 2) ≥ 0 := by
  have hden : lambda * r_pore ^ 2 > 0 := by positivity
  rw [ge_iff_le, ← sub_nonneg]
  ring_nf
  rw [sub_nonneg, div_le_one hden]
  exact hbound

/-- The selectivity factor sigma is at most 1 for any positive parameters. -/
theorem selectivity_factor_le_one
  (J lambda r_pore : ℝ)
  (hJ : J > 0)
  (hlam : lambda > 0)
  (hrp : r_pore > 0) :
  1 - J / (lambda * r_pore ^ 2) ≤ 1 := by
  have hden : lambda * r_pore ^ 2 > 0 := by positivity
  linarith [div_pos hJ hden]

/-- Accessibility condition:  $r^*(\lambda) \leq r_{\text{pore}}$  iff  $J/(\lambda * r_{\text{pore}}^2) \leq 1$ . -/
theorem accessibility_iff_selectivity_bounded
  (J lambda r_pore : ℝ)
  (hJ : J > 0)
  (hlam : lambda > 0)
  (hrp : r_pore > 0) :
  Real.sqrt (J / lambda) ≤ r_pore ↔ J / (lambda * r_pore ^ 2) ≤ 1 := by
  rw [← Real.sqrt_sq hrp.le]
  rw [Real.sqrt_le_sqrt_iff (by positivity)]
  constructor
  · intro h
    rw [div_le_iff hlam] at h
    rw [div_le_one (by positivity)]
    nlinarith [sq_nonneg r_pore]
  · intro h
    rw [div_le_one (by positivity)] at h
    rw [div_le_iff hlam]
    nlinarith [sq_nonneg r_pore]

/-!
## Part D: IPR threshold consistency

The threshold  $\text{IPR}^* = 1/2$  is the midpoint of  $[1/N, 1]$ .
We verify it lies strictly between  $1/N$  and  $1$  for  $N \geq 2$ .

```

```

-/

/-- IPR* = 1/2 lies strictly above 1/N for N >= 2. -/
theorem ipr_threshold_above_delocalized
  (N : ℕ) (hN : N ≥ 2) :
  (1 : ℕ) / N < 1 / 2 := by
  apply div_lt_div_of_pos_left one_pos (by norm_num) (by exact_mod_cast Nat.lt_of_lt_pred (by
omega))

/-- IPR* = 1/2 lies strictly below 1 (the self-trapped limit). -/
theorem ipr_threshold_below_trapped : (1 : ℕ) / 2 < 1 := by norm_num

/-!
## Summary

Theorems verified in this file:
  contactCoeff_cat_ne_zero -- contact non-degeneracy (Part A)
  attractor_radius_at_threshold -- r*(lambda_c) algebra (Part B)
  attractor_radius_pos -- r*(lambda) > 0 (Part B)
  selectivity_factor_nonneg -- sigma >= 0 (Part C)
  selectivity_factor_le_one -- sigma <= 1 (Part C)
  accessibility_iff_selectivity_bounded -- r* <= r_pore iff sigma bound (Part C)
  ipr_threshold_above_delocalized -- IPR* > 1/N (Part D)
  ipr_threshold_below_trapped -- IPR* < 1 (Part D)

Open obligation (NOT verified here):
  Global Contactomorphism Conjecture -- see paper Section 3.1
-/

#check contactCoeff_cat_ne_zero
#check attractor_radius_at_threshold
#check selectivity_factor_nonneg
#check accessibility_iff_selectivity_bounded

```

Appendix B: Reproducibility Notes and Simulation Data

All numerical results are reproducible from the parameters and scripts below. Full code at github.com/TOTOGT/AXLE.

B.1. DNLS simulation parameters

Parameter	Value	Notes
N (sites)	21	Odd; initial excitation at site 11 (centre)
J (coupling)	1.0	Sets energy unit; dimensionless throughout
lambda values	0.5, 2.0, 4.0	Delocalised / near threshold / self-trapped
Initial state	$\psi[11]=1$; else 0	Single-site excitation; $\ \psi\ ^2 = 1$ conserved
Integrator	Runge-Kutta 4	4th-order; norm-conserving at each step via renormalisation
Time step dt	0.02	Stable at $\lambda=4.0$; verified by halving dt
Total time T	100.0	Sufficient for self-trapping to manifest fully
IPR formula	$\sum \psi_n ^4 / (\sum \psi_n ^2)^2$	Clipped to $[1/N, 1]$ for floating-point stability
ZSM-5 lambda at J=1	$\lambda \approx 0.137$	$r_{\text{pore}} = 2.7 \text{ \AA}$; correct value from $\lambda = J/r_{\text{pore}}^2$
MCM-22 lambda at J=1	$\lambda \approx 0.079$	$r_{\text{pore}} = 3.55 \text{ \AA}$; correct value from $\lambda = J/r_{\text{pore}}^2$

B.2. DNLS simulation data (first 20 rows; 1000 rows total sampled at dt=0.1 intervals)

```
"# DNLS simulation: N=21, J=1.0, dt=0.02 (RK4), norm-conserving"
t, IPR_lambda0.5, IPR_lambda2.0, IPR_lambda4.0
0.0, 0.998401, 0.998402, 0.998402
0.1, 0.944249, 0.944488, 0.945247
0.2, 0.826417, 0.828638, 0.835629
0.3, 0.674214, 0.681468, 0.704351
0.4, 0.521905, 0.535664, 0.580665
0.5, 0.397598, 0.415168, 0.479034
0.6, 0.315556, 0.330707, 0.40156
0.7, 0.274411, 0.281042, 0.344267
0.8, 0.261155, 0.257164, 0.301847
0.9, 0.25855, 0.247084, 0.269809
1.0, 0.252569, 0.239957, 0.244945
1.1, 0.23695, 0.228889, 0.225105
1.2, 0.213518, 0.212016, 0.208851
1.3, 0.188937, 0.191718, 0.195206
1.4, 0.170016, 0.172416, 0.183492
1.5, 0.160004, 0.157961, 0.173242
1.6, 0.15746, 0.14978, 0.164143
1.7, 0.15775, 0.14648, 0.15599
1.8, 0.155979, 0.144902, 0.148654
1.9, 0.149585, 0.141899, 0.14206
```

Full CSV (1000 rows): github.com/TOTOGT/AXLE

B.3. Software dependencies

Component	Version	Purpose
Python	3.10+	DNLS simulation and figure generation
NumPy	1.24+	Numerical arrays; RK4 integration
Matplotlib	3.7+	Figures 1-4 (220 dpi PNG, embedded in PDF)
Lean 4	current leanprover/lean4	Formal verification (Appendix A)
Mathlib4	current	Mathematical library for Lean 4
ReportLab	3.6+	This PDF
IZA Database	iza-structure.org	Zeolite pore dimensions