

Table 3. Mean \pm SD $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, and $\delta^{34}\text{S}$ values, size (total length or carapace width in mm) and sample size of all consumers sampled. LA: Louisiana, NC: North Carolina, NJ: New Jersey, MA: Massachusetts. For MA, consumers samples (with exception of Striped Bass) were pooled prior to being run for stable isotopes resulting in no standard deviation.

Site	Species	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	$\delta^{34}\text{S}$	Size (mm)	n
LA	Blue crab	-17.4 ± 1.8	7.5 ± 1.2	8.2 ± 2.2	83.9 ± 43.8	28
	Killifish	-16.0 ± 1.0	8.3 ± 0.7	6.6 ± 2.1	82.7 ± 18	23
	Red drum	-18.5 ± 1.2	11.6 ± 2.1	11.3 ± 2.6	565.3 ± 167.4	12
	Spotted seatrout	-16.8 ± 0.3	15.7 ± 0.4	15.7 ± 1.1	502.5 ± 145	4
NC	Blue crab	-15.6 ± 0.6	8.9 ± 0.5	10.6 ± 1.0	110.7 ± 91.2	8
	Killifish	-16.3 ± 0.3	9.0 ± 0.5	11.1 ± 0.2	65.3 ± 18.7	4
	Red drum	-15.2 ± 1.1	10.5 ± 0.8	9.2 ± 0.8	386.1 ± 129.1	7
	Flounder	-16.4 ± 0.4	11.3 ± 0.1	10.2 ± 1.6	252 ± 80.9	3
	Spotted seatrout	-15.5 ± 4.1	10.2 ± 2.5	9.0 ± 2.9	371.5 ± 85.4	9
NJ	Blue crab	-18.2 ± 1.0	10.3 ± 1.3	10.0 ± 2.7	115.6 ± 15.7	9
	Killifish	-12.4 ± 0.5	7.2 ± 0.3	2.0 ± 1.2	67.8 ± 15.4	9
	Striped bass	-18.7 ± 1.1	14.6 ± 0.6	13.4 ± 1.7	520.7 ± 87.5	10
	Flounder	-20.5 ± 1.3	12.1 ± 1.2	13.6 ± 1.6	289.5 ± 71.3	9
MA	Blue crab	-11.3 ± 0.0	6.5 ± 0.0	5.4 ± 0.0	104.5 ± 13.4	4
	Mummichog	-15.8 ± 0.0	9 ± 0.0	6.4 ± 0.0	44.0 ± 8.5	15
	Striped bass	-18.0 ± 1.6	15.1 ± 1.3	11.4 ± 2.3	94.0 ± 280.0	34
	Flounder	-17.2 ± 0.0	10.8 ± 0	11.7 ± 0.0	197.0 ± 0.0	2