

SI Table 1. The trophic level [mean $\pm$ SD] of each oyster reef-associated consumers and the proportional contribution [median (lower 95% CI, upper 95% CI)] of each basal resource to consumers for each estuary sampled. White shrimp were not collected in NC and therefore not included for that estuary.

Estuary	Consumer	Trophic Level	Benthic microalgae	sPOM	<i>S. alterniflora</i>
NC	Oyster	1.70 $\pm$ 0.03	0.066 (0.003,0.187)	0.89 (0.762,0.98)	0.044 (0.004,0.121)
	Mud crab	1.89 $\pm$ 0.05	0.078 (0.004,0.214)	0.789 (0.651,0.926)	0.133 (0.01,0.278)
	Blue crab	2.29 $\pm$ 0.05	0.158 (0.011,0.379)	0.682 (0.476,0.882)	0.161 (0.021,0.359)
	Pinfish	2.33 $\pm$ 0.05	0.158 (0.006,0.357)	0.707 (0.574,0.841)	0.135 (0.011,0.297)
NI-WB	Oyster	1.92 $\pm$ 0.01	0.933 (0.849,0.986)	0.052 (0.007,0.134)	0.015 (0.001,0.045)
	Mud crab	2.23 $\pm$ 0.02	0.856 (0.739,0.957)	0.094 (0.019,0.187)	0.05 (0.002,0.158)
	White shrimp	2.2 $\pm$ 0.02	0.601 (0.492,0.703)	0.302 (0.222,0.381)	0.097 (0.003,0.226)
	Blue crab	2.63 $\pm$ 0.02	0.729 (0.61,0.841)	0.193 (0.107,0.28)	0.078 (0.005,0.21)
	Pinfish	2.75 $\pm$ 0.02	0.848 (0.752,0.949)	0.13 (0.034,0.226)	0.022 (0.002,0.061)
SI	Oyster	2.10 $\pm$ 0.02	0.81 (0.679,0.939)	0.175 (0.048,0.306)	0.014 (0,0.046)
	Mud crab	2.98 $\pm$ 0.02	0.748 (0.625,0.879)	0.234 (0.101,0.359)	0.019 (0.001,0.066)
	White shrimp	2.20 $\pm$ 0.02	0.41 (0.309,0.513)	0.5 (0.392,0.601)	0.09 (0.001,0.231)
	Blue crab	2.68 $\pm$ 0.02	0.558 (0.458,0.66)	0.405 (0.295,0.502)	0.038 (0.001,0.118)
	Pinfish	3.03 $\pm$ 0.01	0.565 (0.469,0.673)	0.416 (0.308,0.514)	0.018 (0.001,0.056)
GTM	Oyster	2.32 $\pm$ 0.01	0.009 (0.001,0.023)	0.977 (0.937,0.996)	0.015 (0,0.053)
	Mud crab	2.63 $\pm$ 0.03	0.024 (0.003,0.058)	0.929 (0.83,0.99)	0.047 (0.001,0.147)
	White shrimp	2.12 $\pm$ 0.02	0.335 (0.265,0.399)	0.633 (0.573,0.695)	0.032 (0.003,0.082)
	Blue crab	2.56 $\pm$ 0.05	0.081 (0.02,0.173)	0.843 (0.631,0.957)	0.076 (0.003,0.276)
	Pinfish	2.56 $\pm$ 0.03	0.067 (0.007,0.163)	0.889 (0.773,0.981)	0.044 (0.001,0.138)

SI Table 2. Trophic niche size and Sørensen overlap index (proportional values range between 0-1) output from hypervolume analysis for all consumers (blue crabs, mud crabs, oysters, pinfish, and white shrimp) sampled across four estuaries in the SAB. White shrimp were not collected in NC and therefore not included for that estuary.

Consumer	Estuary	Niche size	Sørensen Overlap Index			
			NC	NI-WB	SI	GTM
Oyster	NC	0.88	-	0	0	0.02
	NI-WB	0.15	0	-	0.11	0
	SI	0.68	0	0.11	-	0
	GTM	0.02	0.02	0	0	-
Mud Crab	NC	3.49	-	0	0	0.11
	NI-WB	1.34	0	-	0.17	0
	SI	1.07	0	0.17	-	0
	GTM	0.29	0.11	0	0	-
Blue crab	NC	6.31	-	0		0.24
	NI-WB	1.28	0	-	0.004	0
	SI	1.05	0	0.004	-	0
	GTM	2.47	0.24	0	0	-
Pinfish	NC	6.4	-	0	0	0.15
	NI-WB	0.47	0	-	0	0
	SI	0.29	0	0	-	0
	GTM	1.15	0.15	0	0	-
White Shrimp	NI-WB	1.61	-	-	0	0
	SI	1.19	-	0	-	0.05
	GTM	0.22	-	0	0.05	-