

Table 1. Bulk carbon ($\delta^{13}\text{C}$), nitrogen ($\delta^{15}\text{N}$), and sulfur ($\delta^{34}\text{S}$) stable isotope values (mean $\pm$ SD) and sample sizes (n) for all basal resources (*S. alterniflora*, sPOM, and benthic microalgae) across four estuaries in the SAB. Estuaries included North Carolina (NC), North Inlet–Winyah Bay (NI-WB), Sapelo Island (SI), and Guana Tolomato Matanzas (GTM).

Estuary	Basal resource	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	$\delta^{34}\text{S}$	n
NC	<i>S. alterniflora</i>	-14.9 ± 0.3	3.1 ± 1.3	-3.6 ± 6.6	10
	sPOM	-20.7 ± 1.4	5.6 ± 0.3	11.5 ± 9.3	3
	Benthic microalgae	-18.2 ± 1.6	3.5 ± 1.3	-3.5 ± 6.7	5
NI-WB	<i>S. alterniflora</i>	-14.9 ± 0.2	6 ± 0.9	6.3 ± 6.9	10
	sPOM	-21.1 ± 0.4	4 ± 0.2	-15.3 ± 1	5
	Benthic microalgae	-17.6 ± 0.9	3.6 ± 1.2	3.5 ± 9.7	8
SI	<i>S. alterniflora</i>	-14.9 ± 0.3	5.4 ± 0.8	2.9 ± 6	10
	sPOM	-21.6 ± 0.3	4.6 ± 0.2	-5.8 ± 2.2	10
	Benthic microalgae	-17.6 ± 0.9	3.6 ± 1.2	-14 ± 11.6	8
GTM	<i>S. alterniflora</i>	-14.6 ± 0.2	6.3 ± 0.7	7.6 ± 8.3	10
	sPOM	-22.6 ± 0.3	3.9 ± 0.6	5 ± 3.2	7
	Benthic microalgae	-18.3 ± 0.3	2.8 ± 0.6	-13.7 ± 4.7	6

Table 2. Bulk carbon ($\delta^{13}\text{C}$), nitrogen ($\delta^{15}\text{N}$), and sulfur ($\delta^{34}\text{S}$) stable isotope values (mean $\pm$ SD) and sample sizes (n) for all oyster reef-associated consumers (oyster, mud crab, white shrimp, blue crab, and pinfish) sampled across four estuaries in the SAB. Estuaries included North Carolina (NC), North Inlet–Winyah Bay (NI-WB), Sapelo Island (SI), and Guana Tolomato Matanzas (GTM). White shrimp were not collected in NC and therefore not included for that estuary.

Estuary	Consumer	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	$\delta^{34}\text{S}$	n
NC	Oyster	-18.8 ± 0.2	7.6 ± 0.4	16.5 ± 0.6	10
	Mud crab	-15.5 ± 2	8 ± 1.7	14.8 ± 1.7	8
	Blue crab	-16.1 ± 0	9.1 ± 0	10.8 ± 0	1
	Pinfish	-16.6 ± 1.7	9.3 ± 1.8	12.1 ± 3.5	9
NI-WB	Oyster	-19.1 ± 0.3	8.2 ± 0.3	16.7 ± 0.9	10
	Mud crab	-16.6 ± 0.5	9.2 ± 1.1	15 ± 0.6	8
	White shrimp	-16 ± 0.9	8.9 ± 0.7	7 ± 2.2	10
	Blue crab	-15.3 ± 0.8	10.4 ± 0.4	10.6 ± 2.3	11
	Pinfish	-18.3 ± 1.3	10.8 ± 0.5	13.2 ± 0.9	10
SI	Oyster	-20.2 ± 0.4	7.4 ± 0.3	16.1 ± 0.6	9
	Mud crab	-18.7 ± 0.3	10.4 ± 0.4	15.1 ± 0.6	10
	White shrimp	-16.3 ± 0.5	8.2 ± 0.5	6.8 ± 1.5	9
	Blue crab	-16 ± 0.7	9.6 ± 0.5	10 ± 1.6	10
	Pinfish	-18.4 ± 1.4	10.7 ± 0.4	10.1 ± 1.3	10
GTM	Oyster	-19.3 ± 1.1	8.3 ± 0.6	15.7 ± 1	10
	Mud crab	-18.3 ± 1.1	9.4 ± 0.7	13.5 ± 1.4	7
	White shrimp	-17.7 ± 1.3	7.3 ± 0.4	3.2 ± 1	10
	Blue crab	-14.6 ± 0	9.2 ± 0	10.1 ± 0	1
	Pinfish	-19 ± 1.3	9.1 ± 1.3	10.3 ± 0.1	2