Peak Prediction Using Quantile Regression and Asymmetric Laplace Noise

Abstract

Predicting the location of a peak before it is fully observed is important in many applications. We consider a Gaussian-shaped signal with additive noise and develop a quantile-based method for estimating its amplitude, location, and spread. For fixed location and spread, the amplitude estimator is a weighted τ-quantile of a transformed version of the observations. A τ-aligned asymmetric Laplace noise model provides a probabilistic interpretation of the quantile objective through its likelihood. We study the sequential evolution of the estimated peak location as the observation horizon increases. When the estimate does not stabilize sufficiently early, we propose a two-stage construction of candidate entry–exit intervals based on oscillations in the estimation trajectory. These intervals can be constructed without knowing the true peak location. Monte Carlo simulations evaluate the interval-selection rate and empirical coverage, showing that performance depends on the signal-to-noise ratio. Stabilization is more reliable under the high- and moderate-SNR settings considered, whereas low SNR reduces point-estimation reliability and makes interval coverage more variable. Finally, we illustrate the method using daily COVID-19 cases during the first wave of the pandemic in China, France, and Germany.

Keywords: Peak prediction; quantile regression; asymmetric Laplace noise; sequential estimation; entry–exit intervals; signal-to-noise ratio; COVID-19.

Mathematics subject classification: 62M10, 62G35, 60G35.
JEL classification: C22, C21, C53, I18.

1 Introduction

Predicting peaks in time series data is important across a wide range of applications involving proactive decision-making, risk management, resource allocation, and disaster preparedness. Examples include anticipating disease outbreaks in epidemiology and peak demand in energy systems. Peak prediction is challenging because observations are affected by noise, and the underlying signal may be difficult to identify from partial observations. Measurement errors and random fluctuations can make the location and amplitude of a peak difficult to estimate before it is fully observed. Moreover, real-world time series may exhibit multiple peaks, irregular peak shapes, or other temporal structures that are not adequately described by simple forecasting models.

Popular forecasting approaches, such as Autoregressive Integrated Moving Average (ARIMA) models and Recurrent Neural Networks (RNNs), can be effective for describing temporal dependence and forecasting general patterns, but they are not specifically designed to estimate the timing and magnitude of a future peak. A model that provides a good overall forecast may not necessarily provide an accurate estimate of the location of a peak. There has also been comparatively little theoretical work devoted specifically to peak prediction in time series. One contribution in this direction is the thesis of Stewart et al. (2023), which studies the interplay between trend, seasonality, and noise; see also (Yu et al., 2019; Pei and Shaman, 2017).

Our approach builds on the framework developed in Dermoune et al. (2026). We consider observations available up to a current observation horizon T and a prescribed search horizon $h > T$. The observations are modeled as a Gaussian-shaped signal with amplitude $A > 0$, peak location L, and spread $S > 0$, corrupted by additive noise. Our objective is to estimate these parameters from the observations available up to time T and, in particular, to predict the peak location L before it is fully observed.

Our first contribution is to extend the least-absolute-deviation criterion used in the symmetric case to a general quantile regression criterion indexed by $\tau \in (0, 1)$. As shown in Section 2, for fixed location and spread, the amplitude minimizing this criterion is a weighted τ-quantile of a transformed version of the observations. The weighted-median estimator is recovered as the special case $\tau = 1/2$. We also examine the behavior of the quantile-based objective in the noise-free setting in Section 3.

Our second contribution is a probabilistic interpretation of the estimation criterion. In Section 4, we introduce a τ-aligned asymmetric Laplace noise model whose negative log-likelihood is proportional to the quantile check loss. This connection provides a unified framework for studying the effects of the quantile level τ and the signal-to-noise ratio (SNR) on peak estimation, with the symmetric Laplace model recovered when $\tau = 1/2$. Section 5 investigates the evolution of the sequential peak-location estimator as the observation horizon increases. In particular, we study its stabilization time, defined as the first horizon from which the estimated location remains constant over all subsequent horizons, subject to a minimum plateau-length condition.

Our third contribution addresses situations in which the sequential estimate does not sta-

bilize sufficiently early to provide a reliable point prediction. In Section 6, we develop a data-driven construction of candidate entry and exit times based on the oscillations of the sequential peak-location estimates. These times define candidate intervals for the peak region using only the estimation trajectory, without requiring knowledge of the true peak location L. We then introduce an operational rule for selecting a candidate interval based on whether it contains the current point estimate. The performance of this procedure is assessed through Monte Carlo simulations, including empirical interval-selection rates and coverage probabilities. The true peak location is used only retrospectively to evaluate the selected intervals; the construction itself does not guarantee a prescribed coverage level.

Finally, we illustrate the methodology using daily COVID-19 case counts during the first wave of the pandemic in China (Barmparis and Tsironis, 2020), France, and Germany. The real-world application in Section 7 examines the estimated peak trajectories and the behavior of the proposed entry–exit construction across several quantile levels.

The remainder of the paper is organized as follows. Section 2 introduces the Gaussian peak model and the quantile-based estimation procedure, including the weighted-quantile characterization of the amplitude estimator. Section 3 examines the quantile-based objective in the noise-free setting. Section 4 introduces the τ-aligned asymmetric Laplace noise model and establishes its connection with the quantile loss. Section 5 studies the stabilization of the sequential peak-location estimator, while Section 6 develops the candidate entry–exit intervals and evaluates the operational selection procedure. Section 7 presents the real-world COVID-19 application, including comparisons across quantile levels and tables of estimated peak trajectories.

2 Quantile-Based Estimation Methodology

We consider the problem of estimating the amplitude, location, and spread of a unimodal signal corrupted by noise. The model is:

$$I(t) = A \exp\left(-\frac{(t-L)^2}{S^2}\right) + e(t), \qquad t = 1, \ldots, T. \tag{1}$$

where $A > 0$ is the amplitude, $L > 0$ the peak location, $S > 0$ the spread (scale), and $e(t)$ is additive noise. Given observations $(I(t))_{t=1}^{T}$, our goal is to estimate the underlying parameters using quantile regression, i.e., minimizing the asymmetric loss

$$\rho_\tau(u) = u\left(\tau - \mathbf{1}_{\{u<0\}}\right). \tag{2}$$

for a fixed quantile level $\tau \in (0, 1)$. We define the quantile objective:

$$f_\tau(T, a, l, s) = \frac{1}{T}\sum_{t=1}^{T} \rho_\tau\left(I(t) - a\exp\left(-\frac{(t-l)^2}{s^2}\right)\right). \tag{3}$$

For fixed candidate values of (l, s), the optimal amplitude a minimizes $a \mapsto f_\tau(T, a, l, s)$. This minimizer is a weighted τ-quantile of a transformed version of the data, as described below.

2.1 Weighted Quantile Estimation of Amplitude

Let $w(t) = \exp\left(-\frac{(t-l)^2}{s^2}\right)$ be the weight associated with time point t, and define:

$$x(t) = I(t) \exp\left(\frac{(t-l)^2}{s^2}\right).$$

Then the optimal amplitude a is the weighted τ-quantile of $x(t)$ with weights $w(t)$. The following proposition gives the corresponding characterization of the weighted quantile as the minimizer of the weighted quantile loss. The proof of the following Proposition is the same as in the work of Novoselac (2020).

Proposition 1. *Let $(x(t), w(t))$ be a sequence of real values with positive weights $w(t) > 0$. Then, for any $\tau \in (0, 1)$, the weighted τ-quantile minimizes the weighted quantile loss:*

$$q_\tau = \arg\min_{a \in \mathbb{R}} \sum_{t=1}^{T} w(t)\, \rho_\tau\big(x(t) - a\big),$$

where ρ_τ is the quantile check function defined in (2). To compute q_τ, sort the values $x(t)$ in increasing order, obtaining a permutation π such that

$$x(\pi(1)) \le x(\pi(2)) \le \cdots \le x(\pi(T)).$$

Define the reordered weights $w_\pi(t) = w(\pi(t))$. Let

$$k = \min\left\{ j \in \{1, \ldots, T\} : \sum_{t=1}^{j} w_\pi(t) \ge \tau \sum_{t=1}^{T} w(t) \right\}.$$

Then the weighted τ-quantile satisfies $q_\tau = x(\pi(k))$.

2.2 Quantile-Based Estimation Procedure

We define the following steps to jointly estimate the location $\hat{l}(T)$, amplitude $\hat{a}(T)$, and spread $\hat{s}(T)$:

Algorithm 1 Quantile-based estimation of peak parameters

Require: Time horizon T, upper bound $h \ge T$, quantile level $\tau \in (0, 1)$

1: For each candidate $l \in [1, h]$, find $s^*(l) = \arg\min_s f_\tau(T, a^*(l, s), l, s)$
2: Set $\hat{l}(T) = \arg\min_l f_\tau(T, a^*(l, s^*(l)), l, s^*(l))$
3: Set $\hat{s}(T) = s^*(\hat{l}(T))$
4: Set $\hat{a}(T) = a^*(\hat{l}(T), \hat{s}(T))$

Ensure: Estimated parameters $(\hat{l}(T), \hat{a}(T), \hat{s}(T))$

2.3 Remark on Asymmetry and Robustness

This method generalizes the classic Least Absolute Deviations (LAD) estimator, recovered at $\tau = 0.5$. By adjusting τ, we introduce asymmetry into the estimation loss, enabling the model to better adapt to skewed noise distributions. This makes the approach especially valuable in early-warning systems and epidemic peak detection where overestimation and underestimation have asymmetric consequences.

3 Quantile-Based Objective Function under Noise-Free Conditions

We first consider the idealized scenario where the signal is not corrupted by noise. The model of interest is given by

$$I(t) = A \exp\left(-\frac{(t-L)^2}{S^2}\right), \quad t = 1, \ldots, T,$$

where $A = 15000$, $L = 22$, $S = 10$, and $T \in \{2, 3, 10, 20, 21\}$. The quantile-based estimation procedure is implemented by minimizing the following objective function (3). The amplitude a is computed via a weighted τ-quantile, as described in Section 2.

Since the goal is to predict the location L of the peak before it becomes observable, we restrict the minimization over candidate peak locations $l > T$. For each value of $\tau \in \{0.25, 0.5, 0.75, 0.95\}$, we compute and visualize the objective curve $l \mapsto f_\tau(T, a(T, l), l, s(T, l))$ for the selected values of T (see Figure 1).

4 Quantile-Based Estimation with τ-Aligned Noise

To analyze how quantile-aligned noise affects peak estimation, we introduce a unified noise model based on the *general asymmetric Laplace* (GAL) distribution. This flexible family allows both symmetric and asymmetric noise structures while maintaining the property that the τ-quantile of the noise is at zero.

4.1 Asymmetric Laplace Noise Aligned with the τ-Quantile

We consider an asymmetric Laplace distribution parametrized by a location parameter $m \in \mathbb{R}$, a scale parameter $\lambda > 0$, and an asymmetry parameter $k > 0$. Its density is given by

$$f_{m,\lambda,k}(x) = \frac{\lambda}{k + k^{-1}} \begin{cases} \exp\left(\dfrac{\lambda(x-m)}{k}\right), & x < m, \\ \exp\left(-\lambda k(x-m)\right), & x \geq m. \end{cases}$$




Figure 1: Objective function $l \mapsto f_\tau(T, a, l, s)$ computed for $\tau \in \{0.25, 0.5, 0.75, 0.95\}$ and $T \in \{2, 3, 10, 20, 21\}$, under a noise-free signal. The vertical dashed line indicates the true peak location $L = 22$, and the search is restricted to values $l > T$.

The parameter m is the mode of the distribution, while k controls its asymmetry and λ controls its concentration. The distribution function evaluated at the mode is

$$\mathbb{P}(X < m) = \frac{\lambda}{k + k^{-1}} \int_{-\infty}^{m} \exp\left(\frac{\lambda(x-m)}{k}\right) dx = \frac{k^2}{1+k^2}.$$

Therefore, to align the asymmetric Laplace noise with the τ-quantile regression criterion, we require

$$\mathbb{P}(e(t) < 0) = \tau, \tag{4}$$

and $\tau \in (0,1)$. Since the location of the noise can be fixed at zero, we set $m = 0$. The corresponding asymmetry parameter is then determined by

$$\frac{k^2}{1+k^2} = \tau,$$

or equivalently,

$$k = \sqrt{\frac{\tau}{1-\tau}}. \tag{5}$$

Thus, the τ-aligned asymmetric Laplace noise has density

$$f_{\tau,\lambda}(x) = \frac{\lambda}{k + k^{-1}} \begin{cases} \exp\left(\dfrac{\lambda x}{k}\right), & x < 0, \\ \exp(-\lambda k x), & x \geq 0, \end{cases}$$

where k is given by (5). By construction, the noise satisfies the τ-alignment property (4). In particular, when $\tau = 1/2$, we have $k = 1$, and the asymmetric Laplace distribution reduces to the symmetric Laplace distribution centered at zero. Hence, the symmetric Laplace case is recovered as a special case of the τ-aligned asymmetric Laplace model.

4.2 Simulation under Asymmetric Laplace Noise

We now investigate the behavior of the quantile-based peak estimation procedure when the additive noise follows the τ-aligned asymmetric Laplace distribution introduced in Section 4.1. In the parameterization based on the quantile loss, the density of the noise can be written as

$$f_{\tau,b}(x) = \frac{\tau(1-\tau)}{b} \exp\left(-\frac{1}{b}\rho_\tau(x)\right), \qquad \tau \in (0,1), \quad b > 0. \tag{6}$$

where ρ_τ is the check function defined in (2). Equivalently,

$$f_{\tau,b}(x) = \begin{cases} \dfrac{\tau(1-\tau)}{b} \exp\left(-\dfrac{\tau x}{b}\right), & x \geq 0, \\ \dfrac{\tau(1-\tau)}{b} \exp\left(\dfrac{(1-\tau)x}{b}\right), & x < 0. \end{cases}$$

The normalization follows from

$$\int_{-\infty}^{+\infty} \exp\left(-\frac{1}{b}\rho_\tau(x)\right) dx = \frac{b}{1-\tau} + \frac{b}{\tau} = \frac{b}{\tau(1-\tau)}.$$

Moreover, by construction, the noise satisfies the τ-alignment property (4). This distribution is directly compatible with the quantile-based estimation criterion. Indeed,

$$-\log f_{\tau,b}(e) = -\log\left(\frac{\tau(1-\tau)}{b}\right) + \frac{1}{b}\rho_\tau(e).$$

Therefore, for fixed τ and b, minimizing the negative log-likelihood is equivalent to minimizing the empirical quantile loss. The asymmetric Laplace model thus provides a probabilistic interpretation of the quantile-based peak estimation procedure without introducing a separate estimation criterion. For the simulation study, the true signal is

$$I(t) = A \exp\left(-\frac{(t-L)^2}{S^2}\right),$$

with $A = 1$, $L = 22$, $S = 10$. Since the maximum amplitude of the Gaussian signal is $\max_t I(t) = A$, we define the signal-to-noise ratio as the ratio between the peak amplitude of the signal and the standard deviation of the noise:

$$\mathrm{SNR} = \frac{A}{\sqrt{\mathrm{Var}(e(t))}}. \tag{7}$$

In the simulations, we set SNR = 1000. For each $\tau \in \{0.25, 0.5, 0.75, 0.95\}$, the corresponding asymmetric Laplace noise is generated with asymmetry parameter (5). Thus, the noise variance is calibrated according to

$$\mathrm{Var}(e(t)) = \left(\frac{A}{\mathrm{SNR}}\right)^2. \tag{8}$$

For the τ-aligned asymmetric Laplace distribution, this variance is

$$\mathrm{Var}(e(t)) = b^2 \frac{\tau^2 + (1-\tau)^2}{\tau^2(1-\tau)^2}. \tag{9}$$

Combining (8) and (9) gives

$$b = \frac{A}{\mathrm{SNR}} \frac{\tau(1-\tau)}{\sqrt{\tau^2 + (1-\tau)^2}}. \tag{10}$$

For each τ, the objective function $l \mapsto f_\tau(T, a, l, s)$ is evaluated for the time horizons $T \in \{4, 6, 8, 9, 10\}$. For each candidate peak location l, the scale parameter s is optimized over the prescribed interval, and the amplitude a is determined as the weighted τ-quantile described in Section 2.1. The resulting objective profiles are then compared across the different values of τ. The resulting profiles are displayed in Figure 2. The true peak location $L = 22$ is marked by a vertical dashed line. Each panel corresponds to a different pair (T, τ), and the curves show the behavior of the quantile-loss objective across candidate peak locations.




Figure 2: Objective function profiles $l \mapsto f_\tau(T, a, l, s)$ under *asymmetric Laplace* noise for all combinations of $\tau \in \{0.25, 0.5, 0.75, 0.95\}$ (columns) and $T \in \{4, 6, 8, 9, 10\}$ (rows). Each subfigure shows the loss function for a specific T and τ pair, with the dashed vertical line indicating the true peak location $L = 22$.

5 Stabilization of the Sequential Estimator

We study how the estimated peak location $\hat{l}(T)$ evolves as the observation horizon T increases. We use the model (1) with $h = 30$, $A = 1$, $L = 22$, $S = 10$, τ-aligned asymmetric Laplace noise (Section 4.1), and SNR $\in \{1000, 100, 1\}$. At each $T \in \{4, \ldots, h\}$, $(\hat{l}(T), \hat{a}(T), \hat{s}(T))$ minimizes the objective (3); only $\hat{l}(T)$ is considered below.

5.1 Stabilization time

We define the stabilization time of the path $T \mapsto \hat{l}(T)$ as

$$T^* = \min\{T \in \{4, \ldots, h\} : \ \hat{l}(u) = \hat{l}(T) \text{ for all } u \in \{T, \ldots, h\}\},$$

the first horizon from which the estimate no longer changes. This definition does not use L. To exclude a trivial plateau reduced to the last horizon(s), we require the plateau to contain at least $m = 3$ horizons, $h - T + 1 \geq m$, and set $T^* = \infty$ otherwise. In simulations, where L is known, we also record whether the plateau value equals L, that is, $\hat{l}(T^*) = L$. The operational entry time $T^{(0)}_{\text{entry}}(T)$ of Section 6.4 is a different quantity, defined from the oscillations of the path.

5.2 Large and moderate signal-to-noise ratio

For SNR $= 1000$ and SNR $= 100$, the estimated peak location $\hat{l}(T)$ reaches a plateau whose value equals L in all replicates. The stabilization time satisfies $T^* \leq L$ in all replicates at SNR $= 1000$ and in 99% of replicates at SNR $= 100$. For $\tau = 0.50$ and $M = 200$ independent replicates, Table 1 reports median stabilization times of 12 and 18, respectively. In the single realization used in Section 6.5 (seed 280, $\tau = 0.50$), $T^* = 9$ at SNR $= 1000$. In these regimes the point estimate identifies L after a short transient, and T^* plays the role of an entry time. No exit time is needed.

5.3 Small signal-to-noise ratio

For SNR $= 1$, the path rarely settles. In the same experiment ($\tau = 0.50$, $M = 200$), a plateau of at least $m = 3$ horizons exists in only 34% of the replicates ($T^* < \infty$), with median $T^* = 27$, and $T^* > L$ or $T^* = \infty$ in 93% of them. The plateau value equals L in only 4.5% of all replicates (about 13% of those with a plateau). Without the minimal length ($m = 1$), T^* is always finite, with median $30 = h$, because the last horizon is a trivial plateau; then $\hat{l}(T^*) = L$ in 13.5% of the replicates. Figure 3 shows two replicates: one with finite $T^* = 24$ and one with $T^* = \infty$. In this regime the path alone does not indicate when its current value can be trusted. This motivates the entry times, exit times and the data-driven interval defined in Section 6.1 onwards, which are studied for SNR $= 1$ in Section 6.5. In the finite example the estimate equals L at $T = 22$ but leaves it at $T = 23$, so a single hit of L does not indicate stabilization.

Table 1: Stabilization time T^* (plateau of at least $m = 3$ horizons) for $\tau = 0.50$, based on $M = 200$ replicates. Median T^* is computed over replicates with finite T^*; $T^* = \infty$ is counted as $T^* > L$. The last column is the proportion of all replicates with $\hat{l}(T^*) = L$.

SNR	median T^*	$\mathbb{P}(T^* < \infty)$	$\mathbb{P}(T^* > L)$	$\mathbb{P}(\hat{l}(T^*) = L)$
1000	12	1.00	0.00	1.00
100	18	1.00	0.01	1.00
1	27	0.34	0.93	0.045




Figure 3: Trajectories $T \mapsto \hat{l}(T)$ for $\tau = 0.50$. Top: ten replicates at SNR $= 1000$ and SNR $= 100$. Bottom (SNR $= 1$): a replicate with finite $T^* = 24$ (the smallest finite value among replicates whose plateau equals L), and a replicate with $T^* = \infty$ (no plateau of at least three horizons). The dashed line is $L = 22$ and black dots mark T^*.

6 Entry and Exit Times of the Peak Region

The sequential estimation procedure introduced in the preceding sections provides, for each observation horizon T, an estimate $\hat{l}(T)$ of the peak location. Here, T denotes the current observation horizon, so that the estimator is constructed using only the observations available on $[1, T]$, while the candidate peak location is searched over the fixed horizon $[1, h]$, with $T \leq h$ and $L < h$. The case $T = h$ corresponds to the full-sample estimator and is used in the real-data application of Section 7. Thus, although only the observations $1, \ldots, T$ are used in the estimation, the estimator may identify a peak location beyond the current observation horizon T. For notational simplicity, we write $\hat{l}(T)$ instead of $\hat{l}(T, h)$ when the horizon h is fixed.

The stabilization time defined in Section 5 characterizes the first horizon from which the estimated peak location remains constant over all subsequent horizons, subject to the minimum plateau-length condition introduced there. In this section, we introduce a complementary construction based on the oscillations of the estimated peak location. This construction aims at identifying candidate time intervals describing the region in which the estimated peak location becomes stable, through a data-driven time of entry and a time of exit.

The operational construction is based only on the sequential estimation trajectory $T \longmapsto \hat{l}(T)$. The true peak location L is not required to construct the candidate entry and exit times. In the simulation study, where L is known, it is used only for an ex post assessment of whether a candidate peak region contains the true peak.

6.1 Oscillations of the estimated peak location

We fix a sufficiently large horizon h such that the true peak location satisfies $L < h$. For each observation horizon $T \leq h$, the sequential estimator $\hat{l}(T)$ is constructed using only the observations available on $[1, T]$, while the candidate peak location is searched over $[1, h]$. To characterize the temporal stability of the estimated peak location using only the information available at the current observation horizon T, we consider the oscillations of the estimation path over intervals contained in the observed time range $[1, T]$. For two times t and T satisfying $T_0 \leq t \leq T \leq h$, we define the oscillation of the estimated peak location over the interval $[t, T]$ by

$$\omega_l(t, T) = \max_{t \leq t_1, t_2 \leq T} \left| \hat{l}(t_1) - \hat{l}(t_2) \right|. \tag{11}$$

Thus, $\omega_l(t, T)$ measures the largest variation of the estimated peak location over the observed interval $[t, T]$. In particular, $\omega_l(T, T) = 0$. For a fixed observation horizon T, the function $t \longmapsto \omega_l(t, T)$, with $T_0 \leq t \leq T$, is nonincreasing. Indeed, if $t_1 < t_2 \leq T$, then $[t_2, T] \subset [t_1, T]$, and consequently $\omega_l(t_2, T) \leq \omega_l(t_1, T)$. The strict decreases of this sequence identify the times at which the oscillation level changes when the starting point of the observed interval is moved forward.

6.2 Strict-decrease points of the retrospective oscillation

For a fixed observation horizon $T \leq h$, consider the nonincreasing sequence $t \longmapsto \omega_l(t, T)$, with $t \in \{T_0, \ldots, T\}$. We call an observation time $t < T$ a *strict-decrease point* of the retrospective oscillation if

$$\omega_l(t, T) > \omega_l(t + 1, T).$$

We order the strict-decrease points in decreasing order. Thus, if $t_3 < t_2 < t_1 < T$ are the strict-decrease points, then $t_1 > t_2 > t_3$. The intervals between consecutive strict-decrease points correspond to constant plateaus of the nonincreasing sequence $t \mapsto \omega_l(t, T)$. More precisely, if $U > V$ are two consecutive strict-decrease points, then

$$\omega_l(t, T) = \omega_l(U, T), \qquad t \in \{V + 1, \ldots, U\}.$$

Thus, for each current observation horizon T, the strict-decrease points characterize the successive decreases of the retrospective oscillation computed using only the observations available on $[1, T]$.

6.3 Construction of the candidate exit times

For each current observation horizon $T \leq h$, the candidate exit times are defined from the strict-decrease points of the sequence $t \longmapsto \omega_l(t, T)$. We set $T^{(0)}_{\text{exit}}(T) = T$, and let $T^{(1)}_{\text{exit}}(T) > T^{(2)}_{\text{exit}}(T) > \cdots$ denote the strict-decrease points of $t \mapsto \omega_l(t, T)$ that are strictly smaller than T, ordered in decreasing order. For example, if the strict-decrease points are $t_3 < t_2 < t_1 < T$, then

$$T^{(1)}_{\text{exit}}(T) = t_1, \qquad T^{(2)}_{\text{exit}}(T) = t_2, \qquad T^{(3)}_{\text{exit}}(T) = t_3.$$

The convention $T^{(0)}_{\text{exit}}(T) = T$ provides the initial candidate exit time, for which $\omega_l\left(T^{(0)}_{\text{exit}}(T), T\right) = \omega_l(T, T) = 0$. For each candidate index $k \geq 0$, we define the corresponding oscillation level by

$$\epsilon'_k(T) = \omega_l\left(T^{(k)}_{\text{exit}}(T), T\right).$$

Thus, for each current observation horizon T, the candidate exit times correspond to the successive strict decreases of the retrospective oscillation, while the associated levels $\epsilon'_k(T)$ record the oscillation at those candidate exit times. This construction is based only on the sequential estimation trajectory available up to the current observation horizon T and does not require knowledge of the true peak location L. Operationally, for each fixed T, the candidate exit times are obtained by scanning the sequence $t \longmapsto \omega_l(t, T)$ for its successive strict-decrease points. In numerical implementation, a small tolerance can be used when determining strict decreases.

6.4 Determination of the entry time

For a given current observation horizon $T \leq h$ and a candidate exit time $T_{\text{exit}}^{(k)}(T)$, we first consider all observation times strictly before the candidate exit time. The set

$$E_k(T) = \left\{ t < T_{\text{exit}}^{(k)}(T) : \omega_l \left(t, T_{\text{exit}}^{(k)}(T) \right) \geq \epsilon_k'(T) \right\}, \tag{12}$$

defines the Stage-2 set. If $E_k(T)$ is empty, the candidate does not yield a valid entry time. Otherwise, we define $T_k^*(T) = \max E_k(T)$. The corresponding oscillation level is $\epsilon_k(T) = \omega_l \left(T_k^*(T), T_{\text{exit}}^{(k)}(T) \right)$. We then consider all observation times $t < T_{\text{exit}}^{(k)}(T)$ at which the oscillation equals $\epsilon_k(T)$. The entry time is defined as the smallest such observation horizon:

$$T_{\text{entry}}^{(k)}(T) = \min \left\{ t < T_{\text{exit}}^{(k)}(T) : \omega_l \left(t, T_{\text{exit}}^{(k)}(T) \right) = \epsilon_k(T) \right\}, \tag{13}$$

provided that such an observation time exists. The width associated with the candidate is

$$w_k(T) = T_{\text{exit}}^{(k)}(T) - T_{\text{entry}}^{(k)}(T). \tag{14}$$

The resulting candidate peak region at observation horizon T is therefore represented by the interval

$$\mathcal{R}_k(T) = \left[T_{\text{entry}}^{(k)}(T), T_{\text{exit}}^{(k)}(T) \right].$$

All endpoints are determined entirely from the sequential estimation trajectory available up to the current observation horizon T and from the corresponding retrospective oscillation construction.

6.5 Construction of a data-driven interval for the peak location

We illustrate the construction of the entry–exit intervals introduced in Sections 6.1–6.4 and describe how one of these intervals can be selected using only the observations available at the current horizon. The simulation setting is the one considered in Sections 4 and 5, with $A = 1, L = 22, S = 10, h = 30$, and quantile levels $\tau \in \{0.25, 0.50, 0.75, 0.95\}$. We consider three signal-to-noise ratios, SNR $\in \{1000, 100, 1\}$, corresponding to large, moderate, and small signal-to-noise regimes, respectively. For each current observation horizon T, the two-stage procedure produces a collection of candidate intervals

$$\mathcal{R}_k(T) = \left[T_{\text{entry}}^{(k)}(T), T_{\text{exit}}^{(k)}(T) \right],$$

where k indexes the candidate exit times constructed in Section 6.3. The width of candidate k is

$$w_k(T) = T_{\text{exit}}^{(k)}(T) - T_{\text{entry}}^{(k)}(T).$$

Our objective is to define an operational selection rule that does not require knowledge of the true peak location L. As shown in Section 5, for large and moderate signal-to-noise ratios,

$\hat{l}(T)$ reaches a plateau whose value equals L in all replicates, with $T^* \leq L$ in all replicates at SNR = 1000 and in 99% of replicates at SNR = 100. In these regimes, the entry time alone is sufficient for the single-realization examples considered below. For SNR = 1, by contrast, a plateau of at least three horizons exists in only about one third of the replicates, and $T^* > L$ or $T^* = \infty$ in most of them. The path alone therefore does not indicate when its current value can be trusted. The exit times and the admissible set $\mathcal{K}(T)$ introduced below are therefore studied mainly for SNR = 1. The cases SNR = 1000 and SNR = 100 are also reported as high-SNR benchmark cases.

Structure of the candidate intervals. For a fixed T, the exit times satisfy

$$T_{\text{exit}}^{(k+1)}(T) < T_{\text{exit}}^{(k)}(T), \qquad k \geq 0, \tag{15}$$

whenever consecutive candidates exist, since the nontrivial exit times are the strict-decrease points of $t \mapsto \omega_l(t, T)$, ordered from largest to smallest. The candidate intervals are not necessarily nested: in simulations at SNR = 1, consecutive candidates are nested in only about 30% of the cases and are disjoint in about 16–32% of them (details in the supplementary material). The selection rule below uses only (15).

Operational selection of the interval. At a current horizon T, the sequential estimator $\widehat{l}(T)$ is available, whereas the true peak location L is unknown. We therefore select an interval by requiring it to contain the current point estimate. Define the set of admissible candidate indices by

$$\mathcal{K}(T) = \left\{ k : T_{\text{entry}}^{(k)}(T) \leq \widehat{l}(T) \leq T_{\text{exit}}^{(k)}(T) \right\}. \tag{16}$$

If $\mathcal{K}(T)$ is nonempty, we select the first admissible candidate:

$$\widehat{k}(T) = \min \mathcal{K}(T). \tag{17}$$

Since the exit times decrease with k, this is the admissible candidate with the largest exit time. The rule involves no tie-breaking. The resulting data-driven interval is

$$\widehat{\mathcal{I}}(T) = \left[T_{\text{entry}}^{(\widehat{k}(T))}(T), T_{\text{exit}}^{(\widehat{k}(T))}(T) \right]. \tag{18}$$

If $\mathcal{K}(T)$ is empty, no interval is selected at that horizon, and the procedure can continue as additional observations become available. We define the first operational selection horizon by

$$T_{\text{sel}} = \min \{ T : \mathcal{K}(T) \neq \varnothing \}. \tag{19}$$

Thus, T_{sel} is the earliest observation horizon at which the candidate construction produces an interval consistent with the current estimate. It depends only on whether $\mathcal{K}(T)$ is empty, not on the rule used to choose within $\mathcal{K}(T)$. Both the selected interval and this selection horizon are computed without using L. In contrast, a criterion based on whether the selected

interval contains the observed peak L cannot be used operationally and is therefore reserved for retrospective evaluation.

Choice of the selection rule. An alternative to (17) is to select the narrowest admissible candidate, $\arg\min_{k\in\mathcal{K}(T)} w_k(T)$. The two rules coincide whenever $\mathcal{K}(T) = \{0\}$, which is the case for SNR $= 1000$ and SNR $= 100$ after $T = 22$. For SNR $= 1$, we compared them on $M = 1000$ replicates at $T = 30$, among selected intervals (Table 2). The narrowest rule gives shorter intervals but lower coverage. To account for both, we use the interval score $w + \frac{2}{\alpha}\,\mathrm{dist}(L, \widehat{\mathcal{I}})$ (Winkler, 1972; Gneiting and Raftery, 2007), where w is the width and dist is the distance from L to the interval (0 if L is inside); lower is better. The first-admissible rule has the lower score for all four quantile levels and for $\alpha \in \{0.05, 0.10, 0.20\}$, at $T = 22$, 26 and 30, although the difference is negligible for $\tau = 0.95$ at $T = 30$ and $\alpha = 0.20$. This is a simulation result for the present model, not a general property of the rules.

Table 2: Selection rules at SNR $= 1$, $T = 30$, $M = 1000$, among selected intervals: coverage ($T_{\text{entry}} \le L \le T_{\text{exit}}$), mean width, and interval score ($\alpha = 0.10$). "First" is $\min \mathcal{K}(T)$, "Narrowest" is $\arg\min_{k\in\mathcal{K}(T)} w_k(T)$.

	Coverage		Width		Score	
τ	First	Narrowest	First	Narrowest	First	Narrowest
0.25	0.678	0.610	6.35	5.52	25.0	27.8
0.50	0.611	0.552	6.07	5.39	29.3	31.9
0.75	0.667	0.609	6.01	5.33	23.5	25.2
0.95	0.891	0.849	6.33	5.76	10.2	10.8

Numerical results. Table 3 reports the first operational selection horizon T_{sel} and the corresponding selected interval for each combination of τ and SNR. The interval width is $w_{\widehat{k}(T_{\text{sel}})}(T_{\text{sel}})$.

For SNR $= 100$ and SNR $= 1000$, the first selected interval occurs at $T_{\text{sel}} = 22$ for all four quantile levels. For SNR $= 1$, the first selected horizon varies with τ, taking the values 6, 10, 18, and 9 for $\tau = 0.25$, 0.50, 0.75, and 0.95, respectively. At T_{sel} the set $\mathcal{K}(T_{\text{sel}})$ is a singleton in all twelve cases, so the choice of selection rule has no effect on this table; it matters at later horizons (Table 2). The selected intervals are also substantially narrower in these low-SNR examples. However, an early and narrow interval need not be close to the true peak location: in the present realization, the low-SNR intervals selected at T_{sel} do not contain $L = 22$. This illustrates that the operational criterion guarantees consistency with $\widehat{l}(T)$, not containment of the unknown true peak. Accordingly, $\widehat{\mathcal{I}}(T)$ is a data-driven interval estimate for the peak location. Its construction alone does not establish a confidence level or guarantee

$$\mathbb{P}\Big\{L \in \widehat{\mathcal{I}}(T)\Big\} \ge 1 - \alpha.$$

To interpret this interval as a confidence interval with nominal coverage $1 - \alpha$, an appropriate

Table 3: First operationally selected candidate interval for each quantile level τ and signal-to-noise ratio. The selection horizon is T_{sel}, and the selected candidate index is $\widehat{k}(T_{\mathrm{sel}}) = \min \mathcal{K}(T_{\mathrm{sel}})$. In this realization, $\mathcal{K}(T_{\mathrm{sel}})$ contains a single candidate in every row, so the first-admissible and narrowest rules give the same interval. The last column reports the interval width $w_{\widehat{k}(T_{\mathrm{sel}})}(T_{\mathrm{sel}})$.

τ	SNR	T_{sel}	$\widehat{k}(T_{\mathrm{sel}})$	$\widehat{l}(T_{\mathrm{sel}})$	$T_{\mathrm{entry}}^{(\widehat{k}(T_{\mathrm{sel}}))}(T_{\mathrm{sel}})$	$T_{\mathrm{exit}}^{(\widehat{k}(T_{\mathrm{sel}}))}(T_{\mathrm{sel}})$	$w_{\widehat{k}(T_{\mathrm{sel}})}(T_{\mathrm{sel}})$
0.25	1000	22	0	22	11	22	11
0.25	100	22	0	22	18	22	4
0.25	1	6	0	4	4	6	2
0.50	1000	22	0	22	9	22	13
0.50	100	22	0	22	18	22	4
0.50	1	10	1	8	8	9	1
0.75	1000	22	0	22	12	22	10
0.75	100	22	0	22	16	22	6
0.75	1	18	0	17	17	18	1
0.95	1000	22	0	22	13	22	9
0.95	100	22	0	22	16	22	6
0.95	1	9	0	8	8	9	1

coverage result must be established theoretically or assessed through repeated independent simulation replicates. The single-realization illustration above is intended to demonstrate the operational construction and its behaviour across quantile levels and signal-to-noise regimes, rather than to establish statistical coverage.

Monte Carlo assessment of empirical coverage. We next assess the empirical coverage of the operational intervals using repeated independent simulations. For each combination of $\tau \in \{0.25, 0.50, 0.75, 0.95\}$ and SNR $\in \{1000, 100, 1\}$, we generate $M = 1000$ independent trajectories under the same peak model, with $A = 1$, $L = 22$, $S = 10$, and observation horizon $h = 30$. At each available horizon T, the operational selection rule defined in (16)–(18) is applied separately to each trajectory. The true peak location L is used only after selection to assess whether the selected interval contains it, with $L \in \widehat{\mathcal{I}}(T)$ meaning $T_{\mathrm{entry}}^{(\widehat{k}(T))}(T) \le L \le T_{\mathrm{exit}}^{(\widehat{k}(T))}(T)$. Let $\widehat{\mathcal{I}}_m(T)$ denote the selected interval for replicate m, when an interval is available. We define the interval-selection rate by

$$\widehat{q}_M(T) = \frac{1}{M} \sum_{m=1}^{M} \mathbf{1}\Big\{ \widehat{\mathcal{I}}_m(T) \neq \varnothing \Big\},$$

and the empirical coverage among selected intervals by

$$\widehat{p}_M^{\mathrm{sel}}(T) = \frac{\sum_{m=1}^{M} \mathbf{1}\Big\{L \in \widehat{\mathcal{I}}_m(T)\Big\}}{\sum_{m=1}^{M} \mathbf{1}\Big\{\widehat{\mathcal{I}}_m(T) \neq \varnothing\Big\}},$$

provided that at least one interval is selected. We also report the unconditional success proportion

$$\widehat{p}_M^{\mathrm{all}}(T) = \frac{1}{M} \sum_{m=1}^{M} \mathbf{1}\Big\{L \in \widehat{\mathcal{I}}_m(T)\Big\}.$$

Thus, $\widehat{p}_{\mathrm{all}}(T)$ accounts for both interval-selection failures and selected intervals that do not contain the true peak location. When no interval is selected in any replicate, the conditional coverage is undefined.

Figure 4 displays the empirical coverage among selected intervals as a function of T, while Figure 5 shows the corresponding interval-selection rates. The results reveal a marked dependence on the signal-to-noise ratio. For SNR $= 1000$, no interval is selected in the simulations for $T < 22$, whereas the selection rate and empirical coverage are both equal to one from $T = 22$ onward. For SNR $= 100$, selection is rare at several early horizons. At $T = 22$, the selection rate is between 0.996 and 1.000, and all selected intervals contain L. From $T = 23$ onward, all simulated trajectories yield a selected interval, and the empirical coverage is equal to one. For SNR $= 1000$ and SNR $= 100$ the results do not depend on the selection rule, since $\mathcal{K}(T) = \{0\}$ in (almost) all replicates after $T = 22$.

The low-SNR results are substantially different. At SNR $= 1$, intervals may be selected before $T = 22$, but none of the selected intervals at these earlier horizons contains the true peak location in the simulated sample. At $T = 22$, the interval-selection rate is between 0.634 and 0.711, and the empirical coverage among selected intervals ranges from approximately 0.289 to 0.418, depending on τ. Coverage subsequently increases with the observation horizon, although the rate of increase differs across quantile levels. At $T = 30$, for example, the empirical coverage is approximately 0.678, 0.611, 0.667, and 0.891 for $\tau = 0.25$, 0.50, 0.75, and 0.95, respectively. At this horizon, the interval-selection rate is at least 0.993 for all four quantile levels. Hence, the relatively low coverage at small SNR cannot be attributed solely to failures to select an interval.

Table 4 summarizes the selection rates, empirical coverage probabilities, and mean widths at $T = 30$. In the high- and medium-SNR settings, all selected intervals contain the true peak location in the simulated replicates. Under low SNR, the empirical coverage is lower, despite the high selection rates. The mean widths also differ across the SNR regimes, illustrating that interval width alone does not determine whether an interval contains the true peak.

These findings provide a simulation-based assessment of the operational procedure under the specified model and parameter settings. In particular, they show that an interval can be selected consistently without necessarily achieving high empirical coverage. The results should

therefore be interpreted as empirical coverage probabilities under the considered simulation design, rather than as a general theoretical coverage guarantee.

Table 4: Monte Carlo interval-selection rates, empirical coverage, and mean interval widths at $T = 30$, based on $M = 1000$ independent replicates and the first-admissible rule (17). Coverage means $T_{\text{entry}} \leq L \leq T_{\text{exit}}$. The conditional coverage is calculated among selected intervals, and the unconditional success proportion also accounts for replicates in which no interval is selected.

τ	SNR	$\widehat{q}_M(30)$	$\widehat{p}_M^{\text{sel}}(30)$	$\widehat{p}_M^{\text{all}}(30)$	Mean width
0.25	1000	1.000	1.000	1.000	18.750
0.25	100	1.000	1.000	1.000	12.708
0.25	1	0.998	0.678	0.677	6.350
0.50	1000	1.000	1.000	1.000	18.283
0.50	100	1.000	1.000	1.000	11.992
0.50	1	0.993	0.611	0.607	6.066
0.75	1000	1.000	1.000	1.000	18.799
0.75	100	1.000	1.000	1.000	12.750
0.75	1	0.998	0.667	0.666	6.011
0.95	1000	1.000	1.000	1.000	19.606
0.95	100	1.000	1.000	1.000	14.141
0.95	1	1.000	0.891	0.891	6.326

7 Real-World Application: COVID-19 Case Study

We now demonstrate the application of our quantile-based peak estimation framework to real-world data. We focus on the daily confirmed COVID-19 cases in China, France, and Germany. Figure 6 displays the daily case series for these three countries. The Chinese series is obtained from the Johns Hopkins University Center for Systems Science and Engineering (JHU CSSE) feed via the `COVID19` R package (Guidotti and Ardia, 2020; Guidotti, 2022), while the French and German series are obtained from the European Centre for Disease Prevention and Control (ECDC) open data portal (ECDC, 2020). For each country, we analyze the sequence $T \mapsto \hat{l}(T)$ for the quantile levels $\tau \in \{0.25, 0.5, 0.75, 0.95\}$, using quantile-loss minimization over the Gaussian-shaped parametric peak model. Following Algorithm 1, the candidate peak location ranges over the analysis horizon $h = \max(T)$, so that for each country the search set is $\{1, \ldots, h\}$. We then assess the temporal stabilization of the estimated peak-location path through

$$T_{\text{stab}} = \min\Big\{T : \max_{u \geq T} \hat{l}(u) - \min_{u \geq T} \hat{l}(u) \leq \varepsilon\Big\},$$

the earliest horizon from which all later estimates stay within a range ε (here $\varepsilon = 3$, 5 and 15 for China, France and Germany). This is computed for each country and quantile level. For each country we additionally apply the operational two-stage entry–exit construction of Section 6, which uses the oscillations of the estimated peak-location path to identify a candidate peak-region window $[T_{\text{entry}}^{(k)}, T_{\text{exit}}^{(k)}]$ and whose width is denoted $w_k = T_{\text{exit}}^{(k)} - T_{\text{entry}}^{(k)}$. Because the




Figure 4: Empirical coverage ($T_{\text{entry}} \leq L \leq T_{\text{exit}}$) of the selected operational intervals, using the first-admissible rule, as a function of the observation horizon T, based on $M = 1000$ independent simulated trajectories for each combination of τ and SNR. Coverage is calculated conditionally on interval selection. The horizontal axis gives the observation horizon, and each panel corresponds to one quantile level.




Figure 5: Empirical interval-selection rate as a function of the observation horizon T, based on $M = 1000$ independent simulated trajectories for each combination of τ and SNR. The selection rate is the proportion of replicates for which the operational rule selects an interval, and does not depend on the choice of rule within $\mathcal{K}(T)$. Shaded bands are $\pm$ one Monte Carlo standard error. Each panel corresponds to one quantile level.

observed peak location L is known for each of the three country-level trajectories used here, the resulting Stage-2 candidates can be evaluated retrospectively. As in the rest of the paper, a closed window $[T^{(k)}_{\text{entry}}, T^{(k)}_{\text{exit}}]$ contains L if $T^{(k)}_{\text{entry}} \leq L \leq T^{(k)}_{\text{exit}}$. The candidate of smallest width w_k satisfying this containment criterion is reported for each country and quantile level, with ties broken by the smaller k. Since the time index is discrete, a window may contain L at an endpoint. This minimum-width selection is used only for retrospective evaluation and is not part of the operational procedure when L is unknown. In addition, we apply the operational selection of Section 6.5, which uses only the current estimate $\hat{l}(T)$: at each horizon T we form the admissible set $\mathcal{K}(T)$ of candidates whose interval contains $\hat{l}(T)$ and select $\hat{k}(T) = \min \mathcal{K}(T)$. The observed peak L is used only to assess the resulting intervals. Throughout, L denotes the observed peak day, defined as the day of the maximum daily count (22, 78 and 61 for China, France and Germany), not a true parameter. Throughout this section, the Stage-2 candidates are described by the tuple $(k, \epsilon'_k, T^{(k)}_{\text{exit}}, T^*_k, \epsilon_k, T^{(k)}_{\text{entry}}, w_k)$ with the notation of Section 6: k is the candidate index, ϵ'_k the oscillation level at the candidate exit time, $T^{(k)}_{\text{exit}}$ the exit time, T^*_k the last admissible horizon in the Stage-2 set E_k, ϵ_k the oscillation level attained at T^*_k, $T^{(k)}_{\text{entry}}$ the entry time, and $w_k = T^{(k)}_{\text{exit}} - T^{(k)}_{\text{entry}}$ the width.




Figure 6: Daily confirmed COVID-19 cases in China, France, and Germany (left to right).

7.1 China

For China ($L = 22$), the selected windows (Table 5) are $[15, 26]$, $[15, 24]$, $[22, 25]$ and $[22, 27]$ for $\tau = 0.25, 0.5, 0.75, 0.95$, with widths between 3 and 11. A window containing L is therefore selected at every quantile level, but those for $\tau = 0.75$ and $\tau = 0.95$ begin at L and lie to its right, whereas the others contain L in their interior. For $\tau = 0.5$, the window $[15, 24]$ is close to $(14, 25)$ reported in Dermoune et al. (2026) under the least-absolute-deviation criterion for the same dataset. These results complement the stabilization analysis in Supplementary Table S1: the stabilization time describes the temporal stabilization of the estimated peak-location path, whereas the two-stage construction uses its oscillations to identify candidate windows for the peak region.

Table 5: Retrospective selection of the peak-region candidate for COVID-19 in China. The selected candidate is the narrowest one with $T_{\text{entry}}^{(k)} \le L \le T_{\text{exit}}^{(k)}$.

τ	k	ϵ_k'	$T_{\text{exit}}^{(k)}$	T_k^*	ϵ_k	$T_{\text{entry}}^{(k)}$	w_k	L
0.25	1	1	26	15	3	15	11	22
0.50	2	2	24	15	2	15	9	22
0.75	1	2	25	22	15	22	3	22
0.95	3	6	27	26	9	22	5	22

7.2 France

For France ($L = 78$), the selected windows (Table 6) are $[72, 79]$, $[71, 78]$, $[70, 78]$ and $[78, 85]$ for $\tau = 0.25, 0.5, 0.75, 0.95$, with widths between 7 and 8. A window containing L is therefore selected at every quantile level; only the window for $\tau = 0.25$ contains L in its interior, whereas the others contain L at an endpoint.

Table 6: Retrospective selection of the peak-region candidate for COVID-19 in France. The selected candidate is the narrowest one with $T_{\text{entry}}^{(k)} \le L \le T_{\text{exit}}^{(k)}$.

τ	k	ϵ_k'	$T_{\text{exit}}^{(k)}$	T_k^*	ϵ_k	$T_{\text{entry}}^{(k)}$	w_k	L
0.25	5	13	79	76	13	72	7	78
0.50	5	10	78	72	11	71	7	78
0.75	4	10	78	72	11	70	8	78
0.95	3	5	85	80	6	78	7	78

7.3 Germany

For Germany ($L = 61$), the selected windows (Table 7) are $[60, 62]$, $[59, 61]$, $[57, 61]$ and $[56, 63]$ for $\tau = 0.25, 0.5, 0.75, 0.95$, with widths between 2 and 7. A window containing L is therefore selected at every quantile level; for $\tau = 0.5$ and $\tau = 0.75$ the window ends at L, whereas the others contain L in their interior. These results complement the stabilization analysis in Supplementary Table S3.

Table 7: Retrospective selection of the peak-region candidate for COVID-19 in Germany. The selected candidate is the narrowest one with $T_{\text{entry}}^{(k)} \le L \le T_{\text{exit}}^{(k)}$.

τ	k	ϵ_k'	$T_{\text{exit}}^{(k)}$	T_k^*	ϵ_k	$T_{\text{entry}}^{(k)}$	w_k	L
0.25	7	15	62	60	15	60	2	61
0.50	5	10	61	59	11	59	2	61
0.75	5	7	61	58	11	57	4	61
0.95	6	9	63	56	18	56	7	61

7.4 Admissible candidates $\mathcal{K}(T)$ on the real data

For each country and each τ, we apply at every horizon T the current-horizon construction of Section 6.5 to the path $T \mapsto \hat{l}(T)$ (Tables S1–S3 in the supplement). At the terminal horizon $T = h$ this construction coincides with the two-stage construction used above. As in the rest of the paper, an interval $[a, b]$ contains L if $a \leq L \leq b$; the observed peak day L is 22, 78 and 61 for China, France and Germany.

Size of $\mathcal{K}(T)$. Table 8 lists $\mathcal{K}(h)$ and Table 9 gives the proportion of horizons at which $\mathcal{K}(T)$ is empty, a singleton, or has at least two elements. In contrast with the simulations at SNR = 1000 and 100, where $\mathcal{K}(T) = \{0\}$ after $T = 22$, the set $\mathcal{K}(T)$ here typically contains nontrivial candidates ($k \geq 1$): its largest size is 2 for China and 2 to 4 for France and Germany, and its mean size when nonempty is between 1.1 and 2.5. It is empty at many horizons (31–34% for China, 39–80% for France and Germany); the France and Germany grids start later ($T = 50$ and $T = 40$), so these proportions are not directly comparable across countries. The real paths therefore resemble the low-SNR simulations. The rules $\min \mathcal{K}(T)$ and $\arg\min_k w_k(T)$ select different candidates at 0–38% (China), 0–38% (France) and 6–50% (Germany) of the horizons at which an interval is selected.

Table 8: Admissible set $\mathcal{K}(h)$ at the terminal horizon.

	$\tau = 0.25$	$\tau = 0.50$	$\tau = 0.75$	$\tau = 0.95$
China ($h = 35$)	$\{1\}$	$\{2, 3\}$	$\{2, 3\}$	$\{4, 5\}$
France ($h = 89$)	$\{2, 3\}$	$\{3, 4\}$	$\{1\}$	$\{3, 4, 5, 6\}$
Germany ($h = 75$)	$\{3, 4\}$	$\{3\}$	$\{1, 2\}$	$\{2, 3\}$

Table 9: Proportion of horizons with $\mathcal{K}(T)$ empty, a singleton, or with at least two elements, and largest size of $\mathcal{K}(T)$.

Country	τ	empty	singleton	≥ 2	max $\lvert\mathcal{K}\rvert$	mean $\lvert\mathcal{K}\rvert$ if nonempty
China	0.25	0.31	0.63	0.06	2	1.09
	0.50	0.34	0.22	0.44	2	1.67
	0.75	0.31	0.06	0.63	2	1.91
	0.95	0.34	0.13	0.53	2	1.81
France	0.25	0.68	0.10	0.23	2	1.69
	0.50	0.80	0.03	0.18	2	1.88
	0.75	0.75	0.15	0.10	2	1.40
	0.95	0.80	0.08	0.13	4	2.50
Germany	0.25	0.50	0.14	0.36	2	1.72
	0.50	0.50	0.11	0.39	2	1.78
	0.75	0.50	0.11	0.39	3	1.83
	0.95	0.39	0.00	0.61	4	2.36

Position of the candidates relative to L. We examine $\mathcal{K}(T)$ at $T = L - 3$, $L + 3$ and $L + 10$ (the nearest available horizons: $19, 25, 32$ for China, $75, 81, 88$ for France and $58, 64, 71$

for Germany). For $T < L$, every candidate ends at $T^{(k)}_{\text{exit}}(T) \leq T < L$, so no interval can contain L; this is structural, and none does (0 of 13 candidates; $\mathcal{K}(75)$ is empty for France). For $T > L$, the number of candidates in $\mathcal{K}(T)$ whose interval contains L, pooled over the four values of τ, is 3 of 5 (China), 1 of 3 (France) and 6 of 9 (Germany) at $T = L + 3$, and 5 of 7, 6 of 9 and 6 of 9 at $T = L + 10$. Candidates that miss L are mostly narrow and lie on one side of L, near $\hat{l}(T)$. Table 10 shows, for each country and horizon, the narrowest candidate with $k \geq 1$, width at most 10, and $T^{(k)}_{\text{entry}} \leq L \leq T^{(k)}_{\text{exit}}$ (ties: smaller k, then smaller τ). This choice uses L and is illustrative only. Over all horizons $T > L$, the proportion of candidates covering L has no monotone pattern in k: for $k = 1, 2, 3, \geq 4$ it is $0.82, 0.88, 0.52, 0.42$ (China), $0.25, 0.63, 0.63, 1.00$ (France) and $0.32, 0.64, 0.61, 0.79$ (Germany). These proportions are descriptive, since consecutive horizons are dependent.

Table 10: Illustrative candidates: the narrowest $k \geq 1$ in $\mathcal{K}(T)$ with width at most 10 and $T^{(k)}_{\text{entry}} \leq L \leq T^{(k)}_{\text{exit}}$, at $T = L+3$ and $T = L+10$ ($L = 22, 78, 61$); ties are broken by the smaller k, then the smaller τ. The last column gives the number of candidates in $\mathcal{K}(T)$, pooled over τ, whose interval contains L. The selection uses L and is not operational.

Country	T	τ	k	$\widehat{l}(T)$	$[T^{(k)}_{\text{entry}}, T^{(k)}_{\text{exit}}]$	Width	Covering L
China	25	0.50	2	17	[15, 22]	7	3 of 5
China	32	0.50	2	17	[15, 24]	9	5 of 7
France	81	0.25	2	79	[72, 79]	7	1 of 3
France	88	0.95	3	79	[78, 85]	7	6 of 9
Germany	64	0.50	1	61	[61, 62]	1	6 of 9
Germany	71	0.50	3	64	[61, 65]	4	6 of 9

Operational rule. Table 11 gives the interval selected by $\hat{k}(T) = \min \mathcal{K}(T)$ at the same horizons. It contains L in 4 of 9 cases at $T = L+3$ and in 8 of 12 at $T = L+10$ ($\mathcal{K}(T)$ is empty in three further cases). Several of these intervals contain L only at an endpoint (for example $[13, 22]$, $[78, 85]$ and $[61, 62]$); since the time index is discrete and the intervals are closed, this is a genuine containment. The candidates of Table 10 are not what the operational rule returns in general: the rule often returns a narrow interval on one side of L. This agrees with the low-SNR simulations, where coverage of the selected interval was moderate and increased with T. These are single realizations per country, not coverage probabilities.

Entry and exit times as functions of T. Figure 7 shows, for each country and each τ, the curves $T \mapsto T^{(0)}_{\text{entry}}(T)$ for $T < L$ and $T \mapsto T^{(1)}_{\text{entry}}(T)$, $T \mapsto T^{(1)}_{\text{exit}}(T)$ for $T \geq L$, whether or not the corresponding candidates belong to $\mathcal{K}(T)$. These curves are descriptive: a flat stretch means that the window of the candidate no longer moves as T increases, and a jump means that $\hat{l}$ has changed. The curves before and after L refer to different candidates ($k = 0$ and $k = 1$).

Before the peak, $T^{(0)}_{\text{entry}}(T)$ stays close to the diagonal: it equals $T-1$, so that the window of the $k = 0$ candidate reduces to $[T-1, T]$, at 68%, 63% and 61% of the horizons (pooled over τ)

Table 11: Interval selected by $\hat{k}(T) = \min \mathcal{K}(T)$ at $T = L + 3$ and $T = L + 10$. "Covers L" means $T_{\text{entry}} \leq L \leq T_{\text{exit}}$; a dash means that $\mathcal{K}(T)$ is empty.

Country	T	τ	$\hat{k}$	$\widehat{l}(T)$	$[T_{\text{entry}}, T_{\text{exit}}]$	Width	Covers L
China	25	0.25	1	15	[14, 15]	1	No
	25	0.50	1	17	[16, 24]	8	Yes
	25	0.75	1	20	[13, 22]	9	Yes
	25	0.95	—	—	—	—	—
	32	0.25	1	16	[15, 26]	11	Yes
	32	0.50	2	17	[15, 24]	9	Yes
	32	0.75	2	18	[13, 22]	9	Yes
	32	0.95	4	20	[13, 26]	13	Yes
France	81	0.25	1	79	[79, 80]	1	No
	81	0.50	—	—	—	—	—
	81	0.75	0	81	[80, 81]	1	No
	81	0.95	—	—	—	—	—
	88	0.25	2	82	[81, 84]	3	No
	88	0.50	3	81	[73, 82]	9	Yes
	88	0.75	1	81	[81, 86]	5	No
	88	0.95	3	79	[78, 85]	7	Yes
Germany	64	0.25	1	62	[62, 63]	1	No
	64	0.50	1	61	[61, 62]	1	Yes
	64	0.75	1	60	[59, 63]	4	Yes
	64	0.95	2	58	[56, 59]	3	No
	71	0.25	3	65	[64, 67]	3	No
	71	0.50	2	64	[61, 66]	5	Yes
	71	0.75	1	65	[65, 67]	2	No
	71	0.95	2	61	[60, 65]	5	Yes

for China, France and Germany. The remaining flat stretches correspond to horizons at which $\hat{l}$ is unchanged and do not by themselves indicate stabilization: for France and $\tau = 0.75$, for instance, $\hat{l}(T) = 89 = h$ for $T = 52, \ldots, 56$, the upper bound of the search set.

After the peak, the countries differ. In China the windows of the $k = 1$ candidate become constant over long stretches: $[15, 26]$ from $T = 27$ ($\tau = 0.25$), $[25, 29]$ from $T = 30$ ($\tau = 0.5$), $[22, 25]$ from $T = 26$ ($\tau = 0.75$) and $[28, 30]$ from $T = 31$ ($\tau = 0.95$), up to $h = 35$. Only the first and the third contain L, the latter at its left endpoint. In France and Germany, where only 12 and 15 horizons follow the peak, the entry and exit times keep moving until the end of the sample: the $k = 1$ windows remain close behind the current horizon (the median of $T - T^{(1)}_{\text{exit}}(T)$ over $T \geq L$ is 1 for both countries, against 3 for China), constant stretches last at most 4 horizons in France and 6 in Germany, and none of the terminal $k = 1$ windows contains L.

This is consistent with the interpretation of $k = 1$ as the most recent plateau of $\hat{l}$, which follows T rather than the peak region. For France and Germany, the windows containing L in Tables 6 and 7 correspond to larger indices ($k = 3, \ldots, 7$), which reach back to the peak, whereas in China $k = 1$ already contains L for $\tau = 0.25$ and $\tau = 0.75$. The figure therefore shows when the entry and exit times stop moving, which happens clearly only for China within the sample, but not by itself whether the windows contain L.

8 Conclusion

This paper develops a quantile-based framework for predicting the location of a peak in a Gaussian-shaped time series observed sequentially and corrupted by additive noise. For fixed candidate values of the peak location and spread, the amplitude estimator is characterized as a weighted τ-quantile of a transformed version of the observations. The τ-aligned asymmetric Laplace model provides a probabilistic interpretation of the quantile objective, linking the estimation criterion to a likelihood-based formulation.

We also investigate the evolution of the estimated peak location as the observation horizon increases. The stabilization time describes when the sequential estimate becomes constant over the remaining horizon, subject to a minimum plateau-length condition. The simulations show that this criterion is informative when the signal-to-noise ratio is high or moderate, but becomes less reliable under substantial noise. To address situations in which the point estimate does not stabilize early enough, we introduce a two-stage construction of candidate entry–exit intervals based on the oscillations of the estimation trajectory. The operational selection rule uses the current point estimate and does not require knowledge of the true peak location.

The Monte Carlo experiments show that the performance of the selected intervals depends strongly on the signal-to-noise ratio. Under the high- and moderate-SNR settings considered, the selected intervals achieve full empirical coverage in the simulated replicates once the peak becomes identifiable. Under low SNR, intervals can be selected early without containing the true peak, although empirical coverage generally improves as more observations become avail-




Figure 7: Entry and exit times of the candidates $k = 0$ ($T < L$) and $k = 1$ ($T \geq L$) as functions of the horizon T, for China (top), France (middle) and Germany (bottom) and $\tau = 0.25, 0.5, 0.75, 0.95$ (left to right). Black: $T^{(0)}_{\text{entry}}(T)$ for $T < L$. Green: $T^{(1)}_{\text{entry}}(T)$ (solid) and $T^{(1)}_{\text{exit}}(T)$ (dashed) for $T \geq L$. The vertical and horizontal dashed grey lines mark $T = L$ and the value L (22, 78 and 61). The candidates are plotted whether or not they belong to $\mathcal{K}(T)$.

able. These results also illustrate that a high interval-selection rate does not necessarily imply high coverage. The application to daily COVID-19 cases in China, France, and Germany demonstrates how the proposed construction can be applied to real sequential estimation trajectories. The real-data results also show that the admissible candidate set may be empty at some horizons and that the operationally selected interval need not contain the retrospectively observed peak.

Several limitations suggest directions for further research. The analysis is based on a Gaussian-shaped peak and a specified τ-aligned noise model, so performance under more complex peak shapes, multiple peaks, temporal dependence, or model misspecification requires further investigation. Moreover, the entry–exit intervals are data-driven interval estimates rather than confidence intervals with guaranteed nominal coverage. Establishing theoretical coverage properties, and developing selection rules that balance interval width, coverage, and the possibility that no interval is selected, are important topics for future work.

References

G.D. Barmparis and G.P. Tsironis. Estimating the infection horizon of covid-19 in eight countries with a data-driven approach. *Chaos, Solitons and Fractals*, 135:109842, June 2020. ISSN 0960-0779. doi: 10.1016/j.chaos.2020.109842.

Azzouz Dermoune, Saïd Maanan, Daoud Ounaissi, and Yousri Slaoui. Forecasting the amplitude, the location and the width of the peak of a time series using the weighted median. *Journal of Applied Statistics*, pages 1–29, September 2026. ISSN 1360-0532. doi: 10.1080/02664763.2026.2734150.

European Centre for Disease Prevention and Control. Covid-19 case distribution by country. ECDC Open Data Portal, 2020. URL https://opendata.ecdc.europa.eu/covid19/casedistribution/csv. Accessed 2020-01-15 to 2020-05-31.

Tilmann Gneiting and Adrian E Raftery. Strictly proper scoring rules, prediction, and estimation. *Journal of the American Statistical Association*, 102(477):359–378, March 2007. ISSN 1537-274X. doi: 10.1198/016214506000001437.

Emanuele Guidotti. A worldwide epidemiological database for covid-19 at fine-grained spatial resolution. *Scientific Data*, 9(1), March 2022. ISSN 2052-4463. doi: 10.1038/s41597-022-01245-1.

Emanuele Guidotti and David Ardia. Covid-19 data hub. *Journal of Open Source Software*, 5 (51):2376, July 2020. ISSN 2475-9066. doi: 10.21105/joss.02376.

Vedran Novoselac. The component weighted median absolute deviations problem. *European Journal of Pure and Applied Mathematics*, 13(4):964–976, October 2020. ISSN 1307-5543. doi: 10.29020/nybg.ejpam.v13i4.3808.

Sen Pei and Jeffrey Shaman. Counteracting structural errors in ensemble forecast of influenza outbreaks. *Nature Communications*, 8(1), October 2017. ISSN 2041-1723. doi: 10.1038/s41467-017-01033-1.

Tucker Stewart, Bin Yu, Anderson Nascimento, and Juhua Hu. Enhancing peak network traffic prediction via time-series decomposition, 2023.

Robert L. Winkler. A decision-theoretic approach to interval estimation. *Journal of the American Statistical Association*, 67(337):187–191, March 1972. ISSN 1537-274X. doi: 10.1080/01621459.1972.10481224.

Bin Yu, Giovanna Graciani, Anderson Nascimento, and Juhua Hu. Cost-adaptive neural networks for peak volume prediction with emm filtering. In *2019 IEEE International Conference on Big Data (Big Data)*, pages 4208–4213. IEEE, December 2019. doi: 10.1109/bigdata47090.2019.9006188.

Data Availability Statement The daily COVID-19 case counts analyzed in Section 7 are derived from two publicly available sources. The Chinese series is obtained from the Johns Hopkins University Center for Systems Science and Engineering (JHU CSSE) feed through the `COVID19` R package, covering the period 22 January to 22 March 2020. The French and German series are obtained from the European Centre for Disease Prevention and Control (ECDC) open data portal, covering the periods 15 January to 31 May 2020 and 28 January to 5 July 2020, respectively. All data used in this study are publicly available at the sources cited above and in the bibliography.

Conflict of Interest Statement The authors declare no conflict of interest.

Funding Statement This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.