

Physics 2211: Matter and Interactions

Class III (interactions and momentum)

Today's objective:

Momentum principle and iterations



$$\frac{d\vec{p}}{dt} = \vec{F}_{net}$$

$$d\vec{p} = \vec{F}_{net} dt$$

Physics 2211: *Matter and Interactions*

Class III (interactions and momentum)

Today's objective:

Momentum principle and iterations

Howey Room C203

Office Hours: Weds 7-9 am

E-mail: Flavio.Fenton@physics.gatech.edu

PLUS Sessions: Bagas Widiyanto

Tuesdays and Thursdays from
7:00-8:00 PM in CULC 123.

Physics 221: Matter and Interactions

- **The Momentum Principle:** *The change in momentum of a system is equal to the net interaction (force) acting on the system by the surroundings multiplied by the duration of the interaction*
 - System: *One or more objects of interest*
 - Surroundings: *Everything else in the Universe*
 - Net Force: *The sum of all the interaction (forces)*
- **Newton's Second Law**
 - True for all forces
 - Allows us to quantify the interactions of a system and its surroundings.

$$\frac{d\vec{p}}{dt} = \vec{F}_{net}$$

$$\Delta \vec{p} = \vec{F}_{net} \Delta t$$

■ Applying the momentum principle

1. Chose your system, everything else belongs in the surroundings.
2. Draw a force diagram
 - All forces must be due to interactions with surroundings
3. Chose your time interval
 - Given or estimated
4. Substitute known values into the equation and solve for the unknowns
5. Check units and reasonableness of answer


Physics 221: *Matter and Interactions*

- The momentum principle

$$\Delta \vec{p} = \vec{F}_{net} \Delta t$$

$$\vec{p}_f = \vec{p}_i + \vec{F}_{net} \Delta t$$

Physics of kicking a ball:



What is the system?

What are the surroundings?

What forces are acting on the system?

Important!!: And when?

Yes: $F_{gravitation}$ is all the time




No: $F_{Kicking}$ is NOT all the time!

What if we consider air friction?

Physics 2211: Matter and Interactions

$$\Delta \vec{p} = \vec{F}_{net} \Delta t$$




Note: This equation tells us that:
The direction of change in Momentum,
has to be equal to the Net Force



Physics 2211: Matter and Interactions



$$\Delta \vec{p} = \vec{F}_{net} \Delta t$$

Note: This equation tells us that:
The direction of change in Momentum,
has to be equal to the Net Force



Physics 2211: Matter and Interactions


- Changes in momentum indicate an interaction
- Clicker:** A child rides on a merry-go-round, traveling from location A to location C at a constant speed. What is the direction of the child's change in momentum, between A and C?



9 zero magnitude


Physics 221: Matter and Interactions

- Changes in momentum indicate an interaction
- Clicker:** A child rides on a merry-go-round, traveling from location A to location C at a constant speed. What is the direction of the child's change in momentum, between A and C?



Assume $\vec{V}_A = \langle 0, 5, 0 \rangle$ and $\vec{V}_C = \langle 0, -5, 0 \rangle$, $m=1$
 What is $|\Delta \vec{P}|$ and $\Delta |\vec{P}|$?



$|\langle 0, -10, 0 \rangle| = 10$ and 0 (so not the same!)



$$|\Delta \vec{p}| \neq \Delta |\vec{p}|$$



Physics 2211: Matter and Interactions

- Changes in momentum indicate an interaction
- Clicker:** A child rides on a merry-go-round, traveling from location A to location C at a constant speed. What is the direction of the child's change in momentum, between A and C?



Physics 2211: Matter and Interactions

- Changes in momentum indicate an interaction
- Clicker:** A child rides on a merry-go-round, traveling from location A to location C at a constant speed. What is the direction of the child's change in momentum, between A and C?



9 zero magnitude

Another Example, so let's remember:


- **Applying the momentum principle**

1. Chose your system, everything else belongs in the surroundings.
2. Draw a force diagram
 - All forces must be due to interactions with surroundings
3. Chose your time interval
 - Given or estimated
4. Substitute known values into the equation and solve for the unknowns
5. Check units and reasonableness of answer

Physics 221: Matter and Interactions

On the bucket

- **SRS:** A bucket of chicken is being lifted up a well at a constant speed by a rope. All frictional effects are negligible. In this situation, forces on the bucket of chicken are such that:
 - (1) The upward force by the rope is greater than the downward force of gravity
 - (2) The upward force by the rope is equal to the downward force of gravity.
 - (3) The upward force by the rope is smaller than the downward force of gravity.
 - (4) The upward force by the rope is greater than the sum of the downward force of gravity and a downward force due to the air.
 - (5) None of the above. (The bucket goes up because the rope is being shortened, not because an upward force is exerted on the bucket by the rope).



Physics 221: Matter and Interactions

On the bucket

- **SRS:** A bucket of chicken is being lifted up a well at a constant speed by a rope. All frictional effects are negligible. In this situation, forces on the bucket of chicken are such that:

- (1) The upward force by the rope is greater than the downward force of gravity
- (2) The upward force by the rope is equal to the downward force of gravity.
- (3) The upward force by the rope is smaller than the downward force of gravity.
- (4) The upward force by the rope is greater than the sum of the downward force of gravity and a downward force due to the air.
- (5) None of the above. (The bucket goes up because the rope is being shortened, not because an upward force is exerted on the bucket by the rope).




If the velocity is increasing, what would be the answer?

Physics 221: *Matter and Interactions*


Having fun with $\frac{d\vec{p}}{dt} = 0$

- Explosions?
- Fire?
- Boiling water?



Having fun with $\frac{d\vec{p}}{dt} = 0$

- Fire and Explosions?
- Liquids?
- Motion?



- **SRS:** Which cart(s) experience a net force to the left?

Cart A moves to the left at a constant speed.

Cart B moves to the left, gradually speeding up.

Cart C moves to the left, gradually slowing down.

- (1) A only
- (2) B only
- (3) C only
- (4) A and B
- (5) B and C
- (6) A and C
- (7) A, B, and C

Physics 221: Matter and Interactions

- **SRS:** Which cart(s) experience a net force to the left?

Cart A moves to the left at a constant speed.

Cart B moves to the left, gradually speeding up.

Cart C moves to the left, gradually slowing down.

(1) A only



(2) B only

(3) C only

(4) A and B

(5) B and C

(6) A and C

(7) A, B, and C

Physics 221: *Matter and Interactions*

Q2.2.a

An object is moving in the $+x$ direction.

Which of the following statements about the net force acting on the object could be true?

- A. The net force is in the $+x$ direction
- B. The net force is in the $-x$ direction
- C. The net force is zero

- 1) A only
- 2) B only
- 3) C only
- 4) A and B
- 5) B and C
- 6) A and C
- 7) A, B, and C

Physics 221: *Matter and Interactions*

Q2.2.a

An object is moving in the $+x$ direction.

Which of the following statements about the net force acting on the object could be true?

- A. The net force is in the $+x$ direction
- B. The net force is in the $-x$ direction
- C. The net force is zero

- 1) A only
- 2) B only
- 3) C only
- 4) A and B
- 5) B and C
- 6) A and C
- 7) A, B, and C



Physics 221: *Matter and Interactions*

Q2.3.b

A hockey puck is sliding along the ice with nearly constant momentum $\langle 10, 0, 5 \rangle$ kg m/s when it is suddenly struck by a hockey stick with a force $\langle 0, 0, 2000 \rangle$ N that lasts for only 3 milliseconds (3×10^{-3} s).

What is the new (vector) momentum of the puck?

1) $\langle 10, 0, 11 \rangle$ kg m/s

2) $\langle 0, 0, 6 \rangle$ kg m/s

3) 14.86 kg m/s

4) $\langle 16, 0, 11 \rangle$ kg m/s

5) $\langle 0, 0, 30 \rangle$ kg m/s

Physics 221: *Matter and Interactions*

Q2.3.b

A hockey puck is sliding along the ice with nearly constant momentum $\langle 10, 0, 5 \rangle$ kg m/s when it is suddenly struck by a hockey stick with a force $\langle 0, 0, 2000 \rangle$ N that lasts for only 3 milliseconds (3×10^{-3} s).

What is the new (vector) momentum of the puck?

1) $\langle 10, 0, 11 \rangle$ kg m/s

2) $\langle 0, 0, 6 \rangle$ kg m/s

3) 14.86 kg m/s

4) $\langle 16, 0, 11 \rangle$ kg m/s

5) $\langle 0, 0, 30 \rangle$ kg m/s



What if I ask you next, what would be the momentum 1 second later?
(assume no friction by the ice)

Physics 221: *Matter and Interactions*

Important implications of the Momentum Principle

- If the net force is zero, there is no change in momentum
- Force in one coordinate can not affect the momentum in a different coordinate.

Example: F_x only affects P_x can not affect P_y or P_z

Example2: Object with P_x and force F_y what happens to P_x ?

Nothing

$$p_{fx} = p_{ix} + F_{\text{net},x} \Delta t$$

$$p_{fy} = p_{iy} + F_{\text{net},y} \Delta t$$


$$p_{fz} = p_{iz} + F_{\text{net},z} \Delta t$$

Duck example

Some problems with the HW


- Check for units and make sure they are correct!
- Check for directions (Direction of vectors, magnitudes Vs vectors, which v_{avg} to use , etc...)
- Iterative cases.

Iterative solution



Physics 221: Matter and Interactions

How to solve the projectile motion in a computer?



Iterative solution


- Calculate the net force $\vec{F}_{\text{net,now}}$ acting on the system
 - Update momentum: $\vec{p}_{\text{future}} = \vec{p}_{\text{now}} + \vec{F}_{\text{net,now}} \Delta t$
 - Update position: $\vec{r}_{\text{future}} = \vec{r}_{\text{now}} + \vec{v}_{\text{avg}} \Delta t$
- repeat!

Use:

$$\vec{v}_{\text{avg}} = \vec{p}_{\text{f}} / m$$

Physics 221: Matter and Interactions

How to solve the projectile motion in a computer?



Iterative solution

- Calculate the net force $\vec{F}_{\text{net,now}}$ acting on the system
 - Update momentum: $\vec{p}_{\text{future}} = \vec{p}_{\text{now}} + \vec{F}_{\text{net,now}} \Delta t$
 - Update position: $\vec{r}_{\text{future}} = \vec{r}_{\text{now}} + \vec{v}_{\text{avg}} \Delta t$
- repeat!

Use:

$$\vec{v}_{\text{avg}} = \vec{p}_f / m$$

Physics 221: Matter and Interactions

- **SRS:** *The x-component of momentum of an object is found to increase with time:*

$$t = 0 \text{ s} \quad p_x = 30 \text{ kg m/s}$$

$$t = 1 \text{ s} \quad p_x = 40 \text{ kg m/s}$$

$$t = 2 \text{ s} \quad p_x = 50 \text{ kg m/s}$$

$$t = 3 \text{ s} \quad p_x = 60 \text{ kg m/s}$$

$$\Delta \vec{p} = \vec{F}_{net} \Delta t$$

(1) $F_{net,x} = 0$

(2) $F_{net,x}$ is constant

(3) $F_{net,x}$ is increasing with time

(4) Not enough information

Physics 221: Matter and Interactions

- **SRS:** *The x-component of momentum of an object is found to increase with time:*

$$t = 0 \text{ s} \quad p_x = 30 \text{ kg m/s}$$

$$t = 1 \text{ s} \quad p_x = 40 \text{ kg m/s}$$

$$t = 2 \text{ s} \quad p_x = 50 \text{ kg m/s}$$

$$t = 3 \text{ s} \quad p_x = 60 \text{ kg m/s}$$

$$\Delta \vec{p} = \vec{F}_{net} \Delta t$$

(1) $F_{net,x} = 0$




(2) $F_{net,x}$ is constant

(3) $F_{net,x}$ is increasing with time

(4) Not enough information


Physics 221: Matter and Interactions

- **SRS:** The initial momentum of the kicked ball was $\langle 1.5, 3.5, 0 \rangle \text{ kg m/s}$. The final momentum of the ball was $\langle 1.5, 1.05, 0 \rangle \text{ kg m/s}$. Which graph correctly shows P_y for the ball during this 0.5 s?



Physics 221: Matter and Interactions

- **SRS:** The initial momentum of the kicked ball was $\langle 1.5, 3.5, 0 \rangle \text{ kg m/s}$. The final momentum of the ball was $\langle 1.5, 1.05, 0 \rangle \text{ kg m/s}$. Which graph correctly shows P_y for the ball during this 0.5 s?



What can we say about the force for each case?

Physics 221: *Matter and Interactions*

The Kinematic Equations

$$v = v_0 + at$$

$$\Delta x = v_0 t + \frac{1}{2} at^2$$

$$v^2 = v_0^2 + 2a\Delta x$$

$$\Delta x = \bar{v}t = \frac{1}{2}(v + v_0)t$$

$$\Delta x = vt - \frac{1}{2} at^2$$

You all know these equations, but where did they come from?

Yes! From

$$\frac{d\vec{p}}{dt} = \vec{F}_{net}$$

when $\vec{F}_{Net} = \text{Constant}$

Physics 221: Matter and Interactions

How to solve the projectile motion in a computer?

8. + 0/3 points

MI4 2.1.013. [3220153]

A Ping-Pong ball is acted upon by the Earth, air resistance, and a strong wind. Here are the positions of the ball at several times.

Early time interval:

At $t = 13.40$ s, the position was $\langle 4.65, 3.06, -8.45 \rangle$ m.

At $t = 13.46$ s, the position was $\langle 4.72, 2.87, -9.35 \rangle$ m.

Late time interval:

At $t = 16.40$ s, the position was $\langle 10.77, 11.14, -54.35 \rangle$ m.

At $t = 16.46$ s, the position was $\langle 10.95, 11.65, -55.25 \rangle$ m.

(a) In the early time interval, from $t = 13.40$ s to $t = 13.46$ s, what was the average momentum of the ball? The mass of the Ping-Pong ball is 2.7 grams (2.7×10^{-3} kg). Express your result as a vector.

$\vec{p} =$ $\times \text{ kg} \cdot \text{ m/s}$

(b) In the late time interval, from $t = 16.40$ s to $t = 16.46$ s, what was the average momentum of the ball? Express your result as a vector.

$\vec{p} =$ $\times \text{ kg} \cdot \text{ m/s}$

(c) In the time interval from $t = 13.40$ s (the start of the early time interval) to $t = 16.40$ s (the start of the late time interval), what was the average net force acting on the ball? Express your result as a vector.

$\vec{F}_{\text{net}} =$ $\times \text{ N}$

Physics 221: Matter and Interactions

How to solve the projectile motion in a computer?

8. + 0/3 points

A Ping-Pong ball is acted upon by the Earth, air resistance, and a net force. The position of the ball is recorded at various times.

Early time interval:

At $t = 13.40$ s, the position was $\langle 4.65, 3.06, -8.45 \rangle$ m

At $t = 13.46$ s, the position was $\langle 4.72, 2.87, -9.35 \rangle$ m

Late time interval:

At $t = 16.40$ s, the position was $\langle 10.77, 11.14, -54.35 \rangle$ m

At $t = 16.46$ s, the position was $\langle 10.95, 11.65, -55.25 \rangle$ m

(a) In the early time interval, from $t = 13.40$ s to $t = 13.46$ s, the mass of the Ping-Pong ball is 2.7 grams (2.7×10^{-3} kg).

$\vec{p} =$ \times kg \cdot m/s

(b) In the late time interval, from $t = 16.40$ s to $t = 16.46$ s, your result as a vector.

$\vec{p} =$ \times kg \cdot m/s

(c) In the time interval from $t = 13.40$ s (the start of the early time interval), what was the average net force acting on the ball?

$\vec{F}_{\text{net}} =$ \times N

momentum (initial) < 0.00315, -0.00855, -0.0405 >

momentum (final) < 0.0081, 0.02295, -0.0405 >

$F = dp/dt =$ < 0.00165, 0.0105, 4.25585e-16 >

Run this program

Share or export this program

```
1 GlowScript 1.1 VPython
2 #HW2 Prob 8
3 scene.background=color.white
4 ri=vector(4.65,3.06,-8.45)
5 ti=13.40
6 rf=vector(4.72,2.87,-9.35)
7 tf=13.46
8 ball=sphere(pos=ri,rad=0.2,color=color.yellow)
9 ball.m=2.7e-3
10 deltar=rf-ri
11 deltat=tf-ti
12 v=deltar/deltat
13 p=ball.m*v
14 scene.pause('Click to proceed')
15 print("momentum (initial)",p)
16 pi=p
17 ri=vector(10.77,11.14,-54.35)
18 ti=16.40
19 rf=vector(10.95,11.65,-55.25)
20 tf=16.46
21 deltar=rf-ri
22 deltat=tf-ti
23 v=deltar/deltat
24 p=ball.m*v
25 scene.pause('Click to proceed')
26 print("momentum (final)",p)
27 deltap=p-pi
28 deltat=16.40-13.40
29 dp_dt=deltap/deltat
30 scene.pause('Click to proceed')
31 print("F=dp/dt=",dp_dt)
```

Physics 221: Matter and Interactions

How to solve the projectile motion in a computer?

16. 0/6 points

MI4 2.4.021. [3219763]

Use the approximation that $\vec{v} \approx \vec{p}/m$ for each time step.

You throw a metal block of mass **0.20** kg into the air, and it leaves your hand at time $t = 0$ at location $\langle 0, 2, 0 \rangle$ m with velocity $\langle \mathbf{2.5}, \mathbf{3.5}, 0 \rangle$ m/s. At this low velocity air resistance is negligible. Using the iterative method shown in Section 2.4 with a time step of 0.05 s, calculate step by step the position and velocity of the block at $t = 0.05$ s, $t = 0.10$ s, and $t = 0.15$ s. (Express your answers in vector form.)

$\vec{x}(t = 0.05 \text{ s}) =$ $\times \text{ m}$

$\vec{v}(t = 0.05 \text{ s}) =$ $\times \text{ m/s}$

$\vec{x}(t = 0.10 \text{ s}) =$ $\times \text{ m}$

$\vec{v}(t = 0.10 \text{ s}) =$ $\times \text{ m/s}$

$\vec{x}(t = 0.15 \text{ s}) =$ $\times \text{ m}$

$\vec{v}(t = 0.15 \text{ s}) =$ $\times \text{ m/s}$

Iterative solution

- Calculate the net force $\vec{F}_{\text{net,now}}$ acting on the system
 - Update momentum: $\vec{p}_{\text{future}} = \vec{p}_{\text{now}} + \vec{F}_{\text{net,now}} \Delta t$
 - Update position: $\vec{r}_{\text{future}} = \vec{r}_{\text{now}} + \vec{v}_{\text{avg}} \Delta t$
- repeat!

$$\vec{v}_{\text{avg}} \approx \vec{v}_f$$

$$\vec{v}_{\text{avg}} \approx \frac{\vec{v}_i + \vec{v}_f}{2}$$

Physics 221: Matter and Interactions

How to solve the projectile motion in a computer?

16. 0/6 points

MI4 2.4.021. [3219763]

Use the approximation that $\vec{v} \approx \vec{p}/m$ for each time step.

You throw a metal block of mass **0.20** kg into the air, and it leaves your hand at time $t = 0$ at location $\langle 0, 2, 0 \rangle$ m with velocity $\langle 2.5, 3.5, 0 \rangle$ m/s. At this low velocity air resistance is negligible. Using the iterative method shown in Section 2.4 with a time step of 0.05 s, calculate step by step the position and velocity of the block at $t = 0.05$ s, $t = 0.10$ s, and $t = 0.15$ s. (Express your answers in vector form.)

$\vec{x}(t = 0.05 \text{ s}) =$ ✗ m

$\vec{v}(t = 0.05 \text{ s}) =$ ✗ m/s

$\vec{x}(t = 0.10 \text{ s}) =$ ✗ m

$\vec{v}(t = 0.10 \text{ s}) =$ ✗ m/s

$\vec{x}(t = 0.15 \text{ s}) =$ ✗ m

$\vec{v}(t = 0.15 \text{ s}) =$ ✗ m/s

Run this program

Share or export this program

```
1 GlowScript 1.1 VPython
2 #HW2 Prob 16
3 scene.background=color.yellow
4
5 t=0
6 ball=sphere(pos=vector(0,2,0), rad=0.1, color=color.blue)
7 ball.vel=vector(2.5,3.5,0)
8 ball.m=0.2
9 ball.p=ball.m*ball.vel
10
11
12 deltat=0.05
13 g=9.8
14
15 while t<=0.15:
16     scene.pause('Click to proceed')
17     fnet=vector(0,-ball.m*g,0)
18     ball.p=ball.p + fnet*deltat
19     vavg=ball.p/ball.m
20     print("time",t+deltat,"vavg",vavg)
21     ball.pos=ball.pos + vavg*deltat
22     print("time",t+deltat,"position",ball.pos)
23     t=t+deltat
```

Physics 221: Matter and Interactions

How to solve the projectile motion in a computer?

18. Question Details

MI4 2.4.023. [3219765]

A soccer ball of mass 0.40 kg is rolling with velocity $\langle 0, 0, 3.0 \rangle \text{ m/s}$, when you kick it. Your kick delivers an impulse of magnitude $1.2 \text{ N} \cdot \text{s}$ in the $-x$ direction. The net force on the rolling ball, due to the air and the grass, is 0.20 N in the direction opposite to the direction of the ball's momentum. Using a time step of 0.5 s , find the position of the ball at a time 1.5 s after you kick it, assuming that the ball is at the origin at the moment it is kicked. Use the approximation

$\vec{v}_{\text{avg}} \approx \vec{p}_f / m$. (Express your answer in vector form.)

$\vec{r}_f =$ $\times < -3.97, 0, 3.97 > \text{ m}$

Additional Materials

 [Section 2.4](#)

```
1 GlowScript 1.1 VPython
2 #HW2 Prob 18
3 scene.background=color.yellow
4
5 t=0
6 ball=sphere(pos=vector(0,0,0), rad=0.1, color=color.blue)
7 ball.vel=vector(0,0,2.0)
8 ball.m=0.5
9 ball.p=ball.m*ball.vel
10 impulse=vector(-1.8,0,0)
11 ball.p=ball.p+impulse
12 print("momentum after cick",ball.p)
13 ballpunitary=ball.p/mag(ball.p)
14 print("momentum after cick",ballpunitary)
15 Ffrictionmag=0.26
16 Ffriction=-ballpunitary*Ffrictionmag
17 print("momentum after cick",Ffriction)
18
19 deltat=0.5
20
21 while t<1.5:
22     scene.pause('Click to proceed')
23     ball.p=ball.p+deltat*Ffriction
24     vavg=ball.p/ball.m
25     ball.pos=ball.pos+deltat*vavg
26     print("at time",t+deltat,"position is:",ball.pos)
27     t=t+deltat
28
```