

Discrete Mathematics and Algorithms (CSE 611)

Assignment Set 3

Total Marks: 150

Deadline: October 04, 2013 (Friday), 4:00 pm after class

Each question carries 10 marks.

1. If $F(x)$ denotes the distribution function of a random variable X , then show that

(i) $P(a < X < b) = F(b - 0) - F(a)$, and

(ii) $P(a \leq X \leq b) = F(b) - F(a - 0)$.

2. Show that a function $f(x)$ given by

$$f(x) = \begin{cases} x & 0 < x < 1 \\ k - x & 1 < x < 2 \\ 0 & \text{elsewhere} \end{cases}$$

is a probability density function for a suitable value of the constant k . Calculate the probability that the random variable lies between $\frac{3}{2}$ and $\frac{7}{6}$.

3. The mean of a continuous random variable X is defined by $E(X) = \int_{-\infty}^{\infty} x f(x) dx$, where $f(x)$ is the probability density function of X . If $\mu = E(X)$, we define the variation of X as $Var(X) = E[(X - \mu)^2] = E(X^2) - \mu^2$. Suppose X follows a gamma distribution with parameter $l(> 0)$. Compute $E(X)$ and $Var(X)$.

4. The probability density function of a random variable X is $A \operatorname{sech}(x)$, where $\operatorname{sech}(x)$ is the hyperbolic sec function and it is given by $\operatorname{sech}(x) = \frac{1}{\cosh(x)}$ and $\cosh(x) = \frac{e^x + e^{-x}}{2}$. Find the value of the constant A . Show that $P(X < 1) = \frac{2}{\pi} \tan^{-1}(e)$ and $P(|X| \geq 1) = 2 - \frac{4}{\pi} \tan^{-1}(e)$.

5. Prove that

(i) $P[(A \cap B)|A] = P(B|A)$.

(ii) $P[(B \cup C)|A] = P(B|A) + P(C|A) - P[(B \cap C)|A]$.

6. Interest for money deposited in a savings account is paid at a rate of 0.5 percent per month, with interest compounded monthly. Suppose \$50 is deposited into a savings account each month for a period of five years. What is the total amount in the account four years after the first deposit? Twenty years after the first deposit?

7. Let

$$\begin{aligned} a_r &= \begin{cases} 1, r = 0 \\ 3, r = 1 \\ 2, r = 2 \\ 0, r \geq 3 \end{cases} \\ c_r &= 5^r \text{ for all } r \end{aligned}$$

Given that $c = a * b$, that is c is the convolution of numeric functions a and b . Show that

$$b_r = \frac{25}{42}5^r - \frac{1}{6}(-1)^r + \frac{4}{7}(-1)^r 2^r, r \geq 0.$$

8. Using the generating function, show that solution of the following recurrence relation

$$a_k - 7a_{k-1} + 10a_{k-2} = 3^k$$

with initial conditions $a_0 = 0$ and $a_1 = 1$, is

$$a_k = \frac{8}{3}2^k - \frac{9}{2}3^k + \frac{11}{6}5^r$$

9. Consider an air traffic-control system in which the desired altitude of an aircraft, a_r , is computed by a computer every second and is compared with the actual altitude of the aircraft, b_{r-1} , determined by a tracking radar 1 second earlier. Depending on whether a_r is larger or smaller than b_{r-1} , the altitude of the aircraft will be changed accordingly. Specifically, the change in altitude at the r -th second, $b_r - b_{r-1}$, is proportional to the difference $a_r - b_{r-1}$. That is,

$$b_r - b_{r-1} = K(a_r - b_{r-1})$$

where K is a proportional constant.

(a) Determine b_r , given that $a_r = 1000(\frac{3}{2})^r$, $K = 3$, and $b_0 = 0$.

(b) Determine b_r , given that

$$a_r = \begin{cases} 1000(\frac{3}{2})^r, 0 \leq r \leq 9 \\ 1000(\frac{3}{2})^{10}, r \geq 10 \end{cases}$$

$K = 3$, and $b_0 = 0$.

10. Consider the multiplication of bacteria in a controlled environment. Let a_r denote the number of bacteria there are on the r -th day. We define the rate of growth on the r -th day to be $a_r - 5 \cdot a_{r-1}$. It is known that the rate of growth at r -th day is three times the growth of the $(r - 1)$ -th day. Determine a_r , given that $a_0 = 1$.
11. Let a and b be constants, and m be a positive integer. Then prove that the solution of the recurrence relation

$$T(n) = \begin{cases} b, & n \leq 2 \\ mT(n/2) + an^2, & n > 2 \end{cases}$$

is $T(n) = O(n^{\log_2 m})$.

12. If k is a nonnegative constant, then prove that the recurrence

$$T(n) = \begin{cases} k, & n = 1 \\ 3T(n/2) + kn, & n > 1 \end{cases}$$

has the following solution (for n is a power of 2):

$$T(n) = 3kn^{\log_2 3} - 2kn.$$

13. The sets A and B have m and n elements (respectively) from a linear order. These sets are not necessarily sorted. Also assume that $m \leq n$. Show how to compute $A \cup B$ and $A \cap B$ in $O(n \log m)$ time.
14. Consider the Strassen's matrix multiplication using the divide-and-conquer algorithm, when the order n of the matrices is not power of 2. In this case, explain the Strassen's matrix multiplication for $C = AB$, where $A = \begin{pmatrix} 2 & -1 & 6 \\ 0 & 1 & 8 \\ 1 & 2 & -3 \end{pmatrix}$, and $B = \begin{pmatrix} 0 & -1 & 5 \\ 1 & 1 & 3 \\ -1 & 1 & 0 \end{pmatrix}$.
15. Applying the dynamic programming approach, determine an LCS of the sequences $\langle 1, 0, 0, 1, 0, 1, 0, 1 \rangle$ and $\langle 0, 1, 0, 1, 1, 0, 1, 1, 0 \rangle$. Show, in details, the c and b tables.

Submission Instructions

Copying in assignments leads to award ZERO marks in assignment marks. Also, the source from which you have copied, that source student will be treated under the same rule.

Please submit the assignment in hard copy stating the following at the top:

Discrete Mathematics and Algorithms (CSE 611)

Assignment Set 3

submitted by

Name: XYZ, Roll No: abc