

① GIVEN

* Rigid tank


$$\gg V_{\text{tank}} = 15 \text{ m}^3$$

 \gg separated into 2 sections by membrane each filled w/ $\text{N}_2(\text{g})$
 \gg section A has $\rho_A = 1.6 \text{ kg/m}^3$, $\text{Vol} = V_A$, P_A
 \gg section B $m_B = 6 \text{ kg}$, $\text{vol} = V_B$, P_B
 \gg after membrane punctured $P_{\text{tank}} = 1.8 \text{ kg/m}^3$
 $\gg M \equiv \text{molar mass of Nitrogen } 14.006$
FINDInitial density (kg/m^3) of $\text{N}_2(\text{g})$ @ section BEQUATION

$$\rho = \frac{m}{V}$$

ASSUMPTION

gas is ideal gas. Quasiequilibrium.
closed system.

EFDSOLN

$$\text{initially: } \rho_A = \frac{m_A}{V_A} \dots \textcircled{1}, \rho_B = \frac{m_B}{V_B} \dots \textcircled{2}$$

$$\text{after: } \rho_{\text{tank}} = \frac{m_{\text{tank}}}{V_{\text{tank}}} \dots \textcircled{3}$$

$$\therefore m_{\text{tank}} = m_A + m_B$$

$$\textcircled{3} \Rightarrow \rho_{\text{tank}} = \frac{m_A + m_B}{V_{\text{tank}}}$$

$$m_A = \rho_{\text{tank}} V_{\text{tank}} - m_B = \left(\frac{1.8 \text{ kg}}{\text{m}^3} \right) (15 \text{ m}^3) - 6 \text{ kg} = 21 \text{ kg}$$

$$\therefore V_A = \frac{m_A}{\rho_A} = (21 \text{ kg}) \left(\frac{\text{m}^3}{1.6 \text{ kg}} \right) = 13.125 \text{ m}^3$$

$$V_B = 15 - 13.125 = 1.875 \text{ m}^3$$

$$\rho_B = \frac{m_B}{V_B} = (6 \text{ kg}) \left(\frac{1}{1.875 \text{ m}^3} \right) = \boxed{3.20 \text{ kg/m}^3}$$