Recip-e
Digital medication prescriptions

eHealth DTE testing

Version	Status	Date	Author(s)	Changes
0.1	Draft	13.08.2026	Dries Vanstappen	First version
0.2	Update	02.10.2026	Dries Vanstappen	Added extra field “Dosage elements to use” to Prescriber Use Cases
0.3	Update	09.10.2026	Dries Vanstappen	Added tests for Executors, Patients and Hospitals Minor corrections in syntax and spelling

Reviewers	Name	Reviewed version	Comments
SPOC Klant	Don Leonard		
SPOC K&D			
SPOC GA/TB			
SPOC TPL			
SPOC TPO&P			
SPOC Chain			
Chain Project Leader	Delphine Lanotte		
Chain Service Manager	Sélim Dahmane		

Table of contents

1. INTRODUCTION 4
1.a. Goal of this document 4
2. ACCESS TO EHEALTH DTE 5
3. HOW TO TEST 5
4. PRESCRIBERS 5
4.a. Test-Prescriber-1 5
4.b. Test-Prescriber-2 5
4.c. Test-Prescriber-3 6
4.d. Test-Prescriber-4 7
4.e. Test-Prescriber-5 7
4.f. Test-Prescriber-6 8
4.g. Test-Prescriber-7 8
4.h. Test-Prescriber-8 9
4.i. Test-Prescriber-9 10
4.j. Test-Prescriber-10 10
4.k. Test-Prescriber-11 11
4.l. Test-Prescriber-12 11
5. EXECUTORS 12
5.a. Test-Executor-1 12
5.b. Test-Executor-2 12
6. PATIENTS 12

6.a. Test-Patient-1 12
6.b. Test-Patient-2 12
6.c. Test-Patient-3 13
6.d. Test-Patient-4 13
7. HOSPITALS 14
7.a. Test-Hospital-1 14
7.b. Test-Hospital-2 14
7.c. Test-Hospital-3 14
7.d. Test-Hospital-4 15
7.e. Test-Hospital-5 16
7.f. Test-Hospital-6 16
7.g. Test-Hospital-7 17
7.h. Test-Hospital-8 17
7.i. Test-Hospital-9 18
7.j. Test-Hospital-10 19
7.k. Test-Hospital-11 19
7.l. Test-Hospital-12 20
7.m. Test-Hospital-13 20

1. Introduction

1.a. Goal of this document

This document explains how to test a digital medication prescription setup using the eHealth Digital Testing Environment (DTE).
It describes how to access the DTE and run your tests.
You will find an overview of all tests which are available on the DTE divided per type of actor.
For digital medication prescriptions the 4 possible actors are:

- Prescribers: those who can create a prescription.
- Executors: those who can act upon a prescription.
- Patients: those for whom the prescriptions are created, along with the different mandateholders that can be linked to them.
- Hospitals: Prescribers and Executors acting within the same healthcare organization

If you need any further information or help, please contact the Recip-e team using the form on https://technicalsupport.recip-e.be/
We will answer you as soon as possible.

2. Access to eHealth DTE

The eHealth DTE can be found at https://dte.ehealth.fgov.be
Please contact eHealth to get access.

3. How to test

When you are logged in to the eHealth DTE then you'll find different test projects.
They each have their own title, project description, REST API URL and project related information.
The test projects that are used for testing digital medication prescriptions are "Recip-e Prescriber", "Recip-e Patient", "Recip-e Executor" and "Recip-e Hospital".

You can then use your favorite API platform (for instance Postman) to send the JSONs you created to the REST API URL of the relevant test project.
Or configure your platform to consider the DTE as Recip-e.

4. Prescribers

4.a. Test-Prescriber-1

Action	Create a prescription
Product	Lasix 40 mg tabl. 50 (CNK: 0117-572)
Dosage	Every two days
Prescriber privacy flags	
Pharmacy privacy flags	
Note	Test-Prescriber-1
Scenario	As a prescriber I want to prescribe "Lasix 40 mg tabl. 50" which the patient has to take every other day. I want to add a note "Test-Prescriber-1" to the prescription. I don't want to set provisions for prescribers nor pharmacies.
Dosage elements to use	▪ timing ▪ code

4.b. Test-Prescriber-2

Action	Create a prescription
Product	Medrol 32 mg tabl. 20 (CNK: 0895-540)
Dosage	2 weeks once per day 32 mg, 2 weeks once per day 24 mg, 2 weeks once per day 16 mg, 2 weeks once per day 8mg

Prescriber privacy flags	Only original prescriber may see
Pharmacy privacy flags	All pharmacies may see
Note	Test-Prescriber-2
Scenario	As a prescriber I want to prescribe "Medrol 32 mg tabl. 20". The patient has to take 32mg once per day for two weeks, then 24mg once per day for two weeks, then 16mg once per day for two weeks and finally 8mg once per day for two weeks. I want to add a note "Test-Prescriber-2" to the prescription and set a provision stating that only the original prescriber may view this prescription and that all pharmacists may view this prescription.
Dosage elements to use	▪ sequence ▪ timing ▪ repeat ▪ duration ▪ durationUnit ▪ frequency ▪ period ▪ periodUnit ▪ doseAndRate ▪ doseQuantity

4.c. Test-Prescriber-3

Action	Create a prescription
Product	Methadon 1mg / 1ml oplossing oraal; magistrale bereiding Orale oplossing 80ml - magistrale bereiding • METHADONHYDROCHLORIDE (CNK: 0558387) 1mg / 1ml • CITROENZUUR WATERVRIJ (CNK: 0503037) 1mg / 1ml • BANANEN ESSENCE (CNK: 0561456) 1mg / 1ml • GLYCEROL (CNK: 0525337) 10g / 100ml • GECONSERVEERDE SUIKERSTROOP (CNK: 0584938) 40g / 100ml • WATER GEZUIVERD (CNK: 0507541) 112g / 100ml
Dosage	Daily dose of 20ml During 2 days with a maximum of 4 days.
Prescriber privacy flags	Only original prescriber may see
Pharmacy privacy flags	
Note	Test-Prescriber-3
Scenario	As a prescriber, I want to prescribe the following magistral preparation in a structured way. The patient has to take 20 ml per day for 2 days with a maximum of 4 days. I want to add the note “Test-Prescriber-3” to the prescription and set a provision stating that only the original prescriber may view this prescription. I don't want to set a provision for pharmacies. Methadon 1 mg/1 mL oral solution; compounded preparation Oral syrup 80 mL - magistral preparation • METHADONHYDROCHLORIDE 1mg / 1ml • CITROENZUUR WATERVRIJ 1mg / 1ml • BANANEN ESSENCE 1mg / 1ml • GLYCEROL 10g / 100ml

	• GECONSERVEERDE SUIKERSTROOP 40g / 100ml • WATER GEZUIVERD 112g / 100ml
Dosage elements to use	▪ timing ▪ repeat ▪ duration ▪ durationMax ▪ durationUnit ▪ code ▪ doseAndRate ▪ doseQuantity

4.d. Test-Prescriber-4

Action	Create a prescription
Product	Hukyndra 40 mg inj. opl. s.c. voorgev. pen 6 x 0.4 ml (CNK: 4509-469)
Dosage	Week 0: 80mg, week 2: 40mg, then every two weeks 40mg
Prescriber privacy flags	All prescribers may see
Pharmacy privacy flags	No pharmacy may see
Note	Test-Prescriber-4
Scenario	As a prescriber I want to prescribe "Hukyndra 40 mg inj. opl. s.c. voorgev. pen 6 x 0.4 ml". The patient has to first inject 80mg. two weeks later 40mg and then every 2 weeks 40mg. I want to add the note "Test-Prescriber-4" to the prescription and set a provision stating that all prescribers may see the prescription but no pharmacies may see it.
Dosage elements to use	▪ sequence ▪ timing ▪ repeat ▪ duration ▪ durationUnit ▪ doseAndRate ▪ type ▪ coding ▪ doseQuantity

4.e. Test-Prescriber-5

Action	Create a prescription
Product	Indocollyre 0.1 % oogdruppels opl. druppelfl. 5 ml (CNK: 0435-552)
Dosage	Start one day before the operation until five days after the operation. 4 times per day one drop Operation on 05.12.2026 **For DTE use the following:** Starten 1 dag voor de operatie tot 5 dagen na de operatie – 4 x 1 druppel per dag
Prescriber privacy flags	Only original prescriber may see
Pharmacy privacy flags	No pharmacies may see
Note	Test-Prescriber-5
Scenario	As a prescriber I want to prescribe "Indocollyre 0.1 % oogdruppels opl. druppelfl.

	5 ml". The patient has an operation on 05.12.2026 and has to start the day before the operation until five days after the operation. I want to pass the following instructions to the patient using patientInstruction. "Starten 1 dag voor de operatie tot 5 dagen na de operatie – 4 x 1 druppel per dag" I want to add the note "Test-Prescriber-5" to the prescription and set a provision stating that only the original prescriber may see the prescription and no pharmacy can see it.
Dosage elements to use	▪ patientInstruction

4.f. Test-Prescriber-6

Action	Create a prescription
Product	Durogesic 25 µg/h transderm. pleister sachet 10 (4,2 mg/10,5 cm²) (CNK: 2218-279)
Dosage	Each three days change the patch
Prescriber privacy flags	Only original prescriber and GMD-holder may see
Pharmacy privacy flags	All pharmacies may see
Note	Test-Prescriber-6
Scenario	As a prescriber I want to prescribe "Durogesic 25 µg/h transderm. pleister sachet 10 (4,2 mg/10,5 cm²)". The patient has to change the patch every three days. I want to add the note "Test-Prescriber-6" to the prescription and set a provision which states that only the original prescriber and its GMD-holder may see the prescription and that all pharmacies may see the prescription.
Dosage elements to use	▪ timing ▪ repeat ▪ frequency ▪ period ▪ periodUnit ▪ doseAndRate ▪ type ▪ coding ▪ doseQuantity

4.g. Test-Prescriber-7

Action	Create a prescription
Product	*VOS prescription using VMP-group* bisoprolol oraal 2,5 mg (CNK: 3889)
Dosage	Once per day 2.5mg during twelve weeks
Prescriber privacy flags	Only original prescriber may see
Pharmacy privacy flags	
Note	Test-Prescriber-7
Scenario	As a prescriber I want to prescribe the VMP-group "bisoprolol oraal 2,5 mg". The patient has to take 2.5mg once per day for 12 weeks. I want to add a note "Test-Prescriber-7" to the prescription and set a provision stating that only the original prescriber may see the prescription. I don't want to set a provision for pharmacies.

Dosage elements to use	▪ timing ▪ repeat ▪ duration ▪ durationUnit ▪ frequency ▪ period ▪ periodUnit ▪ doseAndRate ▪ type ▪ coding ▪ doseQuantity ▪ maxDosePerPeriod ▪ numerator ▪ denominator

4.h. Test-Prescriber-8

Action	Create a prescription
Product	*Currently imported from abroad – product normally available in Belgium* Colchicine Opocalcium 1 mg tabl. 20 (CNK: 1127-299)
Dosage	For an acute gout attack: • (with normal kidney function): start within the first 12 hours with 1 mg, followed by 0.5 mg one hour later, and continue on the following days with 0.5 mg 2 to 3 times per day. • Maximum 6 mg during a treatment course. **For DTE use the following:** Voor een acute jichtaanval: • (bij normale nierfunctie): begin binnen de eerste 12 uur met 1 mg, gevolgd door 0,5 mg een uur later, en ga de volgende dagen door met 0,5 mg 2 tot 3 keer per dag. • Max 6mg tijdens een behandelkuur
Prescriber privacy flags	
Pharmacy privacy flags	
Note	Test-Prescriber-8
Scenario	As a prescriber I want to prescribe "Colchicine Opocalcium 1 mg tabl. 20". I want to give the following instructions to the patient using patientInstruction Voor een acute jichtaanval: • (bij normale nierfunctie): begin binnen de eerste 12 uur met 1 mg, gevolgd door 0,5 mg een uur later, en ga de volgende dagen door met 0,5 mg 2 tot 3 keer per dag. • Max 6mg tijdens een behandelkuur I want to add a note "Test-Prescriber-8" to the prescription. I don't want to set any provisions.
Dosage elements to use	▪ patientInstruction

4.i. Test-Prescriber-9

Action	Create a prescription
Product	*Currently imported from abroad – product not available in Belgium* Gutron 5mg **Pass product on as a free text AMPP**
Dosage	Starting dose is 2.5mg three times per day for one week. Every week increase the dose by 2.5 mg with a maximum of 10mg three times a day without an end date.
Prescriber privacy flags	
Pharmacy privacy flags	All pharmacies may see
Note	Test-Prescriber-9
Scenario	As a prescriber I want to prescribe "Gutron 5mg". This product is not available in Belgium. The patient has to take a dose of 2.5mg three times per day for a week. Every following week they have to increase the dose by 2.5mg to a maximum of 10mg three times per day without an end date. I want to add a note "Test-Prescriber-9" to the prescription. I don't want to set a provision for prescribers but I want to set one for pharmacies stating that all pharmacies may see the prescription.
Dosage elements to use	▪ sequence ▪ timing ▪ repeat ▪ duration ▪ durationUnit ▪ code ▪ doseAndRate ▪ type ▪ coding ▪ doseQuantity

4.j. Test-Prescriber-10

Action	Create a prescription
Product	Gasvormige Medicinale Zuurstof Air Products 100 % v/v medicin. gas (samengep.) kader gascilinders 117.7 m3 (CNK: 4005-393)
Dosage	3 liters per minute, 18 hours per day during one month. Add patient instruction “With oxygen humidifier” **For DTE use the following:** Met zuurstofbevochtiger
Prescriber privacy flags	
Pharmacy privacy flags	No pharmacies may see
Note	Test-Prescriber-10
Scenario	As a prescriber I want to prescribe "Gasvormige Medicinale Zuurstof Air Products 100 % v/v medicin. gas (samengep.) kader gascilinders 117.7 m3". The patient has to take three liters per minute for 18 hours per day during one month. I want to pass the following instruction to the patient: "Met zuurstofbevochtiger" I want to add a note "Test-Prescriber-10" to the prescription.

	I don't want to set a provision for prescribers but I want to set one for pharmacies stating that no pharmacy may see the prescription.
Dosage elements to use	▪ patientInstruction ▪ timing ▪ repeat ▪ boundsDuration ▪ duration ▪ durationUnit ▪ frequency ▪ period ▪ periodUnit ▪ doseAndRate ▪ rateQuantity

4.k. Test-Prescriber-11

Action	Create a prescription
Product	Lactulose EG 670 mg/ml siroop 5 l (CNK: 1455-450)
Dosage	Daily before each meal A minimum of 15ml and a maximum of 45ml
Prescriber privacy flags	All prescribers may see
Pharmacy privacy flags	All pharmacies may see
Note	Test-Prescriber-11
Scenario	As a prescriber I want to prescribe "Lactulose EG 670 mg/ml siroop 5 l". The patient has to take at least 15ml and a maximum of 45ml daily before each meal. I want to add a note "Test-Prescriber-11" to the prescription. I want to set a provision stating that all prescribers and all pharmacies may see the prescription.
Dosage elements to use	▪ timing ▪ repeat ▪ when ▪ code ▪ coding ▪ doseAndRate ▪ type ▪ coding ▪ doseRange ▪ low ▪ high

4.l. Test-Prescriber-12

Action	Cancel a prescription
Note	Test-Prescriber-12
Scenario	As a prescriber I want to cancel a prescription.

	I want to add a note “Test-Prescriber-12” to the Task

5. Executors

5.a. Test-Executor-1

Action	Cancel prescription
Note	Test-Executor-1
Scenario	As an executor I want to cancel an active prescription. I want to add a note “Test-Executor-1” to the Task

5.b. Test-Executor-2

Action	Complete prescription
Note	Test-Executor-2
Scenario	As an executor I want to complete an active prescription. I want to add a note “Test-Executor-2” to the Task

6. Patients

6.a. Test-Patient-1

Action	Update Pharmacy PF to "Only specific pharmacies"
Note	Test-Patient-1
Scenario	As a patient I want to update the privacy flags of a prescription so that only a specific pharmacy may see the prescription. I don't want to send an update about the prescriber privacy flag. I want to add a note “Test-Patient-1” to the Task

6.b. Test-Patient-2

Action	Update Prescriber PF to "Only prescriber himself" and Pharmacy PF to "No pharmacy may see"
Note	Test-Patient-2
Scenario	As a patient I want to update the privacy flags of a prescription so that only the original prescriber and no pharmacies may see the prescription. I want to use only 1 FHIR Consent element. I want to add a note “Test-Patient-2” to the Task

6.c. Test-Patient-3

Action	Update Prescriber PF to "All prescribers may see"
Note	Test-Patient-3
Scenario	As a patient I want to update the privacy flags of a prescription so that all prescribers may see the prescription. I don't want to send an update about the pharmacy privacy flag. I want to add a note “Test-Patient-3” to the Task

6.d. Test-Patient-4

Action	Cancel a prescription
Note	Test-Patient-4
Scenario	As a patient I want to cancel a prescription with reason "No longer needed". I want to add a note “Test-Patient-4” to the Task

7. Hospitals

7.a. Test-Hospital-1

Action	Create a prescription
Product	Lasix 40 mg tabl. 50 (CNK: 0117-572)
Dosage	Every two days
Prescriber privacy flags	
Pharmacy privacy flags	
Note	Test-Hospital-1
Scenario	As a prescriber linked to a hospital, I want to prescribe "Lasix 40 mg tabl. 50" which the patient has to take every other day. I want to add a note "Test-Hospital-1" to the prescription. I don't want to set provisions for prescribers nor pharmacies.
Dosage elements to use	▪ timing ▪ code

7.b. Test-Hospital-2

Action	Create a prescription
Product	Medrol 32 mg tabl. 20 (CNK: 0895-540)
Dosage	2 weeks once per day 32 mg, 2 weeks once per day 24 mg, 2 weeks once per day 16 mg, 2 weeks once per day 8mg
Prescriber privacy flags	Only original prescriber may see
Pharmacy privacy flags	All pharmacies may see
Note	Test-Hospital-2
Scenario	As a prescriber linked to a hospital, I want to prescribe "Medrol 32 mg tabl. 20". The patient has to take 32mg once per day for two weeks, then 24mg once per day for two weeks, then 16mg once per day for two weeks and finally 8mg once per day for two weeks. I want to add a note "Test-Hospital-2" to the prescription and set a provision stating that only the original prescriber may view this prescription and that all pharmacists may view this prescription.
Dosage elements to use	▪ sequence ▪ timing ▪ repeat ▪ duration ▪ durationUnit ▪ frequency ▪ period ▪ periodUnit ▪ doseAndRate ▪ doseQuantity

7.c. Test-Hospital-3

Action	Create a prescription

Product	Methadon 1mg / 1ml oplossing oraal; magistrale bereiding Orale oplossing 80ml - magistrale bereiding • METHADONHYDROCHLORIDE (CNK: 0558387) 1mg / 1ml • CITROENZUUR WATERVRIJ (CNK: 0503037) 1mg / 1ml • BANANEN ESSENCE (CNK: 0561456) 1mg / 1ml • GLYCEROL (CNK: 0525337) 10g / 100ml • GECONSERVEERDE SUIKERSTROOP (CNK: 0584938) 40g / 100ml • WATER GEZUIVERD (CNK: 0507541) 112g / 100ml
Dosage	Daily dose of 20ml During 2 days with a maximum of 4 days.
Prescriber privacy flags	Only original prescriber may see
Pharmacy privacy flags	
Note	Test-Hospital-3
Scenario	As a prescriber linked to a hospital, I want to prescribe the following magistral preparation in a structured way. The patient has to take 20 ml per day for 2 days with a maximum of 4 days. I want to add the note “Test-Hospital-3” to the prescription and set a provision stating that only the original prescriber may view this prescription. I don't want to set a provision for pharmacies. Methadon 1 mg/1 mL oral solution; compounded preparation Oral syrup 80 mL - magistral preparation • METHADONHYDROCHLORIDE 1mg / 1ml • CITROENZUUR WATERVRIJ 1mg / 1ml • BANANEN ESSENCE 1mg / 1ml • GLYCEROL 10g / 100ml • GECONSERVEERDE SUIKERSTROOP 40g / 100ml • WATER GEZUIVERD 112g / 100ml
Dosage elements to use	▪ timing ▪ repeat ▪ duration ▪ durationMax ▪ durationUnit ▪ code ▪ doseAndRate ▪ doseQuantity

7.d. Test-Hospital-4

Action	Create a prescription
Product	Hukyndra 40 mg inj. opl. s.c. voorgev. pen 6 x 0.4 ml (CNK: 4509-469)
Dosage	Week 0: 80mg, week 2: 40mg, then every two weeks 40mg
Prescriber privacy flags	All prescribers may see
Pharmacy privacy flags	No pharmacy may see
Note	Test-Hospital-4
Scenario	As a prescriber linked to a hospital, I want to prescribe "Hukyndra 40 mg inj. opl. s.c. voorgev. pen 6 x 0.4 ml". The patient has to first inject 80mg. two weeks later 40mg and then every 2 weeks 40mg. I want to add the note "Test-Hospital-4" to the prescription and set a provision stating that all prescribers may see the prescription but no pharmacies may see

	it.
Dosage elements to use	▪ sequence ▪ timing ▪ repeat ▪ duration ▪ durationUnit ▪ doseAndRate ▪ type ▪ coding ▪ doseQuantity

7.e. Test-Hospital-5

Action	Create a prescription
Product	Indocollyre 0.1 % oogdruppels opl. druppelfl. 5 ml (CNK: 0435-552)
Dosage	Start one day before the operation until five days after the operation. 4 times per day one drop Operation on 05.12.2026 **For DTE use the following:** Starten 1 dag voor de operatie tot 5 dagen na de operatie – 4 x 1 druppel per dag
Prescriber privacy flags	Only original prescriber may see
Pharmacy privacy flags	No pharmacies may see
Note	Test-Hospital-5
Scenario	As a prescriber linked to a hospital, I want to prescribe "Indocollyre 0.1 % oogdruppels opl. druppelfl. 5 ml". The patient has an operation on 05.12.2026 and has to start the day before the operation until five days after the operation. I want to pass the following instructions to the patient using patientInstruction. "Starten 1 dag voor de operatie tot 5 dagen na de operatie – 4 x 1 druppel per dag" I want to add the note "Test-Hospital-5" to the prescription and set a provision stating that only the original prescriber may see the prescription and no pharmacy can see it.
Dosage elements to use	▪ patientInstruction

7.f. Test-Hospital-6

Action	Create a prescription
Product	Durogesic 25 µg/h transderm. pleister sachet 10 (4,2 mg/10,5 cm²) (CNK: 2218-279)
Dosage	Each three days change the patch
Prescriber privacy flags	Only original prescriber and GMD-holder may see
Pharmacy privacy flags	All pharmacies may see
Note	Test-Hospital-6
Scenario	As a prescriber linked to a hospital, I want to prescribe "Durogesic 25 µg/h transderm. pleister sachet 10 (4,2 mg/10,5 cm²)". The patient has to change the patch every three days. I want to add the note "Test-Hospital-6" to the prescription and set a provision which states that only the original prescriber and its GMD-holder may see the

	prescription and that all pharmacies may see the prescription.
Dosage elements to use	▪ timing ▪ repeat ▪ frequency ▪ period ▪ periodUnit ▪ doseAndRate ▪ type ▪ coding ▪ doseQuantity

7.g. Test-Hospital-7

Action	Create a prescription
Product	*VOS prescription using VMP-group* bisoprolol oraal 2,5 mg (CNK: 3889)
Dosage	Once per day 2.5mg during twelve weeks
Prescriber privacy flags	Only original prescriber may see
Pharmacy privacy flags	
Note	Test-Hospital-7
Scenario	As a prescriber linked to a hospital, I want to prescribe the VMP-group "bisoprolol oraal 2,5 mg". The patient has to take 2.5mg once per day for 12 weeks. I want to add a note "Test-Hospital-7" to the prescription and set a provision stating that only the original prescriber may see the prescription. I don't want to set a provision for pharmacies.
Dosage elements to use	▪ timing ▪ repeat ▪ duration ▪ durationUnit ▪ frequency ▪ period ▪ periodUnit ▪ doseAndRate ▪ type ▪ coding ▪ doseQuantity ▪ maxDosePerPeriod ▪ numerator ▪ denominator

7.h. Test-Hospital-8

Action	Create a prescription
Product	*Currently imported from abroad – product normally available in Belgium* Colchicine Opocalcium 1 mg tabl. 20 (CNK: 1127-299)
Dosage	For an acute gout attack: • (with normal kidney function): start within the first 12 hours with 1 mg, followed by 0.5 mg one hour later, and continue on the following days with 0.5 mg 2 to 3 times per day.

	• Maximum 6 mg during a treatment course. **For DTE use the following:** Voor een acute jichtaanval: • (bij normale nierfunctie): begin binnen de eerste 12 uur met 1 mg, gevolgd door 0,5 mg een uur later, en ga de volgende dagen door met 0,5 mg 2 tot 3 keer per dag. • Max 6mg tijdens een behandelkuur
Prescriber privacy flags	
Pharmacy privacy flags	
Note	Test-Hospital-8
Scenario	As a prescriber linked to a hospital, I want to prescribe "Colchicine Opocalcium 1 mg tabl. 20". I want to give the following instructions to the patient using patientInstruction Voor een acute jichtaanval: • (bij normale nierfunctie): begin binnen de eerste 12 uur met 1 mg, gevolgd door 0,5 mg een uur later, en ga de volgende dagen door met 0,5 mg 2 tot 3 keer per dag. • Max 6mg tijdens een behandelkuur I want to add a note "Test-Hospital-8" to the prescription. I don't want to set any provisions.
Dosage elements to use	▪ patientInstruction

7.i. Test-Hospital-9

Action	Create a prescription
Product	*Currently imported from abroad – product not available in Belgium* Gutron 5mg **Pass product on as a free text AMPP**
Dosage	Starting dose is 2.5mg three times per day for one week. Every week increase the dose by 2.5 mg with a maximum of 10mg three times a day without an end date.
Prescriber privacy flags	
Pharmacy privacy flags	All pharmacies may see
Note	Test-Hospital-9
Scenario	As a prescriber linked to a hospital, I want to prescribe "Gutron 5mg". This product is not available in Belgium. The patient has to take a dose of 2.5mg three times per day for a week. Every following week they have to increase the dose by 2.5mg to a maximum of 10mg three times per day without an end date. I want to add a note "Test-Hospital-9" to the prescription. I don't want to set a provision for prescribers but I want to set one for pharmacies stating that all pharmacies may see the prescription.
Dosage elements to use	▪ sequence ▪ timing ▪ repeat ▪ duration ▪ durationUnit

	▪ code ▪ doseAndRate ▪ type ▪ coding ▪ doseQuantity

7.j. Test-Hospital-10

Action	Create a prescription
Product	Gasvormige Medicinale Zuurstof Air Products 100 % v/v medicin. gas (samengep.) kader gascilinders 117.7 m3 (CNK: 4005-393)
Dosage	3 liters per minute, 18 hours per day during one month. Add patient instruction “With oxygen humidifier” **For DTE use the following:** Met zuurstofbevochtiger
Prescriber privacy flags	
Pharmacy privacy flags	No pharmacies may see
Note	Test-Hospital-10
Scenario	As a prescriber linked to a hospital, I want to prescribe "Gasvormige Medicinale Zuurstof Air Products 100 % v/v medicin. gas (samengep.) kader gascilinders 117.7 m3". The patient has to take three liters per minute for 18 hours per day during one month. I want to pass the following instruction to the patient: "Met zuurstofbevochtiger" I want to add a note "Test-Hospital-10" to the prescription. I don't want to set a provision for prescribers but I want to set one for pharmacies stating that no pharmacy may see the prescription.
Dosage elements to use	▪ patientInstruction ▪ timing ▪ repeat ▪ boundsDuration ▪ duration ▪ durationUnit ▪ frequency ▪ period ▪ periodUnit ▪ doseAndRate ▪ rateQuantity

7.k. Test-Hospital-11

Action	Create a prescription
Product	Lactulose EG 670 mg/ml siroop 5 l (CNK: 1455-450)
Dosage	Daily before each meal A minimum of 15ml and a maximum of 45ml
Prescriber privacy flags	All prescribers may see
Pharmacy privacy flags	All pharmacies may see
Note	Test-Hospital-11

Scenario	As a prescriber linked to a hospital, I want to prescribe "Lactulose EG 670 mg/ml siroop 5 l". The patient has to take at least 15ml and a maximum of 45ml daily before each meal. I want to add a note "Test-Hospital-11" to the prescription. I want to set a provision stating that all prescribers and all pharmacies may see the prescription.
Dosage elements to use	▪ timing ▪ repeat ▪ when ▪ code ▪ coding ▪ doseAndRate ▪ type ▪ coding ▪ doseRange ▪ low ▪ high

7.l. Test-Hospital-12

Action	Cancel prescription
Note	Test-Hospital-12
Scenario	As an executor acting within a hospital, I want to cancel an active prescription.

7.m. Test-Hospital-13

Action	Complete prescription
Note	Test-Hospital-13
Scenario	As an executor acting within a hospital, I want to complete an active prescription.