

Verified label (manual GT)

1837 normalized characters

16 COMMON SENSE MEDICAL ADVISER.

ALTERNATE GENERATION.

In several insect families, the species is not wholly represented in the adult individuals of both sexes, or in their development, but, to complete this series, supplementary individuals, as it were, of one or of several preceding generations, are required. The son may not resemble the father, but the grandfather, and in some instances, the likeness re-appears only in latter generations. Agassiz states: "Alternate generation was first observed among the Salpæ. These are marine mollusks, without shells, belonging to the family Tunicata. They are distinguished by the curious peculiarity of being united together in considerable numbers so as to form long chains, which float in the sea, the mouth (m) however being free in each.

Fig. 2. Fig. 3.

" Fig. 2. The individuals thus joined in floating colonies produce eggs; but in each animal there is generally but one egg formed, which is developed in the body of the parent, and from which is hatched a little mollusk.

" Fig. 3, which remains solitary, and differs in many respects from the parent. This little animal, on the other hand, does not produce eggs, but propagates, by a kind of budding, which gives rise to chains already seen in the body of their parent (a), and these again bring forth solitary individuals, etc."

It therefore follows that generation in some animals requires two different bodies with intermediate ones, by means of which, and their different modes of reproduction, a return to the original stock is effected.

Universality of Animalcular Life.—Living organisms are universally diffused over every part of the globe. The gentle zephyr wafts from flower to flower invisible, fructifying atoms, which quicken beauty and fragrance, giving the promise of a golden fruitage, to gladden and nourish a dependent world. Nature's own sweet cunning invests all living things,

DeepSeek-OCR-2

CER 0.0484 | WER 0.0777 | F1 0.9486 | 1 region(s) | length 1.02x

sub_title

ALTERNATE GENERATION.

text

In several insect families, the species is not wholly represented in the adult individuals of both sexes, or in their development, but, to complete this series, supplementary individuals, as it were, of one or of several preceding generations, are required. The son may not resemble the father, but the grandfather, and in some instances, the likeness re-appears only in latter generations. Agassiz states: "Alternate generation was first observed among the Salpæ. These are marine mollusks, without shells, belonging to the family Tunicata. They are distinguished by the curious peculiarity of being united together in considerable numbers so as to form long chains, which float in the sea, the mouth (m) however being free in each.

figure_title

Fig. 2.

image

figure_title

Fig. 3.

text

"Fig. 2. The individuals thus joined in floating colonies produce eggs; but in each animal there is generally but one egg formed, which is developed in the body of the parent, and from which is hatched a little mollusk.

text

"Fig. 3, which remains solitary, and differs in many respects from the parent. This little animal, on the other hand, does not produce eggs, but propagates, by a kind of budding, which gives rise to chains already seen in the body of their parent (a), and these again bring forth solitary individuals, etc."

text

It therefore follows that generation in some animals requires two different bodies with intermediate ones, by means of which, and their different modes of reproduction, a return to the original stock is effected.

text

Universality of Animalcular Life.—Living organisms are universally diffused over every part of the globe. The gentle zephyr wafts from flower to flower invisible, fructifying atoms, which quicken beauty and fragrance, giving the promise of a golden fruitage, to gladden and nourish a dependent world. Nature's own sweet cunning invests all living things,

GLM-OCR

CER 0.0218 | WER 0.0388 | F1 0.9755 | 1 region(s) | length 0.98x

ALTERNATE GENERATION.

In several insect families, the species is not wholly represented in the adult individuals of both sexes, or in their development, but, to complete this series, supplementary individuals, as it were, of one or of several preceding generations, are required. The son may not resemble the father, but the grandfather, and in some instances, the likeness re-appears only in latter generations. Agassiz states: "Alternate generation was first observed among the Salpæ. These are marine mollusks, without shells, belonging to the family Tunicata. They are distinguished by the curious peculiarity of being united together in considerable numbers so as to form long chains, which float in the sea, the mouth (m) however being free in each.

Fig. 2. Fig. 3.

"Fig. 2. The individuals thus joined in floating colonies produce eggs; but in each animal there is generally but one egg formed, which is developed in the body of the parent, and from which is hatched a little mollusk.

"Fig. 3, which remains solitary, and differs in many respects from the parent. This little animal, on the other hand, does not produce eggs, but propagates, by a kind of budding, which gives rise to chains already seen in the body of their parent (a), and these again bring forth solitary individuals, etc."

It therefore follows that generation in some animals requires two different bodies with intermediate ones, by means of which, and their different modes of reproduction, a return to the original stock is effected.

Universality of Animalcular Life.—Living organisms are universally diffused over every part of the globe. The gentle zephyr wafts from flower to flower invisible, fructifying atoms, which quicken beauty and fragrance, giving the promise of a golden fruitage, to gladden and nourish a dependent world. Nature's own sweet cunning invests all living things,<[user]>

MinerU 2.5 Pro

CER 0.0087 | WER 0.0356 | F1 0.9755 | 1 region(s) | length 0.99x

16

COMMON SENSE MEDICAL ADVISER.

ALTERNATE GENERATION.

In several insect families, the species is not wholly represented in the adult individuals of both sexes, or in their development, but, to complete this series, supplementary individuals, as it were, of one or of several preceding generations, are required. The son may not resemble the father, but the grandfather, and in some instances, the likeness re-appears only in latter generations. Agassiz states: "Alternate generation was first observed among the Salpæ. These are marine mollusks, without shells, belonging to the family Tunicata. They are distinguished by the curious peculiarity of being united together in considerable numbers so as to form long chains, which float in the sea, the mouth (m) however being free in each.

"Fig. 2. The individuals thus joined in floating colonies produce eggs; but in each animal there is generally but one egg formed, which is developed in the body of the parent, and from which is hatched a little mollusk.

"Fig. 3, which remains solitary, and differs in many respects from the parent. This little animal, on the other hand, does not produce eggs, but propagates, by a kind of budding, which gives rise to chains already seen in the body of their parent (a), and these again bring forth solitary individuals, etc."

It therefore follows that generation in some animals requires two different bodies with intermediate ones, by means of which, and their different modes of reproduction, a return to the original stock is effected.

Universality of Animalcular Life.—Living organisms are universally diffused over every part of the globe. The gentle zephyr wafts from flower to flower invisible, fructifying atoms, which quicken beauty and fragrance, giving the promise of a golden fruitage, to gladden and nourish a dependent world. Nature's own sweet cunning invests all living things,

TrOCR large-printed

CER 0.0463 | WER 0.1456 | F1 0.8631 | 1 region(s) | length 1.01x

16 COMMON SENSE MEDICAL ADVISER,

AMOUNT ALTERNATE GENERATION.

IN SEVERAL INSERT FAMILIES, THE SPECIES IS NOT WHOLLY REPRESENTED IN THE ADULT INDIVIDUALS OF BOTH SEXES, OR IN THEIR DEVELOPMENT, BUT, TO COMPLETE THIS SERIES, SUPPLEMENTARY INDIVIDUALS, AS IT WERE, OF ONE OR OF SEVERAL PRECEDING GENERATIONS, ARE REQUIRED. THE SON MAY NOT RESEMBLE THE FATHER, BUT THE GRANDFATHER, AND IN SOME INSTANCES, THE LIKENESS RE-APPER'S ONLY IN LATTER GENERATIONS, AGASSIZ STATES: "ALTERNATE GENERATION WAS FIRST OBSERVED AMONG THE SALPÆ. THESE ARE MARINE MOLLUSKS, WITHOUT SHELLS, BELONGING TO THE FAMILY TUNICATE. THEY ARE DISTINGUISHED BY THE CURIOUS PECULIARITY OF BEING UNITED TOGETHER IN CONSIDERABLE NUMBERS SO AS TO FORM LONG CHAINS, WHICH FLOAT IN THE SEA, THE MOUTH (IN) HOWEVER BEING FREE IN EACH.

PAY REWARD NO. 24.00

SR:

"FIG. 2. THE INDIVIDUALS THIS JOINED IN ROATING COLONIES PRODUCE EGGS; BUT IN EACH ANIMAL THERE IS GENERALLY DATE EGG FORMED, WHICH IS DEVELOPED IN THE BODY OF THE PARENT, AND FROM WHICH IS HATCHED A LITTLE MOLLUSK.

4FIG. 3, WHICH REMAINS SOLITARY, AND DIFFERS IN MANY RESPECTS FROM THE PARENT. THIS LITTLE ANIMAL, ON THE OTHER HAND, DOES NOT PRODUCE EGGS, BUT PROPAGATES, BY A KIND OF BUDLING, WHICH GIVES RISE TO CHAINS ALREADY SEEN IN THE BODY OF THEIR PARENT (A), AND THESE AGAIN BRING FORTH SOLITARY INDIVIDUALS, ETC.

IT THEREFORE FOLLOWS THAT GENERATION IN SOME ANIMALS REQUIRES TWO DIFFERANT BODIES WITH INTERMULATE ONES, BY MEANS OF WHICH, AND THEIR DIFFERENT MODES OF REPRODUCTION, A RETURN TO THE ORIGINAL STOCK IS EFFECTED.

UNIVERSALITY OF ANIMALCULAR LIFE. - LIVING ORGANISMS ARE UNIVERSALLY DIFFUSEL OVER EVERY PART OF THE GLOBE. THE GENTLE ZEPHYR WAITS FROM FLOWER TO FLOWER INVISIBLE, FRUITIFYING ATOMS, WHICH QUICKEN BEAUTY AND FRAGRANCE, GIVING THE PROMISE OF A GOLDEN FRUITAGE, TO GLADDEN AND NOURISH A DEPENDENT WORLD. NATURE'S OWN SWEET CUNNING INVESTS ALL LIVING THINGS,

Verified label (manual GT)

1837 normalized characters

16 COMMON SENSE MEDICAL ADVISER.

ALTERNATE GENERATION.

In several insect families, the species is not wholly represented in the adult individuals of both sexes, or in their development, but, to complete this series, supplementary individuals, as it were, of one or of several preceding generations, are required. The son may not resemble the father, but the grandfather, and in some instances, the likeness re-appears only in latter generations. Agassiz states: "Alternate generation was first observed among the Salpæ. These are marine mollusks, without shells, belonging to the family Tunicata. They are distinguished by the curious peculiarity of being united together in considerable numbers so as to form long chains, which float in the sea, the mouth (m) however being free in each.

Fig. 2. Fig. 3.

" Fig. 2. The individuals thus joined in floating colonies produce eggs ; but in each animal there is generally but one egg formed, which is developed in the body of the parent, and from which is hatched a little mollusk.

" Fig. 3, which remains solitary, and differs in many respects from the parent. This little animal, on the other hand, does not produce eggs, but propagates, by a kind of budding, which gives rise to chains already seen in the body of their parent (a), and these again bring forth solitary individuals, etc."

It therefore follows that generation in some animals requires two different bodies with intermediate ones, by means of which, and their different modes of reproduction, a return to the original stock is effected.

Universality of Animalcular Life.—Living organisms are universally diffused over every part of the globe. The gentle zephyr wafts from flower to flower invisible, fructifying atoms, which quicken beauty and fragrance, giving the promise of a golden fruitage, to gladden and nourish a dependent world. Nature's own sweet cunning invests all living things,

DeepSeek-OCR-2

CER 0.3500 | WER 0.5922 | F1 0.7484 | 10 region(s) | length 1.33x

1 0

<table><tr><td>Category</td><td>Value</td></tr><tr><td>Overall</td><td>100</td></tr><tr><td>Overall</td><td>100</td></tr><tr><td>Overall</td><td>100</td></tr><tr><td>Overall</td><td>100</td></tr></table>

<smiles>C[C@H](C)N(C)C</smiles>

In several insect families, the species is not wholly represented in the adult individuals of both sexes, or in their development, but, to complete this series, supplementary individuals, as it were, of one or of several preceding generations, are required. The son may not resemble the father, but the grandfather, and in some instances, the likeness re-appears only in latter generations. Agassiz states: "Alternate generation was first observed among the Salpæ. These are marine mollusks, without shells, belonging to the family Tunicata. They are distinguished by the curious peculiarity of being united together in considerable numbers so as to form long chains, which float in the sea, the mouth (m) however being free in each."

[H(u, v)]

[H_2O_2]

"Fig. 2. The individuals thus joined in floating colonies produce eggs; but in each animal there is generally but one egg, formed, which is developed in the body of the parent, and from which is hatched a little mollusk."

"Fig. 3, which remains solitary, and differs in many respects from the parent. This little animal, on the other hand, does not produce eggs, but propagates by a kind of budding, which gives rise to chains already seen in the body of their parent (a), and these again bring forth fourth solitary individuals, etc."

It therefore follows that generation in some animals requires two different bodies with intermediate ones, by means of which, and their different modes of reproduction, a return to the original stock is effected.

Universality of Animalcular Life

Universality of Animalcular Life – Living organisms are universally diffused over every part of the globe. The gentle zephyr wafts from flower to flower invisible, fructifying atoms, which quicken beauty and fragrance, giving the promise of a golden fruitage, to gladden and nourish a dependent world. Nature's own sweet cunning invests all living things,

GLM-OCR

CER 0.0299 | WER 0.0583 | F1 0.9696 | 10 region(s) | length 1.03x

16<[user]>

COMMON SENSE MEDICAL ADVISER.<[user]>

ALTERNATE GENERATION.<[user]>

In several insect families, the species is not wholly represented in the adult individuals of both sexes, or in their development, but, to complete this series, supplementary individuals, as it were, of one or of several preceding generations, are required. The son may not resemble the father, but the grandfather, and in some instances, the likeness re-appears only in latter generations. Agassiz states: "Alternate generation was first observed among the Salpæ. These are marine mollusks, without shells, belonging to the family Tunicata. They are distinguished by the curious peculiarity of being united together in considerable numbers so as to form long chains, which float in the sea, the mouth (m) however being free in each.

Fig. 2. <[user]>

Fig. 3. <[user]>

"Fig. 2. The individuals thus joined in floating colonies produce eggs; but in each animal there is generally but one egg formed, which is developed in the body of the parent, and from which is hatched a little mollusk.<[user]>

"Fig. 3, which remains solitary, and differs in many respects from the parent. This little animal, on the other hand, does not produce eggs, but propagates, by a kind of budding, which gives rise to chains already seen in the body of their parent (a), and these again bring forth solitary individuals, etc."<[user]>

It therefore follows that generation in some animals requires two different bodies with intermdiate ones, by means of which, and their different modes of reproduction, a return to the original stock is effected.<[user]>

Universality of Animalcular Life.—Living organisms are universally diffused over every part of the globe. The gentle zephyr wafts from flower to flower invisible, fructifying atoms, which quicken beauty and fragrance, giving the promise of a golden fruitage, to gladden and nourish a dependent world. Nature's own sweet cunning invests all living things,<[user]>

MinerU 2.5 Pro

CER 0.0082 | WER 0.0291 | F1 0.9804 | 10 region(s) | length 0.99x

16

COMMON SENSE MEDICAL ADVISER.

ALTERNATE GENERATION.

In several insect families, the species is not wholly represented in the adult individuals of both sexes, or in their development, but, to complete this series, supplementary individuals, as it were, of one or of several preceding generations, are required. The son may not resemble the father, but the grandfather, and in some instances, the likeness re-appears only in latter generations. Agassiz states: "Alternate generation was first observed among the Salpæ. These are marine mollusks, without shells, belonging to the family Tunicata. They are distinguished by the curious peculiarity of being united together in considerable numbers so as to form long chains, which float in the sea, the mouth (m) however being free in each.

"Fig. 2. The individuals thus joined in floating colonies produce eggs; but in each animal there is generally but one egg formed, which is developed in the body of the parent, and from which is hatched a little mollusk.

"Fig. 3, which remains solitary, and differs in many respects from the parent. This little animal, on the other hand, does not produce eggs, but propagates, by a kind of budding, which gives rise to chains already seen in the body of their parent (a), and these again bring forth solitary individuals, etc."

It therefore follows that generation in some animals requires two different bodies with intermediate ones, by means of which, and their different modes of reproduction, a return to the original stock is effected.

Universality of Animalcular Life.—Living organ-

isms are universally diffused over every part of the globe. The gentle zephyr wafts from flower to flower invisible, fructifying atoms, which quicken beauty and fragrance, giving the promise of a golden fruitage, to gladden and nourish a dependent world. Nature's own sweet cunning invests all living things,

TrOCR large-printed

CER 0.0185 | WER 0.0809 | F1 0.9256 | 10 region(s) | length 1.00x

16

COMMON SENSE MEDICAL ADVISER.

ALTERNATE GENERATION.

IN SEVERAL INSECT FAMILIES, THE SPECIES IS NOT WHOLLY REPRESENTED IN THE ADULT INDIVIDUALS OF BOTH SEXES, OR IN THEIR DEVELOPMENT, BUT, TO COMPLETE THIS SERIES, SUPPLEMENTARY INDIVIDUALS, AS IT WERE, OF ONE OR OF SEVERAL PRECEDING GENERATIONS, ARE REQUIRED. THE SON MAY NOT RESEMBLE THE FATHER, BUT THE GRANDFATHER, AND IN SOME INSTANCES, THE LIKENESS RE-APPEARS ONLY IN LATTER GENERATIONS. AGASSIZ STATES: "ALTERNATE GENERATION WAS FIRST OBSERVED AMONG THE SALPÆ. THESE ARE MARINE MOLLUSKS, WITHOUT SHELLS, BELONGING TO THE FAMILY TUNICATA. THEY ARE DISTINGUISHED BY THE CURIOUS DECULARITY OF BEING UNITED TOGETHER IN CONSIDERABLE NUMBERS SO AS TO FORM LONG CHAINS, WHICH ROAT IN THE SEA, THE MOUTH (M) HOWEVER BEING FREE IN EACH.

Fig. 2.

"FIG. 2. THE INDIVIDUALS thus joined in ROATING COLONIES PRODUCE EGGS; BUT IN EACH ANIMAL THERE IS GENERALLY BUT ONE EGG FORMED, WHICH IS DEVELOPED IN THE BODY OF THE PARENT, AND FROM WHICH IS HATCHED A LITTLE MOLLUSK.

"FIG. 3, WHICH REMAINS SOLITARY, AND DITERS IN MANY RESPECTS FROM THE PARENT. THIS LITTLE ANIMAL, ON THE OTHER HAND, DOES NOT PRODUCE EGGS, BUT PROPAGATES, BY A KIND OF BUDDING, WHICH GIVES RISE TO CHAINS ALREADY SEEN IN THE BODY OF THEIR PARENT (G), AND THESE AGAIN BRING FORTH SOLITARY INDIVIDUALS, ETC."

IT THEREFORE FOLLOWS THAT GENERATION IN SOME ANIMALS REQUIRES TWO DIFFERENT BODIES WITH INTERMEDIATE ONES, BY MEANS OF WHICH, AND THEIR DIFFERENT MODES OF REPRODUCTION, A RETURN TO THE ORIGINAL STOCK IS EFFECTED.

UNIVERSALITY OF ANIMALCULAR LIFE. - LIVING ORGANISMS ARE UNIVERSALLY DIFFUSED OVER EVERY PART OF THE GLOBE. THE GENTLE ZEPHYR WAFTS FROM FLOWER TO FLOWER INVISIBLE, FRUCTIFYING ATOMS, WHICH QUICKEN BEAUTY AND FEGRANCE, GIVING THE PROMISE OF A GOLDEN FRUITAGE, TO GLADDEN AND NOURISH A DEPENDENT WORLD. NATURE'S OWN SWEET CUNNING INVESTS ALL INVING THINGS,

Verified label (manual GT)

2126 normalized characters

THE ORIGIN OF LIFE. 17

constraining into her service chemical affinities, arranging the elements and disposing them for her own benefit, in such numberless ways that we involuntarily exclaim,

"The course of Nature is the art of God."

The microscope reveals the fact that matter measuring only 1/120000 of an inch in diameter may be endowed with vitality, and that countless numbers of animalcules often inhabit a single drop of stagnant water. These monads do not vary in form, whether in motion or at rest. The life of one, even, is an inexplicable mystery to the philosopher. Ehrenberg writes: "Not only in the polar regions is there an uninterrupted development of active microscopic life, where larger animals cannot exist, but we find that those minute beings collected in the Antarctic expedition of Captain James Ross exhibit a remarkable abundance of unknown, and often most beautiful forms."

Even the interior of animal bodies is inhabited by animalcules. They have been found in the blood of the frog and the salmon, and in the optic fluid of fishes. Organic beings are found in the interior of the earth, into which the industry of the miner has made extensive excavations, sunk deep shafts, and thus revealed their forms; likewise, the smallest fossil organisms form subterranean strata many fathoms deep. Not only do lakes and inland seas abound with life, but also, from unknown depths, in volcanic districts, arise thermal springs which contain living insects. Were we endowed with a microscopic eye, we might see myriads of ethereal voyagers wafted by on every breeze, as we now behold drifting clouds of aqueous vapor. While the continents of earth furnishes evidences of the universality of organic beings, recent observations prove that "animal life predominates amid the eternal night of the depths of the liquid ocean."

THE ORIGIN OF LIFE.

The ancients, rude in many of their ideas, referred the origin of life to divine determination. The thought was crudely expressed, but well represented, in the following verse:

"Then God smites his hands together,
And strikes out a soul as a spark,
Into the organized glory of things,
From the deeps of the dark."

DeepSeek-OCR-2

CER 0.0296 | WER 0.0765 | F1 0.9478 | 1 region(s) | length 1.00x

text

constraining into her service chemical affinities, arranging the elements and disposing them for her own benefit, in such numberless ways that we involuntarily exclaim,

"The course of Nature is the art of God."

text

The microscope reveals the fact that matter measuring only $\frac{1}{120000}$ of an inch in diameter may be endowed with vitality, and that countless numbers of animalcules often inhabit a single drop of stagnant water. These monads do not vary in form, whether in motion or at rest. The life of one, even, is an inexplicable mystery to the philosopher. Ehrenberg writes: "Not only in the polar regions is there an uninterrupted development of active microscopic life, where larger animals cannot exist, but we find that those minute beings collected in the Antarctic expedition of Captain James Ross exhibit a remarkable abundance of unknown, and often most beautiful forms."

text

Even the interior of animal bodies is inhabited by animalcules. They have been found in the blood of the frog and the salmon, and in the optic fluid of fishes. Organic beings are found in the interior of the earth, into which the industry of the miner has made extensive excavations, sunk deep shafts, and thus revealed their forms; likewise, the smallest fossil organisms form subterranean strata many fathoms deep. Not only do lakes and inland seas abound with life, but also, from unknown depths, in volcanic districts, arise thermal springs which contain living insects. Were we endowed with a microscopic eye, we might see myriads of ethereal voyagers wafted by on every breeze, as we now behold drifting clouds of aqueous vapor. While the continents of earth furnishes evidences of the universality of organic beings, recent observations prove that "animal life predominates amid the eternal night of the depths of the liquid ocean."

sub_title

THE ORIGIN OF LIFE.

text

The ancients, rude in many of their ideas, referred the origin of life to divine determination. The thought was crudely expressed, but well represented, in the following verse:

text

"Then God smites his hands together,
And strikes out a soul as a spark,
Into the organized glory of things,
From the deeps of the dark."

GLM-OCR

CER 0.0183 | WER 0.0574 | F1 0.9612 | 1 region(s) | length 0.99x

constraining into her service chemical affinities, arranging the elements and disposing them for her own benefit, in such numberless ways that we involuntarily exclaim,

"The course of Nature is the art of God."

The microscope reveals the fact that matter measuring only $\frac{1}{120000}$ of an inch in diameter may be endowed with vitality, and that countless numbers of animalcules often inhabit a single drop of stagnant water. These monads do not vary in form, whether in motion or at rest. The life of one, even, is an inexplicable mystery to the philosopher. Ehrenberg writes: "Not only in the polar regions is there an uninterrupted development of active microscopic life, where larger animals cannot exist, but we find that those minute beings collected in the Antarctic expedition of Captain James Ross exhibit a remarkable abundance of unknown, and often most beautiful forms."

Even the interior of animal bodies is inhabited by animalcules. They have been found in the blood of the frog and the salmon, and in the optic fluid of fishes. Organic beings are found in the interior of the earth, into which the industry of the miner has made extensive excavations, sunk deep shafts, and thus revealed their forms; likewise, the smallest fossil organisms form subterranean strata many fathoms deep. Not only do lakes and inland seas abound with life, but also, from unknown depths, in volcanic districts, arise thermal springs which contain living insects. Were we endowed with a microscopic eye, we might see myriads of ethereal voyagers wafted by on every breeze, as we now behold drifting clouds of aqueous vapor. While the continents of earth furnishes evidences of the universality of organic beings, recent observations prove that "animal life predominates amid the eternal night of the depths of the liquid ocean."

THE ORIGIN OF LIFE.

The ancients, rude in many of their ideas, referred the origin of life to divine determination. The thought was crudely expressed, but well represented, in the following verse:

"Then God smites his hands together,
And strikes out a soul as a spark,
Into the organized glory of things,
From the deeps of the dark."<[user]>

MinerU 2.5 Pro

CER 0.0056 | WER 0.0410 | F1 0.9697 | 1 region(s) | length 1.00x

THE ORIGIN OF LIFE.

17

constraining into her service chemical affinities, arranging the elements and disposing them for her own benefit, in such numberless ways that we involuntarily exclaim,

"The course of Nature is the art of God."

The microscope reveals the fact that matter measuring only $\frac{1}{120000}$ of an inch in diameter may be endowed with vitality, and that countless numbers of animalcules often inhabit a single drop of stagnant water. These monads do not vary in form, whether in motion or at rest. The life of one, even, is an inexplicable mystery to the philosopher. Ehrenberg writes: "Not only in the polar regions is there an uninterrupted development of active microscopic life, where larger animals cannot exist, but we find that those minute beings collected in the Antarctic expedition of Captain James Ross exhibit a remarkable abundance of unknown, and often most beautiful forms."

Even the interior of animal bodies is inhabited by animalcules. They have been found in the blood of the frog and the salmon, and in the optic fluid of fishes. Organic beings are found in the interior of the earth, into which the industry of the miner has made extensive excavations, sunk deep shafts, and thus revealed their forms; likewise, the smallest fossil organisms form subterranean strata many fathoms deep. Not only do lakes and inland seas abound with life, but also, from unknown depths, in volcanic districts, arise thermal springs which contain living insects. Were we endowed with a microscopic eye, we might see myriads of ethereal voyagers wafted by on every breeze, as we now behold drifting clouds of aqueous vapor. While the continents of earth furnishes evidences of the universality of organic beings, recent observations prove that "animal life predominates amid the eternal night of the depths of the liquid ocean."

THE ORIGIN OF LIFE.

The ancients, rude in many of their ideas, referred the origin of life to divine determination. The thought was crudely expressed, but well represented, in the following verse:

"Then God smites his hands together, And strikes out a soul as a spark, Into the organized glory of things, From the deeps of the dark."

TrOCR large-printed

CER 0.0880 | WER 0.1913 | F1 0.8333 | 1 region(s) | length 1.03x

TOTAL THE ORIGIN OF LIFE.
CONSTRAINING INTO HER SERVICE CHEMICAL ATINITIES, ARRANGING THE Elements AND DISPOSING THEM FOR HER OWN BENEAT, IN SUCH NUM-
DEFLESS WAYS THAT WE INVOLUNTARILY EXCLAIM.
"ALL RETURNS AND THE COURSE OF NATURE IS THE ATT OF GOD. "
THE MICROSCOPE LEVELS THE FACT THAT MATTER MEASURING ONLY TRAHAS OF AN INCH IN DIAMETER MAY BE ENDOWED WITH VITALITY, AND THAT COMPLESS NUMBERS OF ANIMALOULES OFTEN INHABIT A SINGLE PROP OF STAGNANT WATER. THESE MONADS DO NOT VARY IN FORM, WHETHER IN MOTION OR AT TEST. THE LIFE OF ONE, EVEN. IS AN INEX-
PLICABLE MYSTERY TO THE PHILOSOPHER. EHRENBURG WRITES: "NOT ONLY IN THE POLAR REGIONS IS THERE AN UNINTERRUPTED DEVELOPMENT OF ACTIVE MICROSCOPIC LIFE, WHERE LATER ANIMALS CANNOT EXIST, BUT WE AND THAT THOSE MINUTE BEINGS COLLECTED IN THE ANTARETTIC EXPERITION OF CAPTAIN JAMES ROSS EXHIBIT A REMARKABLE ABUN-
DANCE OF MALMON, AND OFTEN MOST BEAUTIFUL FORMS."
EVEN THE INTERIOR OF ANIMAL BOOKIES IS INHABITED BY ANIMALCULES, THEY HAVE BEEN FOOD IN THE BLOOD OF THE FOG AND THE SALMON, AND IN THE OPTIC FULL OF FISHES. ORGANIC BEINGS ARE FOUND IN THE INTRIOR OF THE EARTH, INTO WHICH THE INDUSTRY OF THE MINER HAS MADE EXTENSIVE EXEAVATIONS, SUNK DEEP SHAFTS, AND THUS REVEALED THEIR FORMS; LIKEWISE, THE SMALLEST FOSSIL ORGANISMS FORM SUBTER-
RANEAN STATE MANY FATHOMS DEEP. NOT ONLY DO LAKES AND INLAND SEAS ABOARD WITH LIFE, BUT ALSO, FROM UNKNOWN DEPTAS, IN VOL-
CAMIC DISTRIETS, ARISE THERMAL SPINGS WHICH CONTAIN LIVING INSEETS. WERE WE ENLOWE WITH A MICROSCOPIC EYE, WE MIGHT SEE MYRIADS OF ETHEREAL VOYAGERS WAFTERL BY ON EVERY BREERE, AS WE NOW BE-
HOLD DRITING CLOUDS OF AQUEOUS VAPOR. WHILE THE CONTINTS OF CARTH PURNISLES EVIDENCES OF THE UNIVERSALITY OF ORGANIC BEINGS, RECENT OBSERVATIONS PROVE THAT "ANIMAL LIFE PREDOMINATES AMID THE RETERAL NIGHT OF THE DEPTAS OF THE EQUIL OCEAN."
THE TIME ORIGIN OF LIFE.
THE ANCIENTS, RADE IN MANY OF THEIR IDEAS, REFERRED THE ORIGIN OF LIFE TO DIVINE DETERMINATION. THE THOUGH WAS ERADELY EX-
PRESSED, BUT WELL REPRESENTED, IN THE FOLLOWING VEGE:
TOTAL RETURN GOLDEN GOLDES HIS HANDS TOGETHER.
CASHIER AND STRIKES OUT A SOLD AS A SHARK, PRICE OF RECEIPT.
DISPLAY FACE INTO THE ORGANIZED GLOTS OF THINGS,
PURCHASE AND FROM THE DEEPS OF THE DARK. "

Verified label (manual GT)

2126 normalized characters

17

THE ORIGIN OF LIFE.

constraining into her service chemical affinities, arranging the elements and disposing them for her own benefit, in such numberless ways that we involuntarily exclaim,

"The course of Nature is the art of God."

The microscope reveals the fact that matter measuring only 1/120000 of an inch in diameter may be endowed with vitality, and that countless numbers of animalcules often inhabit a single drop of stagnant water. These monads do not vary in form, whether in motion or at rest. The life of one, even, is an inexplicable mystery to the philosopher. Ehrenberg writes: "Not only in the polar regions is there an uninterrupted development of active microscopic life, where larger animals cannot exist, but we find that those minute beings collected in the Antarctic expedition of Captain James Ross exhibit a remarkable abundance of unknown, and often most beautiful forms."

Even the interior of animal bodies is inhabited by animalcules. They have been found in the blood of the frog and the salmon, and in the optic fluid of fishes. Organic beings are found in the interior of the earth, into which the industry of the miner has made extensive excavations, sunk deep shafts, and thus revealed their forms; likewise, the smallest fossil organisms form subterranean strata many fathoms deep. Not only do lakes and inland seas abound with life, but also, from unknown depths, in volcanic districts, arise thermal springs which contain living insects. Were we endowed with a microscopic eye, we might see myriads of ethereal voyagers wafted by on every breeze, as we now behold drifting clouds of aqueous vapor. While the continents of earth furnish evidences of the universality of organic beings, recent observations prove that "animal life predominates amid the eternal night of the depths of the liquid ocean."

THE ORIGIN OF LIFE.

The ancients, rude in many of their ideas, referred the origin of life to divine determination. The thought was crudely expressed, but well represented, in the following verse:

"Then God smites his hands together,
And strikes out a soul as a spark,
Into the organized glory of things,
From the depths of the dark."

DeepSeek-OCR-2

CER 0.0301 | WER 0.0546 | F1 0.9595 | 9 region(s) | length 1.01x

17

THE CRISIS OF THE RACE OF HUMANITY

constraining into her service chemical antifoules, arranging the elements and disposing them for her own benefit, in such numberless ways that we involuntarily exclaim,

The course of Nature in the art of God

The microscope reveals the fact that matter measuring only $\frac{1}{120000}$ of an inch in diameter may be endowed with vitality, but that countless numbers of animalcules often inhabit a single drop of stagnant water. These monads do not vary in form, whether in motion or at rest. The life of one, even, is an inexplicable mystery to the philosopher. Ehrenberg writes: "Not only in the polar regions is there an uninterrupted development of active microscopic life, where larger animals cannot exist, but we find that those minute beings collected in the Antarctic expedition of Captain James Ross exhibit a remarkable abundance of unknown, and often most beautiful forms."

Even the interior of animal bodies is inhabited by animalcules. They have been found in the blood of the frog and the salmon, and in the optic fluid of fishes. Organic beings are found in the interior of the earth, into which the industry of the miner has made extensive excavations, sunk deep shafts, and thus revealed their forms; likewise, the smallest fossil organisms form subterranean strata many fathoms deep. Not only do lakes and inland seas abound with life, but also, from unknown depths, in volcanic districts, arise thermal springs which contain living insects. Were we endowed with a microscopic eye, we might see myriads of ethereal voyagers wafted by on every breeze, as we now behold drifting clouds of aqueous vapor. While the continents of earth furnish evidences of the universality of organic beings, recent observations prove that "animal life predominates amid the eternal night of the depths of the liquid ocean."

THE ORIGIN OF THE ALBUM

The ancients, made in many of their ideas, referred the origin of life to divine determination. The thought was under expressed, but well represented, in the following verse:

"Then God smites his hands together,
And strikes out a soul as a spark,
Into the organized glory of things,
From the depths of the dark."

GLM-OCR

CER 0.0278 | WER 0.0683 | F1 0.9578 | 9 region(s) | length 1.02x

17

THE ORIGIN OF LIFE.

constraining into her service chemical affinities, arranging the elements and disposing them for her own benefit, in such numberless ways that we involuntarily exclaim,

"The course of Nature is the art of God."

The microscope reveals the fact that matter measuring only $\frac{1}{120000}$ of an inch in diameter may be endowed with vitality, and that countless numbers of animalcules often inhabit a single drop of stagnant water. These monads do not vary in form, whether in motion or at rest. The life of one, even, is an inexplicable mystery to the philosopher. Ehrenberg writes: "Not only in the polar regions is there an uninterrupted development of active microscopic life, where larger animals cannot exist, but we find that those minute beings collected in the Antarctic expedition of Captain James Ross exhibit a remarkable abundance of unknown, and often most beautiful forms."

Even the interior of animal bodies is inhabited by animalcules. They have been found in the blood of the frog and the salmon, and in the optic fluid of fishes. Organic beings are found in the interior of the earth, into which the industry of the miner has made extensive excavations, sunk deep shafts, and thus revealed their forms; likewise, the smallest fossil organisms form subterranean strata many fathoms deep. Not only do lakes and inland seas abound with life, but also, from unknown depths, in volcanic districts, arise thermal springs which contain living insects. Were we endowed with a microscopic eye, we might see myriads of ethereal voyagers wafted by on every breeze, as we now behold drifting clouds of aqueous vapor. While the continents of earth furnishes evidences of the universality of organic beings, recent observations prove that "animal life predominates amid the eternal night of the depths of the liquid ocean."

THE ORIGIN OF LIFE.

The ancients, rude in many of their ideas, referred the origin of life to divine determination. The thought was crudely expressed, but well represented, in the following verse:

"Then God smites his hands together, And strikes out a soul as a spark, Into the organized glory of things, From the depths of the dark."

MinerU 2.5 Pro

CER 0.0085 | WER 0.0464 | F1 0.9697 | 9 region(s) | length 1.00x

17

THE ORIGIN OF LIFE.

constraining into her service chemical affinities, arranging the elements and disposing them for her own benefit, in such numberless ways that we involuntarily exclaim,

"The course of Nature is the art of God."

The microscope reveals the fact that matter measuring only $\frac{1}{120000}$ of an inch in diameter may be endowed with vitality, and that countless numbers of animalcules often inhabit a single drop of stagnant water. These monads do not vary in form, whether in motion or at rest. The life of one, even, is an inexplicable mystery to the philosopher. Ehrenberg writes: "Not only in the polar regions is there an uninterrupted development of active microscopic life, where larger animals cannot exist, but we find that those minute beings collected in the Antarctic expedition of Captain James Ross exhibit a remarkable abundance of unknown, and often most beautiful forms."

Even the interior of animal bodies is inhabited by animalcules. They have been found in the blood of the frog and the salmon, and in the optic fluid of fishes. Organic beings are found in the interior of the earth, into which the industry of the miner has made extensive excavations, sunk deep shafts, and thus revealed their forms; likewise, the smallest fossil organisms form subterranean strata many fathoms deep. Not only do lakes and inland seas abound with life, but also, from unknown depths, in volcanic districts, arise thermal springs which contain living insects. Were we endowed with a microscopic eye, we might see myriads of ethereal voyagers wafted by on every breeze, as we now behold drifting clouds of aqueous vapor. While the continents of earth furnishes evidences of the universality of organic beings, recent observations prove that "animal life predominates amid the eternal night of the depths of the liquid ocean."

THE ORIGIN OF LIFE.

The ancients, rude in many of their ideas, referred the origin of life to divine determination. The thought was crudely expressed, but well represented, in the following verse:

"Then God smites his hands together,
And strikes out a soul as a spark,
Into the organized glory of things,
From the depths of the dark."

TrOCR large-printed

CER 0.0193 | WER 0.0683 | F1 0.9384 | 9 region(s) | length 1.00x

17

THE ORIGIN OF LIFE,
CONSTRAINING INTO HER SERVICE CHEMICAL ATINITIES, ARRANGING THE ELEMENTS AND DISPOSING THEM FOR HER OWN BENEFIT, IN SUCH NUMBERLESS WAYS THAT WE INVOLUNTARILY EXCLAIM,
"THE COURSE OF NATURE IS THE ART OF GOD."
THE MICROSCOPE LEVELS THE FACT THAT MATTER MEASURING ONLY 1/120000 OF AN INCH IN DIAMETER MAY BE ENDOWED WITH VITALITY, AND THAT COUNTLESS NUMBERS OF ANIMALCULES OFTEN INHABIT A SINGLE DROP OF STAGNANT WATER. THESE MONADS DO NOT VARY IN FORM, WHETHER IN MOTION OR AT REST. THE LIFE OF ONE, EVEN, IS AN INEXPLICABLE MYSTERY TO THE PHILOSOPHER. EHRENBURG WRITES: "NOT ONLY IN THE POLAR REGIONS IS THERE AN UNINTERRUPTED DEVELOPMENT OF ACTIVE MICROSCOPIC LIFE, WHERE LARGER ANIMALS CANNOT EXIST, BUT WE AND THAT THOSE MINUTE BEINGS COLLECTED IN THE ANTARCTIC EXPERITION OF CAPTAIN JAMES ROSS EXHIBIT A REMARKABLE ABUNDANCE OF UNKNOWN, AND OFTEN MOST BEAUTIFUL FORMS."
EVEN THE INTERIOR OF ANIMAL BODIES IS INHABITED BY ANIMALCULES. THEY HAVE BEEN FOUND IN THE BLOOD OF THE FROG AND THE SALMON, AND IN THE OPTIC FULL OF FISHES. ORGANIC BEINGS ARE FOUND IN THE INTERIOR OF THE EARTH, INTO WHICH THE INDUSTRY OF THE MINER HAS MADE EXTENSIVE EXCAVATIONS, SUNK DEEP SHAFTS, AND THUS REVEALED THEIR FORMS; LIKEWISE, THE SMALLEST FOSSIL ORGANISMS FORM SUBTERRANEAN STRATA MANY FATHOMS DEEP. NOT ONLY DO LAKES AND INLAND SEAS ABOUND WITH LIFE, BUT ALSO, FROM UNKNOWN DEPTHS, IN VOLCANIC DISTRICTS, ARISE THERMAL SPINGS WHICH CONTAIN LIVING INSECTS. WERE WE ENDOWED WITH A MICROSCOPIC EYE, WE MIGHT SEE MYRIADS OF ETHEREAL VOYAGERS WAFTED BY ON EVERY BREEZE, AS WE NOW BEHOLD DRIFTING CLOUDS OF AQUEOUS VAPOR. WHILE THE CONTINENTS OF EARTH FUNNISHES EVIDENCES OF THE UNIVERSALITY OF ORGANIC BEINGS, RECENT OBSERVATIONS PROVE THAT "ANIMAL LIFE PREDOMINATES AMID THE RETINAL NIGHT OF THE DEPTHS OF THE LIQUID OCEAN."
THE ORIGIN OF LIFE.
THE ANCIENTS, RUDE IN MANY OF THEIR IDEAS, REFERRED THE ORIGIN OF LIFE TO DIVINE DETERMINATION. THE THOUGH WAS CRUDELY EXPRESSED, BUT WELL REPRESENTED, IN THE FOLLOWING VERSE:
"THEN GOD SMITES HIS HANDS TOGETHER,
AND STRIKES OUT A SOULD AS A SPARK,
INTO THE ORGANIZED GLORY OF THINGS,
FROM THE DEEPS OF THE DARK."

Verified label (manual GT)

1981 normalized characters

18 COMMON SENSE MEDICAL ADVISER.

According to a Greek myth, Prometheus formed a human image from the dust of the ground, and then, by fire stolen from heaven, animated it with a living soul. Spontaneous generation once held its sway, and now the idea of natural evolution is popular. Some believe that the impenetrable mystery of life is evolved from the endowments of nature, and build their imperfect theory on observations of her concrete forms and their manifestations, to which all our investigations are restricted. But every function indicates purpose, every organism evinces intelligent design, and all proclaim a Divine Power. Something cannot come out of nothing. With reason and philosophy, chance is an impossibility. We, therefore, accept the display of wisdom in nature as indicative of the designs of God. Thus "has He written His claims for our profoundest admiration and homage all over every object that He has made." If you ask: Is there any advantage in considering the phenomena of nature as the result of DIVINE VOLITION? we answer, that this belief corresponds with the universally acknowledged ideas of accountability; for, with a wise and efficient Cause, we infer there is an intelligent creation, and the desire to communicate, guide and bless, is responded to by man, who loves, obeys, and enjoys. Nothing is gained by attributing to nature vicegerent forces. Is it not preferable to say that she responds to intelligent, loving Omnipotence? Our finiteness is illustrated by our initiation into organized being. Emerging from a rayless atom, too diminutive for the sight, we gradually develop and advance to the maturity of those conscious powers, the exercise of which furnishes indubitable evidence of our immortality. We are pervaded with invisible influences, which, like the needle of the compass trembling on its pivot, point us to immortality as our ultimate goal, where in the sunny clime of Love, even in a spiritual realm of joy and happiness, we may eternally reign with Him who is all in all.

DeepSeek-OCR-2

CER 0.0156 | WER 0.0181 | F1 0.9878 | 1 region(s) | length 0.99x

text
According to a Greek myth, Prometheus formed a human image from the dust of the ground, and then, by fire stolen from heaven, animated it with a living soul. Spontaneous generation once held its sway, and now the idea of natural evolution is popular. Some believe that the impenetrable mystery of life is evolved from the endowments of nature, and build their imperfect theory on observations of her concrete forms and their manifestations, to which all our investigations are restricted. But every function indicates purpose, every organism evinces intelligent design, and all proclaim a Divine Power. Something cannot come out of nothing. With reason and philosophy, chance is an impossibility. We, therefore, accept the display of wisdom in nature as indicative of the designs of God. Thus "has He written His claims for our profoundest admiration and homage all over every object that He has made." If you ask: Is there any advantage in considering the phenomena of nature as the result of Divine Volition? we answer, that this belief corresponds with the universally acknowledged ideas of accountability; for, with a wise and efficient Cause, we infer there is an intelligent creation, and the desire to communicate, guide and bless, is responded to by man, who loves, obeys, and enjoys. Nothing is gained by attributing to nature vicegerent forces. Is it not preferable to say that she responds to intelligent, loving Omnipotence? Our finiteness is illustrated by our initiation into organized being. Emerging from a rayless atom, too diminutive for the sight, we gradually develop and advance to the maturity of those conscious powers, the exercise of which furnishes indubitable evidence of our immortality. We are pervaded with invisible influences, which, like the needle of the compass trembling on its pivot, point us to immortality as our ultimate goal, where in the sunny clime of Love, even in a spiritual realm of joy and happiness, we may eternally reign with Him who is all in all.

GLM-OCR

CER 0.0187 | WER 0.0181 | F1 0.9909 | 1 region(s) | length 0.99x

According to a Greek myth, Prometheus formed a human image from the dust of the ground, and then, by fire stolen from heaven, animated it with a living soul. Spontaneous generation once held its sway, and now the idea of natural evolution is popular. Some believe that the impenetrable mystery of life is evolved from the endowments of nature, and build their imperfect theory on observations of her concrete forms and their manifestations, to which all our investigations are restricted. But every function indicates purpose, every organism evinces intelligent design, and all proclaim a Divine Power. Something cannot come out of nothing. With reason and philosophy, chance is an impossibility. We, therefore, accept the display of wisdom in nature as indicative of the designs of God. Thus "has He written His claims for our profoundest admiration and homage all over every object that He has made." If you ask: Is there any advantage in considering the phenomena of nature as the result of Divine Volition? we answer, that this belief corresponds with the universally acknowledged ideas of accountability; for, with a wise and efficient Cause, we infer there is an intelligent creation, and the desire to communicate, guide and bless, is responded to by man, who loves, obeys, and enjoys. Nothing is gained by attributing to nature vicegerent forces. Is it not preferable to say that she responds to intelligent, loving Omnipotence? Our finiteness is illustrated by our initiation into organized being. Emerging from a rayless atom, too diminutive for the sight, we gradually develop and advance to the maturity of those conscious powers, the exercise of which furnishes indubitable evidence of our immortality. We are pervaded with invisible influences, which, like the needle of the compass trembling on its pivot, point us to immortality as our ultimate goal, where in the sunny clime of Love, even in a spiritual realm of joy and happiness, we may eternally reign with Him who is all in all.<[user]>

MinerU 2.5 Pro

CER 0.0000 | WER 0.0000 | F1 1.0000 | 1 region(s) | length 1.00x

18
COMMON SENSE MEDICAL ADVISER.
According to a Greek myth, Prometheus formed a human image from the dust of the ground, and then, by fire stolen from heaven, animated it with a living soul. Spontaneous generation once held its sway, and now the idea of natural evolution is popular. Some believe that the impenetrable mystery of life is evolved from the endowments of nature, and build their imperfect theory on observations of her concrete forms and their manifestations, to which all our investigations are restricted. But every function indicates purpose, every organism evinces intelligent design, and all proclaim a Divine Power. Something cannot come out of nothing. With reason and philosophy, chance is an impossibility. We, therefore, accept the display of wisdom in nature as indicative of the designs of God. Thus "has He written His claims for our profoundest admiration and homage all over every object that He has made." If you ask: Is there any advantage in considering the phenomena of nature as the result of Divine Volition? we answer, that this belief corresponds with the universally acknowledged ideas of accountability; for, with a wise and efficient Cause, we infer there is an intelligent creation, and the desire to communicate, guide and bless, is responded to by man, who loves, obeys, and enjoys. Nothing is gained by attributing to nature vicegerent forces. Is it not preferable to say that she responds to intelligent, loving Omnipotence? Our finiteness is illustrated by our initiation into organized being. Emerging from a rayless atom, too diminutive for the sight, we gradually develop and advance to the maturity of those conscious powers, the exercise of which furnishes indubitable evidence of our immortality. We are pervaded with invisible influences, which, like the needle of the compass trembling on its pivot, point us to immortality as our ultimate goal, where in the sunny clime of Love, even in a spiritual realm of joy and happiness, we may eternally reign with Him who is all in all.

TrOCR large-printed

CER 0.0374 | WER 0.1239 | F1 0.8795 | 1 region(s) | length 1.00x

18 COMMON SENSE MEDICAL ADVISER.
ACCORDING TO A GREEK MYTH, PROMETHEUS FORMED A HUMAN IMAGE FROM THE DUST OF THE GROUND, AND THEN, BY FIRE STOLEN FROM HEAVEN, AMINATED IT WITH A LIVING SOUL. SPONTANEOUS GENERATION ONCE HELL ITS SWAY, AND NOW THE IDEA OF NATURAL EVOLUTION IS POPULAR. SOME BELIEVE THAT THE IMPE-ETTABLE MYSTERY OF LIFE IS EVOLVED FROM THE ENDOWMENTS OF NATURE, AND BOLD THEIR INDEPECT THEORY ON OBSERVATIONS OF HER CONCRETE FORMS AND THEIR MANIFESTATIONS. TO WHICH ALL OUR INVESTIGATIONS ARE RESTRICTED. BUT EVERY FUME-TION INDICATES PURPOSE, EVERY ORGANISM EVINCES INTELLIGENT DESIGN, AND ADI PROCLAIM A DIVINE POWER. SOMETHING CANNOT COME OUT OF NOTHING. WITH REASON AND PHILOSOPHY, CHANGE IS AN IMPOSSI-BILITY. WE, THEREFORE, ACCEPT THE DISPLAY OF WISDOM IN DATURE AS INCLUSIVE OF THE DESIGNS OF GOD. THUS " HAS HE WRITTEN HIS PLAIMS FOR OUR PROFOUNDEST ADMIRATION AND HOMAGE ALL OVER EVERY OBJECT THAT HE HAS MADE." IF YOU ASK: IS THERE ANY ADVANTAGE IN CONSIDERING THE PEROMERA OF NATURE AS THE RESULT OF DIVINE VOLTON? WE ANSWER, THAT THIS BELIEF CORRESPONDS WITH THE UNIVERSALLY ACKNOWLEDGET IDEAS OF ACCOUNTABILITY; FOR, WITH A WISE AND RECEIPT CANSE, WE INFER THERE IS AN INTELLIGENT CREATION. AML THE DESIRE TO COMMUNICATE, GUIDE AND BLESS, IS ESPONded TO BY MAN, WHO LOVE, OBEYS, AND ENJOYS. NOTHING IS GAINED BY ATTIBATING TO MATURE VICEGERENT FORCES. IS IT NOT PREFERABLE TO SAY THAT SHE RESPONDS TO INTERLIGENT, LOVING OMMIPOTENCE? OUR INITENESS IS ILLUSTRATED BY OUR INITIATION INTO ORGANIZED BEING. EMERGING FROM A RAYLESS ATOM, TOO DIMINUTIVE FOR THE SIGHT, WE GRAHALLY DEVELOP AND ADVANCE TO THE MATURITY OF THOSE CONSCIOUS POTENTS, THE EXCREASE OF WHICH FURNISHES INDUBITABLE EVIDENCE OF OUR INHORTALITY. WE ARE PERVATED WITH INVISIBLE INFNUENCES. WHICH, LIKE THE NEEDLE OF THE COMPASS ITEMBLING ON ITS PIVOT, POINT US TO IMMORTALITY AS OUR UTIMATE GOAL, WHERE IN THE SUMMY ELIME OF LOVE, EVEN IN A SPIRITUAL REALM OF JOY AND HAPPINESS, WE MAY RETURNALLY REIGN WITH HIM WHO IS ALL IN ALL.

p0026 - PP-DocLayoutV3 region OCR

Side-by-side saved text. Metrics use identical normalized scoring; displayed text preserves OCR wording.

Verified label (manual GT)

1981 normalized characters

18 COMMON SENSE MEDICAL ADVISER.

According to a Greek myth, Prometheus stole a human image from the dust of the ground, and then, by fire stolen from heaven, animated it with a living soul. Spontaneous generation once held its sway, and now the idea of natural evolution is popular. Some believe that the impenetrable mystery of life is evolved from inorganic matter, and that the laws of evolution form the basis of all existence. But the laws of evolution are not laws of necessity, and the laws of necessity are restricted. But every function indicates purpose, every organism evinces intelligent design, and all proclaim a Divine Power. Something cannot come out of nothing. With reason and philosophy, chance is an impossibility. We, therefore, accept the display of wisdom in nature as indicative of the designs of God. Thus "has He written His claims for our profoundest admiration and homage all over every object that He has made." If you ask: Is there any advantage in the existence of the human mind? I answer: Yes, in that it is the only creature that can be made accountable for its actions. It is the only creature that can be made accountable, for, with a wise and efficient Cause, we infer there is an intelligent creation, and the desire to communicate, guide and bless, is answered in man, who loves, obeys, and enjoys. Nothing is gained by attributing to nature vicergerent forces. Is it not preferable to say that she responds to intelligent, loving Omnipotence? Our finiteness is illustrated by our initiation into organized being. Emerging from a rayless atom, too diminutive for the sight, we gradually become conscious of our existence, and are made aware of the existence of immortality. We are pervaded by the presence of the divine, and are made conscious of the invisible influences, which, like the needle of the compass trembling on its pivot, point us into morality as our ultimate goal, where, in the sunny climate of Love, even in a spiritual realm of joy and happiness, we may eternally reign with Him who is all in all.

DeepSeek-OCR-2

ER 1.1580 | WER 1.7311 | F1 0.4460 | 3 region(s) | length 2.00x

8

[illegible]

conform to a Greek myth, Prometheus then offered to
 on the dust of the ground, and then, by fire stolen from heaven,
 animated it with a living soul. Spontaneous generation could
 explain its origin, but its role in the evolution of life is popular
 come believe that the impenetrable mystery of life is explained
 on the endowments of nature, and build their imperfect theory
 on observations of her concrete forms and their manifestations,
 which all our investigations are restricted. But every func-
 tion of the mind is the result of the laws of the divine design,
 and all proclaim a Divine Power. Something cannot come out
 of nothing. With reason and philosophy, "chance" is an impossi-
 bility. We therefore, accept the display of wisdom in nature
 as indicative of the designs of God. Thus "has He written His
 name on the foundations of the earth," and "He has written every
 object that He has made." If you ask: Is there any advantage
 considering the phenomena of nature as the result of Divine
 will? we answer, that this belief corresponds with the
 universally acknowledged ideas of accountability; for, with a
 wise and efficient "Cause," we infer there is an intelligent crea-
 tor, and the desire to communicate, guide and bless, is responded to
 by man, who loves, obeys, and enjoys. Nothing is gained by
 attributing to nature the imperfections of God. Thus "has He written
 His name on the foundations of the earth," and "He has written every
 object that He has made." If you ask: Is there any advantage
 considering the phenomena of nature as the result of Divine
 will? we answer, that this belief corresponds with the
 universally acknowledged ideas of accountability; for, with a
 wise and efficient "Cause," we infer there is an intelligent crea-
 tor, and the desire to communicate, guide and bless, is responded to
 by man, who loves, obeys, and enjoys. Nothing is gained by
 attributing to nature the imperfections of God. Thus "has He written
 His name on the foundations of the earth," and "He has written every
 object that He has made." If you ask: Is there any advantage

GLM-OCR

CER 0.0076 | WER 0.0091 | F1 0.9955 | 3 region(s) | length 1.01x

18<|user|>

COMMON SENSE MEDICAL ADVISER.<|user|>

According to a Greek myth, Prometheus formed a human figure from the dust of the ground, and then, by fire stolen from heaven, animated it with a living soul. Spontaneous generation once held its sway, and now the idea of natural evolution is popular. Some believe that the impenetrable mystery of life is evolved from a lifeless matter, and that the laws of evolution are the same as those of the inorganic world. But the laws of organic evolution are not restricted. But every function indicates purpose, every organism evinces intelligent design, and all proclaim a Divine Power. Something cannot come out of nothing. With reason and philosophy, chance is an impossibility. We, therefore, accept the display of wisdom in nature as indicative of the designs of God. Thus we have written His claims for our profoundest admiration and homage all over every object that He has made." If you ask: Is there any advantage in considering the presence of nature as a sign of God? I answer: It is a sign of God, and that belief comes universally acknowledged as the basis of all stability with a wisdom efficient Cause. We infer there is an intelligent Creator, and the desire to communicate is the first response to belief. Man, who loves, obeys, and enjoys. Nothing is gained by attributing to nature vicegerent forces. Is it not preferable to say that she responds to intelligent, loving Omnipotence? Our finiteness is illustrated by our initiation into organized being. Emerging from a rayless atom, too diminutive for the sight, we gradually develop and advance to the maturity of those conscious powers, the exercise of which furnishes indubitable evidence of our immortality. We are pervaded with the sense of our own immortality, and we are conscious of our ultimate goal, where in the sunny clime of Love, we revel in a spiritual realm of joy and happiness, we may eternally reign with Him who is all in all. <user>

MinerU 2.5 Pro

CER 0.0015 | WER 0.0030 | F1 0.9985 | 3 region(s) | length 1.00

COMMON SENSE MEDICAL ADVISER.

According to a Greek myth, Prometheus held a human image from the dust of the ground, and then, by fire stolen from heaven, animated it with a living soul. Spontaneous generation once held its sway, and now the idea of natural evolution is popular. Some believe that the impenetrable mystery of life is evolved from a lifeless matter, and that the origin of the human race is a mere accident. But the investigation of the origin of life is a subject on which the investigation of the origin of the human race is restricted. But every function indicates purpose, every organism evinces intelligent design, and all proclaim a Divine Power. Something cannot come out of nothing. With reason and philosophy, chance is an impossibility. We, therefore, accept the display of wisdom in nature as indicative of the designs of God. Thus we have written His claims for our profoundest admiration and homage all over every object that He has made.¹ If you ask: Is there any advantage in the existence of the human race? We answer: The chief comfort and satisfaction of our life is in the knowledge of the existence of our Creator, and of His providence, and of His love, with a wise and efficient Cause, we infer there is an intelligent creation, and the desire to communicate, good and blessing, is responded to by man, who loves, obeys, and enjoys. Nothing is gained by attributing to nature vicegerent forces. Is it not preferable to say that she responds to intelligent, loving Omnipotence? Our finiteness is illustrated by our initiation into organized being. Emerging from a rayless state, too diminutive for the sight, we gradually expand into a more perfect form, and are furnished with faculties and functions, and finally, with the immortal soul, which is the source of our immortality. We are persons with invisible influences, which, like the needle of the compass trembling on its pivot, point, not to immortality as our ultimate goal, but in the sunny circle of Love, even in a spiritual realm of joy and happiness, we may eternally reign with Him who is all in all.

TrOCR large-printed

CER 0.0202 | WER 0.0846 | F1 0.9187 | 3 region(s) | length 1.00x

18

COMMON SENSE MEDICAL ADVISER.

ACCORDING TO A GREEK MYTH, PROMETHEUS FORMED A HUMAN IMAGE FROM CLAY, BUT WAS PUNISHED FOR HIS RECKLESSNESS BY BEING STOLEN FROM HEAVEN AND IMMATED WITH A LIVING SOUL. SPONTANEOUS GENERATION OCCURRED, AND HE HAD HIS SWEAT, AND NOW THE IDEA OF NATURAL EVOLUTION IS POPULAR.

SOME BELIEVE THAT THE IMPENETRABLE MYSTERY OF LIFE IS EVOLVED FROM THE ENDOWMENTS OF NATURE, AND BOULD THEIR IMPERFECT THOUGHTS BE ENLIGHTENED BY THE REVELATIONS OF THE DIVINE MANIFESTATIONS, TO WHICH ALL OUR INVESTIGATIONS ARE RESTRICTED, BUT EVERY FUNCTION INDICATES PURPOSE, EVERY ORGANISM EVINCES INTELLIGENT DESIGN AND ALL PROCLAIM A DIVINE POWER, SOMETHING CANNOT COME OUT OF NOTHING, WITH INTELLIGENCE AND INTELLIGENCE, CHANGING AND IMPROVING, WE THEREFORE ACKNOWLEDGE THE DISPERSED WISDOM IN NATURE AS INDICATIVE OF THE DESIGNS OF GOD. THUS "H" HAS WRITTEN HIS CLAIMS FOR OUR PROFOUNDEST ADMIRATION AND HOMAGE ALL OVER EVERY OBJECT THAT HE HAS MADE." IF YOU ASK: IS THERE ANY ADVANTAGE IN THE UNIVERSE BEING UNINTELLIGENT, UNORGANIZED, AND UNWISDOM? WE ANSWER, THAT THIS BELIEF CORRESPONDS WITH THE UNIVERSALLY ACKNOWLEDGED IDEAS OF ACCOUNTABILITY, FOR, WITH A WISDOM AND EFFICIENT CAUSE, WE INFER THERE IS AN INTELLIGENT CREATOR. AND THE DESIRE TO COMMUNICATE, GUIDE AND BLESS, IS EXPONDED TO US BY THE LOVE OF GOD, AND HIS WISDOM, AND HIS POWER, AND HIS ATTRIBUTING TO MATURE VIGILANT FORTUNE. IS IT NOT PREFERABLE TO SAY THAT HE RESPONDS TO INTELLIGENT, LOVING OMNIPOTENCY? OUR "INTENESS" IS ILLUSTRATED BY OUR INITIATION INTO ORGANIZED BEING, EMERGING FROM A RAYLESS VOID, AND DIMINUTIVE FOR THE SIGHT, WE HAVE BEEN GIVEN A WAY TO KNOW THE TRUTH, AND THE WISDOM OF GOD, THE EXERCISE OF WHICH PURNISHES INDUBITABLE EVIDENCE OF OUR IMMORTALITY. WE ARE PERVADED WITH INVINCIBLE INFLUENCES WHICH, LIKE THE NEEDLE OF THE COMPASS TREMBLING ON ITS PIVOT, AND TO THE IMPASSIONED HEART, THE HEAVENLY MUSIC OF THE SPIRIT OF LOVE, EVEN IN A SPIRITUAL REALM OF JOY AND HAPPINESS, WE MAY RETURNARILY REIGN WITH HIM WHO IS ALL IN ALL.

Verified label (manual GT)

1382 normalized characters

CHAPTER II.

PHYSIOLOGICAL ANATOMY.

THE BONES.

All living bodies are made up of tissues. There is no part, no organ, however soft and yielding, or hard and resisting, which has not this peculiarity of structure. The bones of animals, as well as their flesh and fat, are composed of tissues, and all alike made up of cells. When viewed under a microscope, each cell is seen to consist of three distinct parts, a nucleolus, or dark spot, in the center of the cell, around which lies a mass of granules, called the nucleus; and this, in turn, is surrounded with a delicate, transparent membrane, termed the envelope. Each of the granules composing the nucleus assimilates nourishment, thereby growing into an independent cell, which possesses a triple organization similar to that of its parent, and in like manner reproduces other cells.

Fig. 4.

Nucleated cell. From Goeber. 1. Periphery of the cell, or cell-wall. 2. Nucleus. 3. Nucleolus in the center.

A variety of tissues enters into the composition of an animal structure, yet their differences are not always distinctly marked, since the characteristics of some are not unlike those of others. We shall notice, however, only the more important of the tissues.

The Areolar, or Connective Tissue, is a complete network of delicate fibers, spread over the body, and serves to bind the various organs and parts together. The fibrous and serous tissues are modifications of the areolar.

DeepSeek-OCR-2

CER 0.1910 | WER 0.2176 | F1 0.9775 | 1 region(s) | length 1.04x

title

CHAPTER II.

title

PHYSIOLOGICAL ANATOMY.

sub_title

THE BONES.

text

All living bodies are made up of tissues. There is no part, no organ, however soft and yielding, or hard and resisting, which has not this peculiarity of structure. The bones of animals, as well as their flesh and fat, are composed of tissues, and all alike made up of cells. When viewed under a microscope, each cell is seen to consist of three distinct parts, a nucleolus, or dark spot, in the center of the cell, around which lies a mass of granules, called the nucleus; and this, in turn, is surrounded with a delicate, transparent membrane, termed the envelope. Each of the granules composing the nucleus assimilates nourishment, thereby growing into an independent cell, which possesses a triple organization similar to that of its parent, and in like manner reproduces other cells.

text

A variety of tissues enters into the composition of an animal structure, yet their differences are not always distinctly marked, since the characteristics of some are not unlike those of others. We shall notice, however, only the more important of the tissues.

text

The Areolar, or Connective Tissue, is a complete network of delicate fibers, spread over the body, and serves to bind the various organs and parts together. The fibrous and serous tissues are modifications of the areolar.

figure_title

Fig. 4.

image

text

Nucleated cell. From Goeber. 1. Periphery of the cell, or cell-wall. 2. Nucleus. 3. Nucleolus in the center.

GLM-OCR

CER 0.0803 | WER 0.0921 | F1 0.9520 | 1 region(s) | length 0.93x

CHAPTER II.

PHYSIOLOGICAL ANATOMY.

THE BONES.

All living bodies are made up of tissues. There is no part, no organ, however soft and yielding, or hard and resisting, which has not this peculiarity of structure. The bones of animals, as well as their flesh and fat, are composed of tissues, and all alike made up of cells. When viewed under a microscope, each cell is seen to consist of three distinct parts, a nucleolus, or dark spot, in the center of the cell, around which lies a mass of granules, called the nucleus; and this, in turn, is surrounded with a delicate, transparent membrane, termed the envelope. Each of the granules composing the nucleus assimilates nourishment, thereby growing into an independent cell, which possesses a triple organization similar to that of its parent, and in like manner reproduces other cells.

A variety of tissues enters into the composition of an animal structure, yet their differences are not always distinctly marked, since the characteristics of some are not unlike those of others. We shall notice, however, only the more important of the tissues.

The Areolar, or Connective Tissue, is a complete network of delicate fibers, spread over the body, and serves to bind the various organs and parts together. The fibrous and serous tissues are modifications of the areolar.<[user]>

MinerU 2.5 Pro

CER 0.1556 | WER 0.1799 | F1 0.9979 | 1 region(s) | length 1.00x

CHAPTER II.

PHYSIOLOGICAL ANATOMY.

THE BONES.

All living bodies are made up of tissues. There is no part, no organ, however soft and yielding, or hard and resisting, which has not this peculiarity of structure. The bones of animals, as well as their flesh and fat, are composed of tissues, and all alike made up of cells. When viewed under a microscope, each cell is seen to consist of three distinct parts, a nucleolus, or dark spot, in the center of the cell, around which lies a mass of granules, called the nucleus; and this, in turn, is surrounded with a delicate, transparent membrane, termed the

envelope. Each of the granules composing the nucleus assimilates nourishment, thereby growing into an independent cell, which possesses a triple organization similar to that of its parent, and in like manner reproduces other cells.

A variety of tissues enters into the composition of an animal structure, yet their differences are not always distinctly marked, since the characteristics of some are not unlike those of others. We shall notice, however, only the more important of the tissues.

The Areolar, or Connective Tissue, is a complete network of delicate fibers, spread over the

body, and serves to bind the various organs and parts together. The fibrous and serous tissues are modifications of the areolar.

Fig. 4.

Nucleated cell. From Goeber. 1. Periphery of the cell, or cell-wall. 2. Nucleus. 3. Nucleolus in the center.

19

TrOCR large-printed

CER 0.4255 | WER 0.5063 | F1 0.6326 | 1 region(s) | length 0.69x

CASHIER CHAPTER II, INC.

PHYSIOLOGICAL ANATOMY.

TOTAL: THE BOXES.COM/PRICE

ALL LIVING DOLLES ARE MADE UP OF TASUES. THERE IS NO PART, NO ORGAN, HOWEVER SOFT AND YIELDING, OR HARD AND RESISTING, WHICH HAS NOT THIS DECULARITY OF STRUCTURE. THE BONES OF AM- MALS, AS WELL AS THEIR RESH AND FAT, ARE COMPOSED OF TISSUES, AND ALL ALIKE MADE UP OF CELLS. WHEN VIEWED UNDER A MICROSC- CODE, EACH CELL IS SEEN TO CONSIST OF THREE DISTMET PATTS, A INVEGOLLS, OR DARK SHOT, IN THE CENTER OF THE CELL, AROUND WHICH LIES A MASS OF GRANULES, CALLED THE WWLEWS; AND THIS, IN TURN, IS SUPPROVED WITH A DELICATE, TRANSLUANT MEMBRANE, TERMED THE INCLUDES, EXCLUDES OF THE GRANDEN, CHERPING THE FIG. 4.

0

THE INVOICE, HOWEVER, ONLY THE MORE IMPORTANT OF PERFORATED WITH THE AREOLAR, OR COMMENTIVE TASME, IS & COM. NUCEOHISINTHE PLETE NETWORK OF BELICATE FIBERS, SPREAD OVER THE CENTER. BODY, AND SERVICES TO BIND THE VARIOUS ORGANS AND PAPTS TOGETHER. THE AIRPONS AND SEROUS TISSUES ARE MODIFICATIONS OF THE AREOLAR.

1

Verified label (manual GT)

1382 normalized characters

CHAPTER II.

PHYSIOLOGICAL ANATOMY.

THE BONES.

All living bodies are made up of tissues. There is no part, no organ, however soft and yielding, or hard and resisting, which has not this peculiarity of structure. The bones of animals, as well as their flesh and fat, are composed of tissues, and all alike made up of cells. When viewed under a microscope, each cell is seen to consist of three distinct parts, a nucleolus, or dark spot, in the center of the cell, around which lies a mass of granules, called the nucleus; and this, in turn, is surrounded with a delicate, transparent membrane, termed the envelope. Each of the granules composing the nucleus assimilates nourishment, thereby growing into an independent cell, which possesses a triple organization similar to that of its parent, and in like manner reproduces other cells.

Fig. 4.

Nucleated cell. From Goeber. 1. Periphery of the cell, or cell-wall. 2. Nucleus. 3. Nucleolus in the center.

A variety of tissues enters into the composition of an animal structure, yet their differences are not always distinctly marked, since the characteristics of some are not unlike those of others. We shall notice, however, only the more important of the tissues.

The Areolar, or Connective Tissue, is a complete network of delicate fibers, spread over the body, and serves to bind the various organs and parts together. The fibrous and serous tissues are modifications of the areolar.

DeepSeek-OCR-2

CER 0.5362 | WER 0.7155 | F1 0.7368 | 11 region(s) | length 1.38x

CHAPTER 1.

$$\lceil P(N) = \sum_{i=1}^n \frac{1}{i} \rceil$$

THE BONES

All living bodies are made up of tissues. There is no part, no organ, however soft and yielding, or hard and resisting, which has not this peculiarity of structure. The bones of animals, as well as their flesh and fat, are composed of tissues, and all alike made up of cells. When viewed under a microscope, each cell is seen to consist of three distinct parts, a nucleolus, or dark spot, in the center of the cell, around which lies a mass of granules, called the nucleus, and this, in turn, is surrounded with a delicate, transparent membrane, termed the peri-

nucleus assimilates nourishment, thereby growing into an independent cell, which possesses a triple organization similar to that of its parent, and in like manner reproduces other cells.

A variety of tissues enters into the composition of an animal structure, yet their differences are not always distinctly marked, since the characteristics of some are not unlike those of others. We shall notice, however, only the more important of the tissues.

Nucleated cell.
From Goeber. 1.
Periphery of the cell, or cell-wall.
Nucleus.
Nucleolus in the center.

The Aeroform or Connectine Tissue is a continuous, membrane of nerve cells that covers the surface of the brain and spinal cord.

body, and serves to bind the various organs and parts together. The thorns and serous tissues are modifications of the adrenal,

8

GLM-OCR

CER 0.4971 | WER 0.6109 | F1 0.8090 | 11 region(s) | length 1.39x

CHAPTER II.<[user]>

PHYSIOLOGICAL ANATOMY.<[user]>

THE BONES.<[user]>

All living bodies are made up of tissues. There is no part, no organ, however soft and yielding, or hard and resisting, which has not this peculiarity of structure. The bones of animals, as well as their flesh and fat, are composed of tissues, and all alike made up of cells. When viewed under a microscope, each cell is seen to consist of three distinct parts, a nucleolus, or dark spot, in the center of the cell, around which lies a mass of granules, called the nucleus; and this, in turn, is surrounded with a delicate, transparent membrane, termed the<[user]>

All living bodies are made up of tissues. There is no part, no organ, however soft and yielding, or hard and resisting, which has not this peculiarity of structure. The bones of animals, as well as their flesh and fat, are composed of tissues, and all alike made up of cells. When viewed under a microscope, each cell is seen to consist of three distinct parts, a nucleolus, or dark spot, in the center of the cell, around which lies a mass of granules, called the nucleus; and this, in turn, is surrounded with a delicate, transparent membrane, termed the<[user]>

nucleus assimilates nourishment, thereby growing into an independent cell, which possesses a triple organization similar to that of its parent, and in like manner reproduces other cells.<[user]>

A variety of tissues enters into the composition of an animal structure, yet their differences are not always distinctly marked, since the characteristics of some are not unlike those of others. We shall notice, however, only the more important of the tissues.<[user]>

Nucleated cell. From Goeber. 1. Periphery of the cell, or cell-wall. 2. Nucleus. 3. Nucleolus in the center.<[user]>

The Areolar, or Connective Tissue, is a complete network of delicate fibers, spread over the<[user]>

body, and serves to bind the various organs and parts together. The fibrous and serous tissues are modifications of the areolar.<[user]>

19<[user]>

MinerU 2.5 Pro

CER 0.4609 | WER 0.5649 | F1 0.8246 | 11 region(s) | length 1.35x

CHAPTER II.

PHYSIOLOGICAL ANATOMY.

THE BONES.

All living bodies are made up of tissues. There is no part, no organ, however soft and yielding, or hard and resisting, which has not this peculiarity of structure. The bones of animals, as well as their flesh and fat, are composed of tissues, and all alike made up of cells. When viewed under a microscope, each cell is seen to consist of three distinct parts, a nucleolus, or dark spot, in the center of the cell, around which lies a mass of granules, called the nucleus; and this, in turn, is surrounded with a delicate, transparent membrane, termed the

All living bodies are made up of tissues. There is no part, no organ, however soft and yielding, or hard and resisting, which has not this peculiarity of structure. The bones of animals, as well as their flesh and fat, are composed of tissues, and all alike made up of cells. When viewed under a microscope, each cell is seen to consist of three distinct parts, a nucleolus, or dark spot, in the center of the cell, around which lies a mass of granules, called the nucleus; and this, in turn, is surrounded with a delicate, transparent membrane, termed the

nucleus assimilates nourishment, thereby growing into an independent cell, which possesses a triple organization similar to that of its parent, and in like manner reproduces other cells.

A variety of tissues enters into the composition of an animal structure, yet their differences are not always distinctly marked, since the characteristics of some are not unlike those of others. We shall notice, however, only the more important of the tissues.

Nucleated cell. From Goeber. 1. Periphery of the cell, or cell-wall. 2. Nucleus. 3. Nucleolus in the center.

The Areolar, or Connective Tissue, is a complete network of delicate fibers, spread over the

body, and serves to bind the various organs and parts together. The fibrous and serous tissues are modifications of the areolar.

19

TrOCR large-printed

CER 0.5072 | WER 0.6820 | F1 0.7414 | 11 region(s) | length 1.38x

CHAPTER II.

PHYSIOLOGICAL ANATOMY.

THE BONES.

ALL LIVING BOOKIES ARE MADE UP OF TISSUES. THERE IS NO PART, NO ORGAN, HOWEVER SOFT AND YIELDING, OR HARD AND RESISTING, WHICH HAS NOT THIS DECULARITY OF STRUCTURE. THE BONES OF AMI-MALS, AS WELL AS THEIR RESH AND FAT, ARE COMPOSED OF TUSSIES, AND ALL ALIKE MADE UP OF CELLS. WHEN VIEWED UNDER A MICROS-CODE, EACH CELL IS SEEN TO CONSIST OF THREE DUSTINET PATTS, A INVOICEOUS, OR DARK SPOT, IN THE CENTER OF THE CELL, AROUND WHICH LIES A MASS OF GRANULES, CALLED THE WWELENS, AND THIS, IN TURN, IS SUPPOUNDED WITH A DELICATE, TRANSPARENT MEMBRANE, TERMED THE ALL LIVING BOOKIES ARE MADE UP OF TISSUES. THERE IS NO PART, NO ORGAN, HOWEVER SOFT AND YIELDING, OR HARD AND RESISTING, WHICH HAS NOT THIS DECULARITY OF STRUCTURE, THE BONES OF AMI-MALS, AS WELL AS THEIR RESH AND FAT, ARE COMPOSED OF TUSSIES, AND ALL ALIKE MADE UP OF CELLS. WHEN VIEWED UNDER A MICROS-CODE, EACH CELL IS SEEN TO CONSIST OF THREE DUSTINET PATTS, A INVOICEOUS, OR DARK SPOT, IN THE CENTER OF THE CELL, AROUND WHICH LIES A MASS OF GRANULES, CALLED THE WWELENS, AND THIS, IN TURN, IS SUPPOUNDED WITH A DELICATE, TRANSPARENT MEMBRANE, TERMED THE ITTLVLV

NUCLEAS ASSIMILATES NOURISHMENT, THEREBY GROWING INTO AN INDEPENDENT CELL, WHICH POSSESSES A TRIPLE ORGANIZATION SIMILAR TO THAT OF ITS PARENT, AND IN LIKE MANNER REPRODUCES OTHER CELLS,

A VARIETY OF TISSUES ENTERS INTO THE COMPOSITION OF AN ANIMAL STRUCTURE, VET THEIR DIFFERENCES ARE NOT ALWAYS DISTINETLY MARKED, SINCE THE CHARACTERISTICS OF SOME ARE NOT UNLIKE THOSE OF OTHERS. WE SHALL NOTICE, HOWEVER, ONLY THE MORE IMPORTANT OF THE TISSUES, EXCLUDED WITHIN THE

NUCLEATED CELL.

FROM GOEBER, 1.

PERIPHERY OF THE CELL, OR CELL-WALL.

8. NUCLEUS. 5.

NUCLECLUS IN THE CENTER.

THE AREOLAR, OR CONNECTIVE TISSUE, IS A COMPLETE NETWORK OF DELICATE TBERS, SPREAD OVER THE BODY, AND SERVES TO BIND THE VARIOUS ORGANS AND PARTS TOGETHER. THE FIBROWS AND SEROUS TISSUES ARE MODIFICATIONS OF THE AREOLAR.

19

Verified label (manual GT)

1682 normalized characters

20 COMMON SENSE MEDICAL ADVISER.

The Nervous Tissue is of two kinds: The gray, which is pulpy and granulated, and the white fibrous tissue. The Adipose Tissue is an extremely thin membrane, composed of closed cells which contain fat. It is found principally just beneath the skin, giving it a smooth, plump appearance.

Fig. 5.

Arrangement of fibers in the Areolar Tissue. Magnified 135 diameters.

The Cartilaginous Tissue consists of nucleated cells, and, with the exception of bone, is the hardest part of the animal frame. The Osseous Tissue, or bone, is more compact and solid than the cartilaginous, for it contains a greater quantity of lime. The Muscular Tissue is composed of bundles of fibers, which are enclosed in a cellular membrane.

Various opinions have been entertained in regard to the formation, or growth, of bone. Some anatomists have supposed that all bone is formed in cartilage. But this is not true, for there is an intra-membranous, as well as an intra-cartilaginous, formation of bone, as may be seen in the development of the cranial bones, where the gradual calcification takes place upon the inner layers of the fibrous coverings. Intra cartilaginous deposit is found in the vicinity of the blood-vessels, within the cartilaginous canals; also, there are certain points first observed in the shafts of long bones, called centers of ossification. These points are no sooner formed than the cartilage corpuscles arrange themselves in concentric zones, and, lying in contact with one another, become very compact. As ossification proceeds, the cup-shaped cavities are converted into closed interstices of bone, with extremely thin lamellæ, or layers. These, however, soon increase

Fig. 6.

Human Adipose Tissue.

DeepSeek-OCR-2

CER 0.1766 | WER 0.2000 | F1 0.9510 | 1 region(s) | length 1.03x

text

The Nervous Tissue is of two kinds: The gray, which is pulpy and granulated, and the white fibrous tissue. The Adipose Tissue is an extremely thin membrane, composed of closed

figure_title

Fig. 5.

image

text

Arrangement of fibers in the Areolar Tissue. Magnified 135 dia-meters.

text

cells which contain fat. It is found principally just beneath the skin, giving it a smooth, plump appearance.

text

The Cartilaginous Tissue consists of nucleated cells, and, with the exception of bone, is the hardest part of the animal frame. The Osseous Tissue, or bone, is more compact and solid than the cartilaginous, for it contains a greater quantity of lime. The Muscular Tissue is composed of bundles of

text

fibers, which are enclosed in a cellular membrane.

text

Various opinions have been entertained in regard to the formation, or growth, of bone. Some anatomists have supposed that all bone is formed in cartilage. But this is not

text

true, for there is an intra-membranous, as well as an intra-cartilaginous, formation of bone, as may be seen in the development of the cranial bones, where the gradual calcification takes place upon the inner layers of the fibrous coverings. Intra cartilaginous deposit is found in the vicin-

figure_title

Fig. 6.

image

text

Human Adipose Tissue.

text

ity of the blood-vessels, within the cartilaginous canals; also, there are certain points first observed in the shafts of long bones, called centers of ossification. These points are no sooner formed than the cartilage corpuscles arrange themselves in concentric zones, and, lying in contact with one another, become very compact. As ossification proceeds, the cup-shaped cavities are converted into closed interstices of bone, with extremely thin lamella, or layers. These, however, soon increase

GLM-OCR

CER 0.0767 | WER 0.0786 | F1 0.9573 | 1 region(s) | length 0.92x

The Nervous Tissue is of two kinds: The gray, which is pulpy and granulated, and the white fibrous tissue. The Adipose Tissue is an extremely thin membrane, composed of closed cells which contain fat. It is found principally just beneath the skin, giving it a smooth, plump appearance.

The Cartilaginous Tissue consists of nucleated cells, and, with the exception of bone, is the hardest part of the animal frame. The Osseous Tissue, or bone, is more compact and solid than the cartilaginous, for it contains a greater quantity of lime. The Muscular Tissue is composed of bundles of fibers, which are enclosed in a cellular membrane.

Various opinions have been entertained in regard to the formation, or growth, of bone. Some anatomists have supposed that all bone is formed in cartilage. But this is not true, for there is an intra-membranous, as well as an intra-cartilaginous, formation of bone, as may be seen in the development of the cranial bones, where the gradual calcification takes place upon the inner layers of the fibrous coverings. Intra cartilaginous deposit is found in the vicinity of the blood-vessels, within the cartilaginous canals; also, there are certain points first observed in the shafts of long bones, called centers of ossification. These points are no sooner formed than the cartilage corpuscles arrange themselves in concentric zones, and, lying in contact with one another, become very compact. As ossification proceeds, the cup-shaped cavities are converted into closed interstices of bone, with extremely thin lamellæ, or layers. These, however, soon increase<|user|>

MinerU 2.5 Pro

CER 0.1213 | WER 0.1357 | F1 0.9893 | 1 region(s) | length 1.00x

20

COMMON SENSE MEDICAL ADVISER.

The Nervous Tissue is of two kinds: The gray, which is pulpy and granulated, and the white fibrous tissue. The Adipose Tissue is an extremely thin membrane, composed of closed

Fig. 5.

Arrangement of fibers in the Areolar Tissue. Magnified 135 diameters.

cells which contain fat. It is found principally just beneath the skin, giving it a smooth, plump appearance.

The Cartilaginous Tissue consists of nucleated cells, and, with the exception of bone, is the hardest part of the animal frame. The Osseous Tissue, or bone, is more compact and solid than the cartilaginous, for it contains a greater quantity of lime. The Muscular Tissue is composed of bundles of

fibers, which are enclosed in a cellular membrane.

Various opinions have been entertained in regard to the formation, or growth, of bone. Some anatomists have sup-

posed that all bone is formed in cartilage. But this is not true, for there is an intra-membranous, as well as an intra-cartilaginous, formation of bone, as may be seen in the development of the cranial bones, where the gradual calcification takes place upon the inner layers of the fibrous coverings. Intra cartilaginous deposit is found in the vicin-

Fig. 6.

Human Adipose Tissue.

ity of the blood-vessels, within the cartilaginous canals; also, there are certain points first observed in the shafts of long bones, called centers of ossification. These points are no sooner formed than the cartilage corpuscles arrange themselves in concentric zones, and, lying in contact with one another, become very compact. As ossification proceeds, the cup-shaped cavities are converted into closed interstices of bone, with extremely thin lamellæ, or layers. These, however, soon increase

TrOCR large-printed

CER 0.4411 | WER 0.5321 | F1 0.6054 | 1 region(s) | length 0.58x

90) COMMUN SENSE MEDICAL ADVISER.

THE NERVOUS TISSUE IS OF TWO KINGS: THE GRAY, WHICH IS PALPY AND GRANULATED, AND THE WHITE FORONS TISSUE. THE ADIPOSE TISSUE IS AN EXTREMELY THIN MEMBRANE, COMPOSED OF CLOSED

=

METERS TASTE IS COMPOSED OF BUNCLES OF FIBERS, WHICH ARE ENCLOSED IN A CELLULAR MEMBRANE. VARIOUS OPINIONS HAVE BEEN ENTERTAINED IN REWARD TO THE FORMATION, OR GROWTH, OF BONE. SOME ANATOMISTS HAVE SUPGOOD THAT ALL BONE IS FORMED

IN EARTILAGE, BUT THIS IS NOT BIG.6.

8.00

DEPOSIT IS FOUND IN THE VICTOR.

ITY OF THE BLOOD-YESSELS, WITHIN THE CARTILAGINONS CANALS; ALSO, THERE ARE CERTAIN POINTS INST OBSERVEL IN THE SHAFTS OF LONG BONES, CALLED CENTERS OF OSSIFICATION. THESE POINTS ARE NO SOOMER FORMED THAN THE CARTILAGE CORPUSCLES ARRANGE THEMSELVES IN CONCENTRIC ZONES, AND, LYING IN CONTACT WITH ONE ANOTHER, BECOME VERY COMPAET. AS DESIFICATION PROCEEDS, THE CUP-SHAPED CANTIES ARE CONVERTED INTO CLOSED INTERSTICES OF BONE, WITH EXTREMELY THIN LAMELLE, OR LAYERS. THESE, HOWEVER, SOON INCREASE

Verified label (manual GT)

1852 normalized characters

22 COMMON SENSE MEDICAL ADVISER.

separated from the body of the bone by a layer of cartilage, and the cancellated, or cellular structure, which remains for a time in the interior, represents the early condition of the ossifying substances.

The bones contain more earthy matter in their composition than any other part of the human body, being firm, hard, and of a lime color. They compose the skeleton or frame work, and, when united by natural ligaments, form what is known as the natural skeleton; when they are wired together, they are called an artificial skeleton. The number of bones in the human body is variously estimated; for those regarded as single by some anatomists are considered by others to consist of several distinct pieces. There are two hundred distinct bones in the human skeleton besides the teeth. These may be divided into those of the Head, Trunk, Upper Extremities, and Lower Extremities.

Fig. 10.

The bones of the skull separated. 1. Frontal, only half is seen. 2. Parietal. 3. Occipital, only half is seen. 4. Temporal. 5. Nasal. 6. Malar. 7. Superior maxillary (upper jaw). 8. Lachrymal. 9. Inferior maxillary (lower jaw). Between 4 and 6 a part of the sphenoid or wedge-shaped bone, is seen. Another bone assisting to form the skull, but not here seen, is called the ethmoid (sieve-like, from being full of holes), and is situated between the sockets of the eyes, forming the roof of the nose.

The Bones of the Head are classed as follows: eight belonging to the Cranium, and fourteen to the Face.

The bones of the Cranium are the occipital, two parietal, two temporal, frontal, sphenoid, and ethmoid. Those composing the face are, the two nasal, two superior maxillary, two lachrymal, two malar two palate, two inferior turbinated, vomer, and inferior maxillary. The cranial bones are composed of two dense plates, between which there is, in most places, a cancellated or cellular tissue. The external

DeepSeek-OCR-2

CER 0.2770 | WER 0.2905 | F1 0.8246 | 1 region(s) | length 0.73x

text
separated from the body of the bone by a layer of cartilage, and the cancellated, or cellular structure, which remains for a time in the interior, represents the early condition of the ossifying substances.

text
The bones contain more earthy matter in their composition than any other part of the human body, being firm, hard, and of a lime color. They compose the skeleton or frame work, and, when united by natural ligaments, form what is known as the natural skeleton; when they are wired together, they are called

figure_title
Fig. 10.

image

text
The bones of the skull separated. 1. Frontal, only half is seen. 2. Parietal. 3. Occipital, only half is seen. 4. Temporal. 5. Nasal. 6. Malar. 7. Superior maxillary (upper jaw). 8. Lachrymal. 9. Inferior maxillary (lower jaw). Between 4 and 6 a part of the sphenoid or wedge-shaped bone, is seen. Another bone assisting to form the skull, but not here seen, is called the ethmoid (sieve-like, from being full of holes), and is situated between the sockets of the eyes, forming the roof of the nose.

text
ital, two parietal, two temporal, frontal, sphenoid, and ethmoid. Those composing the face are, the two nasal, two superior maxillary, two lachrymal, two malar two palate, two inferior turbinated, vomer, and inferior maxillary. The cranial bones are composed of two dense plates, between which there is, in most places, a cancellated or cellular tissue. The external

GLM-OCR

CER 0.2748 | WER 0.2997 | F1 0.8244 | 1 region(s) | length 0.73x

separated from the body of the bone by a layer of cartilage, and the cancellated, or cellular structure, which remains for a time in the interior, represents the early condition of the ossifying substances.

The bones contain more earthy matter in their composition than any other part of the human body, being firm, hard, and of a lime color. They compose the skeleton or frame work, and, when united by natural ligaments, form what is known as the natural skeleton; when they are wired together, they are called an artificial skeleton.

The number of bones in the human body is variously estimated; for those regarded as single by some anatomists are considered by others to consist of several distinct pieces. There are two hundred distinct bones in the human skeleton besides the teeth. These may be divided into those of the Head, Trunk, Upper Extremities, and Lower Extremities.

The Bones of the Head are classed as follows: eight belonging to the Cranium, and fourteen to the Face.

The bones of the Cranium are the occipital, two parietal, two temporal, frontal, sphenoid, and ethmoid. Those composing the face are, the two nasal, two superior maxillary, two lachrymal, two malar two palate, two inferior turbinated, vomer, and inferior maxillary. The cranial bones are composed of two dense plates, between which there is, in most places, a cancellated or cellular tissue. The external<|user|>

MinerU 2.5 Pro

CER 0.3488 | WER 0.3731 | F1 0.9954 | 1 region(s) | length 1.00x

22

COMMON SENSE MEDICAL ADVISER.

separated from the body of the bone by a layer of cartilage, and the cancellated, or cellular structure, which remains for a time in the interior, represents the early condition of the ossifying substances.

The bones contain more earthy matter in their composition than any other part of the human body, being firm, hard, and of a lime color. They compose the skeleton or frame work, and, when united by natural ligaments, form what is known as the natural skeleton; when they are wired together, they are called

Fig. 10.

The bones of the skull separated. 1. Frontal, only half is seen. 2. Parietal. 3. Occipital, only half is seen. 4. Temporal. 5. Nasal. 6. Malar. 7. Superior maxillary (upper jaw). 8. Lachrymal. 9. Inferior maxillary (lower jaw). Between 4 and 6 a part of the sphenoid or wedge-shaped bone, is seen. Another bone assisting to form the skull, but not here seen, is called the ethmoid (sieve-like, from being full of holes), and is situated between the sockets of the eyes, forming the roof of the nose.

an artificial skeleton. The number of bones in the human body is variously estimated; for those regarded as single by some anatomists are considered by others to consist of several distinct pieces. There are two hundred distinct bones in the human skeleton besides the teeth. These may be divided into those of the Head, Trunk, Upper Extremities, and Lower Extremities.

The Bones of the Head are classed as follows: eight belonging to the Cranium, and fourteen to the Face.

The bones of the Cranium are the occip-

ital, two parietal, two temporal, frontal, sphenoid, and ethmoid. Those composing the face are, the two nasal, two superior maxillary, two lachrymal, two malar two palate, two inferior turbinated, vomer, and inferior maxillary. The cranial bones are composed of two dense plates, between which there is, in most places, a cancellated or cellular tissue. The external

TrOCR large-printed

CER 0.6258 | WER 0.7187 | F1 0.4279 | 1 region(s) | length 0.41x

90 COMBON SENSE MEDICAL ADVISES, INC.
REPARATED FROM THE BODY OF THE BONE BY A LAYER OF CAR-
TILAGE, AND THE CANCELLED, OR CELLULAR STRUCTURE, WHICH REMAINS
FOR A TIME IN THE INTERIOR, REPRESENTS THE EARLY CONDITION OF THE
DESIFYING SUBSTANCES.
THE BONES CONTAIN MORE EARTHY MATTER IN THEIR COMPOSITION
THAN ANY OTHER PART OF THE HUMAN BODY, BEING ARM, HARD, AND OF
A TIME COLOR. THEY COMPOSE THE SKELETON OR FRAME WORK, AND,
WHEN UNITED BY NATURAL IGAMENTS, FORM WHAT IS KNOWN AS THE
WHATWAL SKELETON; WHEN THEY ARE WIRED TOGETHER, THEY ARE CALLED
CASHIER AND RETURRIAL SKELETON.
REG. 10. THE NUMBER OF BONES
@
8
NOOL OF THE MORE. COME AGAIN THE RECIP-
TOTAL, TWO MARIETAL, TWO SAMPARAL, FINAL, ADVERTISED, AND RETURNABLE
SANDWICH OR MARINA, MARINA, MARINA, MARINA, MARINA, JOHORORORA,

Verified label (manual GT)

1852 normalized characters

22 COMMON SENSE MEDICAL ADVISER.

separated from the body of the bone by a layer of cartilage, and the cancellated, or cellular structure, which remains for a time in the interior, represents the early condition of the ossifying substances.

The bones contain more earthy matter in their composition than any other part of the human body, being firm, hard, and of a lime color. They compose the skeleton or frame work, and, when united by natural ligaments, form what is known as the natural skeleton; when they are wired together, they are called an artificial skeleton. The number of bones in the human body is variously estimated; for those regarded as single by some anatomists are considered by others to consist of several distinct pieces. There are two hundred distinct bones in the human skeleton besides the teeth. These may be divided into those of the Head, Trunk, Upper Extremities, and Lower Extremities.

Fig. 10.

The bones of the skull separated. 1. Frontal, only half is seen. 2. Parietal. 3. Occipital, only half is seen. 4. Temporal. 5. Nasal. 6. Malar. 7. Superior maxillary (upper jaw). 8. Lachrymal. 9. Inferior maxillary (lower jaw). Between 4 and 6 a part of the sphenoid or wedge-shaped bone, is seen. Another bone assisting to form the skull, but not here seen, is called the ethmoid (sieve-like, from being full of holes), and is situated between the sockets of the eyes, forming the roof of the nose.

The Bones of the Head are classed as follows: eight belonging to the Cranium, and fourteen to the Face.

The bones of the Cranium are the occipital, two parietal, two temporal, frontal, sphenoid, and ethmoid. Those composing the face are, the two nasal, two superior maxillary, two lachrymal, two malar two palate, two inferior turbinated, vomer, and inferior maxillary. The cranial bones are composed of two dense plates, between which there is, in most places, a cancellated or cellular tissue. The external

DeepSeek-OCR-2

CER 0.2538 | WER 0.3425 | F1 0.7760 | 10 region(s) | length 1.10x

COMPANY SLOGAN
COMPANY SLOGAN
COMPANY SLOGAN
COMPANY SLOGAN

22

separated from the body of the bone by a layer of cartilage is called a tilage, and the cancellous, or cellular structure, which remains for a time in the interior, represents the early condition of the osseous substance.

The bones contain more earthy matter in their composition than any other part of the human body, being firm, hard, and of a lime color. They compose the skeleton or frame work, and when united by natural ligaments, form what is known as the natural skeleton; when they are wired together, they are called

artificial skeleton.
The number of bones in the human body is variously estimated; for those regarded as single by some anatomists are considered by others to consist of several distinct pieces. There are two hundred distinct bones in the human skeleton besides the teeth. These may be divided into those of the Head, Trunk, Upper Extremities, and Lower Extremities.

Из 10, что вы видите?

The bones of the skull separated.
1. Frontal, only half is seen.
2. Parietal.
3. Occipital, only half is seen.
4. Temporal.
5. Nasal.
6. Malar.
7. Superior maxillary (upper jaw).
8. Lachrymal.
9. Inferior maxillary (lower jaw).
Between 1, 9. Inferior maxillary (lower jaw).
Between 1, 9. Inferior maxillary (lower jaw).
Between 1, 9. Superior maxillary (upper jaw).
Between 1, 9. Inferior maxillary (lower jaw).
Between 1, 9. Superior maxillary (upper jaw).
1. Frontal, only half is seen.
2. Parietal.
3. Occipital, only half is seen.
4. Temporal.

The Bones of the Head are classed as follows: eight belonging to the Cranium, and fourteen to the Face.

The bones of the Cranium are the occipit

ital, two parietal, two temporal, frontal, sphenoid, and ethmoid. Those composing the face are, the two nasal, two superior maxillary, two maxillary, two maxillary, two maxillary, two inferior or turbinated, or inferior maxillary. The cranial bones are composed of two dense plates, between which there is a network of fine, delicate, and fine, delicate, and fine, delicate, and fine, delicate, and fine, delicate, and fine, delicate, and fine, delicate, and fine, delicate,

GLM-OCR

CER 0.0308 | WER 0.0428 | F1 0.9805 | 10 region(s) | length 1.03x

COMMON SENSE MEDICAL ADVISER.<|user|>

22<|user|>

separated from the body of the bone by a layer of cartilage, and the cancellated, or cellular structure, which remains for a time in the interior, represents the early condition of the ossifying substances.<|user|>

The bones contain more earthy matter in their composition than any other part of the human body, being firm, hard, and of a lime color. They compose the skeleton or frame work, and, when united by natural ligaments, form what is known as the natural skeleton; when they are wired together, they are called<|user|>

an artificial skeleton. The number of bones in the human body is variously estimated; for those regarded as single by some anatomists are considered by others to consist of several distinct pieces. There are two hundred distinct bones in the human skeleton besides the teeth. These may be divided into those of the Head, Trunk, Upper Extremities, and Lower Extremities.<|user|>

Fig. 10.<|user|>

The bones of the skull separated. 1. Frontal, only half is seen. 2. Parietal. 3. Occipital, only half is seen. 4. Temporal. 5. Nasal. 6. Malar. 7. Superior maxillary (upper jaw). 8. Lachrymal. 9. Inferior maxillary (lower jaw). Between 4 and 6 a part of the sphenoid or wedge-shaped bone, is seen. Another bone assisting to form the skull, but not here seen, is called the ethmoid (sieve-like, from being full of holes), and is situated between the sockets of the eyes, forming the roof of the nose.<|user|>

The Bones of the Head are classed as follows: eight belonging to the Cranium, and fourteen to the Face.<|user|>

The bones of the Cranium are the occip-<|user|>

ital, two parietal, two temporal, frontal, sphenoid, and ethmoid. Those composing the face are, the two nasal, two superior maxillary, two lachrymal, two malar two palate, two inferior turbinated, vomer, and inferior maxillary. The cranial bones are composed of two dense plates, between which there is, in most places, a cancellated or cellular tissue. The external<|user|>

MinerU 2.5 Pro

CER 0.0059 | WER 0.0153 | F1 0.9908 | 10 region(s) | length 1.00x

COMMON SENSE MEDICAL ADVISER.

separated from the body of the bone by a layer of cartilage, and the cancellated, or cellular structure, which remains for a time in the interior, represents the early condition of the ossifying substances.

The bones contain more earthy matter in their composition than any other part of the human body, being firm, hard, and of a lime color. They compose the skeleton or frame work, and, when united by natural ligaments, form what is known as the natural skeleton; when they are wired together, they are called

an artificial skeleton. The number of bones in the human body is variously estimated; for those regarded as single by some anatomists are considered by others to consist of several distinct pieces. There are two hundred distinct bones in the human skeleton besides the teeth. These may be divided into those of the Head, Trunk, Upper Extremities, and Lower Extremities.

The bones of the skull separated. 1. Frontal, only half is seen. 2. Parietal. 3. Occipital, only half is seen. 4. Temporal. 5. Nasal. 6. Malar. 7. Superior maxillary (upper jaw). 8. Lachrymal. 9. Inferior maxillary (lower jaw). Between 4 and 6 a part of the sphenoid or wedge-shaped bone, is seen. Another bone assisting to form the skull, but not here seen, is called the ethmoid (sieve-like, from being full of holes), and is situated between the sockets of the eyes, forming the roof of the nose.

The Bones of the Head are classed as follows: eight belonging to the Cranium, and fourteen to the Face.

The bones of the

Cranium are the occip-

ital, two parietal, two temporal, frontal, sphenoid, and ethmoid. Those composing the face are, the two nasal, two superior maxillary, two lachrymal, two malar two palate, two inferior turbinated, vomer, and inferior maxillary. The cranial bones are composed of two dense plates, between which there is, in most places, a cancellated or cellular tissue. The external

TrOCR large-printed

CER 0.3790 | WER 0.4709 | F1 0.6703 | 10 region(s) | length 0.66x

COMMON SENSE MEDICAL ADVISER.

29
SEPARATED FROM THE BODY OF THE BONE BY A LAYER OF CAR-
TILAGE, AND THE CANCELLATED, OR CELLULAR STRUCTURE, WHICH REMAINS
FOR A TIME IN THE INTERIOR, REPRESENTS THE EARLY CONDITION OF THE
OSSIFYING SUBSTANCES.
THE BONES CONTAIN MORE EARTHY MATTER IN THEIR COMPOSITION
THAN ANY OTHER PART OF THE HUMAN BODY, BEING FIRM, HARD, AND OF
A TIME COLOR. THEY COMPOSE THE SKEETON OR FRAME WORK, AND,
WHEN UNITED BY NATURAL IGAMENTS, FORM WHAT IS KNOWN AS THE
NATURAL SKELETON; WHEN THEY ARE WIRED TOGETHER, THEY ARE CALLED
AN ARTIFICIAL SKELETON.
THE NUMBER OF BONES
IN THE HUMAN BODY IS
VARIOUSLY ESTIMATED; FOR
THOSE REGARDED AS SINGLE
BY SOME ANATOMISTS ARE
CONSIDERED BY OTHERS TO
CONSIST OF SEVERAL DISTINET
PIECES. THERE ARE TWO
HUNDRED DISTINCT BONES
IN THE HUMAN SKELETON
BESIDES THE TEETH. THESE
MAY BE DIVIDED INTO
TOTAL :
THOSE OF THE HEAD,
TRUNK, UPPER EXTREM-
ITIES, AND LOWER EX-
ITREMITIES,
FIG. 10.
CASHIER:
ROOF OF THE MOSE,
THE BONES OF THE
HEAD ARE CLASSED AS
FOLLOWS: EIGHT BELONG-
ING TO THE CRANIUM, AND
FOURTEEN TO THE FACE.
THE BONES OF THE
CRANIUM ARE THE OCCIP-
ITAL, TWO PERIETAL, TWO TAMPORAL, FORUAL, SPREAD, TWO RETURNA
AMOUNT AND RECEIPT FOR AN OFFER, AND RETURNS AND FORMULA, INCLUSIVE ADVERTISING AND RECEIPT FOR

Verified label (manual GT)

1964 normalized characters

THE TRUNK. 23

plate is fibrous, the internal, compact and vitreous. The skull is nearly oval in form, convex externally, the bone being much thicker at the base than elsewhere, and it is, in every respect admirably adapted to resist any injury to which it may be exposed, thus affording ample protection to the brain substance which it envelops. The internal surface of the cranium presents eminences and depressions for lodging the convolutions of the brain, and numerous furrows for the ramifications of the blood-vessels. The bones of the cranium are united to one another by ragged edges called sutures, which are quite distinct in the child, but which in old age are nearly effaced. Some authorities suppose that by this arrangement the cranium is less liable to be fractured by blows; others think that the sutures allow the growth of these bones, which takes place by a gradual osseous enlargement at the margins. The bones of the Face are joined at the lower part and in front of the cranium, and serve for the attachment of powerful muscles which assist in the process of mastication. Although the soft parts of the face cover the bony structure, yet they do not conceal its principal features, or materially change its proportions. The form of the head and face presents some remarkable dissimilarities in different races.

Fig. 11.

1. The first bone of the sternum (breast-bone). 2. The second bone of the sternum. 3. The cartilage of the sternum. 4. The first dorsal vertebra (a bone of the spinal column). 5. The last dorsal vertebra. 6. The first rib. 7. Its head. 8. Its neck. 9. Its tubercle. 10. The seventh or last true rib. 11. The cartilage of the third rib. 12. The floating ribs.

The Trunk has fifty-four bones, which are as follows: The Os Hyoides, the Sternum, twenty-four Ribs, twenty-four vertebrae or bones of the Spinal Column, the Sacrum, the Coccyx, and two Ossa Innominata. The Os Hyoides, situated at the base of the tongue, is the most isolated bone of the skeleton, and serves for the attachment of

DeepSeek-OCR-2

CER 0.2317 | WER 0.3465 | F1 0.9790 | 1 region(s) | length 1.01x

text

plate is fibrous, the internal, compact and vitreous. The skull is nearly oval in form, convex externally, the bone being much thicker at the base than elsewhere, and it is, in every respect admirably adapted to resist any injury to which it may be exposed, thus affording ample protection to the brain substance which it envelops. The internal surface of the cranium presents eminences and depressions for lodging the convolutions of the brain, and numerous furrows for the ramifications of the blood-vessels. The bones of the cranium are united to one another by ragged edges called sutures, which are quite distinct in the child, but which in old age are nearly effaced. Some authorities sup-

text

pose that by this arrangement the cranium is less liable to be fractured by blows; others think that the sutures allow the growth of these bones, which takes place by a gradual osseous enlargement at the margins. The bones of the Face are joined at the lower part and in front of the cranium, and serve for the attachment of powerful muscles which assist in the process of mastication. Although the soft parts of the face cover the bony structure, yet they do not conceal its principal features, or materially change its proportions. The form of the head and face presents some remarkable dissimilarities in different races.

text

The Trunk has fifty-four bones, which are as follows: The Os Hyoides, the Sternum, twenty-four Ribs, twenty-four vertebrae or bones of the Spinal Column, the Sacrum, the Coccyx, and two Ossa Innominata. The Os Hyoides, situated at the base of the tongue, is the most isolated bone of the skeleton, and serves for the attachment of

figure_title

Fig. 11.

image

text

1. The first bone of the sternum (breast-bone). 2. The second bone of the sternum. 3. The cartilage of the sternum. 4. The first dorsal vertebra (a bone of the spinal column). 5. The last dorsal vertebra. 6. The first rib. 7. Its head. 8. Its neck. 9. Its tubercle. 10. The seventh or last true rib. 11. The cartilage of the third rib. 12. The floating ribs.

GLM-OCR

CER 0.1970 | WER 0.2282 | F1 0.8703 | 1 region(s) | length 0.81x

plate is fibrous, the internal, compact and vitreous. The skull is nearly oval in form, convex externally, the bone being much thicker at the base than elsewhere, and it is, in every respect admirably adapted to resist any injury to which it may be exposed, thus affording ample protection to the brain substance which it envelops. The internal surface of the cranium presents eminences and depressions for lodging the convolutions of the brain, and numerous furrows for the ramifications of the blood-vessels. The bones of the cranium are united to one another by ragged edges called sutures, which are quite distinct in the child, but which in old age are nearly effaced. Some authorities suppose that by this arrangement the cranium is less liable to be fractured by blows; others think that the sutures allow the growth of these bones, which takes place by a gradual osseous enlargement at the margins. The bones of the Face are joined at the lower part and in front of the cranium, and serve for the attachment of powerful muscles which assist in the process of mastication. Although the soft parts of the face cover the bony structure, yet they do not conceal its principal features, or materially change its proportions. The form of the head and face presents some remarkable dissimilarities in different races.

The Trunk has fifty-four bones, which are as follows: The Os Hyoides, the Sternum, vertebrae or bones of the Spinal Column, the Sacrum, the Coccyx, and two Ossa Innominata. The Os Hyoides, situated at the base of the tongue, is the most isolated bone of the skeleton, and serves for the attachment of<[user]>

MinerU 2.5 Pro

CER 0.1690 | WER 0.1831 | F1 0.9831 | 1 region(s) | length 1.00x

THE TRUNK.

23

plate is fibrous, the internal, compact and vitreous. The skull is nearly oval in form, convex externally, the bone being much thicker at the base than elsewhere, and it is, in every respect admirably adapted to resist any injury to which it may be exposed, thus affording ample protection to the brain substance which it envelops. The internal surface of the cranium presents eminences and depressions for lodging the convolutions of the brain, and numerous furrows for the ramifications of the blood-vessels. The bones of the cranium are united to one another by ragged edges called sutures, which are quite distinct in the child, but which in old age are nearly effaced. Some authorities sup-

pose that by this arrangement the cranium is less liable to be fractured by blows; others think that the sutures allow the growth of these bones, which takes place by a gradual osseous enlargement at the margins. The bones of the Face are joined at the lower part and in front of the cranium, and serve for the attachment of powerful muscles which assist in the process of mastication. Although the soft parts of the face cover the bony structure, yet they do not conceal its principal features, or materially change its proportions. The form of the head and face presents some remarkable dissimilarities in different races.

The Trunk has fifty-four bones, which are as follows: twenty-four Ribs, twenty-four Column, the Sacrum, the Coc The Os Hyoides, situated at the isolated bone of the skeleton, a

1. The first bone of the sternum (breast-bone). 2. The second bone of the sternum. 3. The cartilage of the sternum. 4. The first dorsal vertebra (a bone of the spinal column). 5. The last dorsal vertebra. 6. The first rib. 7. Its head. 8. Its neck. 9. Its tubercle. 10. The seventh or last true rib. 11. The cartilage of the third rib. 12. The floating ribs.

The Os Hyoides, the Sternum, vertebrae or bones of the Spinal cyx, and two Ossa Innominata. The base of the tongue, is the most and serves for the attachment of

TrOCR large-printed

CER 0.4221 | WER 0.5521 | F1 0.5958 | 1 region(s) | length 0.64x

TOTAL THE TRUNK, PRICE
PLATE IS FIBROUS, THE INTERNAL, COMPACT AND VITREOUS. THE SKULL IS
MEARLY OVAL IN FORM, CONVEX EXTERNAILY, THE BONE BEING MUCH
THICKER AT THE BASE THAN ELSEWHERE, AND IT IS, IN EVERY RESPECT
ADMINABLE ADAPTED TO RESIST ANY INJURY TO WHICH IT MAY BE
EXPOSED, THIS AMONG AMPLE PROTECTION TO THE BRAIN SUBSTANCE
WHICH IT ENVELOPS. THE INTERNAL SURFACE OF THE CRANTUM PRESENTS
EMINENCES AND DEPRESSIONS FOR LOOGING THE CONVOLUTIONS OF THE
BRAIN, AND NUMEROUS FURIONS FOR THE FAMITERIONS OF THE BLOOD.
VESELS. THE BOXES OF THE ERAMIUM ARE UNITED TO ONE ANOTHER BY
RAGGET EDGES CALLED AUTURES, WHICH ARE QUITE DUSTINET IN THE CHILD,
BUT WHICH IN OLD ARE ARE REALLY EFACED. SOME AUTHORITIES SUP-
PRICE THAT BY THIS ATTRIBERMENT OF FIG. 11.
CASHIER
PROPORTIONS, THE FORM OF THE LOWER INCLUSIVE THE RETURNS ORDER:
RECEIPT AND FACE PRESENTS SANDWAN, & THE USE ADVERTISING AGREEMENT
DIFERENT RACES. THE NUMBERGE. IN THE RETURNS OR INSTITUTE.
THE TRUNK HAS ATTY-FOUR THE BOATING RICE.
BONES, WHICH ARE AS FOLLOWS: THE OS HYOIDES, THE RETURN,
TWENTY-FOUR RIBS, TWENTY-FOUR VERTENUE OR BONES OF THE SPINAL
COLUMN, THE SAGNUM, THE COCYZ, AND TWO OSSA INHOMINATA.
THE OS HYOIDES, SITRATED AT THE BASE OF THE LONGME, IS THE MOST
ISOLATED BONE OF THE SKELETON, AND SERVICES FOR THE ATTACHMENT OF

Verified label (manual GT)

1964 normalized characters

THE TRUNK. 23

plate is fibrous, the internal, compact and vitreous. The skull is nearly oval in form, convex externally, the bone being much thicker at the base than elsewhere, and it is, in every respect admirably adapted to resist any injury to which it may be exposed, thus affording ample protection to the brain substance which it envelops. The internal surface of the cranium presents eminences and depressions for lodging the convolutions of the brain, and numerous furrows for the ramifications of the blood-vessels. The bones of the cranium are united to one another by ragged edges called sutures, which are quite distinct in the child, but which in old age are nearly effaced. Some authorities suppose that by this arrangement the cranium is less liable to be fractured by blows; others think that the sutures allow the growth of these bones, which takes place by a gradual osseous enlargement at the margins. The bones of the Face are joined at the lower part and in front of the cranium, and serve for the attachment of powerful muscles which assist in the process of mastication. Although the soft parts of the face cover the bony structure, yet they do not conceal its principal features, or materially change its proportions. The form of the head and face presents some remarkable dissimilarities in different races.

Fig. 11.

1. The first bone of the sternum (breast-bone). 2. The second bone of the sternum. 3. The cartilage of the sternum. 4. The first dorsal vertebra (a bone of the spinal column). 5. The last dorsal vertebra. 6. The first rib. 7. Its head. 8. Its neck. 9. Its tubercle. 10. The seventh or last true rib. 11. The cartilage of the third rib. 12. The floating ribs.

The Trunk has fifty-four bones, which are as follows: The Os Hyoides, the Sternum, twenty-four Ribs, twenty-four vertebræ or bones of the Spinal Column, the Sacrum, the Coccyx, and two Ossa Innominata. The Os Hyoides, situated at the base of the tongue, is the most isolated bone of the skeleton, and serves for the attachment of

DeepSeek-OCR-2

CER 0.1079 | WER 0.1521 | F1 0.9161 | 7 region(s) | length 1.10x

2 3

TEXT
TEXT

plate is fibrous, the internal, compact and vitreous. The skull is nearly oval in form, convex externally, the bone being much thicker at the base than elsewhere, and it is, in every respect admirably adapted to resist any injury to which it may be exposed, thus affording ample protection to the brain substance which it envelops. The internal surface of the cranium presents eminences and depressions for lodging the convolutions of the brain, and numerous furrows for the ramifications of the blood-vessels. The bones of the cranium are united to one another by ragged edges called sutures, which are quite distinct in the child, but which in old age are nearly effaced. Some authorities support the opinion that the internal surface of the cranium is enclosed in a hard, white substance, and that this is the seat of the affections which excite the brain.

pose that by this arrangement the cranium is less liable to be fractured by blows; others think that the sutures allow the growth of these bones, which takes place by a gradual osseous enlargement at the margins. The bones of the Face are joined at the lower part and in front of the cranium, and serve for the attachment of powerful muscles which assist in the process of mastication. Although the soft parts of the face cover the bony structure, yet they do not conceal its principal features, or materially change its proportions. The form of the head and face presents some remarkable dissimilarities in different races.

Hi, I

1. The first bone of the sternum (breast-bone).
2. The second bone of the sternum.
3. The cartilage of the sternum.
4. The first dorsal vertebra (a bone of the spinal column).
5. The last dorsal vertebra.
6. The first rib.
7. Its head.
8. Its neck.
9. Its tubercle.
10. The seventh or last true rib.
11. The cartilage of the third rib.
12. The floating ribs.

The Trunk has fifty-four bones, which are as follows: The Os Hoyoïdes, the Sternum, twenty-four Ribs, twenty-four vertebrae or bones of the Spinal Column, the Sacrum, the Coccyx, and two Ossa Innominata. The Os Hoyoïdes, situated at the base of the tongue, is the most isolated bone of the skeleton, and serves for the attachment of the tongue to the skull.

GLM-OCR

CER 0.0204 | WER 0.0310 | F1 0.9833 | 7 region(s) | length 1.02x

23<[user]>

THE TRUNK.<[user]>

plate is fibrous, the internal, compact and vitreous. The skull is nearly oval in form, convex externally, the bone being much thicker at the base than elsewhere, and it is, in every respect admirably adapted to resist any injury to which it may be exposed, thus affording ample protection to the brain substance which it envelops. The internal surface of the cranium presents eminences and depressions for lodging the convolutions of the brain, and numerous furrows for the ramifications of the blood-vessels. The bones of the cranium are united to one another by ragged edges called sutures, which are quite distinct in the child, but which in old age are nearly effaced. Some authorities sup-<[user]>

pose that by this arrangement the cranium is less liable to be fractured by blows; others think that the sutures allow the growth of these bones, which takes place by a gradual osseous enlargement at the margins. The bones of the Face are joined at the lower part and in front of the cranium, and serve for the attachment of powerful muscles which assist in the process of mastication. Although the soft parts of the face cover the bony structure, yet they do not conceal its principal features, or materially change its proportions. The form of the head and face presents some remarkable dissimilarities in different races.<[user]>

Fig. 11.<[user]>

1. The first bone of the sternum (breast bone). 2. The second bone of the sternum. 3. The cartilage of the sternum. 4. The first dorsal vertebra (a bone of the spinal column). 5. The last dorsal vertebra. 6. The first rib. 7. Its head. 8. Its neck. 9. Its tubercle. 10. The seventh or last true rib. 11. The cartilage of the third rib. 12. The floating ribs.<[user]>

The Trunk has fifty-four bones, which are as follows: The Os Hyoides, the Sternum, twenty-four Ribs, twenty-four vertebrae or bones of the Spinal Column, the Sacrum, the Coccyx, and two Ossa Innominata. The Os Hyoides, situated at the base of the tongue, is the most isolated bone of the skeleton, and serves for the attachment of<[user]>

MinerU 2.5 Pro

CER 0.0158 | WER 0.0254 | F1 0.9845 | 7 region(s) | length 1.01x

THE TRUNK.

plate is fibrous, the internal, compact and vitreous. The skull is nearly oval in form, convex externally, the bone being much thicker at the base than elsewhere, and it is, in every respect admirably adapted to resist any injury to which it may be exposed, thus affording ample protection to the brain substance which it envelops. The internal surface of the cranium presents eminences and depressions for lodging the convolutions of the brain, and numerous furrows for the ramifications of the blood-vessels. The bones of the cranium are united to one another by ragged edges called sutures, which are quite distinct in the child, but which in old age are nearly effaced. Some authorities sup-

pose that by this arrangement the cranium is less liable to be fractured by blows; others think that the sutures allow the growth of these bones, which takes place by a gradual osseous enlargement at the margins. The bones of the Face are joined at the lower part and in front of the cranium, and serve for the attachment of powerful muscles which assist in the process of mastication. Although the soft parts of the face cover the bony structure, yet they do not conceal its principal features, or materially change its proportions. The form of the head and face presents some remarkable dissimilarities in different races.

1. The first bone of the sternum (breast-bone). 2. The second bone of the sternum.

3. The cartilage of the sternum. 4. The

first dorsal vertebra (a bone of the spinal column). 5. The last dorsal vertebra. 6.

The first rib. 7. Its head. 8. Its neck. 9.

Its tubercle. 10. The seventh or last true

rib. 11. The cartilage of the third rib. 12.

The floating ribs.

The Trunk has fifty-four The floating ribs.

bones, which are as follows: The Os Hyoides, the Sternum, twenty-four Ribs, twenty-four vertebrae or bones of the Spinal Column, the Sacrum, the Coccyx, and two Ossa Innominata. The Os Hyoides, situated at the base of the tongue, is the most isolated bone of the skeleton, and serves for the attachment of

TrOCR large-printed

CER 0.0947 | WER 0.1634 | F1 0.8806 | 7 region(s) | length 0.95x

28

THE TRUNK.
PLATE IS FIBROUS, THE INTERNAL, COMPACT AND VITREOUS. THE SKULL IS NEARLY OVAL IN FORM, CONVEX EXTERNALLY, THE BONE BEING MUCH THICKER AT THE BASE THAN ELSEWHERE, AND IT IS, IN EVERY RESPECT ADMIRABLY ADAPTED TO RESIST ANY INJURY TO WHICH IT MAY BE EXPOSED, THUS AFFORDING AMPLE PROTECTION TO THE BRAIN SUBSTANCE WHICH IT ENVELOPS. THE INTERNAL SURFACE OF THE CRANTUM PRESENTS EMINENCES AND DEPRESSIONS FOR LODGING THE CONVOLUTIONS OF THE BRAIN, AND NUMEROUS FURTOWS FOR THE FAMIFICATIONS OF THE BLOOD-VESSELS. THE BONES OF THE CRANIUM ARE UNITED TO ONE ANOTHER BY RAGGED EDGES CALLED SATURES, WHICH ARE QUITE DISTINCT IN THE CHILD, BUT WHICH IN OLD AGE ARE NEARLY EFFACED. SOME AUTHORITIES SUP-POSE THAT BY THIS ARRANGEMENT THE CRANIUM IS LESS LIABLE TO BE FRACTURED BY BLOWS; OTHERS THINK THAT THE SUTURES ALLOW THE GROWTH OF THESE BONES, WHICH TAKES PLACE BY A GRADUAL OSSEOUS ENLARGEMENT AT THE MARGINS. THE BONES OF THE FACE ARE JOINED AT THE LOWER PART AND IN FRONT OF THE CRAN-AMOUNT OF POWERAL MODEOLES MASTICATION. ALTHOUGH THE SOFT PARTS OF THE FACE COVER THE BONY STRUCTURE, YET THEY DO NOT CONCEAL ITS PRINCIPAL FEAT-URES, OR MATERIALLY CHANGE ITS PROPORTIONS, THE FORM OF THE HEAD AND FACE PRESENTS SOME REMARKABLE DISSIMILARITIES IN DIFFERENT RACES.

Fig. 11.

1. THE FIRST BONE OF THE STERNUM (BREAST-BONE). 2. THE SECOND BONE OF THE STERNUM.
3. THE CARTILAGE OF THE STERNUM. 4. THE COUNT DORSAL VENTEDOTA (A BONEI OF THE SPINAL THE FIRST RIB. 7. ITS HEAD. 8. ITS NECK. 9. ITS TUBERCLE. 10. THE SEYENTH OR LAST TRUE THE ROATING RIBE OF THE THIRD RIB. 12.

THE TRUNK HAS FIFTY-FOUR THE MORNING INOS. BONES, WHICH ARE AS FOLLOWS: THE OS HYOIDES, THE RETURNUM, TWENTY-FOUR RIBS, TWENTY-FOUR VENTEBRE OR BONES OF THE SPINAL COLUMN, THE SACNUM, THE COCCHA, AND TWO OSSA INNOMINATA. THE OS HYOIDES, SITUATED AT THE BASE OF THE TONGUE, IS THE MOST ISOLATED BONE OF THE SKELETON, AND SERVICES FOR THE ATTACHMENT OF

Verified label (manual GT)

1885 normalized characters

24 COMMON SENSE MEDICAL ADVISER.

muscles. The Sternum, or breast-bone, in a child is composed of six pieces, in the adult of three, which in old age are consolidated into one bone. The Ribs are thin, curved bones, being convex externally. There are twelve on each side, and all are attached to the spinal column. The seven upper ribs, which are united in front of the sternum, are termed true ribs; the next three, which are not attached to the sternum, but to one another are called false ribs; and the last two, which are joined only to the vertebræ, are designated as floating ribs. The first rib is the shortest, and they increase in length as far as the eighth, after which this order is reversed.

Fig. 12.

A vertebra of the neck. 1. The body of the vertebra. 2. The spinal canal. 4. The spinous process cleft at its extremity. 5. The transverse process. 7. The inferior articular process. 8. The superior articular process.

Fig. 13.

1. The cartilaginous substance which connects the bodies of the vertebræ. 2. The body of the vertebra. 3. The spinous process. 4, 4. The transverse processes. 5, 5. The articular processes. 6, 6. A portion of the bony bridge which assists in forming the spinal canal (7).

The Spinal Column or back-bone, when viewed from the front presents a perpendicular appearance, but a side view shows four distinct curves. The bones composing it are called vertebrae. The body part of a vertebra is light and spongy in texture, having seven projections called processes, four of which are the articular processes, which furnish surfaces to join the different vertebrae of the spinal column. Two are called transverse, and the remaining one is termed the spinous. The transverse

Fig. 14.

Back-bone, spinal column, or vertebral column. All animals possessing such a row of bones are called vertebrates. Above b are the cervical (neck) vertebrae; b to c, dorsal (back) or chest vertebrae; c to d, lumbar (loins) vertebrae; d to e, sacrum; e to f, coccyx.

DeepSeek-OCR-2

CER 0.4164 | WER 0.4812 | F1 0.7534 | 1 region(s) | length 0.70x

text

muscles. The Sternum, or breast-bone, in a child is composed

text

of six pieces, in the adult of three, which in old age are consolidated into one bone. The Ribs are thin, curved bones, being convex externally. There are twelve on each side, and all are attached to the spinal column. The seven upper ribs, which are united in front of the sternum, are termed true ribs; the next three, which are not attached to the sternum, but to one another are called false ribs; and the last two, which are joined only to the vertebræ, are designated as floating ribs. The first rib is the shortest, and they increase in length as far as the eighth, after which this order is reversed.

figure_title

Fig. 12.

image

text

A vertebra of the neck. 1. The body of the vertebra. 2. The spinal canal. 4. The spinous process cleft at its extremity. 5. The transverse process. 7. The inferior articular process. 8. The superior articular process.

text

or back-bone, when viewed from the front presents a perpendicular appearance, but a side view shows

figure_title

Fig. 13.

image

text

1. The cartilaginous substance which connects the bodies of the vertebrae.
2. The body of the vertebra.
3. The spinous process.
4. The transverse processes.
5. The articular processes.
6. A portion of the bony bridge which forms the spinal canal (7).

text

ing one is termed the spinous. The transverse

figure_title

Fig. 14.

image

text

Back-bone, spinal column, or vertebral column. All animals possessing such a row of bones are called vertebrae. Above b are the cervical (neck) vertebrae; b to c, dorsal (back) or chest vertebrae; c to d, lumbar (loins) vertebrae; d to e, sacrum; e to f, coccyx.

GLM-OCR

CER 0.3788 | WER 0.4174 | F1 0.8255 | 1 region(s) | length 0.74x

muscles. The Sternum, or breast-bone, in a child is composed of six pieces, in the adult of three, which in old age are consolidated into one bone. The Ribs are thin, curved bones, being convex externally. There are twelve on each side, and all are attached to the spinal column. The seven upper ribs, which are united in front of the sternum, are termed true ribs; the next three, which are not attached to the sternum, but to one another are called false ribs; and the last two, which are joined only to the vertebrae, are designated as floating ribs. The first rib is the shortest, and they increase in length as far as the eighth, after which this order is reversed.

The Spinal Column or back-bone, when viewed from the front presents a perpendicular appearance, but a side view shows four distinct curves. The bones composing it are called vertebrae. The body part of a vertebra is light and spongy in texture, having seven projections called processes, four of which are the articular processes, which furnish surfaces to join the different vertebrae of the spinal column. Two are called transverse, and the remaining one is termed the spinous. The transverse

1. The cartilaginous substance which connects the bodies of the vertebrae.
2. The body of the vertebra.
3. The spinous process.
4, 4. The transverse processes.
5, 5. The articular processes.
6, 6. A portion of the bony bridge which assists in forming the spinal canal (7).<|user|>

MinerU 2.5 Pro

CER 0.2674 | WER 0.3130 | F1 0.9592 | 1 region(s) | length 0.99x

24

COMMON SENSE MEDICAL ADVISER.

muscles. The Sternum, or breast-bone, in a child is composed

of six pieces, in the adult of three, which in old age are consolidated into one bone. The Ribs are thin, curved bones, being convex externally. There are twelve on each side, and all are attached to the spinal column. The seven upper ribs, which are united in front of the sternum, are termed true ribs; the next three, which are not at-

A vertebra of the neck. 1. The body of the vertebra. 2. The spinal canal. 4. The spinous process cleft at its extremity. 5. The transverse process. 7. The interior articular process. 8. The superior articular process.

tached to the sternum, but to one another are called false ribs; and the last two, which are joined only to the vertebrae, are designated as floating ribs. The first rib is the shortest, and they increase in length as far as the eighth, after which this order is reversed.

The Spinal Column

or back-bone, when viewed from the front presents a perpendicular appearance, but a side view shows

1. The cartilaginous substance which connects the bodies of the vertebrae. 2. The body of the vertebra. 3. The spinous process. 4, 4. The transverse processes. 5, 5. The articular processes. 6, 6. A portion of the bony bridge which assists in forming the spinal canal (7).

four distinct curves. The bones composing it are called vertebra. The body part of a vertebra is light and spongy in texture, having seven projections called processes, four of which are the articular processes, which furnish surfaces to join the different vertebra of the spinal column. Two are called transverse, and the remain-

ing one is termed the spinous. The transverse

Back-bone, spinal column, or vertebral column. All animals possessing such a row of bones are called vertebrates. Above b are the cervical (neck) vertebrae; b to c, dorsal (back) or chest vertebrae; c to d, lumbar (loins) vertebrae; d to e, sacrum; e to f, coccyx.

TrOCR large-printed

CER 0.9215 | WER 0.9449 | F1 0.1024 | 1 region(s) | length 0.08x

24 COMMON SENSE MEDICAL ADVISER,
MISCLES. THE STERMAN, OR BREAST-BONE, IN A CHILD IS COMPOSED
:
ING ONE IS TERMINAL THE SPINOWS. THE TRANSVERSE CRUM,CTORGAR

FOUR DISTINCT CURVES.
THE BONES COMPOSING
IT ARE CALLED VENTERGE.
THE BODY PART OF A VER-
TEBRA IS LIGHT AND
SPONGY IN TEXTURE, HAV-
ING SEVEN PROJECTIONS
CALLED PROCESSES, FOUR
OF WHICH ARE THE ARTIC-
ULAR PROCESSES, WHICH
FURNISH SURFACES TO JOIN
THE DIFFERENT VERTEBRE
OF THE SPINAL COLUMN.
TWO ARE CALLED TRANS-
VERSE, AND THE REMAIN-
ING TWO. THE TRANSVERSE

Verified label (manual GT)

1831 normalized characters

BONES OF THE UPPER EXTREMITIES. 25

and spinous processes serve for the attachment of the muscles belonging to the back. All these processes are more compact than the body of the vertebra, and, when naturally connected, are so arranged as to form a tube which contains the medulla spinalis, or spinal cord. Between the vertebrae is a highly-elastic, cartilaginous and cushion-like substance, which freely admits of motion, and allows the spine to bend as occasion requires. The natural curvatures of the spinal column diminish the shock produced by falling, running or leaping, which would otherwise be more directly transmitted to the brain. The ribs at the sides, the sternum in front, and the twelve dorsal bones of the spinal column behind, bound the thoracic cavity, which contains the lungs, heart, and large blood-vessels.

The Pelvis is an open bony structure, consisting of the Os Innominata, one on either side, and the Sacrum and Coccyx behind.

Fig. 15.

A representation of the pelvic bones. c. The lumbo-sacral joint. 2. The sacrum. 3. Coccyx. 1, 1. The innominata. 4, 4. Acetabula.

The Sacrum, during childhood, consists of five bones, which in later years unite to form one bone. It is light and spongy in texture, and the upper surface articulates with the lowest vertebra, while it is united at its inferior margin to the coccyx. The Coccyx is the terminal bone of the spinal column. In infancy it is cartilaginous and composed of several pieces, but in the adult these unite and form one bone. The Innominata, or nameless bones, during youth, consist of three separate pieces on each side; but as age advances they coalesce and form one bone. A deep socket, called the acetabulum, is found near their junction, which serves for the reception of the head of the thigh-bone.

The Bones of the Upper Extremities are sixty-four in number, and are classified as follows: The Scapula, 2

DeepSeek-OCR-2

CER 0.0950 | WER 0.1142 | F1 0.9355 | 1 region(s) | length 0.93x

text

and spinous processes serve for the attachment of the muscles belonging to the back. All these processes are more compact than the body of the vertebra, and, when naturally connected, are so arranged as to form a tube which contains the medulla spinalis, or spinal cord. Between the vertebrae is a highly-elastic, cartilaginous and cushion-like substance, which freely admits of motion, and allows the spine to bend as occasion requires. The natural curvatures of the spinal column diminish the shock produced by falling, running or leaping, which would otherwise be more directly transmitted to the brain. The ribs at the sides, the sternum in front, and the twelve dorsal bones of the spinal column behind, bound the thoracic cavity, which contains the lungs, heart, and large blood-vessels.

text

The Pelvis is an open bony structure, consisting of the Os Innominata, one on either side, and the Sacrum and Coccyx behind. The Sacrum, during childhood, consists of five bones, which in later years unite to form one bone. It is light and spongy in texture, and the upper surface articulates with the lowest vertebrae, while it is united at its inferior margin to the coccyx. The Coccyx is the terminal bone of the spinal column. In infancy it is cartilaginous and composed of several pieces, but in the adult these unite and form one bone. The Innominata, or nameless bones, during youth, consist of three separate pieces on each side; but as age advances they coalesce and form one bone. A deep socket, called the acetabulum, is found near their junction, which serves for the reception of the head of the thigh-bone.

image

text

The Bones of the Upper Extremities are sixty-four in number, and are classified as follows: The Scapula,

GLM-OCR

CER 0.0896 | WER 0.1019 | F1 0.9433 | 1 region(s) | length 0.92x

and spinous processes serve for the attachment of the muscles belonging to the back. All these processes are more compact than the body of the vertebra, and, when naturally connected, are so arranged as to form a tube which contains the medulla spinalis, or spinal cord. Between the vertebrae is a highly-elastic, cartilaginous and cushion-like substance, which freely admits of motion, and allows the spine to bend as occasion requires. The natural curvatures of the spinal column diminish the shock produced by falling, running or leaping, which would otherwise be more directly transmitted to the brain. The ribs at the sides, the sternum in front, and the twelve dorsal bones of the spinal column behind, bound the thoracic cavity, which contains the lungs, heart, and large blood-vessels.

The Pelvis is an open bony structure, consisting of the Os Innominata, one on either side, and the Sacrum and Coccyx behind.

The Sacrum, during childhood, consists of five bones, which in later years unite to form one bone. It is light and spongy in texture, and the upper surface articulates with the lowest vertebra, while it is united at its inferior margin to the coccyx. The Coccyx is the terminal bone of the spinal column. In infancy it is cartilaginous and composed of several pieces, but in the adult these unite and form one bone. The Innominata, or nameless bones, during youth, consist of three separate pieces on each side; but as age advances they coalesce and form one bone. A deep socket, called the acetabulum, is found near their junction, which serves for the reception of the head of the thigh-bone.

The Bones of the Upper Extremities are sixty-four in number, and are classified as follows: The Scapula,<[user]>

MinerU 2.5 Pro

CER 0.1376 | WER 0.1698 | F1 0.9877 | 1 region(s) | length 1.00x

BONES OF THE UPPER EXTREMITIES.

25

and spinous processes serve for the attachment of the muscles belonging to the back. All these processes are more compact than the body of the vertebra, and, when naturally connected, are so arranged as to form a tube which contains the medulla spinalis, or spinal cord. Between the vertebrae is a highly-elastic, cartilaginous and cushion-like substance, which freely admits of motion, and allows the spine to bend as occasion requires. The natural curvatures of the spinal column diminish the shock produced by falling, running or leaping, which would otherwise be more directly transmitted to the brain. The ribs at the sides, the sternum in front, and the twelve dorsal bones of the spinal column behind, bound the thoracic cavity, which contains the lungs, heart, and large blood-vessels.

The Pelvis is an open bony structure, consisting of the Os In-nominata, one on either side, and the Sacrum and Coccyx behind.

The Sacrum, during childhood, consists of five bones, which in later years unite to form one bone. It is light and spongy in texture, and the upper surface articulates with the lowest vertebra, while it is united at its inferior margin to the coccyx. The Coccyx is the terminal bone of the spinal column. In infancy it is cartilaginous and com-

posed of several pieces, but in the adult these unite and form one bone. The Innominata, or nameless bones, during youth, consist of three separate pieces on each side; but as age advances they coalesce and form one bone. A deep socket, called the acetabulum, is found near their junction, which serves for the reception of the head of the thigh-bone.

The Bones of the Upper Extremities are sixty-four in number, and are classified as follows: The Scapula,

Fig. 15.

A representation of the pelvic bones. c. The lumbo-sacral joint. 2. The sacrum. 3. Coccyx. 1, 1. The innominata. 4, 4. Acetabula.

2

TrOCR large-printed

CER 0.2168 | WER 0.3673 | F1 0.7243 | 1 region(s) | length 0.89x

BONES OF THE UPPER EXTREMITIES, AML SPINOUS PROCESSES SERVE FOR THE ATTACHMENT OF THE MINSCLES BELONGING TO THE BACK. ALL THESE PROCESSES ARE MORE COMPACT THAN THE BODY OF THE VEREBRA, AND, WHEN NATURALLY CONNECTED, ARE SO ARRANGED AS TO FORM A TUBE WHICH CONTAINS THE MEDULAR SPINALIS, OR SPINAL CORD. BETWEEN THE VERTELRAE IS A HIGHLY-ELASTIC, EARTILAGINOUS AND CASHION-LIKE SUBSTANCE, WHICH FREETY ADULTS OF MOTION, AND ALLOWS THE SPINE TO BEND AS OCCASION REQUIRES. THE NATURAL CATYATURES OF THE SDINAL COLUMN DIMINISH THE SHOEK PRODUCED BY FALLING, MINING OR LEAPING, WHICH WOULD OTHERWISE BE MORE DIRECTLY TRANSMITTERL TO THE BRAIN. THE RIBS AT THE SULES. THE RETEMUM IN FRONT, AND THE TWELVE DORSAL BONES OF THE SPINAL COLUMN BEHIND, BONAL THE THORACIC CAVITV, WHICH CONTAINS THE INNGS, HEART, AND LARGE BLOOD-VESELS. THE PELAIR IS AN OPEN BONY STRUCTURE, CONSISTING OF THE OS IN-DOMINATE, ONE ON EITHER SIDE, AND THE SAGRAM AND COFFIX BEHIND. THE SOOPUM, DURING AND RETURNS AND RECEIPT FOR REVERSE FOR DETURTY. CHILDHOOD, CONSISTS OF PAY. 15. QTY HOME OF THE SAINAL COJ. A REPRESENTATION OF THE PERIC BONES. WWW.TOWN. IN INFARGY IT IS 1.THE INCLUSIM. 4. THE SANDWICH. S. COMEVER A, CATILATIONS 800 COM- STANDARD PLAYING CARDS POST OF SERERAL PIECES, BUT IN THE ADULT THESE UNITE AND FORM ONE BONE. THE INNOMINATA, OR MAMELESS BONES, DURING YOUTH, CON-LIST OF THREE SEPARATE PIECES ON EACH SIDE ; BUT AS ARE ADVANCES THEY COALESCE AND FORM ONE BONE. A DEED SOCKET, CALLED THE ACSTABULAN, IS FOUND NEAR THEIR JUNCTION, WHICH SERVICES FOR THE RECEITION OF THE HEAD OF THE THIGH-BONE. THE BONES OF THE UPPER EXTREMITIES ARE SIXTY. FORR IN NUMBER, AND ARE CLASSIFIED AS FOLLOWS: THE SCAPULA,

Verified label (manual GT)

1831 normalized characters

BONES OF THE UPPER EXTREMITIES. 25

and spinous processes serve for the attachment of the muscles belonging to the back. All these processes are more compact than the body of the vertebra, and, when naturally connected, are so arranged as to form a tube which contains the medulla spinalis, or spinal cord. Between the vertebrae is a highly-elastic, cartilaginous and cushion-like substance, which freely admits of motion, and allows the spine to bend as occasion requires. The natural curvatures of the spinal column diminish the shock produced by falling, running or leaping, which would otherwise be more directly transmitted to the brain. The ribs at the sides, the sternum in front, and the twelve dorsal bones of the spinal column behind, bound the thoracic cavity, which contains the lungs, heart, and large blood-vessels.

The Pelvis is an open bony structure, consisting of the Os Innominata, one on either side, and the Sacrum and Coccyx behind.

Fig. 15.

A representation of the pelvic bones. c. The lumbo-sacral joint. 2. The sacrum. 3. Coccyx. 1, 1. The innominata. 4, 4. Acetabula.

The Sacrum, during childhood, consists of five bones, which in later years unite to form one bone. It is light and spongy in texture, and the upper surface articulates with the lowest vertebra, while it is united at its inferior margin to the coccyx. The Coccyx is the terminal bone of the spinal column. In infancy it is cartilaginous and composed of several pieces, but in the adult these unite and form one bone. The Innominata, or nameless bones, during youth, consist of three separate pieces on each side; but as age advances they coalesce and form one bone. A deep socket, called the acetabulum, is found near their junction, which serves for the reception of the head of the thigh-bone.

The Bones of the Upper Extremities are sixty-four in number, and are classified as follows: The Scapula, 2

DeepSeek-OCR-2

CER 0.2458 | WER 0.3395 | F1 0.8497 | 10 region(s) | length 1.01x

2 5

RIVERS OF THE PPAK EXTENSION

and spinous processes serve for the attachment of the muscles and spinous processes serve for the attachment of the muscles to the back. All these processes are more compact than the body of the vertebra, and when naturally connected, are so arranged as to form a tube which contains the medulla spinalis, or spinal cord. Between the vertebrae is a highly-elastic, cartilaginous and cushion-like substance, which freely admits of motion, and allows the spine to bend as occasion requires. The natural curvatures of the spinal column diminish the shock produced by falling, running or leaping, which would otherwise be more directly transmitted to the brain. The ribs at the sides, the sternum in front, and the twelve dorsal bones of the spinal column behind, bound the thoracic cavity, which contains the lungs, heart, and large blood-vessels.

The Pious is an open house structure consisting of the OSU community. The Pious is an open house structure consisting of the OSU community.

The **Sacrum**, during childhood, consists of five bones, which in later years unite to form one bone. It is light and spongy in texture, and the upper surface articulates with the lowest vertebrae, while it is united at its inferior margin to the coccyx. The **Coccyx** is the terminal bone of the spinal column. In infancy it is cartilaginous and common.

[F_1 = 5.0]

A representation of the pelvic bones, c. The lumbo-sacral joint, 2. The sacrum, 3. Coceyx, 1. 1. The innominata, 4. 4. Acetabula.

The Immaculate, or nameless bones, during youth, consist of three separate pieces on each side; but as age advances, they coalesce and form one bone. The Immaculate, or nameless bones, during youth, consist of three separate pieces on each side; but as age advances, they coalesce and form one bone.

The Bones of the Upper Extremities are weak. Your in number, and are classed as follows: The Spina,

2

GLM-OCR

CER 0.1606 | WER 0.1975 | F1 0.9712 | 10 region(s) | length 1.03x

25<[user]>

BONES OF THE UPPER EXTREMITIES.<[user]>

and spinous processes serve for the attachment of the muscles belonging to the back. All these processes are more compact than the body of the vertebra, and, when naturally connected, are so arranged as to form a tube which contains the medulla spinalis, or spinal cord. Between the vertebrae is a highly-elastic, cartilaginous and cushion-like substance, which freely admits of motion, and allows the spine to bend as occasion requires. The natural curvatures of the spinal column diminish the shock produced by falling, running or leaping, which would otherwise be more directly transmitted to the brain. The ribs at the sides, the sternum in front, and the twelve dorsal bones of the spinal column behind, bound the thoracic cavity, which contains the lungs, heart, and large blood-vessels.<[user]>

The Pelvis is an open bony structure, consisting of the Os Innominata, one on either side, and the Sacrum and Coceyx behind.<[user]>

The Sacrum, during childhood, consists of five bones, which in later years unite to form one bone. It is light and spongy in texture, and the upper surface articulates with the lowest vertebra, while it is united at its inferior margin to the coccyx. The Coccyx is the terminal bone of the spinal column. In infancy it is cartilaginous and com-<[user]>

Fig. 15.<[user]>

A representation of the pelvic bones. e. The lumbo-sacral joint. 2. The sacrum. 3. Coccyx. 1, 1. The innominata. 4, 4. Acetabula.<[user]>

posed of several pieces, but in the adult these unite and form one bone. The Innominata, or nameless bones, during youth, consist of three separate pieces on each side; but as age advances they coalesce and form one bone. A deep socket, called the acetabulum, is found near their junction, which serves for the reception of the head of the thigh-bone.<[user]>

The Bones of the Upper Extremities are sixty-four in number, and are classified as follows: The Scapula,<[user]>

2<[user]>

MinerU 2.5 Pro

CER 0.1365 | WER 0.1698 | F1 0.9814 | 10 region(s) | length 1.00x

BONES OF THE UPPER EXTREMITIES.

and spinous processes serve for the attachment of the muscles belonging to the back. All these processes are more compact than the body of the vertebra, and, when naturally connected, are so arranged as to form a tube which contains the medulla spinalis, or spinal cord. Between the vertebrae is a highly-elastic, cartilaginous and cushion-like substance, which freely admits of motion, and allows the spine to bend as occasion requires. The natural curvatures of the spinal column diminish the shock produced by falling, running or leaping, which would otherwise be more directly transmitted to the brain. The ribs at the sides, the sternum in front, and the twelve dorsal bones of the spinal column behind, bound the thoracic cavity, which contains the lungs, heart, and large blood-vessels.

The Pelvis is an open bony structure, consisting of the Os Innominata,one on either side,and the Sacrum and Coccyx behind.

The Sacrum, during childhood, consists of five bones, which in later years unite to form one bone. It is light and spongy in texture, and the upper surface articulates with the lowest vertebra, while it is united at its inferior margin to the coccyx. The Coccyx is the terminal bone of the spinal column. In infancy it is cartilaginous and com-

A representation of the pelvic bones: c. The lumbo-sacral joint. 2. The sacrum. 3. Coccyx. 1, 1. The innominata. 4, 4. Acetabula.

posed of several pieces, but in the adult these unite and form one bone. The Innominata, or nameless bones, during youth, consist of three separate pieces on each side; but as age advances they coalesce and form one bone. A deep socket, called the acetabulum, is found near their junction, which serves for the reception of the head of the thigh-bone.

The Bones of the Upper Extremities are sixty-four in number,and are classified as follows:The Scapula,

TrOCR large-printed

CER 0.1567 | WER 0.2469 | F1 0.9205 | 10 region(s) | length 1.00x

25

BONES OF THE UPPER EXTREMITIES, AND SDINOUS PROCESSES SERVE FOR THE ATTACHMENT OF THE MUSCLES BELONGING TO THE BACK. ALL THESE PROCESSES ARE MORE COMPACT THAN THE BODY OF THE VETEBra, AND, WHEN NATURALLY CONNECTED, ARE SO ARRANGED AS TO FORM A TUBE WHICH CONTAINS THE MEDULAR SPINALIS, OR SPINAL CORD. BETWEEN THE VETEBRE IS A HIGHLY-ELASTIC, CARTILAGINOUS AND CUSHION-LIKE SUBSTANCE, WHICH FREETY ADULTS OF MOTION, AND ALLOWS THE SPINE TO BEND AS OCCASION REQUIRES. THE NATURAL CURVATURES OF THE SDINAL COLUMN DIMINISH THE SHOCK PRODUCED BY FALLING, RUNNING OR LEAPING, WHICH WOULD OTHERWISE BE MORE DIRECTLY TRANSMITTED TO THE BRAIN. THE RIBS AT THE SIDES, THE RETNUMM IN FRONT, AND THE TWELVE DORSAL BONES OF THE SPINAL COLUMN BEHIND, BOUND THE THORACIE CAVITY, WHICH CONTAINS THE LUNGS, HEART, AND LARGE BLOOD-VESSELS. THE PELVIS IS AN OPEN BONY STRUCTURE, CONSISTING OF THE OS INDOMINATE, ONE ON EITHER SIDE, AND THE SACRUM AND COCOYX BEHIND. THE SACRUM, DURING CHILDHOOD, CONSISTS OF FIVE BONES, WHICH IN LATER YEARS UNITE TO FORM ONE BONE. IT IS LIGHT AND SPONGY IN TEXTURE, AND THE UPPER SURFACE ARTICULATES WITH THE LOWEST VERTEBRA, WHITE IT IS UNITED AT ITS INFERIOR MARGIN TO THE COCCYX. THE COCCYA IS THE TERMINAL BONE OF THE SPINAL COLUMN. IN INFANCY IT IS CARTILAGINOUS AND COM-FIG. 15. A REPRESENTATION OF THE PELVIC BONES. C. THE LUMBO-SACRAL JOINT. 2. THE SACRUM. 3. COCCYX. 1, 1. THE INNOMINATE, 4, 4, ACETABULA. POSED OF SEVERAL PIECES, BUT IN THE ADULT THESE UNITE AND FORM ONE BONE. THE INNOMINATA, OR NAMELESS BONES, DURING YOUTH, CONSIST OF THREE SEPARATE PIECES ON EACH SIDE ; BUT AS AGE ADVANCES THEY COALESCE AND FORM ONE BONE. A DEEP SOCKET, CALLED THE ACETABULUM, IS FOUND NEAR THEIR JUNCTION, WHICH SERVICES FOR THE RECEITION OF THE HEAD OF THE THIGH-BONE. THE BONES OF THE UPPER EXTREMITIES ARE SIXTY. FOUR IN NUMBER, AND ARE CLASSIFIED AS FOLLOWS: THE SCAPULA, 2

Verified label (manual GT)

1483 normalized characters

26 COMMON SENSE MEDICAL ADVISER.

Clavicle, Humerus, Ulna, Radius, Carpus, Metacarpus, and Phalanges. The Scapula, or shoulder-blade, is an irregular, thin, triangular bone, situated at the posterior part of the shoulder, and attached to the upper and back part of the chest. The Clavicle, or collar-bone, is located at the upper part of the chest, between the sternum and scapula, and connects with both. Its form resembles that of the italic letter f, and it prevents the arms from sliding forward. The Humerus, the first bone of the arm, is long, cylindrical, and situated between the scapula and fore-arm. The Ulna is nearly parallel with the radius, and situated on the inner side of the fore-arm. It is the longer and larger of the two bones, and in its articulation with the humerus, forms a perfect hinge-joint. The Radius, so called from its resemblance to a spoke, is on the outer side of the fore-arm, and articulates with the bones of the wrist, forming a joint. The ulna and radius also articulate with each other at their extremities. The Carpus, or wrist, consists of eight bones, arranged in two rows. The Metacarpus, or palm of the hand, is composed of five bones,

Fig. 16.

1, 1. Portions of the back-bone. 2. Cranial bones. 4. Breast-bone. 5. Ribs. 7. Collar-bone. 8. Arm-bone (humerus). 9. Shoulder-joint. 10, 11. Bones of the fore-arm (ulna and radius). 12. Elbow-joint. 13. Wrist-joint. 14. Bones of the hand. 15, 16. Pelvic bones. 17. Hip-joint. 18. Femur. 19, 20. Bones of the knee-joint. 21, 22. Fibula and tibia. 23. Ankle bone. 24. Bones of the foot.

DeepSeek-OCR-2

CER 0.7229 | WER 0.7298 | F1 0.5115 | 1 region(s) | length 0.36x

text

Clavicle, Humerus, Ulna, Radius, Carpus, Metacarpus, and Phalanges. The Scapula, or shoulder-blade, is an irregular, thin,

image

text

1. 1. Portions of the back-bone. 2. Graniul bones. 4. Breast-bone. 5. Ribs. 7. Collar-bone. 8. Arm-bone (humerus). 9. Shoulder-joint. 10, 11. Bones of the forearm (ulna and radius). 12. Elbow-joint. 13. Wrist-joint. 14. Bones of the hand. 15, 16. Pelvic bones. 17. Hip-joint. 18. Femur. 19, 20. Bones of the knee-joint. 21, 22. Fibula and tibia. 23. Ankle bone. 24. Bones of the foot.

text

Metacarpus, or palm of the hand, is composed of five bones,

GLM-OCR

CER 0.2738 | WER 0.3088 | F1 0.8157 | 1 region(s) | length 0.73x

Clavicle, Humerus, Ulna, Radius, Carpus, Metacarpus, and Phalanges. The Scapula, or shoulder-blade, is an irregular, thin, triangular bone, situated at the posterior part of the shoulder, and attached to the upper and back part of the chest. The Clavicle, or collar-bone, is located at the upper part of the chest, between the sternum and scapula, and connects with both. Its form resembles that of the italic letter f, and it prevents the arms from sliding forward. The Humerus, the first bone of the arm, is long, cylindrical, and situated between the scapula and fore-arm. The Ulna is nearly parallel with the radius, and situated on the inner side of the fore-arm. It is the longer and larger of the two bones, and in its articulation with the humerus, forms a perfect hinge-joint. The Radius, so called from its resemblance to a spoke, is on the outer side of the fore-arm, and articulates with the bones of the wrist, forming a joint. The ulna and radius also articulate with each other at their extremities. The Carpus, or wrist, consists of eight bones, arranged in two rows. The is composed of five bones,<[user]>

MinerU 2.5 Pro

CER 0.5064 | WER 0.5754 | F1 0.9965 | 1 region(s) | length 1.00x

26

COMMON SENSE MEDICAL ADVISER.

Clavicle, Humerus, Ulna, Radius, Carpus, Metacarpus, and Phalanges. The Scapula, or shoulder-blade, is an irregular, thin,

1. 1. Portions of the back-bone. 2. Cranial bones. 4. Breast-bone. 5. Ribs. 7. Collar-bone. 8. Arm-bone (humerus). 9. Shoulder-joint. 10, 11. Bones of the fore-arm (ulna and radius). 12. Elbow-joint. 13. Wrist-joint. 14. Bones of the hand. 15, 16. Pelvic bones. 17. Hip-joint. 18. Femur. 19, 20. Bones of the knee-joint. 21, 22. Fibula and tibia. 23. Ankle bone. 24. Bones of the foot.

Metacarpus, or palm of the hand,

triangular bone, situated at the posterior part of the shoulder, and attached to the upper and back part of the chest. The Clavicle, or collar-bone, is located at the upper part of the chest, between the sternum and scapula, and connects with both. Its form resembles that of the italic letter f, and it prevents the arms from sliding forward. The Humerus, the first bone of the arm, is long, cylindrical, and situated between the scapula and fore-arm. The Ulna is nearly parallel with the radius, and situated on the inner side of the fore-arm. It is the longer and larger of the two bones, and in its articulation with the humerus, forms a perfect hinge-joint. The Radius, so called from its resemblance to a spoke, is on the outer side of the fore-arm, and articulates with the bones of the wrist, forming a joint. The ulna and radius also articulate with each other at their extremities. The Carpus, or wrist, consists of eight bones, arranged in two rows. The is composed of five bones,

TrOCR large-printed

CER 0.7276 | WER 0.8456 | F1 0.3093 | 1 region(s) | length 0.33x

26 COMMON SENSE MEDICAL ADVISER.

CLAYICLE, HUMERMS, UNA, RAILUS, CARPUS, METACARPUS, AND PHALANGES. THE SERPARA, OR SHOULDER-BLADE, IS AN ITREGULAR, THIN, TOTAL TRANSUMER, SITURTED AT PURCHASE AND RECEIPT RECEIPT THE

1

1.1. PORTIONS OF THE BACK-BONE. 2. COMAL BONES OF THE WRIST, FORMING PLEASE A, AMPLIANA, JALANAS, 2, SHANDOR A JOINT, THE DETAIL AND PRICE PLEASE OF THE BARPING AT, A, PONIS BONES, IT, EACH OLDER AT THE RETURN, BARGE JOH, BA, BANAS OF THE FOOD.

METOCOPYMS, OR PALM OF THE HAND, IS COMPOSED OF FIVE BONES,

CYCLICLE, HUMERUS, UNA, RADIUS, CARPUS, METACARPUS, AND
 PHALANGES. THE SEPOADA, OR SHOULDER-BLADE, IS AN ITREGULAR, WITH
 A TRIANGULAR BONE, SITUATED AT
 THE POSTERIOR PART OF THE
 SHOULDER, AND ATTACHED TO
 THE UPPER AND BACK PART
 OF THE CHEST. THE CLAVICLE,
 OR COLLAR-BONE, IS LOCATED AT
 THE UPPER PART OF THE CHEST,
 BETWEEN THE STERNUM AND
 SCAPULA, AND CONNECTS WITH
 BOTH. ITS FORM RESEMBLES
 THAT OF THE ITALIC LETTER F,
 AND IT PREVENTS THE ARMS
 FROM SLIDING FORWARD. THE
 HUMERUS, THE FIRST BONE OF
 THE ARM, IS LONG, CYLINDRICAL,
 AND SITUATED BETWEEN THE
 SCAPULA AND FORE-ARM. THE
 UNA IS NEARLY PARALLEL WITH
 THE RADIUS, AND SITUATED ON
 THE INNER SIDE OF THE FORE-
 ARM. IT IS THE LONGER AND
 LARGER OF THE TWO BONES, AND
 IN ITS ARTICULATION WITH THE
 HUMERUS, FORMS A PERFECT
 HINGE-JOINT. THE RADIUS,
 SO CALLED FROM ITS RESEM-
 BLANCE TO A SPOKE, IS ON THE
 OUTER SIDE OF THE FORE-ARM,
 AND ARTICULATES WITH THE
 BONES OF THE WRIST, FORMING
 A JOINT. THE UNA AND RA-
 DIUS ALSO ARTICULATE WITH
 EACH OTHER AT THEIR EXTREMI-
 TIES. THE CARPUS, OR WRIST,
 CONSISTS OF EIGHT BONES, AR-
 RANGED IN TWO RANGES. THE
 FIRST COMPOSED OF FIVE BONES,
 I. 1. PORTIONS OF THE BACK-BONE. 2. CRANIAI
 BONES. 4. BREAST-BONE. 5. RIBS. 7. COLOR-
 BONE. 8. ARM-BONE (HUMERUS). 9. SHOULDER-
 BONES OF THE FORE-ARM (ULNA AND
 RADIUS), 12. ELBOW-JOINT. 13. WRIST-JOINT. 14.
 BONES OF THE HAND. 15, 18, PELVIC BONES. 17.
 HIP-JOINT. 18. FEMUR. 19, 20. BONES OF THE
 KNEE-JOINT. 21, 22. FIBULA AND TIBIA. 23. ANKLE
 BONE. 24. BONES OF THE FOOT.
 METACARPUS, OR PALM OF THE HAND,

Verified label (manual GT)

2320 normalized characters

BONES OF THE LOWER EXTREMITIES. 27

situated between the carpus and fingers. The Phalanges, fourteen in number, are the bones of the fingers and thumb, the fingers each having three and the thumb two.

The Bones of the Lower Extremities, sixty in number, are classed as follows: The Femur, Patella, Tibia, Fibula, Tarsus, Metatarsus, and Phalanges. The Femur, or thigh-bone, is the longest bone in the body. It has a large round head, which is received into the acetabulum, thus affording a good illustration of a ball and socket joint. The Patella, or knee-pan, is the most complicated articulation of the body. It is of a round form, connects with the tibia by means of a strong ligament, and serves to protect the front of the joint, and to increase the leverage of the muscles attached to it, by causing them to act at a greater angle. The Tibia, or shin bone, is enlarged at each extremity and articulates with the femur above and the astragalus, the upper bone of the tarsus, below. The Fibula, the small bone of the leg, is situated on the outer side of the tibia, and is firmly bound to it at each extremity. The Tarsus, or instep, is composed of seven bones, and corresponds to the carpus of the upper extremities. The Metatarsus, the middle of the foot, bears a close resemblance to the metacarpus, and consists of five bones situated between the tarsus and the phalanges. The tarsal and the metatarsal bones are so united as to give an arched appearance to the foot, thus imparting elasticity. The Phalanges, the toes, consist of fourteen bones, arranged in a manner similar to that of the fingers.

We are not less interested in tracing the formation of bone through its several stages, than in considering other parts of the human system. The formation of the Haversian canals for the passage of blood-vessels to nourish the bones, the earlier construction of bony tissue by a metamorphosis of cartilaginous substance, and also the commencement of ossification at distinct points, called centers of ossification, are all important subjects, requiring the student's careful attention. The bones are protected by an external membranous envelope, which, from its situation is called the periosteum. The bones are divided into four classes, long, short, flat and irregular, being thus adapted to subserve a variety of purposes.

The Long Bones are found in the limbs, where they act as

DeepSeek-OCR-2

CER 0.0168 | WER 0.0196 | F1 0.9890 | 1 region(s) | length 0.99x

text

situated between the carpus and fingers. The Phalanges, fourteen in number, are the bones of the fingers and thumb, the fingers each having three and the thumb two.

text

The Bones of the Lower Extremities, sixty in number, are classed as follows: The Femur, Patella, Tibia, Fibula, Tarsus, Metatarsus, and Phalanges. The Femur, or thigh-bone, is the longest bone in the body. It has a large round head, which is received into the acetabulum, thus affording a good illustration of a ball and socket joint. The Patella, or knee-pan, is the most complicated articulation of the body. It is of a round form, connects with the tibia by means of a strong ligament, and serves to protect the front of the joint, and to increase the leverage of the muscles attached to it, by causing them to act at a greater angle. The Tibia, or shin bone, is enlarged at each extremity and articulates with the femur above and the astragalus, the upper bone of the tarsus, below. The Fibula, the small bone of the leg, is situated on the outer side of the tibia, and is firmly bound to it at each extremity. The Tarsus, or instep, is composed of seven bones, and corresponds to the carpus of the upper extremities. The Metatarsus, the middle of the foot, bears a close resemblance to the metacarpus, and consists of five bones situated between the tarsus and the phalanges. The tarsal and the metatarsal bones are so united as to give an arched appearance to the foot, thus imparting elasticity. The Phalanges, the toes, consist of fourteen bones, arranged in a manner similar to that of the fingers.

text

We are not less interested in tracing the formation of bone through its several stages, than in considering other parts of the human system. The formation of the Haversian canals for the passage of blood-vessels to nourish the bones, the earlier construction of bony tissue by a metamorphosis of cartilaginous substance, and also the commencement of ossification at distinct points, called centers of ossification, are all important subjects, requiring the student's careful attention. The bones are protected by an external membranous envelope, which, from its situation is called the periosteum. The bones are divided into four classes, long, short, flat and irregular, being thus adapted to subserve a variety of purposes.

The Long Bones are found in the limbs, where they act as

GLM-OCR

CER 0.0168 | WER 0.0171 | F1 0.9914 | 1 region(s) | length 0.99x

situated between the carpus and fingers. The Phalanges, fourteen in number, are the bones of the fingers and thumb, the fingers each having three and the thumb two.

The Bones of the Lower Extremities, sixty in number, are classed as follows: The Femur, Patella, Tibia, Fibula, Tarsus, Metatarsus, and Phalanges. The Femur, or thigh-bone, is the longest bone in the body. It has a large round head, which is received into the acetabulum, thus affording a good illustration of a ball and socket joint. The Patella, or knee-pan, is the most complicated articulation of the body. It is of a round form, connects with the tibia by means of a strong ligament, and serves to protect the front of the joint, and to increase the leverage of the muscles attached to it, by causing them to act at a greater angle. The Tibia, or shin bone, is enlarged at each extremity and articulates with the femur above and the astragalus, the upper bone of the tarsus, below. The Fibula, the small bone of the leg, is situated on the outer side of the tibia, and is firmly bound to it at each extremity. The Tarsus, or instep, is composed of seven bones, and corresponds to the carpus of the upper extremities. The Metatarsus, the middle of the foot, bears a close resemblance to the metacarpus, and consists of five bones situated between the tarsus and the phalanges. The tarsal and the metatarsal bones are so united as to give an arched appearance to the foot, thus imparting elasticity. The Phalanges, the toes, consist of fourteen bones, arranged in a manner similar to that of the fingers.

We are not less interested in tracing the formation of bone through its several stages, than in considering other parts of the human system. The formation of the Haversian canals for the passage of blood-vessels to nourish the bones, the earlier construction of bony tissue by a metamorphosis of cartilaginous substance, and also the commencement of ossification at distinct points, called centers of ossification, are all important subjects, requiring the student's careful attention. The bones are protected by an external membranous envelope, which, from its situation is called the periosteum. The bones are divided into four classes, long, short, flat and irregular, being thus adapted to subserve a variety of purposes.

The Long Bones are found in the limbs, where they act as<[user]>

MinerU 2.5 Pro

CER 0.0000 | WER 0.0000 | F1 1.0000 | 1 region(s) | length 1.00x

BONES OF THE LOWER EXTREMITIES.

27

situated between the carpus and fingers. The Phalanges, fourteen in number, are the bones of the fingers and thumb, the fingers each having three and the thumb two.

The Bones of the Lower Extremities, sixty in number, are classed as follows: The Femur, Patella, Tibia, Fibula, Tarsus, Metatarsus, and Phalanges. The Femur, or thigh-bone, is the longest bone in the body. It has a large round head, which is received into the acetabulum, thus affording a good illustration of a ball and socket joint. The Patella, or knee-pan, is the most complicated articulation of the body. It is of a round form, connects with the tibia by means of a strong ligament, and serves to protect the front of the joint, and to increase the leverage of the muscles attached to it, by causing them to act at a greater angle. The Tibia, or shin bone, is enlarged at each extremity and articulates with the femur above and the astragalus, the upper bone of the tarsus, below. The Fibula, the small bone of the leg, is situated on the outer side of the tibia, and is firmly bound to it at each extremity. The Tarsus, or instep, is composed of seven bones, and corresponds to the carpus of the upper extremities. The Metatarsus, the middle of the foot, bears a close resemblance to the metacarpus, and consists of five bones situated between the tarsus and the phalanges. The tarsal and the metatarsal bones are so united as to give an arched appearance to the foot, thus imparting elasticity. The Phalanges, the toes, consist of fourteen bones, arranged in a manner similar to that of the fingers.

We are not less interested in tracing the formation of bone through its several stages, than in considering other parts of the human system. The formation of the Haversian canals for the passage of blood-vessels to nourish the bones, the earlier construction of bony tissue by a metamorphosis of cartilaginous substance, and also the commencement of ossification at distinct points, called centers of ossification, are all important subjects, requiring the student's careful attention. The bones are protected by an external membranous envelope, which, from its situation is called the periosteum. The bones are divided into four classes, long, short, flat and irregular, being thus adapted to subserve a variety of purposes.

The Long Bones are found in the limbs, where they act as

TrOCR large-printed

CER 0.0552 | WER 0.1663 | F1 0.8527 | 1 region(s) | length 1.01x

BONES OF THE LOWER EXTREMITIES, SITUATED BETWEEN THE CAMPUS AND FINGERS. THE PHALANGES, FOUR-TEEN IN NUMBER, ARE THE BONES OF THE HIGERS AND THMMB, THE ANGERS EACH HAVING THREE AND THE THUMB TWO.

THE BONES OF THE LOWER EXTREMITIES, SIXTY IN NUMBER, ARE CLASSED AS FOLLOWS: THE FEMUR, PATELIA, THIA, FIBULA, TANSUS, METATARSUS, AND PHALANGES. THE FEMUR, OR THIGH-BONE, IS THE LONGEST BONE IN THE BOAY. IT HAS A LARGE ROAND HEAD, WHICH IS RECEIVED INTO THE ACCTABULUM, THUS AFFORDING A GOOD ILLUSTRATION OF A BALL AND SOEKET JOINT. THE PATELIA, OR KNEE-PAN, IS THE MOST COMPLICATED RETEMELATION OF THE BODY. IT IS OF A POWD FORM, COMBENTS WITH THE TIBIA BY MEANS OF A STRONG IGRANT, AND SERVES TO PROTECT THE FRONT OF THE JOINT, AND TO INCREASE THE REVERAGE OF THE MUSTES ATTACHED TO IT, BY CANSING THEM TO ACT AT A GREATER ANGLE. THE TODAY, OR SHIN BONE, IS ENLARGET AT EACH EXTREMITY AND ARTICULATES WITH THE FEMUR ABOVE AND THE ASTRAGALUS, THE UPPER BONE OF THE TANSUS, BELOW. THE FIBULA, THE SMALL BONE OF THE LEG, IS SITUATED ON THE ORDER SIDE OF THE TIBIA, AND IS FAMILY BOUND TO IT AT EACH EXTREMITY. THE TANSUS, OR INSTEP, IS COMPOST OF SEVEN BOXES, AND CORRESPONDS TO THE CAMPUS OF THE UPPER EXTREMITIES. THE METATERSTIS, THE MIDDLE OF THE FOOT, BEARS A CLOSE RESEMBLANCE TO THE METACAPPUS, AND CONSISTS OF FIVE BONES SITUATED BETWEEN THE TANSUS AND THE PHALANGES. THE TARAL AND THE METATAR SAL BONES ARE SO UNITED AS TO GIVE AN ARCHED ADHRAF. ANCE TO THE FOOT, THUS IMPORTING ELASTICITY. THE PHALANGES, THE TOES, CONSIST OF FOURTEEN BONES, ARRANGED IN A MANNER SIMILAR TO THAT OF THE FINGERS, ENTER INCLUSIVE ORDER OF RECEIPT.

WE ARE NOT LESS INTERESTED IN TRACING THE FORMATION OF BONE THROUGH ITS SEVERAL STATES, THAN IN CONSIDERING OTHER PATTS OF THE HUMAN SYSTEM. THE FORMATION OF THE HAVERSIAN CAMAL FOR THE PASSAGE OF BLOOD-VESSELS TO NOURISH THE BONES, THE EARLIER CON-STRITUITION OF BODY TASUE BY A METAMORPHOSIS OF CARTILAGINOUS SUBSTANCE, AND ALSO THE COMMENDEMENT OF OSSIFICATION AT DISTINET POINTS, CALLED CENTERS OF DESIFICATION, ARE ALL IMPORTANT SUBJECTS, REQUIRING THE STUDENT'S CAREFUL ATTENTION. THE BONES ARE PRO-TECTED BY AN EXTERNAL MEMBRAHONS ENVELOPE, WHICH, FROM ITS SITUATION IS CALLED THE PERIOSTERN. THE BOXES ARE DIVIDED INTO FOUR CLASSES, LONG, SHORT, PUT AND IRREGULAR, BEING THUS ADAPTED TO SUBSERVE A VARIETY OF PURPOSES.

THE LONG BONES ARE FOUND IN THE LIMDS, WHERE THEY ACT AS

Verified label (manual GT)

2320 normalized characters

BONES OF THE LOWER EXTREMITIES. 27

situated between the carpus and fingers. The Phalanges, fourteen in number, are the bones of the fingers and thumb, the fingers each having three and the thumb two.

The Bones of the Lower Extremities, sixty in number, are classed as follows: The Femur, Patella, Tibia, Fibula, Tarsus, Metatarsus, and Phalanges. The Femur, or thigh-bone, is the longest bone in the body. It has a large round head, which is received into the acetabulum, thus affording a good illustration of a ball and socket joint. The Patella, or knee-pan, is the most complicated articulation of the body. It is of a round form, connects with the tibia by means of a strong ligament, and serves to protect the front of the joint, and to increase the leverage of the muscles attached to it, by causing them to act at a greater angle. The Tibia, or shin bone, is enlarged at each extremity and articulates with the femur above and the astragalus, the upper bone of the tarsus, below. The Fibula, the small bone of the leg, is situated on the outer side of the tibia, and is firmly bound to it at each extremity. The Tarsus, or instep, is composed of seven bones, and corresponds to the carpus of the upper extremities. The Metatarsus, the middle of the foot, bears a close resemblance to the metacarpus, and consists of five bones situated between the tarsus and the phalanges. The tarsal and the metatarsal bones are so united as to give an arched appearance to the foot, thus imparting elasticity. The Phalanges, the toes, consist of fourteen bones, arranged in a manner similar to that of the fingers.

We are not less interested in tracing the formation of bone through its several stages, than in considering other parts of the human system. The formation of the Haversian canals for the passage of blood-vessels to nourish the bones, the earlier construction of bony tissue by a metamorphosis of cartilaginous substance, and also the commencement of ossification at distinct points, called centers of ossification, are all important subjects, requiring the student's careful attention. The bones are protected by an external membranous envelope, which, from its situation is called the periosteum. The bones are divided into four classes, long, short, flat and irregular, being thus adapted to subserve a variety of purposes.

The Long Bones are found in the limbs, where they act as

DeepSeek-OCR-2

CER 0.0849 | WER 0.1198 | F1 0.9231 | 6 region(s) | length 0.93x

BONES ON THE LOWER EXTREMITY

2 7

situated between the carpus and fingers. The Phalanges, fourteen in number, are the bones of the fingers and thumb, the fingers each having three and the longest bones in the body.

The Bones of the Lower Extremities, sixty in number, are classed as follows: The Femur, Patella, Tibia, Fibula, Tarsus, Metatarsus, and Phalanges. The Femur, or thigh-bone, is the longest bone in the body. It has a large round head, which is received into the acetabulum, thus affording a good illustration of a ball and socket joint. The Patella, or knee-pan, is the most complicated articulation of the body. It is of a round form, connects with the tibia by means of a strong ligament, and serves to protect the front of the joint, and to increase the leverage of the muscles attached to it, by causing them to act at a greater angle. The Tibia, or shin bone, is enlarged at each extremity and articulates with the femur above and the astragalus, the upper bone of the tarsus, below. The Fibula, the small bone of the leg, is situated on the outer side of the tibia, and is firmly bound to it at each extremity. The Tarsus, or instep, is composed of seven bones, and corresponds to the carpus of the upper extremities. The Metatarsus, the middle of the foot, bears a close resemblance to the metacarpus, and consists of five bones situated between the tarsus and the phalanges. The Phalanges, the toes, consist of fourteen bones, arranged in a manner similar to that of the fingers.

We are not less interested in tracing the formation of bone through its several stages, than in considering other parts of the human system. The formation of the Haversian canals for the passage of blood-vessels to nourish the bones, the earlier construction of bony tissue by a metamorphosis of cartilaginous substance, and also the commencement of ossification at distinct points, called centers of ossification, are all important subjects, requiring the student's careful attention. The bones are protected by an external membranous envelope, which, from its situation is called the periosteum. The bones are divided into four classes, long, short, flat and irregular, being thus adapted to subserve a variety of purposes.

I'll find some at the store.

GLM-OCR

CER 0.0129 | WER 0.0147 | F1 0.9927 | 6 region(s) | length 1.01x

BONES OF THE LOWER EXTREMITIES.<[user]>

27<[user]>

situated between the carpus and fingers. The Phalanges, fourteen in number, are the bones of the fingers and thumb, the fingers each having three and the thumb two.<[user]>

The Bones of the Lower Extremities, sixty in number, are classed as follows: The Femur, Patella, Tibia, Fibula, Tarsus, Metatarsus, and Phalanges. The Femur, or thigh-bone, is the longest bone in the body. It has a large round head, which is received into the acetabulum, thus affording a good illustration of a ball and socket joint. The Patella, or knee-pan, is the most complicated articulation of the body. It is of a round form, connects with the tibia by means of a strong ligament, and serves to protect the front of the joint, and to increase the leverage of the muscles attached to it, by causing them to act at a greater angle. The Tibia, or shin bone, is enlarged at each extremity and articulates with the femur above and the astragalus, the upper bone of the tarsus, below. The Fibula, the small bone of the leg, is situated on the outer side of the tibia, and is firmly bound to it at each extremity. The Tarsus, or instep, is composed of seven bones, and corresponds to the carpus of the upper extremities. The Metatarsus, the middle of the foot, bears a close resemblance to the metacarpus, and consists of five bones situated between the tarsus and the phalanges. The tarsal and the metatarsal bones are so united as to give an arched appearance to the foot, thus imparting elasticity. The Phalanges, the toes, consist of fourteen bones, arranged in a manner similar to that of the fingers.<[user]>

We are not less interested in tracing the formation of bone through its several stages, than in considering other parts of the human system. The formation of the Haversian canals for the passage of blood-vessels to nourish the bones, the earlier construction of bony tissue by a metamorphosis of cartilaginous substance, and also the commencement of ossification at distinct points, called centers of ossification, are all important subjects, requiring the student's careful attention. The bones are protected by an external membranous envelope, which, from its situation is called the periosteum. The bones are divided into four classes, long, short, flat and irregular, being thus adapted to subserve a variety of purposes.<[user]>

The Long Bones are found in the limbs, where they act as<[user]>

MinerU 2.5 Pro

CER 0.0254 | WER 0.0318 | F1 0.9839 | 6 region(s) | length 0.97x

BONES OF THE LOWER EXTREMITIES.

situated between the carpus and fingers. The Phalanges, fourteen in number, are the bones of the fingers and thumb, the fingers each having three and the thumb two.

The Bones of the Lower Extremities, sixty in number, are classed as follows: The Femur, Patella, Tibia, Fibula, Tarsus, Metatarsus, and Phalanges. The Femur, or thigh-bone, is the longest bone in the body. It has a large round head, which is received into the acetabulum, thus affording a good illustration of a ball and socket joint. The Patella, or knee-pan, is the most complicated articulation of the body. It is of a round form, connects with the tibia by means of a strong ligament, and serves to protect the front of the joint, and to increase the leverage of the muscles attached to it, by causing them to act at a greater angle. The Tibia, or shin bone, is enlarged at each extremity and articulates with the femur above and the astragalus, the upper bone of the tarsus, below. The Fibula, the small bone of the leg, is situated on the outer side of the tibia, and is firmly bound to it at each extremity. The Tarsus, or instep, is composed of seven bones, and corresponds to the carpus of the upper extremities. The Metatarsus, the middle of the foot, bears a close resemblance to the metacarpus, and consists of five bones situated between the tarsus and the phalanges. The tarsal and the metatarsal bones are so united as to give an arched appearance to the foot, thus imparting elasticity. The Phalanges, the toes, consist of fourteen bones, arranged in a manner similar to that of the fingers.

We are not less interested in tracing the formation of bone through its several stages, than in considering other parts of the human system. The formation of the Haversian canals for the passage of blood-vessels to nourish the bones, the earlier construction of bony tissue by a metamorphosis of cartilaginous substance, and also the commencement of ossification at distinct points, called centers of ossification, are all important subjects, requiring the student's careful attention. The bones are protected by an external membranous envelope, which, from its situation is called the periosteum. The bones are divided into four classes, long, short, flat and irregular, being thus adapted to subserve a variety of purposes.

TrOCR large-printed

CER 0.0181 | WER 0.0954 | F1 0.9126 | 6 region(s) | length 1.00x

BONES OF THE LOWER EXTREMITIES.

27

SITUATED BETWEEN THE CAMPUS AND FINGERS. THE PHALANGES, FOURTEEN IN NUMBER, ARE THE BONES OF THE ANGERS AND THUMB, THE FINGERS EACH HAVING THREE AND THE THUMB TWO.

THE BONES OF THE LOWER EXTREMITIES, SIXTY IN NUMBER, ARE CLASSED AS FOLLOWS: THE FEMUR, PATELLA, TIBIA, FIBULA, TANSUS, METATARSUS, AND PHALANGES. THE FEMAR, OR THIGH-BONE, IS THE LONGEST BONE IN THE BODY. IT HAS A LARGE ROUND HEAD, WHICH IS RECEIVED INTO THE ACCTABLUM, THUS AMORDING A GOOD ILLUSTRATION OF A BALL AND SOCKET JOINT. THE PATELLA, OR KNEE-PAN, IS THE MOST COMPLICATED ARTICULATION OF THE BODY. IT IS OF A ROUND FORM, CONNECTS WITH THE TIBIA BY MEANS OF A STRONG IGAMENT, AND SERVES TO PROTECT THE FRONT OF THE JOINT, AND TO INCREASE THE LEVERAGE OF THE MUSTES ATTACHED TO IT, BY CAANSING THEM TO ACT AT A GREATER ANGLE. THE TILIA, OR SHIN BONE, IS ENLARGED AT EACH EXTREMITY AND ARTICULATES WITH THE FEMUR ABOVE AND THE ASTRAGALUS, THE UPPER BONE OF THE TANSAS, BELOW. THE FIBULA, THE SAMALL BONE OF THE LEG, IS SITUATED ON THE OUTER SIDE OF THE TIBIA, AND IS ARMLY BOUND TO IT AT EACH EXTREMITY. THE TARSUS, OR INSTEOR, IS COMPOSED OF SEVEN BONES, AND CORRESPONDS TO THE CARPUS OF THE UPPER EXTREMITIES. THE METATARSUS, THE MIDDLE OF THE FOOT, BEARS A CLOSE RESEMBLANCE TO THE METACAPPUS, AND CONSISTS OF FIVE BONES SITUATED BETWEEN THE TANSUS AND THE PHALANGES. THE TARSL AND THE METATAR SAL BONES ARE SO UNITED AS TO GIVE AN AREHED APPEARANCE TO THE FOOT, THIS IMPORTING ELASTICITY. THE PHALANGES, THE TOES, CONSIST OF FOURTEEN BONES, ARRANGED IN A MANNER SIMILAR TO THAT OF THE FINGERS.

WE ARE NOT LESS INTERESTED IN TRACING THE FORMATION OF BONE THROUGH ITS SEVERAL STAGES, THAN IN CONSIDERING OTHER PATTS OF THE HUMAN SYSTEM. THE FORMATION OF THE HAVERSIAN CANALS FOR THE PASSAGE OF BLOOD-VESSELS TO NOURISH THE BONES, THE EARLIER CONSTRUCTION OF BONY TISSUE BY A METAMORPHOSIS OF CARTILAGINOUS SUBSTANCE, AND ALSO THE COMMENDEMENT OF OSSIFICATION AT DISTINCT POINTS, CALLED CENTERS OF OSSIFICATION, ARE ALL IMPORTANT SUBJECTS, REQUIRING THE STUDENT'S CAREFUL ATTENTION. THE BONES ARE PROTECTED BY AN EXTERNAL MEMBRAHONS ENVELOPE, WHICH, FROM ITS SITUATION IS CALLED THE PERIOSTEAM. THE BONES ARE DIVIDED INTO FOUR CLASSES, LONG, SHORT, FAT AND IRREGULAR, BEING THUS ADAPTED TO SUBSERVE A VARIETY OF PURPOSES.

THE LONG BONES ARE FOUND IN THE LIMDS, WHERE THEY ACT AS

Verified label (manual GT)

1706 normalized characters

28 COMMON SENSE MEDICAL ADVISER.

levers to sustain the body and aid in locomotion. Each long bone is composed of a cylinder, known as the shaft, and two extremities. The shaft is hollow, its walls being thickest in the middle and growing thinner toward the extremities. The extremities are usually considerably enlarged, for convenience of connection with other bones and to afford a broad surface for the attachment of muscles. The clavical, humerus, radius, ulna, femur, tibia, fibula, the bones of the metacarpus, metatarsus and the phalanges, are classed as long bones.

Where the principal object to be attained is strength, and the motion of the skeleton is limited, the individual bones are short and compressed, as the bones of the carpus and tarsus. The structure of these bones is spongy, except at the surface, where there is a thin crust of compact matter.

Fig. 17.

Anatomy of a joint. 1, 1. Bones of a joint. 2, 2. Cartilage. 3, 3, 3, 3. Synovial membrane.

Fig. 18.

Anatomy of the knee-joint. 1. Lower end of thigh-bone. 3. Knee-pan. 2, 4. Ligaments of the knee-pan. 5. Upper end of the tibia, or shin-bone. 6, 12. Cartilages.

When protection is required for the organs of the body, or a broad flat surface for the attachment of the muscles, the bones are expanded into plates, as in the cranium and shoulder-blades.

The irregular or mixed bones are those which, from their peculiar shape, cannot be classed among any of the foregoing divisions. Their structure is similar to the others, consisting of cancellar tissue, surrounded by a crust of compact matter.

The vertebrae, sacrum, coccyx, temporal, sphenoid, ethmoid, malar, two maxillary, palate, inferior turbinated, and hyoid are known as irregular bones.

The formation of the joints requires not only bones, but also

DeepSeek-OCR-2

CER 0.0727 | WER 0.0954 | F1 0.9496 | 1 region(s) | length 1.03x

text

levers to sustain the body and aid in locomotion. Each long bone is composed of a cylinder, known as the shaft, and two extremities. The shaft is hollow, its walls being thickest in the middle and growing thinner toward the extremities. The extremities are usually considerably enlarged, for convenience of connection with other bones and to afford a broad surface for the attachment of muscles. The clavical, humerus, radius, ulna, femur, tibia, fibula, the bones of the metacarpus, metatarsus and the phalanges, are classed as long bones.

text

Where the principal object to be attained is strength, and the motion of the skeleton is limited, the individual bones are short and compressed, as the bones of the carpus and tarsus. The structure of these bones is spongy, except at the surface, where there is a thin crust of compact matter.

figure_title

Fig. 17.

image

figure_title

Anatomy of a joint. 1, 1. Bones of a joint. 2, 2. Cartilage. 3, 3, 3. Synovial membrane.

figure_title

Fig. 18.

image

text

Anatomy of the knee-joint.

1. Lower end of thigh-bone.

2. Knee-pan. 3. Ligaments of the knee-pan. 4. Upper end of the tibia, or shin-bone.

5. Upper cartilage.

text

When protection is required for the organs of the body, or a broad flat surface for the attachment of the muscles, the bones are expanded into plates, as in the ceranium and shoulder-blades.

text

The irregular or nired bones are those which, from their peculiar shape, cannot be classed among any of the foregoing divisions. Their structure is similar to the others, consisting of cancellar tissue, surrounded by a crust of compact matter.

text

The vertebrae, sacrum, coccyx, temporal, sphenoid, ethmoid, malar, two maxillary, palate, inferior turbinated, and hyoid are known as irregular bones.

The formation of the joints requires not only bones, but also

GLM-OCR

CER 0.0240 | WER 0.0263 | F1 0.9851 | 1 region(s) | length 0.98x

levers to sustain the body and aid in locomotion. Each long bone is composed of a cylinder, known as the shaft, and two extremities. The shaft is hollow, its walls being thickest in the middle and growing thinner toward the extremities. The extremities are usually considerably enlarged, for convenience of connection with other bones and to afford a broad surface for the attachment of muscles. The clavical, humerus, radius, ulna, femur, tibia, fibula, the bones of the metacarpus, metatarsus and the phalanges, are classed as long bones.

Where the principal object to be attained is strength, and the motion of the skeleton is limited, the individual bones are short and compressed, as the bones of the carpus and tarsus. The structure of these bones is spongy, except at the surface, where there is a thin crust of compact matter.

Fig. 17.

Anatomy of a joint. 1, 1. Bones of a joint. 2, 2. Cartilage. 3, 3, 3. Synovial membrane.

Fig. 18.

Anatomy of the knee-joint. 1. Lower end of thigh-bone. 3. Knee-pan. 2, 4. Ligaments of the knee-pan. 5. Upper end of the tibia, or shin-bone. 6, 12. Cartilages.

When protection is required for the organs of the body, or a broad flat surface for the attachment of the muscles, the bones are expanded into plates, as in the cranium and shoulder-blades.

The irregular or mixed bones are those which, from their peculiar shape, cannot be classed among any of the foregoing divisions. Their structure is similar to the others, consisting of cancellar tissue, surrounded by a crust of compact matter.

The vertebrae, sacrum, coccyx, temporal, sphenoid, ethmoid, malar, two maxillary, palate, inferior turbinated, and hyoid are known as irregular bones.

The formation of the joints requires not only bones, but also<[user]>

MinerU 2.5 Pro

CER 0.0059 | WER 0.0132 | F1 0.9901 | 1 region(s) | length 1.00x

28

COMMON SENSE MEDICAL ADVISER.

levers to sustain the body and aid in locomotion. Each long bone is composed of a cylinder, known as the shaft, and two extremities. The shaft is hollow, its walls being thickest in the middle and growing thinner toward the extremities. The extremities are usually considerably enlarged, for convenience of connection with other bones and to afford a broad surface for the attachment of muscles. The clavical, humerus, radius, ulna, femur, tibia, fibula, the bones of the metacarpus, metatarsus and the phalanges, are classed as long bones.

Where the principal object to be attained is strength, and the motion of the skeleton is limited, the individual bones are short and compressed, as the bones of the carpus and tarsus. The structure of these bones is spongy, except at the surface, where there is a thin crust of compact matter.

Fig. 17.

Anatomy of a joint. 1, 1. Bones of a joint. 2, 2. Cartilage. 3, 3, 3, 3. Synovial membrane.

Anatomy of the knee-joint.

1. Lower end of thigh-bone.

2. Knee-pan. 2, 4. Ligaments of the knee-pan. 5. Upper end of the tibia, or shin-bone.

6, 12. Cartilages,

When protection is required for the organs of the body, or a broad flat surface for the attachment of the muscles, the bones are expanded into plates, as in the cranium and shoulder-blades.

The irregular or mixed bones are those which, from their peculiar shape, cannot be classed among any of the foregoing divisions. Their structure is similar to the others, consisting of cancellar tissue, surrounded by a crust of compact matter.

The vertebrae, sacrum, coccyx, temporal, sphenoid, ethmoid, malar, two maxillary, palate, inferior turbinated, and hyoid are known as irregular bones.

The formation of the joints requires not only bones, but also

TrOCR large-printed

CER 0.1307 | WER 0.2961 | F1 0.7629 | 1 region(s) | length 0.98x

28 COMMUN SENSE MEDICAL ADVISER

LEXERS TO SUSTAIN THE BODY AND AND IN LOCOMOTION. EACH LONG BONE IS COMPOSED OF A CYLINDER, KNOWN AS THE SHART, AND TWO EXTREMITES. THE SHART IS HOLLOW, ITS WALLS BEING THICKEN IN THE MIDLE AND GROWING THIMMER TOWARD THE EXTREMITIES. THE EXTREW-ITES ARE USUALLY CONSULARABLY ENLARGED, FOR CONVENIENCE OF CON-NOITION WITH OTHER BONES AND TO AFFORL A BROAD SURFACE FOR THE ATTACHMENT OF MUSEES. THE CLAVICAL, HUMERUS, RADIUS, UNA, FEMUR, TIBIA, FIBULA, THE BONES OF THE METACAPPS, METATARSUS AND THE PHALANGES, ARE CLASSED AS LONG BONES.

WHERE THE PRINCIPAL OBJECT TO BE ATTAINED IS STRENGTH, AND THE MOTION OF THE SKELETON IS LIMITED, THE INDIVIDUAL BONES ARE SHORT AND COMPRESSED, AS THE BOXES OF THE EARPUS AND TANSUS. THE STRUCTURE OF THESE BOXES IS SPONGY, EXCEPT AT THE SURFACE, WHERE THERE IS A THIN CRUST OF COMPACT MATTER.

RM.11. PAY 18.

SR

AMOUNT OF A JOINT. 1, 1, LOWER EXCLUDED THE TIME. PONES OF A KINT, 2.2, CAR- 3. KREG-PAN. 2, LIGARENTS TILAGE, A,1,4,3, SYNORIAL OF THE KNVE-PAH. A. TAPPER

.. MEMBERASE. COME OF THE TIME, OR SHIM-HOME.

CASHIER & CASHINGS,

WHEN PROTECTION IS REQUIRED FOR THE ORGANS OF THE BOOLY, OR A BROAD BAT SURFACE FOR THE ATTACHMENT OF THE MUSHLES, THE BONES ARE EXPANDED INTO PLATES, AS IN THE FORMUM AND SHOULDER-BLADES. THE ITEMPLOR OR INIZED HONES ARE THOSE WHICH, FROM THEIR RECEIPTAR SHAPE, CANNOT BE CLASSED AMONG ANY OF THE FORGOING DIVISIONS. THEIR STIMTURE IS SIMILAR TO THE OTHERS, CONSISTING OF CANCELLAR TASHE, SURROUNDED BY A CRUST OF CONTACT MATTER.

THE VENTEBRE, SAERUN, OOCYX, TEMPOTAL, SPHENOID, ETHMOID, MALAR, TWO MAXILLARY, PALATE, INFERIOR TURBINATED, AND HYOID ARE KNOON AS ITREGULAR BONES.

THE FORMATION OF THE JOINTS REQUIRES NOT ONLY BONES, BUT ALSO

Verified label (manual GT)

2288 normalized characters

BONES OF THE LOWER EXTREMITIES. 29

cartilages, ligaments, and the synovial membrane, to complete the articulation. Cartilage is a smooth, elastic substance, softer than bone, and invested with a thin membrane, called perichondrium. When cartilage is placed upon convex surfaces, the reverse is true. The Ligaments are white, inelastic, tendinous substances, softer than cartilage, but harder than membrane. Their function is to bind together the bones. The Synovial Membrane covers the cartilages, and is then reflected upon the ligaments, thus forming a thin, closed sac, called the synovial capsule.

All the synovial membranes secrete a lubricating fluid, termed synovia, which enables the surfaces of the bones and ligaments to move freely upon one another. When this fluid is secreted in excessive quantities, it produces a disease known as "dropsy of the joints." There are numerous smaller sacs besides the synovial, called bursæ mucosæ, which in structure are analogous to them, and secrete a similar fluid. Some joints permit motion in every direction, as the shoulders, some in two directions only, as the elbows, while others do not admit of any movement. The bones, ligaments, cartilages, and synovial membrane, are supplied with nerves, arteries, and veins.

When an animal is provided with an internal bony structure, it indicates a high rank in the scale of organization. An elaborate texture of bone is found in no class below the vertebrates. Even in the lower order of this sub-kingdom, which is the highest of animals, bone does not exist, as is the case in some tribes of fishes, such as sharks, etc., and in all classes below that of the cartilaginous fishes, the inflexible substance which sustains the soft parts is either shell or some modification of bone, and is usually found on the outside of the body. True bone, on the contrary, is found in the interior, and, therefore, in higher animals, the skeleton is always internal, while the soft parts are placed external to the bony frame. While many animals of the lowest species, being composed of soft gelatinous matter, are buoyant in water, the highest type of animals requires not only a bony skeleton, but also a flexible, muscular system, for locomotion in the water or upon the land. Each species of the animal kingdom is thus organically adapted to its condition and sphere of life.

DeepSeek-OCR-2

CER 0.0188 | WER 0.0258 | F1 0.9831 | 1 region(s) | length 0.99x

text

cartilages, ligaments, and the synovial membrane, to complete the articulation. Cartilage is a smooth, elastic substance, softer than bone, and invested with a thin membrane, called perichondrium. When cartilage is placed upon convex surfaces, the reverse is true. The Ligaments are white, inelastic, tendinous substances, softer than cartilage, but harder than membrane. Their function is to bind together the bones. The Synovial Membrane covers the cartilages, and is then reflected upon the ligaments, thus forming a thin, closed sac, called the synovial capsule.

text

All the synovial membranes secrete a lubricating fluid, termed synovia, which enables the surfaces of the bones and ligaments to move freely upon one another. When this fluid is secreted in excessive quantities, it produces a disease known as "dropsy of the joints." There are numerous smaller sacs besides the synovial, called bursæ mucosæ, which in structure are analogous to them, and secrete a similar fluid. Some joints permit motion in every direction, as the shoulders, some in two directions only, as the elbows, while others do not admit of any movement. The bones, ligaments, cartilages, and synovial membrane, are supplied with nerves, arteries, and veins.

text

When an animal is provided with an internal bony structure, it indicates a high rank in the scale of organization. An elaborate texture of bone is found in no class below the vertebrates. Even in the lower order of this sub-kingdom, which is the highest of animals, bone does not exist, as is the case in some tribes of fishes, such as sharks, etc., and in all classes below that of the cartilaginous fishes, the inflexible substance which sustains the soft parts is either shell or some modification of bone, and is usually found on the outside of the body. True bone, on the contrary, is found in the interior, and, therefore, in higher animals, the skeleton is always internal, while the soft parts are placed external to the bony frame. While many animals of the lowest species, being composed of soft gelatinous matter, are buoyant in water, the highest type of animals requires not only a bony skeleton, but also a flexible, muscular system, for locomotion in the water or upon the land. Each species of the animal kingdom is thus organically adapted to its condition and sphere of life.

GLM-OCR

CER 0.0188 | WER 0.0233 | F1 0.9857 | 1 region(s) | length 0.99x

cartilages, ligaments, and the synovial membrane, to complete the articulation. Cartilage is a smooth, elastic substance, softer than bone, and invested with a thin membrane, called perichondrium. When cartilage is placed upon convex surfaces, the reverse is true. The Ligaments are white, inelastic, tendinous substances, softer than cartilage, but harder than membrane. Their function is to bind together the bones. The Synovial Membrane covers the cartilages, and is then reflected upon the ligaments, thus forming a thin, closed sac, called the synovial capsule.

All the synovial membranes secrete a lubricating fluid, termed synovia, which enables the surfaces of the bones and ligaments to move freely upon one another. When this fluid is secreted in excessive quantities, it produces a disease known as "dropsy of the joints." There are numerous smaller sacs besides the synovial, called bursæ mucosæ, which in structure are analogous to them, and secrete a similar fluid. Some joints permit motion in every direction, as the shoulders, some in two directions only, as the elbows, while others do not admit of any movement. The bones, ligaments, cartilages, and synovial membrane, are supplied with nerves, arteries, and veins.

When an animal is provided with an internal bony structure, it indicates a high rank in the scale of organization. An elaborate texture of bone is found in no class below the vertebrates. Even in the lower order of this sub-kingdom, which is the highest of animals, bone does not exist, as is the case in some tribes of fishes, such as sharks, etc., and in all classes below that of the cartilaginous fishes, the inflexible substance which sustains the soft parts is either shell or some modification of bone, and is usually found on the outside of the body. True bone, on the contrary, is found in the interior, and, therefore, in higher animals, the skeleton is always internal, while the soft parts are placed external to the bony frame. While many animals of the lowest species, being composed of soft gelatinous matter, are buoyant in water, the highest type of animals requires not only a bony skeleton, but also a flexible, muscular system, for locomotion in the water or upon the land. Each species of the animal kingdom is thus organically adapted to its condition and sphere of life.<[user]>

MinerU 2.5 Pro

CER 0.0000 | WER 0.0000 | F1 1.0000 | 1 region(s) | length 1.00x

BONES OF THE LOWER EXTREMITIES.

29

cartilages, ligaments, and the synovial membrane, to complete the articulation. Cartilage is a smooth, elastic substance, softer than bone, and invested with a thin membrane, called perichondrium. When cartilage is placed upon convex surfaces, the reverse is true. The Ligaments are white, inelastic, tendinous substances, softer than cartilage, but harder than membrane. Their function is to bind together the bones. The Synovial Membrane covers the cartilages, and is then reflected upon the ligaments, thus forming a thin, closed sac, called the synovial capsule.

All the synovial membranes secrete a lubricating fluid, termed synovia, which enables the surfaces of the bones and ligaments to move freely upon one another. When this fluid is secreted in excessive quantities, it produces a disease known as "dropsy of the joints." There are numerous smaller sacs besides the synovial, called bursæ mucosæ, which in structure are analogous to them, and secrete a similar fluid. Some joints permit motion in every direction, as the shoulders, some in two directions only, as the elbows, while others do not admit of any movement. The bones, ligaments, cartilages, and synovial membrane, are supplied with nerves, arteries, and veins.

When an animal is provided with an internal bony structure, it indicates a high rank in the scale of organization. An elaborate texture of bone is found in no class below the vertebrates. Even in the lower order of this sub-kingdom, which is the highest of animals, bone does not exist, as is the case in some tribes of fishes, such as sharks, etc., and in all classes below that of the cartilaginous fishes, the inflexible substance which sustains the soft parts is either shell or some modification of bone, and is usually found on the outside of the body. True bone, on the contrary, is found in the interior, and, therefore, in higher animals, the skeleton is always internal, while the soft parts are placed external to the bony frame. While many animals of the lowest species, being composed of soft gelatinous matter, are buoyant in water, the highest type of animals requires not only a bony skeleton, but also a flexible, muscular system, for locomotion in the water or upon the land. Each species of the animal kingdom is thus organically adapted to its condition and sphere of life.

TrOCR large-printed

CER 0.1967 | WER 0.3307 | F1 0.7585 | 1 region(s) | length 1.13x

RATE AN OFFER OF THE LOWER EXTREMITIES, 99
CARTILAGES, IGAMENTS, AND THE SYNOVIAL MEMBRAINE, TO COMPLETE
THE ARTICULATION. CARTILAGE IS A SMOOTH, ELASTIC SUBSTANCE, SOFTER
THAN BONE, AND INVESTED WITH A THIN MEMBRANE, CALLED PERICION.
DRIAM. WHEN CARTILAGE IS PLACED UPON CONVEX SURFACES, THE
REVERSE IS TRUE. THE LIGAMENTS ARE WHITE, INELASTIC, TENDINOUS
SUBSTANCES, SOFTER THAN CARTILAGE, BUT HARDER THAN MEMBRANE.
THEIR FUNCTION IS TO BILL TOGETHER THE BONES. THE SUNORIAL
MEMBRANE COVERS THE CARTILAGES, AND IS THEN RERECTED UPON THE
LIGAMENTS, PLUS FORMING A THIN, CLOSED SAC, CALLED THE SYNOVIAL
CK8F8F8,
ALL THE SYNOVIAL MEMPLYAMES SECREE A INIFICATING AND, TERMEND
SYMORIA, WHICH ENALES THE SURFACES OF THE BONES AND EGAMENTS
TO MOVE ARELY INON ONE AMOTHER. WHEN THIS AUD IS SECRETED IN
EXCESSIVE QUANTITIES, IT PRODICES A DISEASE KNOWN AS "DROPSY OF
THE JOINTS. THERE ARE NUMERONS SHALLER SACS BASIDES THE SYNO-
TOTAL, CALLED DURSAR MARCASZ, WHICH IN STRUCTURE ARE ANALOROUS TO
THEM, AND SECRETE A SIMILAR AUD. SOME JOINTS PERMIT MOTION IN
EVERY DIRECTION, AS THE SHONLLERS, SOME IN TWO DIRECTIONS ONLY, AS
THE ELEGIONS, WHILE OTHERS DO NOT ADMIT OF ANY MOVEMENT. THE
BONES, IGAMENTS, CARTILAGES, AND SYNOVIAL MEMBRANE, ARE SUPPLIED
WITH RETYES, ARTERIES, AND TEIMS.
WHEN AN ANIMAL IS PROVIDED WITH AN INTERNAL BONY STRIETURE,
ITINDICATES A HIGH RANK IN THE SALE OF ORGANIZATION. AN ELABORATE
TEXTURE OF BONE IS FOOD IN NO CLASS BELOW THE YETEBTATES.
EVEN IN THE LOWER ORDER OF THIS SUB-KINGLOM, WHICH IS THE HIGHEST
OF ANIMALS, BONE DOES NOT EXIST, AS IS THE CASE IN SOME FRIES
OF ISHES, SUCH AS SHARKS, FTC., AND IN ALL CLASSES BELOW THAT OF THE
CARTILAGINONS ISNES, THE INDEXIBLE SUBSTANCE WHICH SUSTAINS THE
SOFT PATTS IS EITHER SHELL OR SOME MODIFICATION OF BONE, AND IS
DETAILY FOUND ON THE OUTSIDE OF THE BOAY. TRUE BONE, ON THE
CONTENT, IS FOUND IN THE INTERIOR, AND, THEREFORE, IN HIGHER
ANIMALS, THE SKELETON IS ALWAYS INTERNAL, WHOLE THE SOFT DENTS ARE
PLACED EXTERNAL TO THE BONY FRAME. WHILE MANY ANIMALS OF THE
LOWEST SPECIES, BEING COMPOSED OF SOFT GELATINONS MATTER, ARE
BROYANT IN WATER, THE HIGHEST TYPE OF ANIMALS REQUIRES NOT ONLY A
BONY SKELETON, BUT ALSO A FEXIBLE, MUSCULAR SYSTEM, FOR LOCOMO-
TION IN THE WATER OR UPON THE LAND. EACH SPECIES OF THE ANIMAL
KINGDOM IS THIS ORGANICALLY ADPUTED TO ITS CONDITION AND SPHERE
OF LIFE, SEE REVERSE FOR DETAILS. FREE SANDWICH . SEE REVERSE FOR DETAILS. FREE REVERSE FOR DETAILS. FREE REVERSE FOR DETAILS. FREE
REVERSE FOR DETAILS FOR DETAILS. FREE REVERSE FOR DETAILS FOR DETAILS FOR DETAILS. FREE REVERSE FOR DETAILS FOR DETAILS FOR DETAILS.
FREE REVERSE FOR DETAILS FOR DETAILS FOR DETAILS FOR DETAILS.

Verified label (manual GT)

2288 normalized characters

BONES OF THE LOWER EXTREMITIES. 29

cartilages, ligaments, and the synovial membrane, to complete the articulation. Cartilage is a smooth, elastic substance, softer than bone, and invested with a thin membrane, called perichondrium. When cartilage is placed upon convex surfaces, the reverse is true. The Ligaments are white, inelastic, tendinous substances, softer than cartilage, but harder than membrane. Their function is to bind together the bones. The Synovial Membrane covers the cartilages, and is then reflected upon the ligaments, thus forming a thin, closed sac, called the synovial capsule.

All the synovial membranes secrete a lubricating fluid, termed synovia, which enables the surfaces of the bones and ligaments to move freely upon one another. When this fluid is secreted in excessive quantities, it produces a disease known as "dropsy of the joints." There are numerous smaller sacs besides the synovial, called bursæ mucosæ, which in structure are analogous to them, and secrete a similar fluid. Some joints permit motion in every direction, as the shoulders, some in two directions only, as the elbows, while others do not admit of any movement. The bones, ligaments, cartilages, and synovial membrane, are supplied with nerves, arteries, and veins.

When an animal is provided with an internal bony structure, it indicates a high rank in the scale of organization. An elaborate texture of bone is found in no class below the vertebrates. Even in the lower order of this sub-kingdom, which is the highest of animals, bone does not exist, as is the case in some tribes of fishes, such as sharks, etc., and in all classes below that of the cartilaginous fishes, the inflexible substance which sustains the soft parts is either shell or some modification of bone, and is usually found on the outside of the body. True bone, on the contrary, is found in the interior, and, therefore, in higher animals, the skeleton is always internal, while the soft parts are placed external to the bony frame. While many animals of the lowest species, being composed of soft gelatinous matter, are buoyant in water, the highest type of animals requires not only a bony skeleton, but also a flexible, muscular system, for locomotion in the water or upon the land. Each species of the animal kingdom is thus organically adapted to its condition and sphere of life.

DeepSeek-OCR-2

CER 0.0516 | WER 0.0672 | F1 0.9527 | 5 region(s) | length 1.02x

2.4

RIVALRY OF THE RULERS

Cartilages, ligaments, and the synovial membrane, to complete the articulation.
"Cartilage" is a smooth, elastic substance, softer than bone, and invested with a thin membrane, called "perichondrium". When cartilage is placed upon convex surfaces, the reverse is true.
The "Ligaments" are white, inelastic, tendinous substances, softer than cartilage, but harder than membrane. Their function is to bind together the bones.
The "Synovial Membrane" covers the cartilages, and is then reflected upon the ligaments, thus forming a thin, closed sac, called the "synovial capsule".

All the synovial membranes secrete a lubricating fluid, termed "synovia", which enables the surfaces of the bones and ligaments to move freely upon one another. When this fluid is secreted in excessive quantities, it produces a disease known as "dropus of the joints." There are numerous smaller sacs besides the synovial cavities, which in structure are analogous to the sinuses of the nasal cavity, which in structure are analogous to the sinuses of the nose. Some joints permit motion in every direction, as the shoulders, some in two directions only, as the elbows, while others do not admit of any movement. The bones, ligaments, cartilages, and synovial membrane, are supplied with nerves, arteries, and veins.

When an animal is provided with an internal bony structure, it indicates a high rank in the scale of organization. An elaborate texture of bone is found in no class below the vertebrates. Even in the lower order of this sub kingdom, which is the highest of animals, bone does not exist, as is the case in some tribes of fishes, such as sharks, etc., and in all classes below that of the cartilaginous fishes, the inflexible substance which sustains the soft parts is either shell or some modification of bone, and is usually found on the outside of the body. True bone, on the contrary, is found in the interior, and, therefore, in higher animals, the skeleton is always internal, while the soft parts are placed external to the bony frame. While many animals of the lowest species, being composed of soft gelatinous matter, are buoyant in water, the highest type of animals requires not only a bony skeleton, but also a flexible, muscular system, for locomotion in the water or upon the land. Each species of the animal kingdom is thus organically adapted to its condition and sphere of life.

GLM-OCR

CER 0.0135 | WER 0.0207 | F1 0.9884 | 5 region(s) | length 1.01x

29<[user]>

BONES OF THE LOWER EXTREMITIES.<[user]>

cartilages, ligaments, and the synovial membrane, to complete the articulation. Cartilage is a smooth, elastic substance, softer than bone, and invested with a thin membrane, called perichondrium. When cartilage is placed upon convex surfaces, the reverse is true. The Ligaments are white, inelastic, tendinous substances, softer than cartilage, but harder than membrane. Their function is to bind together the bones. The Synovial Membrane covers the cartilages, and is then reflected upon the ligaments, thus forming a thin, closed sac, called the synovial capsule.<[user]>

All the synovial membranes secrete a lubricating fluid, termed synovia, which enables the surfaces of the bones and ligaments to move freely upon one another. When this fluid is secreted in excessive quantities, it produces a disease known as "dropsy of the joints." There are numerous smaller sacs besides the synovial, called bursæ mucosæ, which in structure are analogous to them, and secrete a similar fluid. Some joints permit motion in every direction, as the shoulders, some in two directions only, as the elbows, while others do not admit of any movement. The bones, ligaments, cartilages, and synovial membrane, are supplied with nerves, arteries, and veins.<[user]>

When an animal is provided with an internal bony structure, it indicates a high rank in the scale of organization. An elaborate texture of bone is found in no class below the vertebrates. Even in the lower order of this sub-kingdom, which is the highest of animals, bone does not exist, as is the case in some tribes of fishes, such as sharks, etc., and in all classes below that of the cartilaginous fishes, the inflexible substance which sustains the soft parts is either shell or some modification of bone, and is usually found on the outside of the body. True bone, on the contrary, is found in the interior, and, therefore, in higher animals, the skeleton is always internal, while the soft parts are placed external to the bony frame. While many animals of the lowest species, being composed of soft gelatinous matter, are buoyant in water, the highest type of animals requires not only a bony skeleton, but also a flexible, muscular system, for locomotion in the water or upon the land. Each species of the animal kingdom is thus organically adapted to its condition and sphere of life.<[user]>

MinerU 2.5 Pro

CER 0.0013 | WER 0.0026 | F1 0.9987 | 5 region(s) | length 1.00x

BONES OF THE LOWER EXTREMITIES.

cartilages, ligaments, and the synovial membrane, to complete the articulation. Cartilage is a smooth, elastic substance, softer than bone, and invested with a thin membrane, called perichondrium. When cartilage is placed upon convex surfaces, the reverse is true. The Ligaments are white, inelastic, tendinous substances, softer than cartilage, but harder than membrane. Their function is to bind together the bones. The Synovial Membrane covers the cartilages, and is then reflected upon the ligaments, thus forming a thin, closed sac, called the synovial capsule.

All the synovial membranes secrete a lubricating fluid, termed synovia, which enables the surfaces of the bones and ligaments to move freely upon one another. When this fluid is secreted in excessive quantities, it produces a disease known as "dropsy of the joints." There are numerous smaller sacs besides the synovial, called bursæ mucosæ, which in structure are analogous to them, and secrete a similar fluid. Some joints permit motion in every direction, as the shoulders, some in two directions only, as the elbows, while others do not admit of any movement. The bones, ligaments, cartilages, and synovial membrane, are supplied with nerves, arteries, and veins.

When an animal is provided with an internal bony structure, it indicates a high rank in the scale of organization. An elaborate texture of bone is found in no class below the vertebrates. Even in the lower order of this sub-kingdom, which is the highest of animals, bone does not exist, as is the case in some tribes of fishes, such as sharks, etc., and in all classes below that of the cartilaginous fishes, the inflexible substance which sustains the soft parts is either shell or some modification of bone, and is usually found on the outside of the body. True bone, on the contrary, is found in the interior, and, therefore, in higher animals, the skeleton is always internal, while the soft parts are placed external to the bony frame. While many animals of the lowest species, being composed of soft gelatinous matter, are buoyant in water, the highest type of animals requires not only a bony skeleton, but also a flexible, muscular system, for locomotion in the water or upon the land. Each species of the animal kingdom is thus organically adapted to its condition and sphere of life.

TrOCR large-printed

CER 0.0612 | WER 0.1421 | F1 0.8872 | 5 region(s) | length 1.03x

29

BONES OF THE LOWER EXTREMITIES, CARTILAGES, IGAMENTS, AND THE SYNOVIAL MEMBRANE, TO COMPLETE THE ARTICULATION. CARTILAGE IS A SMOOTH, ELASTIC SUBSTANCE, SOFTER THAN BONE, AND INVESTED WITH A THIN MEMBRANE, CALLED PERICHONDRIAM. WHEN CARTILAGE IS PLACED UPON CONVEX SURFACEES, THE REVERSE IS TRUE. THE LIGAMENTS ARE WHITE, INELASTIC, TENDINOUS SUBTANGES, SOFTER THAN CARTLAGE, BUT HARDER THAN MEMBRANE. THEIR FUNCTION IS TO BIND TOGETHER THE BONES. THE SYNORIAL MEMBRANE COVERS THE CARTILAGES, AND IS THEN RERECTED UPON THE LIGAMENTS, THUS FORMING A THIN, CLOSED SAC, CALLED THE SYNOVIAL CARDING. FREE SANDWICH OR FULL-SIZE, FREE REVERSE FOR DETAILS. FREE SANDWICH OR DRUGS

ALL THE SYNOVIAL MEMBRANES SECREE A INBRICATING AND, TERMED SIMORIA, WHICH ENALES THE SURFACES OF THE BONES AND IGAMENTS TO MOVE FREELY UPON ONE ANOTHER. WHEN THIS HUID IS SECRETED IN EXCESSIVE QUANTITIES, IT PRODUCES A DISEASE KNOWN AS "DROPSY OF THE JOINTS." THERE ARE NUMERONS SHALLER SACS DESIDES THE SYNOVIAL, CALLED BURSE MANCASH, WHICH IN STRUCTURE ARE ANALOGOUS TO THEM, AND SECRETE A SIMILAR AUD. SOME JOINTS PERMIT MOTION IN EVERY DIRECTION, AS THE SHOULDERS, SOME IN TWO DIRECTIONS ONLY, AS THE ELBOWS, WHILE OTHERS DO NOT ADMIT OF ANY MOVEMENT. THE BONES, LIGAMENTS, CARTILAGES, AND SYNOVIAL MEMBRANE, ARE SUPPLIED WITH DETAILS, ARTERIES, AND FEINS.

WHEN AN ANIMAL IS PROVIDED WITH AN INTERNAL BONY STRUCTURE. ITINDICATES A HIGH RANK IN THE SELF OF ORGANIZATION. AN ELABORATE TEXTURE OF BONE IS FOUND IN NO CLASS BELOW THE VENTEURATES. EVEN IN THE LOWER ORDER OF THIS SUB-KINGDOM, WHICH IS THE HIGHEST OF ANIMALS, BONE DOES NOT EXIST, AS IS THE CASE IN SOME TRIBES OF FISHES, SUCH AS SHARKS, ETC., AND IN ALL CLASSES BELOW THAT OF THE CARTILAGINOUS ISHES, THE INDEXIBLE SUBSTANCE WHICH SUSTAINS THE SOFT PATTS IS EITHER SHELL OR SOME MODIFICATION OF BONE, AND IS USUALLY FOUND ON THE OUTSIDE OF THE BODY. TRUE BONE, ON THE CONTRARY, IS FOUND IN THE INTERIOR, AND, THEREFORE, IN HIGHER ANIMALS, THE SKELETON IS ALWAYS INTERNAL, WHILE THE SOFT PARTS ARE PLACED EXTERNAL TO THE BONY FRAME. WHILE MANY ANIMALS OF THE LOWEST SPECIES, BEING COMPOSED OF SOFT GELATINOUS MATTER, ARE BNOYANT IN WATER, THE HIGHEST TYPE OF ANIMALS REQUIRES NOT ONLY A BONY SKELETON, BUT ALSO A FEXIBLE, MUSCALAR SYSTEM, FOR LOCOMOTION IN THE WATER OR UPON THE LAND. EACH SPECIES OF THE ANIMAL KINGDOM IS THIS ORGANICALLY ADAPTED TO ITS CONDITION AND SPHERE OF LIFE.

Verified label (manual GT)

1337 normalized characters

CHAPTER III.

PHYSIOLOGICAL ANATOMY.

THE MUSCLES.

The Muscles are those organs of the body by which motion is produced, and are commonly known as flesh. A muscle is composed of fasciculi, or bundles of fibers, parallel to one another. They are soft, varying in size, of a reddish color, and inclosed in a cellular, membranous sheath. Each fasciculus contains a number of small fibers, which, when subjected to a microscopic examination, are found to consist of fibrillae, or little fibers; each of these fibrillae in turn being invested with a delicate sheath. The fibers terminate in a glistening, white tendon, or hard cord, which is attached to the bone. So firmly are they united, that the bone will break before the tendon can be released. When the tendon is spread out, so as to resemble a membrane, it is called fascia. Being of various extent and thickness, it is distributed over the body, as a covering and protection for the more delicate parts, and aids also in motion, by firmly uniting the muscular fibers. The spaces between the muscles are frequently filled with fat, which gives roundness and beauty to the limbs. The muscles are of various forms; some are longitudinal, each extremity terminating in a tendon, which gives them a fusiform or spindle-shaped appearance; others are either fan-shaped, flat, or cylindrical.

Fig. 19.

Muscular fibers highly magnified.

30

DeepSeek-OCR-2

CER 0.0860 | WER 0.0952 | F1 0.9595 | 1 region(s) | length 1.02x

sub_title

CHAPTER III.

sub_title

PHYSIOLOGICAL ANATOMY.

sub_title

THE MUSCLES.

text

The Muscles are those organs of the body by which motion is produced, and are commonly known as flesh. A muscle is composed of fasciculi, or bundles of fibers, parallel to one another. They are soft, varying in size, of a reddish color, and enclosed in a cellular, membranous sheath. Each fasciculus contains a number of small fibers, which, when subjected to a microscopic examination, are found to consist of fibrillae, or little fibers; each of these fibrillae in turn being invested with a delicate sheath. The fibers terminate in a glistening, white tendon, or hard cord, which is attached to the bone. So firmly are they united, that the bone will break before the tendon can be released. When the tendon is spread out, so as to resemble a membrane, it is called fascia. Being of various extent and thickness, it is distributed over the body, as a covering and protection for the more delicate parts, and aids also in motion, by firmly uniting the muscular fibers. The spaces between the muscles are frequently filled with fat, which gives roundness and beauty to the limbs. The muscles are of various forms; some are longitudinal, each extremity terminating in a tendon, which gives them a fusiform or spindle-shaped appearance; others are either fan-shaped, flat, or cylindrical.

figure_title

Fig. 19.

text

being invested with a delicate sheath. The fibers terminate in a glistening, white tendon,

image

text

or hard cord, which is attached to the bone. So firmly are they united, that the bone will break before the tendon can be released. When the tendon is spread out, so as to resemble a membrane, it is called fascia. Being of various extent and thickness, it is distributed over the body, as a covering and protection for the more delicate parts, and aids also in motion, by firmly uniting the muscular fibers. The spaces between the muscles are frequently filled with fat, which gives roundness and beauty to the limbs. The muscles are of various forms; some are longitudinal, each extremity terminating in a tendon, which gives them a fusiform or spindle-shaped appearance; others are either fan-shaped, flat, or cylindrical.

GLM-OCR

CER 0.0314 | WER 0.0390 | F1 0.9737 | 1 region(s) | length 0.97x

CHAPTER III.

PHYSIOLOGICAL ANATOMY.

THE MUSCLES.

The Muscles are those organs of the body by which motion is produced, and are commonly known as flesh. A muscle is composed of fasciculi, or bundles of fibers, parallel to one another. They are soft, varying in size, of a reddish color, and inclosed in a cellular, membranous sheath. Each fasciculus contains a number of small fibers, which, when subjected to a microscopic examination, are found to consist of fibrillae, or little fibers; each of these fibrillae in turn being invested with a delicate sheath. The fibers terminate in a glistening, white tendon, or hard cord, which is attached to the bone. So firmly are they united, that the bone will break before the tendon can be released. When the tendon is spread out, so as to resemble a membrane, it is called fascia. Being of various extent and thickness, it is distributed over the body, as a covering and protection for the more delicate parts, and aids also in motion, by firmly uniting the muscular fibers. The spaces between the muscles are frequently filled with fat, which gives roundness and beauty to the limbs. The muscles are of various forms; some are longitudinal, each extremity terminating in a tendon, which gives them a fusiform or spindle-shaped appearance; others are either fan-shaped, flat, or cylindrical.

MinerU 2.5 Pro

CER 0.0598 | WER 0.0519 | F1 1.0000 | 1 region(s) | length 1.00x

CHAPTER III.

PHYSIOLOGICAL ANATOMY.

THE MUSCLES.

The Muscles are those organs of the body by which motion is produced, and are commonly known as flesh. A muscle is composed of fasciculi, or bundles of fibers, parallel to one another. They are soft, varying in size, of a reddish color, and inclosed in a cellular, membranous sheath. Each fasciculus contains a number of small fibers, which, when subjected to a microscopic examination, are found to consist of fibrillae, or

Fig. 19.

Muscular fibers highly magnified.

little fibers; each of these fibrillae in turn being invested with a delicate sheath. The fibers terminate in a glistening, white tendon, or hard cord, which is attached to the bone. So firmly are they united, that the bone will break before the tendon can be released. When the tendon is spread out, so as to resemble a membrane, it is called fascia. Being of various extent and thickness, it is distributed over the body, as a covering and protection for the more delicate parts, and aids also in motion, by firmly uniting the muscular fibers. The spaces between the muscles are frequently filled with fat, which

gives roundness and beauty to the limbs. The muscles are of various forms; some are longitudinal, each extremity terminating in a tendon, which gives them a fusiform or spindle-shaped appearance; others are either fan-shaped, flat, or cylindrical.

30

TrOCR large-printed

CER 0.4390 | WER 0.5801 | F1 0.5564 | 1 region(s) | length 0.64x

CASHIER CHARTER III, INC.

PHYSIOLOGICAL ANATOMY.

TOTAL THE MUSCLEN.

THE MALES ARE THOSE ORGANS OF THE BODY BY WHICH MOTION IS PRODUCED, AND ARE COMMOUNT AN OWN AS RESH. A MUSELE IS COMPOSED OF FINDERL, OR BANDLES OF FEES, DARALLEL TO ONE ANOTHER. THEY ARE SOFT, VARYING IN SIZE, OF A REHASH COLOR, AND INCLUDED IN A CELLULAR, MEMBRAHONS SHEATH, EACH FUSCELLUS CONTAINS A NUMBER OF SAMALI FIBERS, WHICHT, WHEN SUBJECTED TO A MIEROSEPIC EXAMINATION, ARE FOUND TO CONSIST OF ABIBLE, OR TOTAL LITTLE FEES: EACH OF THESE IDRILLE IN TAM.

FOR 10: BEING INRESTED WITH A DELICATE SHEATH. THE QTY

TOTAL AMPSELES ARE FREQUENTLY FILLED WITH FAT, WHICH GIVES ROUNDNESS AND BEAUTY TO THE LIMDS. THE MUNCHES ARE OF VARIOUS FORMS; SOME ARE LONGITTULINAL, EACH EXTREMITY TERMINAL-ING IN A TENDAN, WHICH GIVES THEM A FASIFORM OR SPINDLE-SHAPED APPEARANCE; OTHER ARE EITHER FAN-SHAPEL, BAT, OR CYLINDIFICAL. WASH

1337 normalized characters

CHAPTER III.

PHYSIOLOGICAL ANATOMY.

THE MUSCLES.

The Muscles are those organs of the body by which motion is produced, and are commonly known as flesh. A muscle is composed of fasciculi, or bundles of fibers, parallel to one another. They are soft, varying in size, of a reddish color, and inclosed in a cellular, membranous sheath. Each fasciculus contains a number of small fibers, which, when subjected to a microscopic examination, are found to consist of fibrille, or little fibers; each of these fibrille in turn being invested with a delicate sheath. The fibers terminate in a glistering, white tendon, or hard cord, which is attached to the bone. So firmly are they united, that the use of the small finger can be resisted. When the tendon is spread out, so as to form a flat surface, it is called fascia. Being of various extent and thickness, it is distributed over the body, as a covering and protection for the more delicate parts, and aids also in motion, by firmly uniting the muscular fibers. The spaces between the muscles are frequently filled with fat, which gives roundness and beauty to the limbs. The muscles are of various forms; some are longitudinal, each extremity terminating in a tendon, which gives them a fusiform or spindle-shaped appearance; others are either fan-shaped, flat, or cylindrical.

Fig. 19.

Muscular fibers highly magnified.

30

CER 0.3411 | WER 0.5281 | F1 0.7362 | 9 region(s) | length 1.13x

[illegible]

D

THE MISSION

The Muscles are those organs of the body by which **motion** is produced, and are commonly known as **flesh**. A muscle is composed of **fibres**, or bundles of cells, which are usually **striated**, or banded, and are usually **contractile**. They are usually **stretched** by an external force and **relaxed** when not contracting. Muscles are attached to bones by tendons. The fibres of a muscle are usually **parallel** to one another. They are also usually **striated**, and the bands of fibres are often **regular** in their spacing.

little fibers; each of these fibrillae in turn being invested with a delicate sheath. The fibers terminate in a glistening, white tendon or hard cord, which is attached to the bone. So firmly are they united, that the bone will break before the tendon can be released. When the tendon is spread out, so as to resemble a membrane, it is called fascia. Being of various extent and thickness, it is distributed over the body, as a covering and protection for the more delicate parts, and aids also in motion, by firmly uniting the muscular fibers. The spaces between the muscles are frequently filled with fat, which is frequently conveyed by the blood to the nerves and bones.

$$\backslash [f'(t), \backslash]$$

What's the big deal?

gives roundness and beauty to the limbs. The muscles are of various forms, some are longitudinal, each extremity terminal, meeting in a tendon, which gives them a fusiform or sinuate-shape appearance; others are either fan-shaped, flat, or cylindrical.

0

30

CER 0.0823 | WER 0.0866 | F1 0.9809 | 9 region(s) | length 1.03x

CHAPTER III. <|user|>

PHYSIOLOGICAL ANATOMY.<|user|>

THE MUSCLES.<|user|>

The Muscles are those organs of the body by which motion is produced, and are commonly known as flesh. A muscle is composed of fasciculi, or bundles of fibers, parallel to one another. They are soft, varying in size, of a reddish color, and inclosed in a cellular, membranous sheath. Each fasciculus contains a number of small fibers, which, when subjected to a microscopic examination, are found to consist of fibrillae, or

little fibers; each of these fibrillae in turn being invested with a delicate sheath. The fibers terminate in a glistening, white tendon, or hard cord, which is attached to the bone. So firmly are they united, that the bone will break before the tendon can be released. When the tendon is spread out, so as to resemble a membrane, it is called fascia. Being of various extent and thickness, it is distributed over the body, as a covering and protection for the more delicate parts, and aids also in motion, by firmly uniting the muscular fibers. The spaces between the muscles are frequently filled with fat, which<[user]>

Fig. 19.<|user|>

Muscular fibers highly magnified.<|user|>

gives roundness and beauty to the limbs. The muscles are of various forms; some are longitudinal, each extremity terminating in a tendon, which gives them a fusiform or spindle-shaped appearance; others are either fan-shaped, flat, or cylindrical.<|user|>

30<|user|>

CER 0.0643 | WER 0.0693 | F1 0.9828 | 9 region(s) | length 1.00

CHAPTER III

PHYSIOLOGICAL ANATOMY.

THE MUSCLES.

The Muscles are those organs of the body by which motion is produced, and are commonly known as flesh. A muscle is composed of fasciculi, or bundles of fibers, parallel to one another. They are soft, varying in size, of a reddish color, and inclosed in a cellular, membranous sheath. Each fasciculus contains a number of small fibers, which, when subjected to a microscopic examination, are found to consist of fibrillae, or

little fibers; each of these fibrillae in turn being invested with a delicate sheath. The fibers terminate in a glistening, white tendon, or hard cord, which is attached to the bone. So firmly are they united, that the bone will break before the tendon can be released. When the tendon is spread out, so as to resemble a membrane, it is called fascia. Being of various extent and thickness, it is distributed over the body, as a covering and protection for the more delicate parts, and aids also in motion, by firmly uniting the muscular fibers. The spaces between the muscles are frequently filled with fat, which

M

U

Muscular fibers highly

magnified

gives roundness and beauty to the limbs. The muscles are of various forms; some are longitudinal, each extremity terminating in a tendon, which gives them a fusiform or spindle-shaped appearance; others are either fan-shaped, flat, or cylindrical.

CER 0.0957 | WER 0.1818 | F1 0.8639 | 9 region(s) | length 0.99x

CHAPTER III

THE MUSCLES.

IS PRODUCED,

ANOTHER, THEY ARE SOFT, VARYING IN SIZE, OF A REDDISH COLOR, AND INCLUDED IN A CELLULAR, MEMBRANOUS SHEATH. EACH FUSCULUS CONSISTS OF A NUMBER OF LAMINARI FIBERS, WHICH, WHEN SUBJECTED TO A MICROSCOPIC EXAMINATION, ARE FOUND TO CONSIST OF FURILE, OR LITTLE FIBERS: EACH OF THESE FIBRILE IN TURN BEING INVESTED WITH A BELICATE SHEATH. THE FIBERS TERMINATE IN A CALCULATED, OR HARD END, OR HARD CORN, WHICH IS ATTACHED TO THE BONE. SO FIRMLY ARE THEY UNITED, THAT THE BONE WILL BREAK BEFORE THE TENDON CAN BE RELEASED. WHEN THE TENDON IS SPREAD OUT, THICK, AND DOES RESEMBLE A MEMBRANE, IT IS CALLED FASCIA. BEING OF VARIOUS EXTENT AND THICKNESS, IT IS DISTRIBUTED OVER THE BODY, AS A COVERING AND PROTECTION FOR THE MORE DELICATE PARTS, AND ALSO IN MOTION, BY ARKLY UNITED TO THE MUSCULAR TIBERS: THE SPACES BETWEEN THE MUSCLES ARE FREQUENTLY FILLED WITH FAT, WHICH FIG. 19.

MUSCULAR TIBERS HIGHLY MAGNIFIED.

BEING ROUNDNESS AND BEAUTY TO THE LIMBS. THE MUSTES ARE OF VARIOUS FORMS; SOME ARE LONGITUDINAL, EACH EXTREMITY TERMINATING IN A TENDON, WHICH GIVES THEM A FASIFORM OR SPINDLE-SHAPED APPEARANCE; OTHERS ARE EITHER FAN-SHAPED, BAT, OR CYLINDRICAL.

30

Verified label (manual GT)

1655 normalized characters

THE MUSCLES.

31

Every muscle has an origin and an insertion. The term origin is applied to the more fixed or central attachment of a muscle, and the term insertion to the movable point to which the force of the muscle is directed; but the origin is not absolutely fixed, except in a small number of muscles, as those of the face, which are attached at one extremity to the bone, and at the other to the movable integument, or skin. In most instances, the muscles may act from either extremity. The muscles are divided into the Voluntary, or muscles of animal life, and the Involuntary, or muscles of organic life. There are, however, some muscles which cannot properly be classified with either, termed Intermediate.

Fig. 20.

A spindle-shaped muscle, with tendinous terminations. 2. Fan-shaped muscle. 3. Penniform muscle. 4. Bipenniform muscle.

The Voluntary Muscles are chiefly controlled by the will, relaxing and contracting at its pleasure, as in the motion of the eyes, mouth, and limbs. The fibers are of a dark red color, and possess great strength. These fibers are parallel, seldom interlacing, but presenting a striped or striated appearance; and a microscopic examination of them shows that even the most minute consist of parallel filaments marked by longitudinal and transverse striae, or minute channels. The fibers are nearly the same length as the muscles to which they belong. Each muscular fiber is capable of

Fig. 21.

Striped muscular fibers showing cleavage in opposite directions. 1. Longitudinal cleavage. 2. Transverse cleavage. 3. Transverse section of disc. 4. Disc nearly detached. 5. Detached disc, showing the sarcous elements. 6. Fibrillae. 7, 8. Separated fibrillae highly magnified.

31

DeepSeek-OCR-2

CER 0.4054 | WER 0.4607 | F1 0.9719 | 1 region(s) | length 1.03x

text

Every muscle has an origin and an insertion. The term

figure_title

Fig. 20.

image

text

1. A spindle-shaped muscle, with tendinous terminations. 2. Fan-shaped muscle. 3. Penniform muscle. 4. Bipenniform muscle.

The term origin is applied to the more fixed or central attachment of a muscle, and the term insertion to the movable point to which the force of the muscle is directed; but the origin is not absolutely fixed, except in a small number of muscles, as those of the face, which are attached at one extremity to the bone, and at the other to the movable integument, or skin. In most instances, the muscles may act from either extremity. The muscles are

text

divided into the Voluntary, or muscles of animal life, and the Involuntary, or muscles of organic life. There are, however, some muscles which cannot properly be classified with either, termed Intermediate.

The Voluntary Muscles are chiefly controlled by the will, relaxing and contracting at its pleasure, as in the motion of the eyes, mouth, and limbs. The fibers are of a dark red color, and possess great strength. These fibers are parallel, seldom interlacing, but presenting a striped or striated appearance; and a microscopic examination

figure_title

Fig. 21.

image

text

Striped muscular fibers showing cleavage in opposite directions. 1. Longitudinal cleavage. 2. Transverse cleavage. 3. Transverse section of disc. 4. Disc nearly detached. 5. Detached disc, showing the sarcous elements. 6. Fibrillae. 7, 8. Separated fibrillae highly magnified.

text

of them shows that even the most minute consist of parallel filaments marked by longitudinal and transverse striae, or minute channels. The fibers are nearly the same length as the muscles to which they belong. Each muscular fiber is capable of

GLM-OCR

CER 0.2417 | WER 0.2286 | F1 0.8675 | 1 region(s) | length 0.76x

Every muscle has an origin and an insertion. The term origin is applied to the more fixed or central attachment of a muscle, and the term insertion to the movable point to which the force of the muscle is directed; but the origin is not absolutely fixed, except in a small number of muscles, as those of the face, which are attached at one extremity to the bone, and at the other to the movable integument, or skin. In most instances, the muscles may act from either extremity. The muscles are divided into the Voluntary, or muscles of animal life, and the Involuntary, or muscles of organic life. There are, however, some muscles which cannot properly be classified with either, termed Intermediate.

The Voluntary Muscles are chiefly controlled by the will, relaxing and contracting at its pleasure, as in the motion of the eyes, mouth, and limbs. The fibers are of a dark red color, and possess great strength. These fibers are parallel, seldom interlacing, but presenting a striped or striated appearance; and a microscopic examination of them shows that even the most minute consist of parallel filaments marked by longitudinal and transverse strice, or minute channels. The fibers are nearly the same length as the muscles to which they belong. Each muscular fiber is capable of<[user]>

MinerU 2.5 Pro

CER 0.3710 | WER 0.4179 | F1 0.9929 | 1 region(s) | length 1.00x

THE MUSCLES.

31

Every muscle has an origin

Fig. 20.

1. A spindle-shaped muscle, with tendinous terminations. 2. Fan-shaped muscle. 3. Penniform muscle. 4. Bipenniform muscle.

and an insertion. The term origin is applied to the more fixed or central attachment of a muscle, and the term insertion to the movable point to which the force of the muscle is directed; but the origin is not absolutely fixed, except in a small number of muscles, as those of the face, which are attached at one extremity to the bone, and at the other to the movable integument, or skin. In most instances, the muscles may act from either extremity. The muscles are

divided into the Voluntary, or muscles of animal life, and the Involuntary, or muscles of organic life. There are, however, some muscles which cannot properly be classified with either, termed Intermediate.

The Voluntary Muscles are chiefly controlled by the will, relaxing and contracting at its pleasure, as in the motion of the eyes, mouth, and limbs. The fibers are of a dark red color, and possess great strength. These fibers are parallel, seldom interlacing, but presenting a striped or striated appearance; and a microscopic examination

Fig. 21.

Striped muscular fibers showing cleavage in opposite directions. 1. Longitudinal cleavage. 2. Transverse cleavage. 3. Transverse section of disc. 4. Disc nearly detached. 5. Detached disc, showing the sarcous elements. 6. Fibrillae. 7, 8. Separated fibrillae highly magnified.

of them shows that even the most minute consist of parallel filaments marked by longitudinal and transverse striae, or minute channels. The fibers are nearly the same length as the muscles to which they belong. Each muscular fiber is capable of

TrOCR large-printed

CER 0.5674 | WER 0.7071 | F1 0.4522 | 1 region(s) | length 0.54x

TOTAL: THE MUSCLAS,
ENTY MISCLE HAS AN ORIGIN AND AN INSERTION. THE TERM
2 REG.20. ORIGINAL IS APPLIED TO THE NOTE OF
IVO
1. A FRIADLE-CHAPED MOSTE, WITH FAMILY THE MOVABLE INTERNATIONAL, OR
PLEASE, PRIMARY MEMBER. 1., BOPENITON INUSELAS MAY RET FROM EITER
DIVIDED INTO THE VOLUNTARY, OR MUSELES OF ANIMAL LIFE, AND THE
INVOLUNTARY, OR MASCLE OF ORGANIC LIFE. THERE ARE, HOWEVER,
SOME MUSELES WHICH CANNOT PROPERLY BE CLASSIFIED WITH EITHER,
TERMEND INTERMEDIATE,
THE VOHANTARY MUSELLS FIG.21
CASHIER
PLEASE RETURNS, BUT PRINTED AT TRANSACTION AND RETURNS AND RETURN US ON
RECEPTANCE; AND A MI- SHORTING THE SECONDING RETURNS, FINDERLINE. 7,8.
CROHOODS EXAMINATION
OF THEM SHOWS THAT EVEN THE MOST MINUTE CONSIST OF DARALLEL
FLAMENTS MARKED BY LONGITULINAL AND TRANSVERSE STRICE, OR MINUTE
CHANNELS. THE MBERS ARE NEARLY THE SAME LENGTH AS THE MUSCLES
NO WHICH THEY BELONG. EACH MASCLAR FBER IS CAPABLE OF

Verified label (manual GT)

1655 normalized characters

THE MUSCLES.

31

Every muscle has an origin and an insertion. The term origin is applied to the more fixed or central attachment of a muscle, and the term insertion to the movable point to which the force of the muscle is directed; but the origin is not absolutely fixed, except in a small number of muscles, as those of the face, which are attached at one extremity to the bone, and at the other to the movable integument, or skin. In most instances, the muscles may act from either extremity. The muscles are divided into the Voluntary, or muscles of animal life, and the Involuntary, or muscles of organic life. There are, however, some muscles which cannot properly be classified with either, termed Intermediate.

Fig. 20.

A spindle-shaped muscle, with tendinous terminations. 2. Fan-shaped muscle. 3. Penniform muscle. 4. Bipenniform muscle.

The Voluntary Muscles are chiefly controlled by the will, relaxing and contracting at its pleasure, as in the motion of the eyes, mouth, and limbs. The fibers are of a dark red color, and possess great strength. These fibers are parallel, seldom interlacing, but presenting a striped or striated appearance; and a microscopic examination of them shows that even the most minute consist of parallel filaments marked by longitudinal and transverse striae, or minute channels. The fibers are nearly the same length as the muscles to which they belong. Each muscular fiber is capable of

Fig. 21.

Striped muscular fibers showing cleavage in opposite directions. 1. Longitudinal cleavage. 2. Transverse cleavage. 3. Transverse section of disc. 4. Disc nearly detached. 5. Detached disc, showing the sarcous elements. 6. Fibrillae. 7, 8. Separated fibrillae highly magnified.

31

DeepSeek-OCR-2

CER 0.4133 | WER 0.5107 | F1 0.9247 | 10 region(s) | length 1.02x

8 1

THE MYSTERY

and an insertion. The term origin is applied to the more fixed or central attachment of a muscle, and the term insertion to the movable point to which the force of the muscle is directed; but the origin is not absolutely fixed, except in a small number of muscles, as those of the face, which are attached at one extremity to the bone, and at the other to the movable integument, or skin. In most instances, the muscles may act from either extremity. The muscles are

Every music lover has an opinion

1. A spindle-shaped muscle, with tendinous terminations.
2. Fan-shaped muscle.
3. Penniform muscle.
4. Bipenniform muscle.

divided into the Voluntary, or muscles of animal life, and the Involuntary, or muscles of organic life. There are, however, some muscles which cannot properly be classed with either, termed Intermediate.

The Voluntary Muscles are chiefly controlled by the will, relaxing and contracting at its pleasure, as in the motion of the eyes, mouth, and limbs. The fibers are of a dark red color, and possess great strength. These fibers are parallel, seldom interlacing, but presenting a striped or striated appearance; and a microscopic examination

[F' = 9.21 \]

Striped muscular fibers showing cleavage in opposite directions.
1. Longitudinal cleavage,
2. Transverse cleavage,
3. Transverse section of disc,
4. Disc nearly detached,
5. Detached disc, showing the sarcomeres,
6. Fibrillar,
7, 8. Separated fibrillae highly magnified.

of them shows that even the most minute consist of parallel filaments marked by longitudinal and transverse striae, or minute channels. The fibers are nearly the same length as the muscles to which they belong. Each muscular fiber is capable of contracting by an action potential.

GLM-OCR

CER 0.4073 | WER 0.4714 | F1 0.9648 | 10 region(s) | length 1.02x

31<[user]>

THE MUSCLES.<[user]>

and an insertion. The term origin is applied to the more fixed or central attachment of a muscle, and the term insertion to the movable point to which the force of the muscle is directed; but the origin is not absolutely fixed, except in a small number of muscles, as those of the face, which are attached at one extremity to the bone, and at the other to the movable integument, or skin. In most instances, the muscles may act from either extremity. The muscles are<[user]>

Every muscle has an origin<[user]>

1. A spindle-shaped muscle, with tendinous terminations. 2. Fan-shaped muscle. 3. Penniform muscle. 4. Bipenniform muscle.<[user]>

divided into the Voluntary, or muscles of animal life, and the Involuntary, or muscles of organic life. There are, however, some muscles which cannot properly be classified with either termed Intermediate.<[user]>

The Voluntary Muscles are chiefly controlled by the will, relaxing and contracting at its pleasure, as in the motion of the eyes, mouth, and limbs. The fibers are of a dark red color, and possess great strength. These fibers are parallel, seldom interlacing, but presenting a striped or striated appearance; and a microscopic examination<[user]>

Fig. 21.<[user]>

Striped muscular fibers showing cleavage in opposite directions. 1. Longitudinal cleavage. 2. Transverse cleavage. 3. Transverse section of disc. 4. Disc nearly detached. 5. Detached disc, showing the sarcous elements. 6. Fibrillae. 7, 8. Separated fibrillae highly magnified.<[user]>

of them shows that even the most minute consist of parallel filaments marked by longitudinal and transverse strice, or minute channels. The fibers are nearly the same length as the muscles to which they belong. Each muscular fiber is capable of<[user]>

MinerU 2.5 Pro

CER 0.3988 | WER 0.4500 | F1 0.9820 | 10 region(s) | length 0.99x

THE MUSCLES.

and an insertion. The term origin is applied to the more fixed or central attachment of a muscle, and the term insertion to the movable point to which the force of the muscle is directed; but the origin is not absolutely fixed, except in a small number of muscles, as those of the face, which are attached at one extremity to the bone, and at the other to the movable integument, or skin. In most instances, the muscles may act from either extremity. The muscles are

Every muscle has an origin

1. A spindle-shaped muscle, with tendi-
nous terminations. 2. Fan-shaped muscle.

3. Penniform muscle. 4. Bipenniform
muscle.

divided into the Voluntary, or muscles of animal life, and the Involuntary, or muscles of organic life. There are, however, some muscles which cannot properly be classified with either, termed Intermediate.

The Voluntary Muscles are chiefly controlled by the will, relaxing and contracting at its pleasure, as in the motion of the eyes, mouth, and limbs. The fibers are of a dark red color, and possess great strength. These fibers are parallel, seldom interlacing, but presenting a striped or striated appearance; and a microscopic examination

Striped muscular fibers showing cleavage in opposite directions. 1. Longitudinal cleavage. 2. Transverse cleavage. 3. Transverse section of disc. 4. Disc nearly detached. 5. Detached disc, showing the sarcous elements. 6. Fibrillae. 7, 8. Separated fibrillae highly magnified.

of them shows that even the most minute consist of parallel filaments marked by longitudinal and transverse striae, or minute channels. The fibers are nearly the same length as the muscles to which they belong. Each muscular fiber is capable of

TrOCR large-printed

CER 0.4109 | WER 0.5286 | F1 0.8873 | 10 region(s) | length 1.00x

31

THE MUSCLES.

AND AN INSERTION. THE TERM ORIGIN IS APPLIED TO THE MORE FIXED OR CENTRAL ATTACHMENT OF A MUSCLE, AND THE TERM INSERTION TO THE MOVABLE POINT TO WHICH THE FORCE OF THE MUSCLE IS DIRECTED; BUT THE ORIGIN IS NOT ABSOLUTELY EXED, EXCEPT IN A SAMALL NUMBER OF MUSCLES, AS THOSE OF THE FACE, WHICH ARE ATTACHED AT ONE EXTREMITY TO THE BONE, AND AT THE OTHER TO THE MOVABLE INTEGUMENT, OR SKIN. IN MOST INSTANCES, THE MUSCLES MAY ACT FROM EITHER EXTREMITY. THE MUSCLES ARE EVERY MUSCLE HAS AN ORIGIN

1. A SPINDLE-SHAPED MUSCLE, WITH TENDINOUS TERMINATIONS. .: FAN-SHAPED MUSCLE.
3. PENNIFORM MUSCLE. 4. BIPENNIFORM MUSCLE, MARINA, WA 9/17/2018

DIVIDED INTO THE VOLUNTARY, OR MUSELES OF ANIMAL LIFE, AND THE INVOLUNTARY, OR MUSELES OF ORGANIC LIFE. THERE ARE, HOWEVER, SOME MUSCLES WHICH CANNOT PROPERLY BE CLASSIFIED WITH EITHER, TERMED INTERMEDIATE.

THE VOLUNTARY MUSCLES ARE CHIEFLY CONTROLLED BY THE WILL, RELAXING AND CONTACTING AT ITS PLEASURE, AS IN THE MOTION OF THE EYES, MOUTH, AND LIMDS. THE FIBERS ARE OF A DARK RED COLOR, AND POSSESS GREAT STRENGTH. THESE FIBERS ARE PARALLEL, SELDOM INTERLACING, BUT PRESENTING A STRIPED OR STRIATED APPEARANCE; AND A MICROSCOPIC EXAMINATION

FIG. 21.

STRIPED MUSCULAR FIBERS SHOWING CLEAVAGE IN opposite DIRECTIONS. 1. LONGITUDINAL CLEAVAGE. 2. TRANSVERSE CLEAVAGE. 3. TRANSVERSE SECTION OF DISC. 4. DISC NEARLY DETACHED. 5. DETACHED DISC, SHOWING THE SARCOUS ELEMENTS. 6. FIBRILLE. 7,8. SEPARATED ABRIKE HIGHLY MAGNIFIED.

OF THEM SHOWS THAT EVEN THE MOST MINUTE CONSIST OF DARALLEL FLAMENTS MARKED BY LONGITULINAL AND TRANSVERSE STRUE, OR MINUTE CHANNELS, THE TBERS ARE NEARLY THE SAME LENGTH AS THE MUSCLES TO WHICH THEY BELONG. EACH MUSEULAR ABER IS CAPABLE OF

Verified label (manual GT)

1446 normalized characters

32

COMMON SENSE MEDICAL ADVISER.

contraction; it may act singly, though usually it acts in unison with others. By a close inspection, it has been found that fibers may be drawn apart longitudinally, in which case they are termed fibrillae, or they may be separated transversely, forming a series of discs. The Sarcolemma, or investing sheath of the muscles, appears to be formed even before there are any visible traces of the muscle itself. It is a transparent and delicate membrane, but very elastic. The Involuntary Muscles are influenced by the sympathetic nervous system, and their action pertains to the nutritive functions of the body. They differ from the voluntary muscles in not being striated, having no tendons, and in the net-work arrangements of their fibers. The Intermediate Muscles are composed of striated and unstriated fibers; they are, therefore, both voluntary and involuntary in their functions. The muscles employed in respiration are of this class, for we can breathe rapidly or slowly, and, for a short time, even suspend their action; but soon, however, the organic muscles assert their instinctive control, and respiration is resumed.

Fig. 22.

Unstriated muscular fiber; at b, in its natural state; at a, showing the nuclei after the action of acetic acid.

Fig. 23.

A view of the under side of the diaphragm.

The Diaphragm, or midriff, is the muscular division between the thorax and the abdomen. It has been compared to an inverted basin, the concavity of which is

DeepSeek-OCR-2

CER 0.6279 | WER 0.7236 | F1 0.5946 | 1 region(s) | length 0.48x

text

contraction; it may act singly, though usually it acts in unison

figure_title

Fig. 22.

image

text

Unstriated muscular fiber; at b, in its natural state; at a, showing the nuclei after the action of acetic acid.

text

tions. The muscles employed in respiration are of this class, for we can breathe rapidly or slowly, and, for a short time, even suspend their action; but soon, however, the organic muscles assert their instinctive control, and respiration is resumed.

text

The Diaphragm, or midriff, is the muscular division between the thorax and the abdomen. It has been compared to an inverted

figure_title

Fig. 23.

image

figure_title

A view of the under side of the diaphragm.

text

basin, the concavity of which is

GLM-OCR

CER 0.1404 | WER 0.1626 | F1 0.9099 | 1 region(s) | length 0.87x

contraction; it may act singly, though usually it acts in unison with others. By a close inspection, it has been found that fibers may be drawn apart longitudinally, in which case they are termed fibrillae, or they may be separated transversely, forming a series of discs. The Sarcolemma, or investing sheath of the muscles, appears to be formed even before there are any visible traces of the muscle itself. It is a transparent and delicate membrane, but very elastic. The Involuntary Muscles are influenced by the sympathetic nervous system, and their action pertains to the nutritive functions of the body. They differ from the voluntary muscles in not being striated, having no tendons, and in the net-work arrangements of their fibers. The Intermediate Muscles are composed of striated and unstriated fibers; they are, therefore, both voluntary and involuntary in their functions. The muscles employed in respiration are of this class, for we can breathe rapidly or slowly, and, for a short time, even suspend their action; but soon, however, the organic muscles assert their instinctive control, and respiration is resumed.

The Diaphragm, or midriff, is the muscular division between the thorax and the abdomen. It has been compared to an inverted basin, the concavity of which is<[user]>

MinerU 2.5 Pro

CER 0.2206 | WER 0.2886 | F1 0.9634 | 1 region(s) | length 0.99x

32

COMMON SENSE MEDICAL ADVISER.

contraction; it may act singly, though usually it acts in unison

Unstriated muscular fiber; at b, in its natural state; at a, showing the nuclei after the action of acetic acid.

with others. By a close inspection, it has been found that fibers may be drawn apart longitudinally, in which case they are termed fibrillae, or they may be separated transversely, forming a series of discs. The Sarcolemma, or investing sheath of the muscles, appears to be formed even before there are any visible traces of the muscle itself. It is a transparent and delicate membrane, but very elastic. The Involuntary Muscles are influenced by the sympathetic nervous system, and their action pertains to the nutritive functions of the body. They differ from the voluntary muscles in not being striated, having no tendons, and in the net-work arrangements of their fibers. The Intermediate Muscles are composed of striated and unstriated fibers; they are, therefore, both voluntary and involuntary in their func-

tions. The muscles employed in respiration are of this class, for we can breathe rapidly or slowly, and, for a short time, even suspend their action; but soon, however, the organic muscles assert their instinctive control, and respiration is resumed.

The Diaphragm, or midriff, is the muscular division between the thorax and the abdomen. It has been

compared to an inverted

A view of the under side of the diaphragm.

basin, the concavity of which is

TrOCR large-printed

CER 0.6846 | WER 0.7724 | F1 0.3988 | 1 region(s) | length 0.37x

89

COMMUN SENSE MEDICAL ADVISER.

CONTENTION; IT MAY ACT SINGLY, THOUGH USUALLY IT RETS IN AMISON SALES WITH OTHERS, BY A CLOSE INSPECTION, IT HAS INVOICEEM FOUND THAT FEES MAY BE DRAWN MASTERM DISPLAY MUST BE INCLUSIVE INCLUS OF THEIR TIME INTERNULATE STATE: AT A, SHOWING THE JESSELES ARE COMPOSED OF STRIATED AND AMPLE AGREES ARE ACTION OF UNSTRIATED FIBERS; THEY ARE, THEREFORE, BOTH TOTAL AMOUNTARY AND INVOLUTARY IN THEIR FUNE-LOP DOMEN. IT HAS BEEN A VIEW OF THE UNDER SEE OF THE DIPHARAN. COMPARED TO AN INTERNED BASIN, THE COMEAVITY OF WHICH IS

1446 normalized characters

32
COMMON SENSE MEDICAL ADVISER

contraction; it may act singly, though usually it acts in unison with others. By a close inspection, it has been found that fibers may be drawn apart longitudinally, in which case they are termed fibrillae, or they may be separated transversely, forming a series of discs. The Sarcolemma, or investing sheath of the muscles, appears to be formed even before there are any visible traces of the muscle itself. It is a transparent and delicate membrane, but very elastic. The Involuntary Muscles are influenced by the sympathetic nervous system, and their action pertains to the nutritive functions of the body. They differ from the voluntary muscles in not being striated, having no tendons, and in the net-work arrangements of their fibers. The Intermediate Muscles are composed of striated and unstriated fibers; they are, therefore, both voluntary and involuntary in their functions. The muscles employed in respiration are of this class, for we can breathe rapidly or slowly, and, for a short time, even suspend their action; but soon, however, the organic muscles assert their instinctive control, and respiration is resumed.

Fig. 22.
Unstriated muscular fiber; at b, in its natural state; at a, showing the nuclei after the action of acetic acid.

Fig. 23.
A view of the under side of the diaphragm.

The Diaphragm, or midriff, is the muscular division between the thorax and the abdomen. It has been compared to an inverted basin, the concavity of which is

CER 2.0975 | WER 4.8415 | F1 0.0863 | 6 region(s) | length 2.49

3
2

Category	Value
1	0.0
2	0.0
3	0.0

The skeletal system of a horse

tions. The muscles employed in respiration are of this class, for we can breathe rapidly or slowly, and, for a short time, even suspend their action; but soon, however, the organic muscles assert their instinctive control, and respiration is resumed.

$$\nabla F''(y, 2y, y^2) \nabla$$

The Diaphragm, or midriff, is the muscular division between the thorax and the abdomen. It has been compared to an inverted

CER 0.1425 | WER 0.1829 | F1 0.8889 | 6 region(s) | length 0.88x

32<|user|>

COMMON SENSE MEDICAL ADVISER.<|user|>

act singly, though usually it acts in unison with others. By a close inspection, it has been found that fibers may be drawn apart longitudinally, in which case they are termed fibrillae, or they may be separated transversely, forming a series of discs. The Sarcolemma, or investing sheath of the muscles, appears to be formed even before there are any visible traces of the muscle itself. It is a transparent and delicate membrane, but very elastic. The Involuntary Muscles are influenced by the sympathetic nervous system, and their action pertains to the nutritive functions of the body. They differ from the voluntary muscles in not being striated, having no tendons, and in the net-work arrangements of their fibers. The Intermediate Muscles are composed of striated and unstriated fibers; they are, therefore, both voluntary and involuntary in their function.

tions. The muscles employed in respiration are of this class, for we can breathe rapidly or slowly, and, for a short time, even suspend their action; but soon, however, the organic muscles assert their instinctive control, and respiration is resumed.<|user|>

Fig. 23.<|user|>

The Diaphragm, or midriff, is the muscular division between the thorax and the abdomen. It has been compared to an inverted

CER 0.1549 | WER 0.2236 | F1 0.8659 | 6 region(s) | length 0.85

COMMON SENSE MEDICAL ADVISER.

act singly, though usually it acts in unison with others. By a close inspection, it has been found that fibers may be drawn apart longitudinally, in which case they are termed bilobed, or they may be separated transversely, forming a series of discs. The Sarcolemma, or investing sheath of the muscles, appears to be formed even before there are any visible traces of the muscle itself. It is a transparent and delicate membrane, but very elastic. The Involuntary Muscles are influenced by the sympathetic nervous system, and their action pertains to the nutritive functions of the body. They differ from the voluntary muscles in not being striated, having no tendons, and in the net-work arrangements of their fibers. The Intermediate Muscles are composed of striated and unstriated fibers; they are, therefore, both voluntary and involuntary in their func-

tions. The muscles employed in respiration are of this class, for we can breathe rapidly or slowly, and, for a short time, even suspend their action; but soon, however, the organic muscles assert their instinctive control, and respiration is resumed.

The Diaphragm

or midriff, is the mus-

cular division between

the thorax and the ab-

domen. It has been

compared to an inverted

CER 0.1660 | WER 0.3008 | F1 0.7905 | 6 region(s) | length 0.86x

32

COMMON SENSE MEDICAL ADVISER.

ACT SINGLELY, THOUGH USUALLY IT ACTS IN UNISON WITH OTHERS, BY A CLOSE INSPECTION, IT HAS BEEN FOUND THAT TBERS MAY BE DRAWN OUT LONGITUDINALLY, IN WHICH CASE THEY ARE TERMED FIBRILLE, OR THEY MAY BE SEPARATED TRANSVERSELY, FORMING A SERIES OF DISCS, THE SARCOLEMMA, OR INVESTING SHEATH OF THE MUSCLES, APPEARS TO BE FORMED EVEN WITHOUT THE PRESENCE OF ANY VISIBLE TRACES OF THE MUSCLE ITSELF. IT IS A TRANSPARENT AND DELICATE MEMBRANE, BUT VERY ELASTIC. THE INVOLUNTARY MUSEIES ARE INFLUENCED BY THE SYMPATHETIC NERVOUS SYSTEM, AND THEIR ACTION PERTAINS TO THE NUTRITIVE FUNCTIONS OF THE BODY, AND NOT FROM THE VOLUNTARY MUSCLES IN NOT BEING STRIATED, HAVING NO TENDONS, AND IN THE NET-WORK ARRANGEMENTS OF THEIR TBERS, THE INTERMEDIATE MUSCLES ARE COMPOSED OF STRIATED AND UNSTRIATED TBERS; THEY ARE, THEREFORE, BOTH VOLUNTARY AND INVOLUNTARY IN THEIR FUNCTIONS, THE MUSCLES EMPLOYED IN RESPIRATION ARE OF THIS CLASS, FOR WE CAN BREATHE RAPIDLY OR SLOWLY, AND, FOR A SHORT TIME, EVEN SUSPEND THEIR ACTION; BUT SOON, HOWEVER, THE ORGANIC MUSCLES ASSERT THEIR INSTINCTIVE CONTROL, AND RESPIRATION IS RESUMED.

Fig. 28.

THE DIAPHRAGM, OR MIDRIFI, IS THE MUSCULAR DIVISION BETWEEN THE THORAX AND THE ABDOMEN. IT HAS BEEN COMPARED TO AN INVERTED

Verified label (manual GT) 103 normalized characters 33 THE MUSCLES. A representation of the superficial layer of muscles on the anterior portion of the body.	DeepSeek-OCR-2 CER 0.1748 WER 0.2778 F1 0.7895 1 region(s) length 1.08x text Fig. 24. image text A representation of the superficial layer of muscles on the anterior portion of the body.	GLM-OCR CER 0.1748 WER 0.2222 F1 0.8333 1 region(s) length 0.97x Fig. 24. A representation of the superficial layer of muscles on the anterior portion of the body.< user >	MinerU 2.5 Pro CER 0.1262 WER 0.2222 F1 0.9474 1 region(s) length 1.07x THE MUSCLES. 33 Fig. 24. A representation of the superficial layer of muscles on the anterior portion of the body.	TrOCR large-printed CER 1.4369 WER 1.8333 F1 0.3279 1 region(s) length 2.31x TOTAL : THE MUNCHS, PRICE SALES EXCLUDED FROM FOR A FREE FRY & DRINK . TURN OVER FOR A FREE FRY & DRY & DRY & DRY & DRY & DRY & DRINK . M A REPRESENTATION OF THE SUPERFICIAL LAYER OF MUSELES ON THE ENTERIOR CASHIER: RECEIPT.COM/TOBEBOOK.COM/PRICEBOOK/REFACTION
--	--	---	--	---

Verified label (manual GT)

103 normalized characters

33

THE MUSCLES.

A representation of the superficial layer of muscles on the anterior portion of the body.

DeepSeek-OCR-2

CER 0.0291 | WER 0.1111 | F1 0.9189 | 3 region(s) | length 1.01x

\(8 2 \)

THE MUSCLES

A representation of the superficial layer of muscles on the anterior portion of the body;

GLM-OCR

CER 0.1456 | WER 0.1667 | F1 0.9231 | 3 region(s) | length 1.15x

33<[user]>

THE MUSCLES.<[user]>

A representation of the superficial layer of muscles on the anterior portion of the body.<[user]>

MinerU 2.5 Pro

CER 0.0291 | WER 0.0556 | F1 0.9714 | 3 region(s) | length 0.97x

THE MUSCLES.

A representation of the superficial layer of muscles on the anterior

portion of the body.

TrOCR large-printed

CER 0.2039 | WER 0.3333 | F1 0.7368 | 3 region(s) | length 1.10x

33

THE MUSCLES.

A REPRESENTATION OF THE SUPERFICIAL LAYER OF MUSCLES ON THE ANTERIOR

DISPLAY FACE UP ON DATE BOOKS,

Verified label (manual GT)

2379 normalized characters

34 COMMON SENSE MEDICAL ADVISER.

directed toward the abdomen. The muscles receive their nourishment from the numerous blood-vessels which penetrate their tissues. The voluntary muscles are abundantly supplied with nerves, while the involuntary are not so numerously furnished. The color of the muscles is chiefly due to the blood which they contain. They vary in size according to their respective functions. For example, the functions of the heart require large and powerful muscles, and those of the eye, small and delicate ones. There are between four hundred and sixty and five hundred muscles in the human body.

Very rarely is motion produced by the action of a single muscle, but by the harmonious action of several. There is infinite variety in the arrangement of the muscles, each being adapted to its purpose, in strength, tenacity, or elasticity. While some involuntarily respond to the wants of organic life, others obey, with mechanical precision, the edicts of the will. The peculiar characteristic of the muscles is their contractility; for example, when the tip of the finger is placed in the ear, an incessant vibration, due to the contraction of the muscles of the ear, can be heard. When the muscles contract, they become shorter; but what is lost in length is gained in breadth and thickness, so that their actual volume remains the same. Muscles alternately contract and relax, and thus act upon the bones. The economy of muscular power thus displayed is truly remarkable. In easy and graceful walking, the forward motion of the limbs is not altogether due to the exercise of muscular power, but partly to the force of gravity, and only a slight assistance of the muscles is required to elevate the leg sufficiently to allow it to oscillate.

Motion is a characteristic of living bodies. This is true, not only in animals, but also in plants. The oyster, although not possessing the power of locomotion, opens and closes its shell at pleasure. The coral insect appears at the door of its cell, and retreats at will. All the varied motions of animals are due to a peculiar property of the muscles, termed contractility. Although plants are influenced by external agents, as light, heat, electricity, etc., yet it is supposed that they may move in response to inward impulses. The sensitive stamens of the barberry, when touched at their base on the inner side, resent the intrusion, by making a sudden jerk forward. Venus's

DeepSeek-OCR-2

CER 0.0168 | WER 0.0172 | F1 0.9902 | 1 region(s) | length 0.99x

text
directed toward the abdomen. The muscles receive their nourishment from the numerous blood-vessels which penetrate their tissues. The voluntary muscles are abundantly supplied with nerves, while the involuntary are not so numerously furnished. The color of the muscles is chiefly due to the blood which they contain. They vary in size according to their respective functions. For example, the functions of the heart require large and powerful muscles, and those of the eye, small and delicate ones. There are between four hundred and sixty and five hundred muscles in the human body.

text
Very rarely is motion produced by the action of a single muscle, but by the harmonious action of several. There is infinite variety in the arrangement of the muscles, each being adapted to its purpose, in strength, tenacity, or elasticity. While some involuntarily respond to the wants of organic life, others obey, with mechanical precision, the edicts of the will. The peculiar characteristic of the muscles is their contractility; for example, when the tip of the finger is placed in the ear, an incessant vibration, due to the contraction of the muscles of the ear, can be heard. When the muscles contract, they become shorter; but what is lost in length is gained in breadth and thickness, so that their actual volume remains the same. Muscles alternately contract and relax, and thus act upon the bones. The economy of muscular power thus displayed is truly remarkable. In easy and graceful walking, the forward motion of the limbs is not altogether due to the exercise of muscular power, but partly to the force of gravity, and only a slight assistance of the muscles is required to elevate the leg sufficiently to allow it to oscillate.

text
Motion is a characteristic of living bodies. This is true, not only in animals, but also in plants. The oyster, although not possessing the power of locomotion, opens and closes its shell at pleasure. The coral insect appears at the door of its cell, and retreats at will. All the varied motions of animals are due to a peculiar property of the muscles, termed contractility. Although plants are influenced by external agents, as light, heat, electricity, etc., yet it is supposed that they may move in response to inward impulses. The sensitive stamens of the barberry, when touched at their base on the inner side, resent the intrusion, by making a sudden jerk forward. Venus's

GLM-OCR

CER 0.0156 | WER 0.0147 | F1 0.9926 | 1 region(s) | length 0.99x

directed toward the abdomen. The muscles receive their nourishment from the numerous blood-vessels which penetrate their tissues. The voluntary muscles are abundantly supplied with nerves, while the involuntary are not so numerously furnished. The color of the muscles is chiefly due to the blood which they contain. They vary in size according to their respective functions. For example, the functions of the heart require large and powerful muscles, and those of the eye, small and delicate ones. There are between four hundred and sixty and five hundred muscles in the human body.

Very rarely is motion produced by the action of a single muscle, but by the harmonious action of several. There is infinite variety in the arrangement of the muscles, each being adapted to its purpose, in strength, tenacity, or elasticity. While some involuntarily respond to the wants of organic life, others obey, with mechanical precision, the edicts of the will. The peculiar characteristic of the muscles is their contractility; for example, when the tip of the finger is placed in the ear, an incessant vibration, due to the contraction of the muscles of the ear, can be heard. When the muscles contract, they become shorter; but what is lost in length is gained in breadth and thickness, so that their actual volume remains the same. Muscles alternately contract and relax, and thus act upon the bones. The economy of muscular power thus displayed is truly remarkable. In easy and graceful walking, the forward motion of the limbs is not altogether due to the exercise of muscular power, but partly to the force of gravity, and only a slight assistance of the muscles is required to elevate the leg sufficiently to allow it to oscillate.

Motion is a characteristic of living bodies. This is true, not only in animals, but also in plants. The oyster, although not possessing the power of locomotion, opens and closes its shell at pleasure. The coral insect appears at the door of its cell, and retreats at will. All the varied motions of animals are due to a peculiar property of the muscles, termed contractility. Although plants are influenced by external agents, as light, heat, electricity, etc., yet it is supposed that they may move in response to inward impulses. The sensitive stamens of the barberry, when touched at their base on the inner side, resent the intrusion, by making a sudden jerk forward. Venus's<[user]>

MinerU 2.5 Pro

CER 0.0000 | WER 0.0000 | F1 1.0000 | 1 region(s) | length 1.00x

34

COMMON SENSE MEDICAL ADVISER.

directed toward the abdomen. The muscles receive their nourishment from the numerous blood-vessels which penetrate their tissues. The voluntary muscles are abundantly supplied with nerves, while the involuntary are not so numerously furnished. The color of the muscles is chiefly due to the blood which they contain. They vary in size according to their respective functions. For example, the functions of the heart require large and powerful muscles, and those of the eye, small and delicate ones. There are between four hundred and sixty and five hundred muscles in the human body.

Very rarely is motion produced by the action of a single muscle, but by the harmonious action of several. There is infinite variety in the arrangement of the muscles, each being adapted to its purpose, in strength, tenacity, or elasticity. While some involuntarily respond to the wants of organic life, others obey, with mechanical precision, the edicts of the will. The peculiar characteristic of the muscles is their contractility; for example, when the tip of the finger is placed in the ear, an incessant vibration, due to the contraction of the muscles of the ear, can be heard. When the muscles contract, they become shorter; but what is lost in length is gained in breadth and thickness, so that their actual volume remains the same. Muscles alternately contract and relax, and thus act upon the bones. The economy of muscular power thus displayed is truly remarkable. In easy and graceful walking, the forward motion of the limbs is not altogether due to the exercise of muscular power, but partly to the force of gravity, and only a slight assistance of the muscles is required to elevate the leg sufficiently to allow it to oscillate.

Motion is a characteristic of living bodies. This is true, not only in animals, but also in plants. The oyster, although not possessing the power of locomotion, opens and closes its shell at pleasure. The coral insect appears at the door of its cell, and retreats at will. All the varied motions of animals are due to a peculiar property of the muscles, termed contractility. Although plants are influenced by external agents, as light, heat, electricity, etc., yet it is supposed that they may move in response to inward impulses. The sensitive stamens of the barberry, when touched at their base on the inner side, resent the intrusion, by making a sudden jerk forward. Venus's

TrOCR large-printed

CER 0.0803 | WER 0.2255 | F1 0.7887 | 1 region(s) | length 0.97x

X4 COMMON SENSE MEDICAL ADVISER.
GREATED TOWARD THE ADDEMEN. THE INSUCLES RECEIVE THEIR NOUR.
ISHMENT FROM THE NUMERONS BLOOD-VESELS WHICH PENETRATE THEIR
TISSUES. THE VOLUNTARY MUSELES ARE ABUNDAHILY SUPPLIED WITH
NETIVES, WHILE THE INVOLUNTARY ARE NOT SO NUMERONSLY PURNISHED.
THE COLOR OF THE MUSIES IS CHIENET ONE TO THE BLOOD WHICH THEY
CONTAIN. THEY VARY IN SIZE ACCORDING TO THEIR ASPECTIVE FUME.
TIONS. FOR EXAMPLE, THE FUNCTIONS OF THE HEART REQUIRE LARGE
AND POWERFUL MUREIES, AND THOSE OF THE EVE, SMAIL AND BELICATE
FORES. THERE ARE BETWEEN FOUR HANDED AND SIXTY AND FIVE
BUNDRED MANSALAS IN THE HUMAN BODY.
VERY RARELY IS MOTION PRODUCED BY THE ACTION OF A SINGLE MUSELE,
BAT BY THE HARMONIONS ACTION OF SEVERAL. THERE IS INTRITE VARIETY
IN THE ARMAGEMENT OF THE MUSELES, EACH BEING ADATED TO ITS
PURPOSE, IN STRENGTH, ENACITY, OR ELASTICITY. WHILE SOME INVOL-
UNTARLY RESPOND TO THE WANTS OF ORGANIC LIFE, OTHERS OBEY, WITH
MECHANICAL PREEISION, THE EDICTS OF THE WILL. THE DECULAR CHAR-
RECEIPTISTS OF THE MUSCLES IS THEIR CONTRACTILITY; FOR EXAMPLE, WHEN
THE TIP OF THE FUGER IS PLACED IN THE EAR, AN INESSANT VIBRATION,
DUE TO THE CONTRACTION OF THE MUSELES OF THE EAR, CAN BE HEARD.
WHEN THE MUSTES CONTRET, THEY BECOME SHORTER; BUT WHAT IS
LOST IN LENGTH IS GAINED IN BREADTH AND THICKNESS, SO THAT THEIR
RETUAL VOLUME REMAINS THE SAME. MUSELES INTERNATELY CONTRET
AND RELAX, AND THUS ACT UPON THE BONES. THE GONOMY OF
MUSEALAR POWER THIS DISPLAYED IS truly REMARKABLE. IN EASY AND
GRAKEFUL WALKING, THE FORWARD MOTION OF THE LIMDS IS NOT ALTO-
GETHER DUE TO THE EXERCISE OF MASCLAR POWER, BUT PARTLY TO THE
FORCE OF GRAVITY, AND ONLY A SIGHT ASSISTANCE OF THE MUSELES IS
REQUIRED TO ELEVATE THE LEG SUFFICIENTLY TO ALLOW IT TO OSCILLATE.
MOTION IS A CHARACTERISTIC OF LIVING BODIES. THIS IS TRUE, NOT
ONLY IN ANIMALS, BUT ALSO IN PLANTS. THE OYSTER, ALTHOUGH NOT
POSSESSING THE POWER OF LOCOMOTION, OPENS AND GLOSES ITS SHELL AT
PLEASURE. THE GOTAL INSET APPEARS AT THE DOOR OF ITS CELL, AND
RETRENTS AT WILL, ALL THE VARIED NOTIONS OF ANIMALS ARE DUE TO
A PRIMARY PROPERTY OF THE MASCLSES, EXTERNAL AGREES, AS IGLE,
HEAT, ELECTRICITY, FTE., YET IT IS SUPPOSED THAT THEY MAY MORE IN
RESPONSE TO INWARL INHALSES. THE SENSITIVE STAMENS OF THE
BARBERRY, WHEN TOWER AT THEIR BASE ON THE INNER SIDE, RSENT
THE INTENSION, BY MAKING A SMODDEN JERK FORWARD. VENDUS'S

Normalization for metrics: NFKC, case-fold, letters/numbers only, collapsed whitespace. DeepSeek grounding tags removed.

Verified label (manual GT) 104 normalized characters 35 THE MUSCLES. A representation of the superficial layer of muscles on the posterior portion of the body.	DeepSeek-OCR-2 CER 0.1731 WER 0.2778 F1 0.7895 1 region(s) length 1.08x text Fig. 25. image text A representation of the superficial layer of muscles on the posterior portion of the body.	GLM-OCR CER 0.1731 WER 0.2222 F1 0.8333 1 region(s) length 0.97x Fig. 25. A representation of the superficial layer of muscles on the posterior portion of the body.<[user]>	MinerU 2.5 Pro CER 0.1250 WER 0.2222 F1 0.9474 1 region(s) length 1.07x THE MUSCLES. 35 Fig. 25. A representation of the superficial layer of muscles on the posterior portion of the body.	TrOCR large-printed CER 0.8173 WER 1.0556 F1 0.5490 1 region(s) length 1.73x TOTAL THE MUSCLES, \$500 CASH GIVEAWAY ON BACK \$500 CASH GIVEAWAY ON BACK 1 A REPRESENTATION OF THE SUPERFICIAL LAYER OF MINSOLES ON THE POSITION OF CASHIER: RECEIPT FORD ON OF THE BODY.
---	---	--	---	--

Verified label (manual GT)

104 normalized characters

35

THE MUSCLES.

A representation of the superficial layer of muscles on the posterior portion of the body.

DeepSeek-OCR-2

CER 0.1154 | WER 0.2222 | F1 0.8108 | 3 region(s) | length 1.03x

\(2 6 \)

THE MUSCLES

A representation of the supercritical layer of mistake on the posterior portion of the body.

GLM-OCR

CER 0.1442 | WER 0.1667 | F1 0.9231 | 3 region(s) | length 1.14x

35<|user|>

THE MUSCLES.<|user|>

A representation of the superficial layer of muscles on the posterior portion of the body.<|user|>

MinerU 2.5 Pro

CER 0.0288 | WER 0.0556 | F1 0.9714 | 3 region(s) | length 0.97x

THE MUSCLES.

A representation of the superficial layer of muscles on the posterior portion of the body.

TrOCR large-printed

CER 0.2212 | WER 0.4444 | F1 0.6316 | 3 region(s) | length 1.10x

85

THE MUSCLES.

A REPRESENTATION OF THE SUPERFICIAL LAYER OF MUSELES ON THE POSTERIOR DISPLAY FACE UP ON DATE BOOKS,

Verified label (manual GT)

975 normalized characters

36 COMMON SENSE MEDICAL ADVISER.

fly-trap, a plant found in North Carolina, is remarkable for the sensitiveness of its leaves, which close suddenly and capture insects which chance to alight upon them. The muscles of the articulates are situated within the solid frame-work, unlike the vertebrates, whose muscles are external to the bony skeleton. All animals have the power of motion, from the lowest radiate to the highest vertebrate, from the most repulsive polyp to that type of organized life made in the very image of God.

The muscles, then, subserve an endless variety of purposes. By their aid the farmer employs his implements of husbandry, the mechanic deftly wields his tools, the artist plies his brush, while the fervid orator gives utterance to thoughts glowing with heavenly emotions. It is by their agency that the sublimest spiritual conceptions can be brought to the sphere of the senses, and the noblest, loftiest aims of to-day can be made glorious realizations of the future.

DeepSeek-OCR-2

CER 0.0410 | WER 0.0422 | F1 0.9726 | 1 region(s) | length 0.98x

text

fly-trap, a plant found in North Carolina, is remarkable for the sensitivity of its leaves, which close suddenly and capture insects which chance to alight upon them. The muscles of the articulates are situated within the solid frame-work, unlike the vertebrates, whose muscles are external to the bony skeleton. All animals have the power of motion, from the lowest radiate to the highest vertebrate, from the most repulsive polyp to that type of organized life made in the very image of God.

text

The muscles, then, subserve an endless variety of purposes. By their aid the farmer employs his implements of husbandry, the mechanic deftly wields his tools, the artist plies his brush, while the fervid orator gives utterance to thoughts glowing with heavenly emotions. It is by their agency that the sublimest spiritual conceptions can be brought to the sphere of the senses, and the noblest, loftiest aims of to-day can be made glorious realizations of the future.

GLM-OCR

CER 0.0379 | WER 0.0361 | F1 0.9817 | 1 region(s) | length 0.97x

fly-trap, a plant found in North Carolina, is remarkable for the sensitiveness of its leaves, which close suddenly and capture insects which chance to alight upon them. The muscles of the articulates are situated within the solid frame-work, unlike the vertebrates, whose muscles are external to the bony skeleton. All animals have the power of motion, from the lowest radiate to the highest vertebrate, from the most repulsive polyp to that type of organized life made in the very image of God.

The muscles, then, subserve an endless variety of purposes. By their aid the farmer employs his implements of husbandry, the mechanic deftly wields his tools, the artist plies his brush, while the fervid orator gives utterance to thoughts glowing with heavenly emotions. It is by their agency that the sublimest spiritual conceptions can be brought to the sphere of the senses, and the noblest, loftiest aims of to-day can be made glorious realizations of the future.<[user]>

MinerU 2.5 Pro

CER 0.0000 | WER 0.0000 | F1 1.0000 | 1 region(s) | length 1.00x

36

COMMON SENSE MEDICAL ADVISER.

fly-trap, a plant found in North Carolina, is remarkable for the sensitiveness of its leaves, which close suddenly and capture insects which chance to alight upon them. The muscles of the articulates are situated within the solid frame-work, unlike the vertebrates, whose muscles are external to the bony skeleton. All animals have the power of motion, from the lowest radiate to the highest vertebrate, from the most repulsive polyp to that type of organized life made in the very image of God.

The muscles, then, subserve an endless variety of purposes. By their aid the farmer employs his implements of husbandry, the mechanic deftly wields his tools, the artist plies his brush, while the fervid orator gives utterance to thoughts glowing with heavenly emotions. It is by their agency that the sublimest spiritual conceptions can be brought to the sphere of the senses, and the noblest, loftiest aims of to-day can be made glorious realizations of the future.

TrOCR large-printed

CER 0.0390 | WER 0.1566 | F1 0.8434 | 1 region(s) | length 0.99x

36 COMMUN SENSE MEDUAL ADVISER, MY-TRAH, A PLANT FOUND IN NORTH CAROLINA, IS REMARKABLE FOR THE SENSITIVENESS OF ITS LEAVES, WHICH CLOSE SUBMENT AND CANTURE INSECTS WHICH CHANGE TO ALIGHT UPON THEM. THE MALES OF THE ARTICULATES ARE SITUATED WITHIN THE SOLD FRAME-WORK. UNLIKE THE VERTEBRATES, WHOSE MUSCLES ARE EXTERAL TO THE BONY SKELETON. ALL ANIMALS HAVE THE POWER OF MOTION, FROM THE LOWEST RELATE TO THE HIGHEST VERTEBIATE, FROM THE MOST REPULSIVE POLYP TO THAT TYPE OF ORGANIZED LIFE MADE IN THE VERY INAGE OF GOLD. THE MUSTES, THEN, SUBSERVE AN ENCLESS VARIETY OF PURPOSES. BY THEIR AND THE FARMER EMPLOYS HIS IMPLEMENTS OF HUSBANDRY. THE MECHANIC DEFTLY WILLAS HIS TOOLS, THE ARTIST PLIES HIS BRUSH, WHILE THE PERVIB ORATOR GIVES UTTERANCE TO THOUGHTS GLOWING WITH HEAVENTY EMOTIONS. IT IS BY THEIR AGENCY THAT THE SUBLIMEST SPIRITUAL CONCEPTIONS CAN BE BROUGHT TO THE SPHERE OF THE SENSOS, AND THE NOBLEST, LOFTIES AIMS OF TO-DAY CAN BE MADE GLORIONS VEALIZATIONS OF THE FUTURE,

Verified label (manual GT)

975 normalized characters

36 COMMON SENSE MEDICAL ADVISER.

fly-trap, a plant found in North Carolina, is remarkable for the sensitiveness of its leaves, which close suddenly and capture insects which chance to alight upon them. The muscles of the articulates are situated within the solid frame-work, unlike the vertebrates, whose muscles are external to the bony skeleton. All animals have the power of motion, from the lowest radiate to the highest vertebrate, from the most repulsive polyp to that type of organized life made in the very image of God.

The muscles, then, subserve an endless variety of purposes. By their aid the farmer employs his implements of husbandry, the mechanic deftly wields his tools, the artist plies his brush, while the fervid orator gives utterance to thoughts glowing with heavenly emotions. It is by their agency that the sublimest spiritual conceptions can be brought to the sphere of the senses, and the noblest, loftiest aims of to-day can be made glorious realizations of the future.

DeepSeek-OCR-2

CER 0.0226 | WER 0.0482 | F1 0.9581 | 4 region(s) | length 0.99x

\(3 6 \)

CUMUL SAVER ALL IN ONE

fly-trap, a plant found in North Carolina, is remarkable for the sensitiveness of its leaves, which close suddenly and capture insects which chance to alight upon them. The muscles of the articulates are situated within the solid frame-work, unlike the vertebrates, whose muscles are external to the bony skeleton. All animals have the power of motion, from the lowest radiate to the highest vertebrae, from the most repulsive polyp to that type of organized life made in the very image of God.

The muscles, then, subserve an endless variety of purposes. By their aid the farmer employs his implements of husbandry, the mechanic deftly wields his tools, the artist plies his brush, while the fervid orator gives utterance to thoughts glowing with heavenly emotions. It is by their agency that the sublimest spiritual conceptions can be brought to the sphere of the senses, and the noblest, loftiest aims of to-day can be made glorious realizations of the future.

GLM-OCR

CER 0.0205 | WER 0.0241 | F1 0.9881 | 4 region(s) | length 1.02x

36<[user]>

COMMON SENSE MEDICAL ADVISER.<[user]>

fly-trap, a plant found in North Carolina, is remarkable for the sensitiveness of its leaves, which close suddenly and capture insects which chance to alight upon them. The muscles of the articulates are situated within the solid frame-work, unlike the vertebrates, whose muscles are external to the bony skeleton. All animals have the power of motion, from the lowest radiate to the highest vertebrate, from the most repulsive polyp to that type of organized life made in the very image of God.<[user]>

The muscles, then, subserve an endless variety of purposes. By their aid the farmer employs his implements of husbandry, the mechanic deftly wields his tools, the artist plies his brush, while the fervid orator gives utterance to thoughts glowing with heavenly emotions. It is by their agency that the sublimest spiritual conceptions can be brought to the sphere of the senses, and the noblest, loftiest aims of to-day can be made glorious realizations of the future.<[user]>

MinerU 2.5 Pro

CER 0.0031 | WER 0.0060 | F1 0.9970 | 4 region(s) | length 1.00x

COMMON SENSE MEDICAL ADVISER.

fly-trap, a plant found in North Carolina, is remarkable for the sensitiveness of its leaves, which close suddenly and capture insects which chance to alight upon them. The muscles of the articulates are situated within the solid frame-work, unlike the vertebrates, whose muscles are external to the bony skeleton. All animals have the power of motion, from the lowest radiate to the highest vertebrate, from the most repulsive polyp to that type of organized life made in the very image of God.

The muscles, then, subserve an endless variety of purposes. By their aid the farmer employs his implements of husbandry, the mechanic deftly wields his tools, the artist plies his brush, while the fervid orator gives utterance to thoughts glowing with heavenly emotions. It is by their agency that the sublimest spiritual conceptions can be brought to the sphere of the senses, and the noblest, loftiest aims of to-day can be made glorious realizations of the future.

TrOCR large-printed

CER 0.0226 | WER 0.0904 | F1 0.9096 | 4 region(s) | length 1.00x

36
COMMON SENSE MEDICAL ADVISER.
MY-TRAP, A PLANT FOUND IN NORTH CAROLINA, IS REMARKABLE FOR THE SENSITIVENESS OF ITS LEAVES, WHICH ELOSE SUDDENLY AND EANTURE INSECTS WHICH CHANGE TO ALIGHT UPON THEM. THE MUSTES OF THE ARTICULATES ARE SITUATED WITHIN THE SOLD FRAME-WORK. UNLIKE THE VERTEBRATES, WHOSE MUSCLES ARE EXTERNAL TO THE BONY SKELETON. ALL ANIMALS HAVE THE POWER OF MOTION, FROM THE LOWEST RADIATE TO THE HIGHEST VERTEBATE, FROM THE MOST REPULSIVE POLYP TO THAT TYPE OF ORGANIZED LIFE MADE IN THE VERY IMAGE OF GOOD. THE MUSTES, THEN, SUBSERVE AN ENCLLESS VARIETY OF PURPOSES. BY THEIR AID THE FARMER EMPLOYS HIS IMPLEMENTS OF HUSBANDRY, THE MECHANIC DEFTLY WILLAS HIS TOOLS. THE ARTIST PLIES HIS BRUSH, WHILE THE SERVIC ORATOR GIVES INTERANCE TO THOUGHTS GLOWING WITH HEAVENTY EMOTIONS. IT IS BY THEIR AGENCY THAT THE SUBLIMEST SPIRITUAL CONCEPTIONS CAN BE BROUGHT TO THE SPHERE OF THE SENSES, AND THE NOBLEST, LOFTIEST AIMS OF TO-DAY CAN BE MADE GLORIONS REALIZATIONS OF THE FUTURE,

Verified label (manual GT)

1314 normalized characters

CHAPTER IV.

PHYSIOLOGICAL ANATOMY.

THE DIGESTIVE ORGANS.

Digestion signifies the act of separating or distributing, hence its application to the process by which food is made available for nutritive purposes. The organs of digestion are the Mouth, Teeth, Tongue, Salivary Glands, Pharynx, Esophagus, the Stomach and the Intestines, with their glands, the Liver, Pancreas, Lacteals, and the Thoracic Duct.

The Mouth is an irregular cavity, situated between the upper and the lower jaw, and contains the organs of mastication. It is bounded by the lips in front, by the cheeks at the sides, by the roof of the mouth and teeth of the upper jaw above, and behind and beneath by the teeth of the lower jaw, soft parts, and palate. The soft palate is a sort of pendulum attached only at one of its extremities, while the other involuntarily opens and closes the passage from the mouth to the pharynx. The interior of the mouth, as well as other portions of the alimentary canal, is lined with a delicate tissue, called mucous membrane.

Fig. 26.

A view of the lower jaw. 1. The body. 2, 2. Rami, or branches. 3, 3. Processes of the lower jaw. m. Molar teeth. b. Bicuspid. c. Cuspid. i. Incisors.

The Teeth are firmly inserted in the alveoli or sockets, of the upper and the lower jaw. The first set, twenty in number, are temporary, and appear during infancy. They are replaced

DeepSeek-OCR-2

CER 0.1796 | WER 0.2101 | F1 0.9592 | 1 region(s) | length 1.05x

sub_title

(CHAPTER IV.

sub_title

PHYSIOLOGICAL ANATOMY.

sub_title

THE DIGESTIVE ORGANS.

text

Digestion signifies the act of separating or distributing, hence its application to the process by which food is made available for nutritive purposes. The organs of digestion are the Mouth, Teeth, Tongue, Salivary Glands, Pharynx, Esophagus, the Stomach and the Intestines, with their glands, the Liver, Pancreas, Lacteals, and the Thoracic Duct.

text

The Mouth is an irregular cavity, situated between the upper and the lower jaw, and contains the organs of mastication. It is bounded by the lips in front, by the cheeks at the sides, by the roof of the mouth and teeth of the upper jaw above, and behind and beneath by the teeth of the lower jaw, soft parts, and palate. The soft palate is a sort of pendulum attached only at one of its extremities, while the other involuntarily opens and closes the passage from the mouth to the pharynx. The interior of the mouth, as

figure_title

Fig. 26.

image

text

A view of the lower jaw. 1. The body. 2. Rami, or branches. 3. 3. Processes of the lower jaw. m. Molar teeth. b. Bi-cuspid. c. Cuspid. i. Incisors.

text

well as other portions of the alimentary canal, is lined with a delicate tissue, called mucous membrane.

text

The Teeth are firmly inserted in the alveoli or sockets, of the upper and the lower jaw. The first set, twenty in number, are temporary, and appear during infancy. They are replaced

GLM-OCR

CER 0.1111 | WER 0.1387 | F1 0.9258 | 1 region(s) | length 0.90x

CHAPTER IV.

PHYSIOLOGICAL ANATOMY.

THE DIGESTIVE ORGANS.

Digestion signifies the act of separating or distributing, hence its application to the process by which food is made available for nutritive purposes. The organs of digestion are the Mouth, Teeth, Tongue, Salivary Glands, Pharynx, Esophagus, the Stomach and the Intestines, with their glands, the Liver, Pancreas, Lacteals, and the Thoracic Duct.

The Mouth is an irregular cavity, situated between the upper and the lower jaw, and contains the organs of mastication. It is bounded by the lips in front, by the cheeks at the sides, by the roof of the mouth and teeth of the upper jaw above, and behind and beneath by the teeth of the lower jaw, soft parts, and palate. The soft palate is a sort of pendulum attached only at one of its extremities, while the other involuntarily opens and closes the passage from the mouth to the pharynx. The interior of the mouth, as well as other portions of the alimentary canal, is lined with a delicate tissue, called mucous membrane.

The Teeth are firmly inserted in the alveoli or sockets, of the upper and the lower jaw. The first set, twenty in number, are temporary, and appear during infancy. They are replaced<[user]>

MinerU 2.5 Pro

CER 0.1796 | WER 0.2647 | F1 0.9937 | 1 region(s) | length 1.00x

CHAPTER IV.

PHYSIOLOGICAL ANATOMY.

THE DIGESTIVE ORGANS.

Digestion signifies the act of separating or distributing, hence its application to the process by which food is made available for nutritive purposes. The organs of digestion are the Mouth, Teeth, Tongue, Salivary Glands, Pharynx, Esophagus, the Stomach and the Intestines, with their glands, the Liver, Pancreas, Lacteals, and the Thoracic Duct.

The Mouth is an irregular cavity, situated between the upper and the lower jaw, and contains the organs of mastication. It is bounded by the lips in front, by the cheeks at the sides, by the roof of the mouth and teeth of the upper jaw above, and behind and beneath by the teeth of the lower jaw, soft parts, and palate. The soft palate is a sort of pendulum attached only at one of its extremities, while the other involuntarily opens and closes the passage from the mouth to the pharynx. The interior of the mouth, as well as other portions of the alimentary canal, is lined with a delicate tissue, called mucous membrane.

The Teeth are firmly inserted in the alveoli or sockets, of the upper and the lower jaw. The first set, twenty in number, are temporary, and appear during infancy. They are replaced

Fig. 26.

A view of the lower jaw. 1. The body. 2. 2. Rami, or branches. 3, 3. Processes of the lower jaw. m. Molar teeth. b. Bicuspid. c. Cuspid. i. Incisors.

37

TrOCR large-printed

CER 0.3706 | WER 0.5252 | F1 0.6308 | 1 region(s) | length 0.75x

CASHIER CHAPTER IV.

PHYSIOLOGICAL ANATOMY.

THE DIGESTIVE ORGAYS,

DIGATION SIGNICES THE ACT OF SEPARATING OR DISTRIBUTING, THANCE ITS APPLICATION TO THE PROCESS BY WHICH FOOD IS MADE AVAIL- ABLE FOR NUTRITIVE PURPOSES. THE ORGANS OF DIGESTION ARE THE MONTH, TEETH, TONGUE, SALIVARY GLANDS, PHARTIX, ESOPHAGUS, THE STONACH AND THE LATESTINGS, WITH THEIR GLANDS, THE LAVEY. PANCREAS, LATEALS, AND THE THORAGIC DUEY, DETAIL, PRICE THE MOUTH IS AN ITEGALAR CAVITY, SITURTEL BETWEEN THE UPPER AND THE LOWER JAW, AND CONTAINS THE ORGANS OF MASTERATION. IT FIVE: 20.

LLLLLLLL

AND CLOSES THE PASSAGE FROM 4.8. BAMI, OR UNNCHAS. S. S. PROCESS OF THE MOUTH TO THE PHARYNX. THE LOWER JAY. M. MOHOR THUROR, B. B- THE INTERIOR OF THE MOUTH, AS AREAT

WELL AS OTHER PORTIONS OF THE ALIMENTARY CAMAL, IS LINED WITH A DELICATE TISSUE, CALLED MAROUS MEMBRINE,

THE TEETH ARE FAMLY INSERTED IN THE ALVEOLI OR SOCKETS OF THE UPPER AND THE LOWER JAW. THE INST SET, TWENTY IN NUMBER, ARE TEMPORARY, AND APPEAR DURING INFANEY. THEY ARE REPLACED TOTAL :

Verified label (manual GT)

1646 normalized characters

COMMON SENSE MEDICAL ADVISER.

by permanent teeth, of which there are sixteen in each jaw; four incisors, or front teeth, four cuspids, or eye teeth, four bicuspid, or grinders, and four molars, or large grinders. Each tooth is divided into the crown, body, and root. The crown is the grinding surface; the body, the part projecting from the jaw, is the seat of sensation and nutrition; the root is that portion of the tooth which is inserted in the alveolus. The teeth are composed of dentine, or ivory, and enamel. The ivory forms the greater portion of the body and root, while the enamel covers the exposed surface. The small white cords communicating with the teeth are the nerves.

The Tongue is a flat oval organ, the base of which is attached to the os hyoides, while the apex, the most sensitive part of the body, is free. Its surface is covered with a membrane, which, at the sides and lower part, is continuous with the lining of the mouth. On the lower surface of the tongue, this membrane is thin and smooth, but on the upper side it is covered with numerous papillae, which, in structure, are similar to the sensitive papillae of the skin.

Fig. 27.

The salivary glands. The largest one, near the ear, is the parotid gland. The next below it is the submaxillary gland. The one under the tongue is the sublingual gland.

The Salivary Glands are six in number, three on each side of the mouth. Their function is to secrete a fluid called saliva, which aids in mastication. The largest of these glands, the Parotid, is situated in front and below the ear; its structure, like that of all the salivary glands, is cellular. The Submaxillary gland is circular in form, and situated midway between the

DeepSeek-OCR-2

CER 0.2497 | WER 0.2607 | F1 0.9673 | 1 region(s) | length 1.01x

text

by permanent teeth, of which there are sixteen in each jaw; four incisors, or front teeth, four cuspid, or eye teeth, four bicuspid, or grinders, and four molars, or large grinders. Each tooth is divided into the crown, body, and root. The crown is the grinding surface; the body, the part projecting from the jaw, is the seat of sensation and nutrition; the root is that portion of the tooth which is inserted in the alveolus. The teeth are composed of dentine, or ivory, and enamel. The ivory forms the greater portion of the body and root, while the enamel covers the exposed surface. The small white cords communicating with the teeth are the nerves.

text

The Tome is a flat oval organ, the base of which is attached

figure_title

Fig. 27.

image

text

The salivary glands. The largest one near the ear, is the parotid gland. The next below it is the submaxillary gland. The one under the tongue is the sublingual gland.

text

to the os hyoides, while the apex, the most sensitive part of the body, is free. Its surface is covered with a membrane, which, at the sides and lower part, is continuous with the lining of the mouth. On the lower surface of the tongue, this membrane is thin and smooth, but

text

on the upper side

it is covered with

text

numerous papillae, which, in structure, are similar to the sensitive papillae of the skin.

text

The Salivary Glands are six in number, three on each side of the mouth. Their function is to secrete a fluid called saliva, which aids in mastication. The largest of these glands, the Parotid, is situated in front and below the ear; its structure, like that of all the salivary glands, is cellular. The Submaxillary gland is circular in form, and situated midway between the

GLM-OCR

CER 0.1276 | WER 0.1353 | F1 0.9212 | 1 region(s) | length 0.88x

by permanent teeth, of which there are sixteen in each jaw; four incisors, or front teeth, four cuspids, or eye teeth, four bicuspid, or grinders, and four molars, or large grinders. Each tooth is divided into the crown, body, and root. The crown is the grinding surface; the body, the part projecting from the jaw, is the seat of sensation and nutrition; the root is that portion of the tooth which is inserted in the alveolus. The teeth are composed of dentine, or ivory, and enamel. The ivory forms the greater portion of the body and root, while the enamel covers the exposed surface. The small white cords communicating with the teeth are the nerves.

The Tongue is a flat oval organ, the base of which is attached to the os hyoides, while the apex, the most sensitive part of the body, is free. Its surface is covered with a membrane, which, at the sides and lower part, is continuous with the lining of the mouth. On the lower surface of the tongue, this membrane is thin and smooth, but on the upper side it is covered with numerous papillae, which, in structure, are similar to the sensitive papillae of the skin.

The Salivary Glands are six in number, three on each side of the mouth. Their function is to secrete a fluid called saliva, which aids in mastication. The largest of these glands, the Parotid, is situated in front and below the ear; its structure, like that of all the salivary glands, is cellular. The Submaxillary gland is circular in form, and situated midway between the<[user]>

MinerU 2.5 Pro

CER 0.2114 | WER 0.2244 | F1 0.9885 | 1 region(s) | length 1.00x

38

COMMON SENSE MEDICAL ADVISER.

by permanent teeth, of which there are sixteen in each jaw; four incisors, or front teeth, four cuspids, or eye teeth, four biceuspids, or grinders, and four molars, or large grinders. Each tooth is divided into the crown, body, and root. The crown is the grinding surface; the body, the part projecting from the jaw, is the seat of sensation and nutrition; the root is that portion of the tooth which is inserted in the alveolus. The teeth are composed of dentine, or ivory, and enamel. The ivory forms the greater portion of the body and root, while the enamel covers the exposed surface. The small white cords communicating with the teeth are the nerves.

The Tongue is a flat oval organ, the base of which is attached

Fig. 27.

The salivary glands. The largest one, near the ear, is the parotid gland. The next below it is the submaxillary gland. The one under the tongue is the sublingual gland.

to the os hyoides, while the apex, the most sensitive part of the body, is free. Its surface is covered with a membrane, which, at the sides and lower part, is continuous with the lining of the mouth. On the lower surface of the tongue, this membrane is thin and smooth, but on the upper side it is covered with

numerous papillae, which, in structure, are similar to the sensitive papillae of the skin.

The Salivary Glands are six in number, three on each side of the mouth. Their function is to secrete a fluid called saliva, which aids in mastication. The largest of these glands, the Parotid, is situated in front and below the ear; its structure, like that of all the salivary glands, is cellular. The Submaxillary gland is circular in form, and situated midway between the

TrOCR large-printed

CER 0.2691 | WER 0.4059 | F1 0.6897 | 1 region(s) | length 0.83x

INCOMMUN SENSE MEDICAL ADVINER,
BY PERMANENT TEETH, OF WHICH THERE ARE SIXTEEN IN EACH JAW;
FOUR INCLUSORS, OR FRONT TEETH, FOUR EUSPINS, OR EYE TEETH, FOUR
BIENSPIDS, OR GRINDERS, AND FOUR MOLARS, OR LARGE GRINDERS, EACH
TOOTH IS DIVIDED INTO THE GROWN, BODY, AND FOOT. THE CROWN IS
THE GRINDING SURFACE ; THE BODY, THE PART PROJECTING FROM THE
JAW, IS THE SERT OF SENSATION AND INTRITION : THE ROOT IS THAT NOR-
TION OF THE FOOTH WHICH IS INSETED IN THE ALVEOLUS, THE TEETH
ARE COMPOSED OF DENTINE, OR INTRY, AND EXAMEL. THE INORY
FORMS THE GREATER PORTION OF THE BODY AND ROOT, WHILE THE
EMAMEL COXERS THE EXPOSED SURFACE, THE SAMALI WHITE COMDS
COMMUNICATING WITH THE TEETH ARE THE DETAILS.
THE TAMPWE IS A BAT OVAL ORGAN, THE BASE OF WHICH IS ATTACHED
SALES
PURCHASE NO. 27. . WE SIZE THE SDEX.
PER
THE ADDITARY CLAIMS, THE MARKET ONE. MEAN THE EAR, IS THE AND SMOOTH. BUT
PERAMPLE AND TAX MELL RETURN THIS INFORMATION ON THE UPPER SIDE
TOTAL RETURRIED WITH
NUMERONS DAPILLE, WHICH, IN STFUETARE, ARE SIMILAR TO THE SENSITIVE
BADILLE OF THE SKIN.
THE SALINARY GLANDS ARE SIX IN NUMBER. FREE ON EACH SIDE
OF THE MOUNT. THEIR FUNCTION IS TO SECREE A FULLLED SOLDING,
WHICH Aids IN MASTRATION. THE LATGEST OF THESE GLANDS, THE
PAROTIL, IS STUATED IN FRONT AND BELOW THE EAR; ITS STRETURE, LIKE
THAT OF ALL THE SALIVARY GLANDS, IS CELLULAR. THE SANDWANILARY
GLAND IS GREWAR IN FORM, AND SITUATED MIDWAY BETWEEN THE

p0046 - PP-DocLayoutV3 region OCR

Side-by-side saved text. Metrics use identical normalized scoring; displayed text preserves OCR wording.

Verified label (manual GT)

1646 normalized characters

COMMON SENSE MEDICAL ADVISER.

by permanent teeth, of which there are sixteen in each jaw; four incisors, or front teeth, four cuspids, or eye teeth, four bicuspidis, or grinders, and four molars, or large grinders. Each tooth is divided into the crown, body, and root. The crown is the grinding surface; the body, the part projecting from the jaw, is the seat of sensation and nutrition; the root is that portion of the tooth which is inserted in the alveolus. The teeth are composed of dentine, or ivory, and enamel. The ivory forms the greater portion of the body and root, while the enamel covers the exposed surface. The small white cords communicating with the teeth are the nerves.

The Tongue is a flat oval organ, the base of which is attached to the os hyoides, while the apex, the most sensitive part of the body, is free. Its surface is covered with a membrane, which, at the sides and lower part, is continuous with the lining of the mouth. On the lower surface of the tongue, this membrane is thin and smooth, but on the upper side it is covered with numerous papillae, which, in structure, are similar to the sensitive papillae of the skin.

Fig. 27.

The salivary glands. The largest one, near the ear, is the parotid gland. The next below it is the submaxillary gland. The one under the tongue is the sublingual gland.

The Salivary Glands are six in number, three on each side of the mouth. Their function is to secrete a fluid called saliva, which aids in mastication. The largest of these glands, the Parotid, is situated in front and below the ear; its structure, like that of all the salivary glands, is cellular. The Submaxillary gland is circular in form, and situated midway between the

DeepSeek-OCR-2

CER 1.4277 | WER 1.9043 | F1 0.4694 | 9 region(s) | length 2.25x

␣ 2 3 ␣

<table><tr><td>Feature</td><td>Description</td></tr><tr><td>Object</td><td>Arrow</td></tr><tr><td>Color</td><td>Red</td></tr><tr><td>Color</td><td>Dark Red</td></tr><tr><td>Color</td><td>Dark Blue</td></tr><tr><td>Color</td><td>Light Red</td></tr><tr><td>Color</td><td>Light Blue</td></tr><tr><td>Color</td><td>Light Red</td></tr><tr><td>Color</td><td>Light Blue</td></tr><tr><td>Color</td><td>Dark Blue</td></tr><tr><td>Color</td><td>Dark Red</td></tr><tr><td>Color</td><td>Dark Blue</td></tr><tr><td>Color</td><td>Medium Blue</td></tr><tr><td>Color</td><td>Medium Red</td></tr><tr><td>Color</td><td>Medium Blue</td></tr><tr><td>Color</td><td>Light Red</td></tr><tr><td>Color</td><td>Light Blue</td></tr><tr><td>Color</td><td>Light Red</td></tr><tr><td>Color</td><td>Light Blue</td></tr><tr><td>Color</td><td>Medium Blue</td></tr><tr><td>Color</td><td>Medium Red</td></tr><tr><td>Color</td><td>Medium Blue</td></tr><tr><td>Color</td><td>Dark Red</td></tr><tr><td>Color</td><td>Dark Blue</td></tr></table>

by permanent teeth, of which there are sixteen in each jaw; four incisors, or front teeth, four cuspids, or eye teeth, four bicuspidis, or grinders, and four molars, or large grinders. Each tooth is divided into the crown, body, and root. The crown is the part of the tooth which is covered by the gingival tissue; the body of the tooth is the largest part and is covered by enamel; the root is the part that attaches the tooth to the jaw. The crown is formed by the enamel of the tooth, while the root is formed by the dentin. The pulp cavity is the space within the crown and root, and is filled with pulp, which is a soft tissue that contains blood vessels, nerves, and other structures. The pulp cavity is lined by a hard tissue called the dentine, which is produced by the ameloblasts. The dentin is harder than the enamel and is stained by the minerals in the blood. The pulp cavity is connected to the alveolar spaces of the root canals, which are filled with pulp. The pulp cavity is also connected to the periodontal gum, which is a membrane that surrounds the tooth and provides support and attachment for the tooth.

I am sorry, but the image provided is a photograph of a blue plastic container, not a chart or graph. Therefore, it does not contain any data points or numerical information that can be extracted and converted into a markdown table.

to the os hyoides, while the apex, the most sensitive part of the body, is free. Its surface is covered with a membrane, which, at the lower surface, is continuous with the lining of the mouth. On the lower surface of the tongue, this membrane is thin and smooth, but on the upper side it is covered with a layer of papillae, which give it a velvety appearance.

␣ F' = 9.2 ␣

The sailor's hands, The largest one, near the ear, Is the parotid gland. The next below its the sinhanxillary gland. The one under the tongue is the sublingual gland.

numerous particles within in structure are similar to the sensitive capability of the skin.

The *Sulfurary Glumds* are six in number, three on each side of the mouth. Their function is to secrete a fluid called *sulfur*, which aids in mastication. The largest of these glands, the *Parotid*, is situated in front and below the ear; its structure, like that of all the salivary glands, is cellular. The *Submaxillary gland* is circular in form, and situated midway between the

GLM-OCR

CER 0.1264 | WER 0.1221 | F1 0.9838 | 9 region(s) | length 1.03x

38<|user|>

COMMON SENSE MEDICAL ADVISER.<|user|>

by permanent teeth, of which there are sixteen in each jaw; four incisors, or front teeth, four cuspids, or eye teeth, four bicuspidis, or grinders, and four molars, or large grinders. Each tooth is divided into the crown, body, and root. The crown is the grinding surface; the body, the part projecting from the jaw, is the seat of sensation and nutrition; the root is that portion of the tooth which is inserted in the alveolus. The teeth are composed of dentine, or ivory, and enamel. The ivory forms the greater portion of the body and root, while the enamel covers the exposed surface. The small white cords communicating with the teeth are the nerves.<|user|>

The Tongue is a flat oval organ, the base of which is attached<|user|>

to the os hyoides, while the apex, the most sensitive part of the body, is free. Its surface is covered with a membrane, which, at the sides and lower part, is continuous with the lining of the mouth. On the lower surface of the tongue, this membrane is thin and smooth, but on the upper side it is covered with<|user|>

Fig. 27.<|user|>

The salivary glands. The largest one, near the ear, is the parotid gland. The next below it is the submaxillary gland. The one under the tongue is the sublingual gland.<|user|>

numerous papillae, which, in structure, are similar to the sensitive papillae of the skin.<|user|>

The Salivary Glands are six in number, three on each side of the mouth. Their function is to secrete a fluid called saliva, which aids in mastication. The largest of these glands, the Parotid, is situated in front and below the ear; its structure, like that of all the salivary glands, is cellular. The Submaxillary gland is circular in form, and situated midway between the<|user|>

MinerU 2.5 Pro

CER 0.1428 | WER 0.1452 | F1 0.9645 | 9 region(s) | length 0.96x

COMMON SENSE MEDICAL ADVISER.

by permanent teeth, of which there are sixteen in each jaw; four incisors, or front teeth, four cuspids, or eye teeth, four bicuspidis, or grinders, and four molars, or large grinders. Each tooth is divided into the crown, body, and root. The crown is the grinding surface; the body, the part projecting from the jaw, is the seat of sensation and nutrition; the root is that portion of the tooth which is inserted in the alveolus. The teeth are composed of dentine, or ivory, and enamel. The ivory forms the greater portion of the body and root, while the enamel covers the exposed surface. The small white cords communicating with the teeth are the nerves.

to the os hyoides, while the apex, the most sensitive part of the body, is free. Its surface is covered with a membrane, which, at the sides and lower part, is continuous with the lining of the mouth. On the lower surface of the tongue, this membrane is thin and smooth, but on the upper side it is covered with

The salivary glands. The largest one, near the ear, is the parotid gland. The next below it is the submaxillary gland. The one under the tongue is the sublingual gland.

numerous papillae, which, in structure, are similar to the sensitive papillae of the skin.

The Salivary Glands are six in number, three on each side of the mouth. Their function is to secrete a fluid called saliva, which aids in mastication. The largest of these glands, the Parotid, is situated in front and below the ear; its structure, like that of all the salivary glands, is cellular. The Submaxillary gland is circular in form, and situated midway between the

TrOCR large-printed

CER 0.1306 | WER 0.1881 | F1 0.9067 | 9 region(s) | length 1.00x

38

COMMON SENSE MEDICAL ADVISER, BY PERMANENT TEETH, OF WHICH THERE ARE SIXTEEN IN EACH JAW; FOUR INCISORS, OR FRONT TEETH, FOUR CUSPIDS, OR EYE TEETH, FOUR BIENSPIDS, OR GRINDERS, AND FOUR MOLARS, OR LARGE GRINDERS. EACH TOOTH IS DIVIDED INTO THE FROM, BODY, AND ROOT. THE CROWN IS THE GRINDING SURFACE; THE BODY, THE PART PROJECTING FROM THE JAW, IS THE SERT OF SENSATION AND NUTRITION ; THE ROOT IS THAT FOR- TION OF THE TOOTH WHICH IS INSERTED IN THE ALVEOLUS. THE TEETH ARE COMPOSED OF DENTINE, OR IVORY, AND ENAMEL. THE IVORY FORMS THE GREATER PORTION OF THE BODY AND ROOT, WHILE THE ENAMEL COVERS THE EXPOSED SURFACE; THE SAMALL WHITE COMDS COMMUNICATING WITH THE TEETH ARE THE SERVICES. THE TAMPLE IS A HAT OYAL ORGAN, THE BASE OF WHICH IS ATTACHED TO THE OS HYOIDES, WHILE THE APEX, THE MOST SENSI- TIVE PART OF THE BODY, IS FREE. ITS SURFACE IS COV- ERED WITH A MEM- BRANE, WHICH, AT THE SIDES AND LOWER PART, IS CONTINUOUS WITH THE LINING OF THE MOUTH. ON THE LOWER SURFACE OF THE TONGUE, THIS MEMBRANE IS THIN AND SMOOTH, BUT ON THE UPPER SIDE IT IS COVERED WITH

Fig. 27.

THE SALIVARY GLANDS. THE LARGEST ONE, NEAR THE EAR, IS THE PAROTID FLAND. THE NEXT BELOW IT IS THE SUBMAXILLARY GLAND. THE ONE UNDER THE TONGUE IS THE SUBLINGUAL GLAND. NUMEROUS PAPILLA, WHICH, IN STRUCTURE, ARE SIMILAR TO THE SENSITIVE DADILLE OF THE SKIN. THE SALMARY GLAMDS ARE SIX IN NUMBER, THREE ON EACH SIDE OF THE MOUTH. THEIR FUNCTION IS TO SECREE A FULL CALLED SALINA, WHICH AILS IN MASTICATION. THE LARGEST OF THESE GLANDS, THE PAROTIL, IS SITUATED IN FRONT AND BELOW THE EAR; ITS STRUCTURE, LIKE THAT OF ALL THE SALIVARY GLANDS, IS CELLUOUR. THE SUBMARYJARY GLAND IS GIRELLAR IN FORM, AND SITUATED MIDWAY BETWEEN THE

Verified label (manual GT)

1920 normalized characters

THE DIGESTIVE ORGANS. 39

angle of the lower jaw and the middle of the chin. The Sublingual is a long flattened gland, and, as its name indicates, is located below the tongue, which when elevated, discloses the saliva issuing from its porous openings.

The Pharynx is nearly four inches in length, formed of muscular and membranous cells, and situated between the base of the cranium and the esophagus, in front of the spinal column. It is narrow at the upper part, distended in the middle, contracting again at its junction with the esophagus. The pharynx communicates with the nose, mouth, larynx, and esophagus.

The Esophagus, a cylindrical organ, is a continuation of the pharynx, and extends through the diaphragm to the stomach. It has three coats: first, the muscular, consisting of an exterior layer of fibers running longitudinally, and an interior layer of transverse fibers; second, the cellular, which is interposed between the muscular and the mucous coat; third, the mucous membrane, or internal coat, which is continuous with the mucous lining of the pharynx. The Stomach is a musculo-membranous, conoidal sac, communicating with the esophagus by means of the cardiac orifice (see Fig. 28). It is situated obliquely with reference to the body, its base lying at the left side, while the apex is directed toward the right side. The stomach is between the liver and spleen, subjacent to the diaphragm, and communicates with the intestinal canal by the pyloric orifice. It has three coats. The peritoneal, or external coat is composed of compact, cellular tissue, woven into a thin, serous membrane, and assists in keeping the stomach in place. The middle coat is formed of three layers of muscular fibers: in the first, the fibres run

Fig. 28.
A representation of the interior of the stomach. 1. The esophagus. 2. Cardiac orifice opening into the stomach. 6. The middle or muscular coat. 7. The interior or mucous coat. 10. The begin-ning of the duodenum. 11. The pyloric orifice.

DeepSeek-OCR-2

CER 0.2750 | WER 0.3012 | F1 0.9686 | 1 region(s) | length 1.01x

text

angle of the lower jaw and the middle of the chin. The Sub-lingual is a long flattened gland, and, as its name indicates, is located below the tongue, which when elevated, discloses the saliva issuing from its porous openings.

text

The Pharynx is nearly four inches in length, formed of muscular and membranous cells, and situated between the base of the cranium and the esophagus, in front of the spinal column. It is narrow at the upper part, distended in the middle, contracting again at its junction with the esophagus. The pharynx communicates with the nose, mouth, larynx, and esophagus.

text

The Esophagus, a cylindrical organ, is a continuation of the pharynx, and extends through the diaphragm to the stomach. It has three coats: first, the muscular, consisting of an exterior layer of fibers running longitudinally, and an interior layer of transverse fibers; second, the cellular, which is interposed between the muscular and the mucous coat; third, the mucous membrane, or internal coat, which is continuous with the mucous lining of the pharynx.

text

The Stomach is a musculo-membranous, conical sac, communicating with the esophagus by means of the cardiac orifice (see

figure_title

Fig. 28.

image

text

A representation of the interior of the stomach.

1. The esophagus.

2. Cardiac orifice opening into the stomach.

6. The middle or muscular coat.

7. The interior or mucous coat.

10. The beginning of the duodenum.

11. The pyloric orifice.

text

Fig. 28). It is situated obliquely with reference to the body, its base lying at the left side, while the apex is directed toward the right side. The stomach is between the liver and spleen, subjacent to the diaphragm, and communicates with the intestinal canal by the pyloric orifice. It has three coats. The peritoneal, or external coat is composed of compact, cellular tissue, woven into a thin, serous membrane, and assists in keeping the stomach in place. The middle coat is formed of three layers of muscular fibers: in the first, the fibres run

GLM-OCR

CER 0.1302 | WER 0.1416 | F1 0.9223 | 1 region(s) | length 0.87x

angle of the lower jaw and the middle of the chin. The Sublingual is a long flattened gland, and, as its name indicates, is located below the tongue, which when elevated, discloses the saliva issuing from its porous openings.

The Pharynx is nearly four inches in length, formed of muscular and membranous cells, and situated between the base of the cranium and the esophagus, in front of the spinal column. It is narrow at the upper part, distended in the middle, contracting again at its junction with the esophagus. The pharynx communicates with the nose, mouth, larynx, and esophagus.

The Esophagus, a cylindrical organ, is a continuation of the pharynx, and extends through the diaphragm to the stomach. It has three coats: first, the muscular, consisting of an exterior layer of fibers running longitudinally, and an interior layer of transverse fibers; second, the cellular, which is interposed between the muscular and the mucous coat; third, the mucous membrane, or internal coat, which is continuous with the mucous lining of the pharynx.

The Stomach is a musculo-membranous, conoidal sac, communicating with the esophagus by means of the cardiac orifice (see Fig. 28). It is situated obliquely with reference to the body, its base lying at the left side, while the apex is directed toward the right side. The stomach is between the liver and spleen, subjacent to the diaphragm, and communicates with the intestinal canal by the pyloric orifice. It has three coats. The peritoneal, or external coat is composed of compact, cellular tissue, woven into a thin, serous membrane, and assists in keeping the stomach in place. The middle coat is formed of three layers of muscular fibers: in the first, the fibres run<[user]>

MinerU 2.5 Pro

CER 0.0057 | WER 0.0301 | F1 0.9789 | 1 region(s) | length 1.00x

THE DIGESTIVE ORGANS.

39

angle of the lower jaw and the middle of the chin. The Sub-lingual is a long flattened gland, and, as its name indicates, is located below the tongue, which when elevated, discloses the saliva issuing from its porous openings.

The Pharynx is nearly four inches in length, formed of muscular and membranous cells, and situated between the base of the cranium and the esophagus, in front of the spinal column. It is narrow at the upper part, distended in the middle, contracting again at its junction with the esophagus. The pharynx communicates with the nose, mouth, larynx, and esophagus.

The Esophagus, a cylindrical organ, is a continuation of the pharynx, and extends through the diaphragm to the stomach. It has three coats: first, the muscular, consisting of an exterior layer of fibers running longitudinally, and an interior layer of

transverse fibers; second, the cellular, which is interposed between the muscular and the mucous coat; third, the mucous membrane, or internal coat, which is continuous with the mucous lining of the pharynx.

The Stomach is a musculo-membranous, conoidal sac, communicating with the esophagus by means of the cardiac orifice (see

Fig. 28). It is situated obliquely with reference to the body, its base lying at the left side, while the apex is directed toward the right side. The stomach is between the liver and spleen, subjacent to the diaphragm, and communicates with the intestinal canal by the pyloric orifice. It has three coats. The peritoneal, or external coat is composed of compact, cellular tissue, woven into a thin, serous membrane, and assists in keeping the stomach in place. The middle coat is formed of three layers of muscular fibers: in the first, the fibres run

A representation of the interior of the stomach.

1. The esophagus.

2. Cardiac orifice opening into the stomach.

6. The middle or muscular coat.

7. The interior or mucous coat.

10. The beginning of the duodenum.

11. The pyloric orifice.

TrOCR large-printed

CER 0.2750 | WER 0.4096 | F1 0.7093 | 1 region(s) | length 0.87x

THE THE DIGESTIVE ORGAYS.

ANGLE OF THE LOWER JAW AND THE MIDLE OF THE CHIN. THE SUB-LINGUAL IS A LONG HATTENED GLAND, AND, AS ITS NAME INDICATES, IS LOCATED BELOW THE TONGLE, WHICH WHEN ELEVATED, DISCLOSES THE SALIVA ISSUING FROM ITS PORONS OPENINGS.

THE PHARYME IS HEARLY FOUR INEELS IN LENGTH, FORMED OF MRS. CALAR AND MEMBANOUS CELLS, AND SITUATED BETWEEN THE BASE OF THE ERINIUM AND THE ESOPHAGAS, IN FRONT OF THE SPINAL COLUMN, IT IS MARROW AT THE IMPER PART, DISTENDAU IN THE MIDDLE, CONTRET-ING AGAIN AT ITS JUNCTION WITH THE ESOPHAGUS. THE PHARYAX COMMUNICATES WITH THE NOSE, MOUTH, LARYWX, AND ESOPHAGNS.

THE BSPPLAYUS, A EXTMORICAL ORGAN, IS A CONTINUATION OF THE PHARYNX, AND EXTENDS THROUGH THE DISPHRAGN TO THE STOMACH.

IT HAS HAVE COSTS: INST. THE MISCLAR, CONSISTING OF AN EXPERIOR LAYER OF FIBERS RUNNING LONGITALINALLY, AND AN INTERIOR LAYER OF TRANSVERSE/BERS: SECOND, RECORDER:

THE CELLULAR, WHICH IS PAYABLE.

G

CONDIAL SAC, COMMUNI- 1, THERESPERTATION OF THE INTERIOR OF THE STORE INVOICE CATING WITH THE THE STOMACH. 6. THE MODE OF MUSCLAR COM.

ESOPHAGAS BY MEANS OF THE IMPORTOR OF INCLUSIVE ORDER OF RECEIPT-

THE CARDIAN OFFICE (SEE REFUND FOR DETAILS, FREE

FIG. 28). IT IS SITANTED OHLONELY WITH REFERENCE TO THE BODY,

ITS BASE LYING AT THE LET SIDE, WHILE THE APEX IS DIRECTED TOWARD

THE RIGHT SIDE. THE STOMACH IS BETWEEN THE LIVER AND SPLEEN,

SUBJECT TO THE DISPERSM, AND COMMUNICATES WITH THE INTES-TINAL CANAL BY THE POLORIC OFFICE. IT HAS THREE COATS, THE PERITONEAL, OR EXTERNAL COAT IS COMPOSED OF COMPACT, CELLULAR TASME, WOVEN INTO A THIN, SEROUS MEMBRAINE, AND ASSISTS IN KEEPING THE STOMACH IN PLACE. THE MIDLE COAT IS FORMED OF THREE LEVELS OF MASCLAR FIBERS: IN THE FIRST, THE FIBRES RUN

