More of the protein the child already makes: from a BUB1B genotype to drug hypotheses in mosaic variegated aneuploidy

Track 2 report · Rare Disease, Real Kid: MVA Hackathon 2026
Santiago Marin Martinez (Hugging Face / GitHub: smarin00) · 24 October 2026
Code, results and every number in this report: https://github.com/smarin00/mva-hackathon-2026

Summary. This child's mosaic variegated aneuploidy (MVA) is explained by two *BUB1B* alleles, found in Track 1 (rank points 100/100). The stop-gain p.Leu737* is degraded by nonsense-mediated decay (NMD), so every BubR1 molecule the child has comes from the missense allele p.Asn1002Lys. Four lines of evidence indicate that this allele makes a working but unstable protein: a protein language model calibrated against six experimentally measured MVA alleles, the AlphaFold structure, an independent predictor (AlphaMissense) and the published biology of the domain. SpliceAI excludes a splicing effect. **The child already makes a working protein; there just isn't enough of it.** Because chromosome segregation fails at about 6% of normal BubR1 and is near normal at about 13%, raising the existing protein about two-fold is a realistic therapeutic goal.

A direction-aware screen of Open Targets, followed by documented curation, gives three approved-drug hypotheses. **Niacin** (NAD^+ → SIRT2 → BubR1 K668) is the lead, followed by **arimoclomol** (chaperone induction; FDA-approved from age 2) and **dasatinib + quercetin** (senescent-cell clearance, downstream). The same screen gives a **do-no-harm list** of 31 compounds, many of them oncology drugs, that would push the wrong way. A check against current FDA labels shows that niacin and arimoclomol could continue alongside the chemotherapy this child would receive for an MVA-associated cancer, whereas dasatinib should not.

For other families, a **variant atlas** maps all 19,950 possible BubR1 missense changes and all 1,050 possible premature stops to a mechanism and a therapy direction. Without any annotation it recovers the experimentally defined map of the protein. It agrees with ClinVar benign variants it never saw (1 of 33 in its damaging bin, against 34% expected; $p = 2 \times 10^{-5}$). The pipeline then extends the atlas to the six other genes linked to MVA. Everything runs from one case file with `make all`, and a clean rerun reproduced every result. All findings are hypotheses; a validation plan in patient cells, with go/no-go criteria, is given in Section 5.




Figure 1. From genotype to drug hypotheses. The truncating allele makes no protein, so the phenotype rests on the missense allele. Evidence that this allele is destabilised, not inactive, turns the problem into one of protein quantity. That defines the therapeutic goal, which drives the drug screen, the do-no-harm list, the cancer-treatment check and the resource for other families.

For families: what this report says, in plain words

Every cell needs a quality-control step when it divides, to make sure each new cell gets exactly one copy of every chromosome. A protein called BubR1 is central to that step. In MVA caused by changes in the *BUB1B* gene, some cells end up with too many or too few chromosomes, which causes the condition's features and raises the risk of some childhood cancers.

Everyone has two copies of *BUB1B*. In this child, one copy carries a change that stops the cell from making any protein from it. The other copy makes BubR1 that most likely works but falls apart faster than it should, so there is too little of it. Laboratory studies of other MVA families showed that when this kind of unstable BubR1 is brought back to normal amounts, the quality-control step works again, and that cells need only a fraction of the normal amount to divide correctly.

That points to a strategy: not replacing the protein, but helping the cell keep more of the protein it already makes. We looked for existing, approved medicines that act on the cell's protein-protection systems. We found three worth testing in the laboratory, and a list of medicines that would push in the wrong direction, which may be useful to the child's doctors now. None of this has been tested in this child's cells yet, and it is not a treatment recommendation. It is a set of specific, testable ideas, together with the experiments that would show whether they are right.

1. Case and genotype

Allele	GRCh38	cDNA	Protein	Consequence
1	chr15:40209701 T>G	c.2210T>G	p.Leu737*	stop-gain, exon 17 of 23
2	chr15:40220612 T>G	c.3006T>G	p.Asn1002Lys	missense, pseudokinase domain

Transcript ENST00000287598 (MANE Select); BubR1, 1,050 amino acids (UniProt O60566).

Both variants were called in the provided VCF and again from the raw reads by DeepVariant, and Exomiser ranked *BUB1B* first (Track 1). **Phase is unresolved.** The variants are 10.9 kb apart, beyond the reach of short-read pairs, and read-backed phasing found no shared phase block. The diagnosis needs them in trans (in cis, the other copy would be wild type). The evidence for trans is phenotypic: MVA with constitutional aneuploidy arises from biallelic *BUB1B* loss, and MVA1 patients typically carry one truncating and one missense allele [1]. Parental testing or long-read sequencing would settle it.

2. Mechanism

2.1 p.Leu737* is a null allele

From the Ensembl transcript structure (`bin/nmd_check.py`), the premature stop lies in exon 17 of 23, **747 nt upstream of the last exon-exon junction**. That is about 15 times the 50-nt boundary beyond which transcripts are degraded by NMD [2, 3]. Consistent with this, truncating MVA alleles are not detectable as protein in patient cells, and their transcripts are reduced or absent [4]. SpliceAI [5] predicts no splice effect (all delta scores ≤ 0.03), so exon skipping does not rescue the allele.

The point matters because the truncated protein would not be useless if it were made. BubR1's C-terminal kinase domain is a catalytically inactive pseudokinase that the checkpoint does not need: a truncation at residue 731 fully rescues checkpoint signalling and chromosome alignment [6]. A product ending at residue 736 would keep every checkpoint-essential feature (KEN1, the TPR domain, the GLEBS motif that binds Bub3) [7]. NMD, not the loss of sequence, makes this allele null. **All of the child's BubR1 therefore comes from the p.Asn1002Lys allele.**

2.2 p.Asn1002Lys most likely destabilises BubR1

Prior from the literature. The characterised MVA missense alleles in and against the pseudokinase domain act on protein *quantity*, not activity. I909T, L1012P and R727C turn over about two-fold faster, depend on HSP90, are cleared by the proteasome and are present at 2–6-fold lower levels in patient cells. Raising I909T or L1012P back to normal levels fully restores checkpoint function [4]. p.Asn1002Lys lies in the same domain, ten residues from L1012P.




Figure 2. Three predictors tested against six alleles with measured stability [4]. (a) ESM-2 separates the destabilising from the stable alleles completely, and the patient allele falls between the two best-characterised destabilisers. Shading marks the calibrated bins used by the atlas. (b) AlphaMissense calls the patient allele likely pathogenic but does not separate the measured classes. (c) Burial does not separate them either. Sources: `findings/track2/esm_scores.tsv`, `alphamissense/summary.txt`, `structure_context.tsv`.

Line 1: a protein language model, calibrated. ESM-2 (650M) masked-marginal scores [8, 9] have no scale on their own, so we read them against six MVA alleles whose stability was measured experimentally [4] (`bin/esm_variant_score.py`):

Variant	Measured class	ESM-2 log-likelihood ratio	Percentile in pseudokinase domain
L1012P	destabilising	−6.94	12.8
N1002K	**patient**	**−5.81**	**18.2**
I909T	destabilising	−5.71	18.8
L844F	destabilising	−3.33	34.8
R727C	destabilising	−2.80	40.4
Q921H	stable	−0.24	78.7
R550Q	stable	+0.37	87.3

The model separates the two classes completely, and the patient allele falls between L1012P and I909T, the two alleles for which HSP90 dependence and rescue by overexpression were shown directly. With six alleles, a perfect 4-versus-2 split arises by chance with $p = 1/15 \approx 0.07$: suggestive, not conclusive. ESM-2 scores evolutionary constraint, not folding. It serves as a stability proxy here because the pseudokinase domain has no catalytic role.

Line 2: structure. In the AlphaFold model (AF-O60566-F1) [10], N1002 is confidently placed (pLDDT 91) and buried (relative solvent accessibility 0.16) in a pocket lined by Trp973, Phe977, Trp978 and Val998. Its side-chain amide lies within hydrogen-bonding distance (≤ 3.5 Å) of three backbone atoms, Trp978 N and O and Val998 O, bridging two segments of the fold. Lysine is longer and positively charged: it cannot reproduce this

donor-acceptor bridge, and it puts a charge into a pocket lined by aromatic side chains. We also report a negative result: burial alone does not separate the measured classes, because the stable control Q921H is buried too (Figure 2c). The structure supplies a physical mechanism, not independent discrimination.

Line 3: an independent predictor. AlphaMissense [11], trained on primate population frequency and structural context rather than on these alleles, scores p.Asn1002Lys 0.92 (*likely pathogenic*). The two predictors therefore agree on the patient allele. Unlike ESM-2, however, AlphaMissense does **not** reproduce the measured classes. It calls two verified destabilisers likely benign (R727C 0.24, L844F 0.29), leaves a third ambiguous (I909T 0.45) and scores the stable control Q921H (0.49) above three of the four. Across all 6,173 missense changes it scores, the two predictors correlate moderately (Spearman $|\rho| = 0.65$; 0.80 in the TPR domain). Our reading is that a general pathogenicity score, tuned to clearly damaging variants, underrates mild hypomorphic stability alleles, which is the class MVA1 missense alleles belong to. That is why ESM-2 calibrated on measured stability is the primary line here; with six alleles this remains indicative.

Line 4: splicing excluded. SpliceAI predicts no splice effect (all delta scores ≤ 0.02, in *BUB1B* and in the overlapping *PAK6* read-through annotation).

Conclusion. No single line is proof; together they make a stability defect the most likely mechanism. The decisive experiment is a cycloheximide chase in patient cells (Section 5).

2.3 Disease mechanism and therapeutic goal

Loss or gain of function	**Loss of function**: one null allele plus one hypomorphic (destabilised) allele
Pathway disrupted	**Spindle assembly checkpoint.** BubR1, with Bub3, Mad2 and Cdc20, forms the mitotic checkpoint complex, which holds back the APC/C–Cdc20 ubiquitin ligase until every kinetochore is attached; BubR1 also stabilises kinetochore-microtubule attachments
Downstream consequence	Premature anaphase and chromosome missegregation, giving **mosaic aneuploidy** across tissues; cancer predisposition (embryonal rhabdomyosarcoma, Wilms tumour, leukaemia [1, 12]); in BubR1-hypomorphic mice, early senescence and progeroid features [13, 14]

The quantitative key. Segregation is very sensitive to the BubR1 dose: about 6% of normal causes missegregation in most cells, while about 13% has virtually no effect [4]. Biallelic null alleles have not been reported and are presumably lethal [1]. MVA1 therefore sits in a narrow band of residual BubR1, and **a roughly two-fold increase of the existing, functional protein may be enough to cross the threshold.** The goal is to raise the steady-state level of a working but unstable protein; there is no enzyme activity to restore.

Bulk sequencing depth was also profiled for aneuploidy. Chromosomes 16, 17 and 19 show 1.3–2.3% higher depth, depending on the GC model, but chromosome GC content predicts the shift under all three GC-correction models tested (Spearman ρ 0.48–0.69), and the B-allele frequency shows no matching signal. These shifts are reported as unconfirmed (`findings/aneuploidy/`). This is the expected result for MVA: aneuploidy differs from cell to cell and averages out in bulk DNA.

3. From mechanism to candidates

3.1 A direction-aware screen, then curation

Candidate identification combined an automated screen with documented curation:

1. **Direction-aware screen** (`bin/opentargets_candidates.py`). A 13-gene panel around BubR1 dosage (`config/genes/bub1b.yaml`) records, for each gene, the direction that would help the child and the paper behind it: SIRT2 *up*, HSF1 *up*, HSP90 *do not inhibit*, and so on. The panel was crossed with every drug and clinical candidate in Open Targets (API 26.9.0, data 26.09) using the ChEMBL action type, and each drug-target pair was classified automatically: *in the desired direction*, *against it* (do-no-harm) or *review*.
2. **Named-drug lookup** for agents that act on a pathway rather than through one annotated target (niacin, nicotinamide riboside, arimoclomol, valproate, senolytics).
3. **Curation** of every result against the mechanism, and against pediatric safety and approval status taken from primary regulatory sources.

The screen is informative mainly for what it rules out (Figure 3). **No drug acts directly on BUB1B, SIRT2 or HSF1**, so the leads below enter through named-drug lookup and the literature. Its 15 compounds in the desired direction were all curated out (12 HDAC inhibitors, two BCL-xL inhibitors and the CBP inhibitor PRI-724; Section 3.3). Its 31 compounds acting against the child's biology form the do-no-harm list (Section 3.4). The screen also exposed database errors that curation corrected: arimoclomol is still listed at phase 3 despite FDA approval, and several dietary supplements (nicotinamide riboside, quercetin, fisetin) carry "approved" flags.




Figure 3. The screen, gene by gene. Compounds in Open Targets acting on each panel gene, split by whether they act in the direction that would help (right) or against it (left), and what curation concluded. Salt forms are merged into one compound. Source: `findings/track2/opentargets/ot_target_drugs.tsv` .

3.2 Candidates

	Candidate	Route to more BubR1	Evidence for BubR1 specifically	Approval	Main risks
1	**Niacin** (nicotinic acid)	NAD^+ → SIRT2 → deacetylation of BubR1 K668 → less degradation	**Direct, in vivo** [14]	FDA-approved (dyslipidaemia)	Flushing; liver toxicity at high dose; no pediatric NAD^+ data
2	**Arimoclomol**	Amplifies the HSF1 heat-shock response → chaperone support for a misfolding-prone protein	Mechanism-class extrapolation	FDA 2024 (Niemann-Pick C, with miglustat), age ≥ 2 [15]; not EU-approved [16]	Hypersensitivity; no BubR1 data
3	**Dasatinib + quercetin**	Downstream: clears senescent cells	Genetic clearance in BubR1 progeroid mice [13]	Dasatinib FDA-approved, incl. pediatric Ph+ ALL	Marrow suppression, pleural effusion; treats consequences, not cause

1. Niacin: the lead. BubR1 levels fall when NAD^+ and SIRT2 activity fall. SIRT2 deacetylates BubR1 at K668, counteracting the acetyltransferase CBP, and raises BubR1 abundance in vivo. The NAD^+ precursor NMN raised BubR1 levels in mice, and more SIRT2 **extended the median lifespan of BubR1-hypomorphic mice** [14], the closest animal model of the dosage loss in MVA. p.Asn1002Lys is full length and keeps K668, so the pathway still has its substrate. Niacin is the approved NAD^+ precursor of choice: at 750–1,000 mg/day it raised blood NAD^+ up to eight-fold in adults with mitochondrial myopathy [17]. It enters NAD^+ synthesis through NAPRT, bypassing NAMPT, and unlike nicotinamide it does not feed back to inhibit sirtuins. Nicotinamide riboside, a dietary supplement rather than an approved medicine, is a better-tolerated alternative with pediatric precedent: 25 mg/kg/day for four months in 24 patients with ataxia-telangiectasia, 17 of them under 18 [18]. *Gap:* SIRT2-mediated stabilisation was shown for wild-type BubR1; whether it rescues a destabilised mutant is the first thing to test. *Caveat:* SIRT2 can also deacetylate a second site, K250, whose acetylation protects BubR1 in mitosis [19]. The in vivo net effect of more SIRT2 was more BubR1 [14], but validation should measure BubR1 through mitosis as well as at steady state.

2. Arimoclomol. It amplifies HSF1-driven induction of HSP70-family chaperones under stress, supporting proteins that are prone to misfold. The neighbouring allele L1012P is a verified chaperone-dependent client [4]. HSF1 also induces HSP90, which should help an HSP90 client, but that is a prediction. Arimoclomol is approved in the US for children from age 2 [15], which shortens any path to a trial; the EU's CHMP gave a negative opinion in July 2026 [16].

3. Dasatinib + quercetin. Clearing p16Ink4a-positive senescent cells in BubR1 progeroid mice delayed ageing phenotypes in eye, muscle and fat, even when started late [13]. That study used genetic clearance; dasatinib + quercetin is the senolytic regimen studied in humans. This treats consequences rather than cause and carries real toxicity, so it ranks last.

3.3 Considered and not ranked: HDAC inhibitors

HDAC2 and HDAC3 remove a protective acetylation at BubR1 K250 at the end of mitosis [20], which is why the screen returned 12 HDAC inhibitors in the desired direction. Six are listed as approved in Open Targets, including givinostat, which the FDA approved in 2024 for Duchenne muscular dystrophy from age 6 [21]; valproate, a weak class I inhibitor with long pediatric use, entered through named-drug lookup. We read the three primary papers in full, and they argue against ranking any of these. K250 acetylation exists only in mitosis and sets *when* BubR1 is destroyed, not how much there is: an acetylation-deficient allele left overall BubR1 levels unchanged [22]. The direct effect of keeping K250 acetylated is a longer metaphase. In a checkpoint that is too short that might help, but HDAC inhibitors also use mitotic arrest to kill dividing cells [19, 20], and the oncology HDAC inhibitors cause thrombocytopenia. The route enters the validation plan as an exploratory arm (mitotic timing and missegregation in patient cells), not as a candidate.

Rejected at curation. PRI-724 is listed as a CBP inhibitor, the desired direction, but it blocks the CBP–β-catenin interaction rather than CBP's acetyltransferase activity, so it cannot act on BubR1 K668. Navitoclax and obatoclax (BCL-xL inhibitors) cause thrombocytopenia and are not justified in a child. High-dose nicotinamide inhibits sirtuins. Proteasome inhibitors rescued mutant BubR1 in vitro [4] but are too toxic systemically; they remain a proof of mechanism.

3.4 Do-no-harm list

This may be the output most useful to the child's doctors today, because several of these agents are in oncology use or trials and the child is cancer-predisposed.

Class	Compounds	Why it is expected to harm	Examples
HSP90 inhibitors	9	Mutant BubR1 is an HSP90 client; HSP90 inhibition depleted it in patient-derived cells [4]	ganetespib, tanespimycin, luminespib, onalespib
NAMPT inhibitors	2	Deplete NAD^+, opposing candidate 1	daporinad (trialled in childhood leukaemia), teglarinad
Aurora B inhibitors	18	Impair error correction in an already weakened checkpoint	barasertib, tozasertib, danusertib
MPS1 (TTK) inhibitors	2	Weaken the spindle assembly checkpoint directly	BAY-1161909, BAY-1217389

The HSP90 count includes CD24Fc, a fusion protein that ChEMBL annotates as an HSP70/HSP90 inhibitor. Only the HSP90 entry rests on a direct measurement in MVA patient cells; the others follow from the mechanism.

3.5 Compatibility with cancer treatment

MVA1 predisposes to embryonal rhabdomyosarcoma, Wilms tumour and leukaemia [1, 12]. A medicine this child might take for years has to be compatible with the chemotherapy the child would receive if one of these cancers develops. We checked each candidate against the standard regimens, VAC and VAC/VI for rhabdomyosarcoma [23, 24] and EE-4A and DD-4A for Wilms tumour [25], using the current US prescribing information (FDA labels via openFDA [26]; `make oncology`). Three checks are run for each pair: one label names

the other drug; one drug inhibits or induces CYP3A4 while the other is a CYP3A4 substrate; or both labels warn of the same organ toxicity. Every flag carries the label sentences it rests on (`findings/track2/oncology/compatibility.tsv`).




Figure 4. Candidates against the chemotherapy for MVA-associated cancers. Each cell lists the checks that flagged the pair; shading darkens with the number of flags. Label versions and dates: `findings/track2/oncology/labels.tsv`.

The check separates the candidates. Niacin's only flags are liver warnings shared with dactinomycin and cyclophosphamide (both carry hepatic veno-occlusive disease), which liver-enzyme monitoring addresses. Arimoclomol raises no flag; its label states that it neither inhibits nor induces CYP3A4/5. Dasatinib inhibits CYP3A4, which clears vincristine, irinotecan and doxorubicin, and it overlaps with every regimen drug on marrow suppression. Valproate overlaps on marrow and liver, a further reason not to rank it. So the two candidates that act on the BubR1 dose, niacin and arimoclomol, could continue alongside cancer treatment, while dasatinib, which treats consequences, should be stopped before it. The flags are prompts for clinical review, not verdicts, and the absence of a flag is not proof of safety. No leukaemia regimen was modelled, because the leukaemia subtype in MVA is not specified.

4. A resource for other families

4.1 The BUB1B variant atlas

MVA1 patients typically carry one truncating and one missense allele [1], so each new family faces the two questions this report answered for one child. The atlas (`bin/variant_atlas.py`, `findings/track2/atlas/`) answers them for every possible variant.

It recovers the functional map unaided. ESM-2 sees only the sequence, yet the most constrained features are exactly the checkpoint-essential ones [7]: KEN1 (mean −11.2, more constrained than 98.6% of positions), KEN2 (−9.0), the GLEBS motif (−7.8) and the TPR domain (−6.9). The linker at residues 450–650, which AlphaFold models as disordered (median pLDDT 34), is nearly unconstrained (−1.0). Experiment, evolution and structure agree.

Truncating variants. Stops at codons 1–969 are predicted null by NMD. Stops at **codons 970–1,050 (the last 81 codons) escape NMD and keep every checkpoint-essential feature**, so by analogy with the functional 731X truncation [6] they may act as mild or tolerated alleles. A truncating *BUB1B* variant should not be assumed null without checking its position.

Missense variants. All 19,950 changes are binned by their calibrated ESM-2 score and mapped to a therapy direction. A stability defect, where dose-raising applies, covers 2,190 changes (11%). A defect in an essential functional site, where more protein may not compensate, covers 3,174 (16%); likely tolerated covers 4,001 (20%); and 10,585 (53%) remain uncertain. The uncertain share comes from linker regions without characterised function and from the wide gap between only six calibration anchors. The atlas is a triage tool, not a classifier.




Figure 5. BUB1B variant atlas. Top: positional constraint (ESM-2, mean over 19 substitutions, 9-residue window), with the calibration alleles and the patient's missense. Bottom: the NMD fate of a premature stop at each codon, with the patient's stop. Figure as committed in `findings/track2/atlas/atlas_figure.png`.

4.2 Independent check against ClinVar

`bin/clinvar_validate.py` maps every *BUB1B* missense SNV in ClinVar (file date 2026-10-04) [27] to the atlas, excluding the six calibration alleles. ClinVar holds 1,494 *BUB1B* missense records, 97% of them uncertain or conflicting.




Figure 6. Atlas bins for ClinVar variants the atlas never saw. The reference is every missense change reachable by a single nucleotide substitution (the changes ClinVar can contain). Benign variants avoid the damaging bin (one-sided binomial $p = 2 \times 10^{-5}$) and are enriched in the tolerated bin ($p = 1 \times 10^{-4}$). Unique variants are counted. Source: `findings/track2/clinvar_validation.tsv`.

Of 33 unique benign or likely benign variants, one falls in the damaging bin (3%, against 34% of reachable changes) and 19 in the tolerated bin (58%, against 26%). Some benign assertions use in-silico evidence, so the test is not fully independent. The pathogenic side cannot be tested: ClinVar holds a single pathogenic missense outside the calibration set (Q467H, one submitter), which the atlas places in its indeterminate bin.

What ClinVar is missing. It lists L1012P, L844F and R727C, experimentally verified destabilising MVA alleles [4], as variants of uncertain significance (VUS), and this child's p.Asn1002Lys as a VUS too. Calibrated computational evidence of this kind, together with the published functional data, is what variant-classification guidelines can use to move such alleles out of uncertainty. Of the 1,427 unique ClinVar VUS, the atlas places 419 as likely tolerated and 107 in the stability-defect class, where the strategy proposed here would apply if they occurred with a null allele. AlphaMissense agrees on part of this triage: 12% of VUS records are called damaging and 28% tolerated by both predictors, a higher-confidence subset.

4.3 The other genes that cause MVA

Six other genes are linked to MVA in the HPO annotations. The pipeline finds them itself, as the numbered OMIM series of the Track 2 disease and its Orphanet group, and builds the same constraint and NMD maps for each (`make related-genes`; `bin/related_genes.py`). It returned CEP57 (MVA2), TRIP13 (MVA3), CENATAC (MVA4) and MAD1L1 (MVA7), plus BUB1 and BUB3 from the Orphanet MVA group. Two of these, CENATAC and MAD1L1, had been missed by our hand-made list. Without calibration alleles these atlases report constraint and NMD geometry only, not mechanism; adding measured alleles to a gene's file switches on the same calibrated reading as for *BUB1B*.




Figure 7. Constraint and NMD maps for the six related MVA genes, built without manual input from `findings/genes/<GENE>/atlas_missense.tsv` and `atlas_truncating.tsv` (full figures with UniProt features in the same folders).

5. Proposed validation

In patient-derived cells (a lymphoblastoid line or fibroblasts), with healthy controls:

Step	Experiment	Go / no-go
1	BubR1 protein level (western blot)	Confirms the premise if well below 50% of control
2	Cycloheximide chase	Confirms instability if p.Asn1002Lys BubR1 turns over faster than wild type
3	Niacin or nicotinamide riboside, arimoclomol; dose series; BubR1 at steady state and in mitosis	**Go** if BubR1 rises above about 13% of control
4	Checkpoint (nocodazole arrest) and missegregation (live imaging or interphase FISH)	**Go** if missegregation falls measurably
5	HSP90 inhibitor as positive control	Tests the do-no-harm prediction: BubR1 should fall further
6	Exploratory: valproate at 0.3–0.7 mM; acetyl-K250 BubR1, time from nuclear-envelope breakdown to anaphase, missegregation	A longer mitosis without fewer errors, or mitotic death, rules the route out

A positive result in steps 3–4 would justify a discussion with the child's clinicians. Steps 1–2 would also provide the functional evidence needed to reclassify p.Asn1002Lys.

6. How far each claim can be trusted

Claim	Evidence	Strength	What would refute it
p.Leu737* makes no protein	NMD geometry (747 nt); truncating MVA alleles absent as protein [4]	Strong	Stable mutant transcript in patient RNA
The two alleles are in trans	Phenotype; biallelic loss in MVA1 [1]	Indirect	Parental testing or long reads showing cis
p.Asn1002Lys destabilises BubR1	Calibrated ESM-2 ($p \approx 0.07$), AlphaMissense agreement, structural H-bond bridge, domain prior	Moderate	Normal turnover in a cycloheximide chase
A two-fold rise would be enough	Dose-response in cells [4]	Moderate (other alleles, cell lines)	No fall in missegregation when BubR1 is raised
Niacin raises mutant BubR1	SIRT2–K668 route in mice, on wild-type BubR1 [14]	Weak to moderate	No rise in patient cells
Arimoclomol helps	Mechanism class; L1012P is chaperone-dependent [4]	Weak	No rise in patient cells
HSP90 inhibitors harm	Depletion of mutant BubR1 in patient cells [4]	Strong	—
Aurora B, MPS1, NAMPT inhibitors harm	Mechanism	Moderate	—

What we tested ourselves. Every predictor was tested against measured data before being used, and negative results are reported: burial does not separate the measured classes, and AlphaMissense does not either. Guards stop the run rather than analyse the wrong residue when the reference residue or isoform does not match. Fifteen tests on synthetic data run on every push (GitHub Actions). A clean clone, run end to end with `make all` (about 15 hours on a 16-core server), reproduced the Track 1 submission and every Track 2 table exactly. The aneuploidy estimates agreed within 0.03 percentage points; read placement by bwa-mem varies with thread batching, but no call changed.

7. Scalability and reuse

We went deep on one child's mechanism rather than broad across drug databases; the breadth comes from the pipeline.

- **One case file, one command.** `params.yaml` holds the data paths, HPO terms and route. `make all` runs Track 1 from raw reads or from the VCF, then Track 2 on the top-ranked candidate, then the related genes. Track 2 needs no patient data and can be rerun from `config/cases/mva1_proband.yaml` alone. Versions, container digests and database releases are recorded.
- **Gene-agnostic by design.** NMD geometry, structural context, calibrated language-model scoring, the direction-aware screen, the ClinVar check and the oncology check all read a gene file. For a new gene, a minimal file is generated from UniProt; adding a domain map, measured alleles, a pathway panel or an oncology block switches on the corresponding analysis. That curation, not the code, is the real work for a new gene, and it is explicit and reviewable.
- **The direction-aware screen** turns a mechanism into a candidate list and a do-no-harm list in one pass, which a target-to-drug lookup does not.
- **Any disorder with a handful of functionally characterised variants** can be calibrated the same way, which covers many recessive loss-of-function disorders where the residual protein is what matters.

8. Limitations

- Phase is inferred from the phenotype, not observed.
- The stability mechanism rests on four supporting lines (n = 6 calibration; burial not discriminative; a general predictor that does not separate the measured classes; a literature prior), not on a measurement in this child's cells.
- The atlas's missense bins rest on six calibration alleles; 53% of missense changes remain uncertain, and the pathogenic side could not be tested against ClinVar.
- No candidate has BubR1-specific evidence in human cells. Niacin's link runs through wild-type BubR1 in mice; arimoclomol's is a mechanism-class extrapolation; senolytics act downstream. Pediatric dosing and safety of niacin at NAD^+-raising doses are unknown.
- The oncology check reads label text: flags are prompts for review, and the leukaemia regimen was not modelled.
- Bulk sequencing cannot measure mosaic aneuploidy at the levels expected in MVA.
- All drug and target annotations come from public databases at one release; regulatory status was checked against primary sources on 19–20 October 2026.

9. Methods, data handling and AI use

Methods, commands and versions are in the repository (`README.md`, `Makefile`, `docs/decisions.md`, `docs/reference_data.md`); the figures in this report are drawn from the committed result tables by `scripts/report_figures.py`. Track 1 used Nextflow 26.04 with DeepVariant 1.10.0, Exomiser 15.1.0 (data 2602) and Ensembl VEP 116 on GRCh38. Track 2 used Ensembl 116, UniProt O60566, the AlphaFold DB model, ESM-2 650M, SpliceAI 1.3.1, AlphaMissense, ClinVar (file date 2026-10-04), HPO annotations (2026-09-01), Open Targets 26.09 and openFDA drug labels.

Patient data were analysed only on a dedicated server and never entered the repository. All data and genotype-level derived files will be deleted by 23 November 2026 (`docs/DELETION.md`).

AI assistance: Anthropic, Claude via claude.ai, consumer Pro plan (Claude Opus 5.5); model training disabled; no outputs rated. It was used for pipeline design, code, literature search, drafting of the gene knowledge file and of this report. All analyses were run, and all results and literature claims reviewed, by the author. No genome-scale data were shared (`docs/llm_usage.md`).

References

1. Hanks S. et al. Constitutional aneuploidy and cancer predisposition caused by biallelic mutations in BUB1B. *Nat Genet* 36, 1159–1161 (2004).
2. Nagy E., Maquat L.E. A rule for termination-codon position within intron-containing genes: when nonsense affects RNA abundance. *Trends Biochem Sci* 23, 198–199 (1998).
3. Lindeboom R.G.H., Supek F., Lehner B. The rules and impact of nonsense-mediated mRNA decay in human cancers. *Nat Genet* 48, 1112–1118 (2016).
4. Suijkerbuijk S.J.E. et al. Molecular causes for BUBR1 dysfunction in the human cancer predisposition syndrome mosaic variegated aneuploidy. *Cancer Res* 70, 4891–4900 (2010).
5. Jaganathan K. et al. Predicting splicing from primary sequence with deep learning. *Cell* 176, 535–548 (2019).
6. Suijkerbuijk S.J.E. et al. The vertebrate mitotic checkpoint protein BUBR1 is an unusual pseudokinase. *Dev Cell* 22, 1321–1329 (2012).
7. BubR1 TPR domain supports the mitotic checkpoint by promoting MCC formation and MCC-APC/C interaction. *Commun Biol* (2026). doi:10.1038/s42003-026-10384-9 (domain boundaries).
8. Lin Z. et al. Evolutionary-scale prediction of atomic-level protein structure with a language model. *Science* 379, 1123–1130 (2023).
9. Meier J. et al. Language models enable zero-shot prediction of the effects of mutations on protein function. *NeurIPS* 34 (2021).
10. Jumper J. et al. Highly accurate protein structure prediction with AlphaFold. *Nature* 596, 583–589 (2021).
11. Cheng J. et al. Accurate proteome-wide missense variant effect prediction with AlphaMissense. *Science* 381, eadg7492 (2023).
12. Atlas of Genetics and Cytogenetics in Oncology and Haematology: BUB1B (atlasgeneticsoncology.org/gene/854).
13. Baker D.J. et al. Clearance of p16Ink4a-positive senescent cells delays ageing-associated disorders. *Nature* 479, 232–236 (2011).
14. North B.J. et al. SIRT2 induces the checkpoint kinase BubR1 to increase lifespan. *EMBO J* 33, 1438–1453 (2014).
15. US FDA. FDA approves first treatment for Niemann-Pick disease, type C (20 September 2024); Miplyffa (arimoclomol) prescribing information.
16. Zevra Therapeutics. Update on the EU regulatory submission for arimoclomol (negative CHMP opinion, 24 July 2026).
17. Pirinen E. et al. Niacin cures systemic NAD^+ deficiency and improves muscle performance in adult-onset mitochondrial myopathy. *Cell Metab* 31, 1078–1090 (2020).
18. Veenhuis S.J.G. et al. Nicotinamide riboside improves ataxia scores and immunoglobulin levels in ataxia telangiectasia. *Mov Disord* (2021). doi:10.1002/mds.28788
19. Choi E. et al. BubR1 acetylation at prometaphase is required for modulating APC/C activity and timing of mitosis. *EMBO J* 28, 2077–2089 (2009).
20. Park I. et al. HDAC2/3 binding and deacetylation of BubR1 initiates spindle assembly checkpoint silencing. *FEBS J* 284, 4035–4050 (2017).
21. US FDA. Duvyzat (givinostat) prescribing information (approved 21 March 2024).
22. Park I. et al. Loss of BubR1 acetylation causes defects in spindle assembly checkpoint signaling and promotes tumor formation. *J Cell Biol* 202, 295–309 (2013).
23. Arndt C.A.S. et al. Vincristine, actinomycin, and cyclophosphamide compared with vincristine, actinomycin, and cyclophosphamide alternating with vincristine, topotecan, and cyclophosphamide for intermediate-risk rhabdomyosarcoma: Children's Oncology Group study D9803. *J Clin Oncol* 27, 5182–5188 (2009).
24. Hawkins D.S. et al. Addition of vincristine and irinotecan to vincristine, dactinomycin, and cyclophosphamide does not improve outcome for intermediate-risk rhabdomyosarcoma: a report from the Children's Oncology Group. *J Clin Oncol* 36, 2770–2777 (2018).
25. Green D.M. et al. Comparison between single-dose and divided-dose administration of dactinomycin and doxorubicin for patients with Wilms' tumor: a report from the National Wilms' Tumor Study Group. *J Clin Oncol* 16, 237–245 (1998).
26. US FDA prescribing information via the openFDA drug label API; set IDs, versions and dates in `findings/track2/oncology/labels.tsv`.
27. Landrum M.J. et al. ClinVar: improving access to variant interpretations and supporting evidence. *Nucleic Acids Res* 46, D1062–D1067 (2018).