

PRODUCT MONOGRAPH

^{PR}TEVA-ATOMOXETINE

(atomoxetine hydrochloride capsules)

10, 18, 25, 40 and 60 mg atomoxetine (as atomoxetine hydrochloride)

Selective Norepinephrine Reuptake Inhibitor
for Attention-Deficit/Hyperactivity Disorder (ADHD)

Teva Canada Limited
30 Novopharm Court
Toronto, Ontario
M1B 2K9

Date of Preparation:
May 11, 2012

Submission Control No: 155184

Table of Contents

<u>PART I: HEALTH PROFESSIONAL INFORMATION.....</u>	<u>3</u>
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE	3
CONTRAINDICATIONS	4
WARNINGS AND PRECAUTIONS	5
ADVERSE REACTIONS	13
DRUG ABUSE AND DEPENDENCE	21
DRUG INTERACTIONS	22
DOSAGE AND ADMINISTRATION	24
OVERDOSAGE	27
ACTION AND CLINICAL PHARMACOLOGY	28
STORAGE AND STABILITY	32
DOSAGE FORMS, COMPOSITION AND PACKAGING	32
<u>PART II: SCIENTIFIC INFORMATION</u>	<u>34</u>
PHARMACEUTICAL INFORMATION	34
CLINICAL TRIALS	35
DETAILED PHARMACOLOGY	42
TOXICOLOGY	45
REFERENCES	48
<u>PART III: CONSUMER INFORMATION.....</u>	<u>52</u>

PrTEVA-ATOMOXETINE®
(atomoxetine hydrochloride capsules)

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Clinically Relevant Nonmedicinal Ingredients
Oral	10, 18, 25, 40, 60 mg capsules	Starch <i>For a complete listing see Dosage Forms, Composition and Packaging section.</i>

INDICATIONS AND CLINICAL USE

TEVA-ATOMOXETINE (atomoxetine hydrochloride) is indicated for the treatment of Attention-Deficit/Hyperactivity Disorder (ADHD) in children 6 years of age and over, adolescents, and adults.

A diagnosis of ADHD (DSM-IV) implies the presence of hyperactive-impulsive or inattentive symptoms that cause impairment and that were present before age 7 years. The symptoms must be persistent, must be more severe than is typically observed in individuals at a comparable level of development, must cause clinically significant impairment, e.g., in social, academic, or occupational functioning, and must be present in 2 or more settings, e.g., school (or work) and at home. The symptoms must not be better accounted for by another mental disorder. For the Inattentive type, at least 6 of the following symptoms must have persisted for at least 6 months: lack of attention to details/careless mistakes, lack of sustained attention, poor listener, failure to follow through on tasks, poor organization, avoids tasks requiring sustained mental effort, loses things, easily distracted, forgetful. For the Hyperactive-Impulsive type, at least 6 of the following symptoms must have persisted for at least 6 months: fidgeting/squirming, leaving seat, inappropriate running/climbing, difficulty with quiet activities, “on the go,” excessive talking, blurting answers, can’t wait turn, intrusive. For a Combined Type diagnosis, both inattentive and hyperactive-impulsive criteria must be met.

Special Diagnostic Considerations

The specific etiology of ADHD is unknown, and there is no single diagnostic test. Adequate diagnosis requires the use not only of medical but also of special psychological, educational, and social resources. Learning may or may not be impaired. The diagnosis must be based upon a complete history and evaluation of the patient and not solely on the presence of the required

number of DSM-IV characteristics.

Need for Comprehensive Treatment Program

TEVA-ATOMOXETINE is indicated as an integral part of a total treatment program for ADHD that may include other measures (psychological, educational, and social) for patients with this syndrome. Drug treatment may not be indicated for all patients with this syndrome. Drug treatment is not intended for use in the patient who exhibits symptoms secondary to environmental factors and/or other primary psychiatric disorders, including psychosis.

Appropriate educational placement is essential in children and adolescents with this diagnosis and psychosocial intervention is often helpful. When remedial measures alone are insufficient, the decision to prescribe drug treatment medication will depend upon the physician's assessment of the chronicity and severity of the patient's symptoms.

Pediatrics (<6 years of age)

The safety and efficacy of atomoxetine hydrochloride in pediatric patients less than 6 years of age have not been established.

CONTRAINDICATIONS

- *Hypersensitivity:* TEVA-ATOMOXETINE (atomoxetine hydrochloride) is contraindicated in patients known to be hypersensitive to atomoxetine or other constituents of the product (*see* WARNINGS).
- *Monoamine Oxidase Inhibitors:* TEVA-ATOMOXETINE should not be taken with monoamine oxidase inhibitors (MAOI), or within 2 weeks after discontinuing MAOI. Treatment with MAOI should not be initiated within 2 weeks after discontinuing TEVA-ATOMOXETINE. With other drugs that affect brain monoamine concentrations, there have been reports of serious, sometimes fatal, reactions (including hyperthermia, rigidity, myoclonus, autonomic instability with possible rapid fluctuations of vital signs, and mental status changes that include extreme agitation progressing to delirium and coma) when taken in combination with MAOI. Some cases presented with features resembling neuroleptic malignant syndrome. Such reactions may occur when these drugs are given concurrently or in close proximity.
- *Narrow Angle Glaucoma:* In clinical trials, atomoxetine hydrochloride use was associated with an increased risk of mydriasis and therefore its use is not recommended in patients with narrow angle glaucoma.
- Symptomatic cardiovascular disease.
- Moderate to severe hypertension.
- Advanced arteriosclerosis.

- Uncontrolled hyperthyroidism.

WARNINGS AND PRECAUTIONS

POTENTIAL ASSOCIATION WITH THE OCCURRENCE OF BEHAVIOURAL AND EMOTIONAL CHANGES, INCLUDING SELF-HARM

Pediatric Placebo-Controlled Clinical Trial Data: An increased risk over placebo for suicide-related events in children and adolescents taking atomoxetine hydrochloride, was identified in a pooled analysis of placebo-controlled trials of 6-18 weeks duration. Of 1357 patients who received atomoxetine hydrochloride, 5 (0.37%) had reports of **suicidal ideation** compared to 0% of 851 patients who received placebo. In addition, one **suicide attempt** (overdose) was identified, which occurred in an atomoxetine hydrochloride patient. No completed suicides occurred during these trials. (See also WARNINGS AND PRECAUTIONS, Special Populations, *Pediatrics, 6-18 years of age.*)

Post-Marketing Data: There have been very rare reports of suicidal ideation, suicidal attempts, suicidal depression and completed suicides in children, adolescents and adults (see ADVERSE REACTIONS, Post-Market Adverse Drug Reactions, Tables 5 and 6).

ADHD and its related co-morbidities may be associated with increased risk of suicidal ideation and/or behavior. Rigorous clinical monitoring for suicidal ideation or other indicators of potential for suicidal behavior is advised in patients of all ages. This includes monitoring for agitation-type of emotional and behavioral changes, and clinical worsening.

Screening Patients for Bipolar Disorder

Particular care should be taken in treating ADHD in patients with comorbid bipolar disorder because of concern for possible induction of a mixed/manic episode in patients at risk for bipolar disorder. Prior to initiating treatment with Atomoxetine, patients with comorbid depressive symptoms should be adequately screened to determine if they are at risk for bipolar disorder; such screening should include a detailed psychiatric history, including a family history of suicide, bipolar disorder, and depression.

Emergence of New Psychotic or Manic Symptoms

Treatment emergent psychotic or manic symptoms, e.g., hallucinations, delusional thinking, or mania in children and adolescents without a prior history of psychotic illness or mania can occur with atomoxetine use at usual doses. If such symptoms occur, consideration should be given to a possible causal role of atomoxetine, and discontinuation of treatment should be considered. In a pooled analysis of multiple short-term, placebo-controlled studies, such symptoms occurred in about 0.2% (4 patients with events out of 1939 exposed to atomoxetine for several weeks at usual doses) of atomoxetine-treated patients compared to 0 out of 1056 placebo-treated patients.

Pre-Existing Psychosis:

Administration of medications for ADHD may also exacerbate symptoms of behavior disturbance and thought disorder in patients with a pre-existing psychotic disorder.

Hepatic Effects

Severe Liver Injury: Post-marketing reports indicate that atomoxetine hydrochloride can cause severe liver injury in rare cases. Although no evidence of liver injury was detected in clinical trials of about 6000 patients, there have been rare cases in postmarketing experience of clinically significant liver injury that were considered probably or possibly related to atomoxetine hydrochloride. In one patient, liver injury, manifested by elevated hepatic enzymes (up to 40 X upper limit of normal [ULN]) and jaundice (bilirubin up to 12 X ULN), recurred upon rechallenge and was followed by recovery upon drug discontinuation, providing evidence that atomoxetine hydrochloride likely caused the liver injury. Such reactions may occur several months after therapy is started, but laboratory abnormalities may continue to worsen for several weeks after drug is stopped. Because of probable under-reporting, it is impossible to provide an accurate estimate of the true incidence of these events. The patient described above recovered from his liver injury, and did not require a liver transplant. However, severe liver injury due to any drug may potentially progress to acute liver failure resulting in death or the need for a liver transplant.

TEVA-ATOMOXETINE should be discontinued in patients with jaundice or laboratory evidence of liver injury, and should not be restarted. Laboratory testing to determine liver enzyme levels should be done upon the first symptom or sign of liver dysfunction (e.g., pruritus, dark urine, jaundice, right upper quadrant tenderness, or unexplained "flu-like" symptoms). (*See also* CONSUMER INFORMATION).

Allergic Events

Although uncommon, allergic reactions, including angioneurotic reactions, rash, angioneurotic edema, and urticaria, have been reported in patients taking atomoxetine hydrochloride.

Effects on Growth

Chronic open-label treatment studies (up to 3 years) indicate minimal, if any, long-term effects of atomoxetine hydrochloride on weight and height compared with normal growth curves.

In general, the weight and height gain of pediatric patients treated with atomoxetine hydrochloride lags behind that predicted by normative population data for about the first 9-12 months of treatment. Subsequently, weight gain rebounds and at about 3 years of treatment, patients treated with atomoxetine hydrochloride have gained 17.9 kg on average, 0.5 kg more than predicted by their baseline data. After about 12 months, gain in height stabilizes, and at 3 years, patients treated with atomoxetine hydrochloride have gained 19.4 cm on average, 0.4 cm less than predicted by their baseline data.

This growth pattern was generally similar regardless of pubertal status at the time of treatment initiation. Patients who were pre-pubertal at the start of treatment (girls ≤ 8 years old, boys ≤ 9 years old) gained an average of 2.1 kg and 1.2 cm less than predicted after three years. Patients who were pubertal (girls > 8 to ≤ 13 years old, boys > 9 to ≤ 14 years old) or late pubertal (girls > 13 years old, boys > 14 years old) had average weight and height gains that were close to or exceeded those predicted after three years of treatment.

Growth followed a similar pattern in both extensive and poor metabolizers (EMs, PMs). PMs treated for at least 2 years gained an average of 2.4 kg and 1.1 cm less than predicted, while EMs gained an average of 0.2 kg and 0.4 cm less than predicted.

During acute treatment studies (up to 9 weeks), atomoxetine hydrochloride-treated patients lost an average of 0.4 kg, while placebo patients gained an average of 1.5 kg. In a controlled trial that randomized patients to placebo or 1 of 3 atomoxetine doses, 1.3%, 7.1%, 19.3%, and 29.1% of patients lost at least 3.5% of their body weight in the placebo, 0.5, 1.2, and 1.8 mg/kg/day atomoxetine hydrochloride dose groups, respectively. During acute treatment studies, atomoxetine hydrochloride -treated patients grew an average of 0.9 cm while placebo-treated patients grew an average of 1.1 cm. Whether final adult height or weight is affected by treatment with atomoxetine hydrochloride is unknown. Growth should be monitored during treatment with atomoxetine hydrochloride.

Cardiovascular

Pre existing Cardiovascular and Cerebral Vascular Conditions

TEVA-ATOMOXETINE should be used with caution in patients with hypertension, tachycardia, congenital long QT syndrome, or cardiovascular or cerebrovascular disease because it can increase blood pressure and heart rate. Patients should be screened for pre-existing or underlying cardiovascular or cerebrovascular conditions before initiation of treatment with TEVA-ATOMOXETINE and monitored for new conditions of the heart or brain during the course of treatment. Pulse and blood pressure should be measured at baseline, following TEVA-ATOMOXETINE dose increases, and periodically while on therapy.

In pediatric placebo-controlled trials, atomoxetine hydrochloride -treated subjects experienced a mean increase in heart rate of about 6 beats/minute compared with placebo subjects. At the final study visit before drug discontinuation, 3.1% (16/518) of atomoxetine hydrochloride -treated subjects had heart rate increases of at least 25 beats/minute and a heart rate of at least 110 beats/minute, compared with 0.3% (1/338) of placebo subjects. 1.9% (10/518) of atomoxetine hydrochloride -treated subjects and 0.3% (1/338) of placebo subjects had a heart rate increase of at least 25 beats/minute and a heart rate of at least 110 beats/minute on 2 or more occasions. Tachycardia was identified as an adverse event for 0.5% (3/657) of these pediatric subjects compared with 0.0% (0/408) of placebo subjects. The mean heart rate increase in extensive metabolizer (EM) patients was 5.8 beats/minute, and in poor metabolizer (PM) patients 9.7 beats/minute.

Atomoxetine hydrochloride -treated pediatric subjects experienced mean increases of about 1.5

mm Hg in systolic and diastolic blood pressures compared with placebo. At the final study visit before drug discontinuation, 5.2% (26/496) of atomoxetine hydrochloride -treated pediatric subjects had high systolic blood pressure measurements compared with 2.4% (8/327) of placebo subjects. High systolic blood pressures were measured on 2 or more occasions in 6.0% (30/496) of atomoxetine hydrochloride -treated subjects and 2.4% (8/327) of placebo subjects. At the final study visit before drug discontinuation, 3.0% (15/506) of atomoxetine hydrochloride -treated pediatric subjects had high diastolic blood pressure measurements compared with 0.9% (3/330) of placebo subjects. High diastolic blood pressures were measured on 2 or more occasions in 4.2% (21/506) of atomoxetine hydrochloride -treated subjects and 0.9% (3/330) of placebo subjects. (High systolic and diastolic blood pressure measurements were defined as those exceeding the 95th percentile, stratified by age, gender, and height percentile - National High Blood Pressure Education Working Group on Hypertension Control in Children and Adolescents.)

In adult placebo-controlled trials, atomoxetine hydrochloride -treated subjects experienced a mean increase in heart rate of 5 beats/minute compared with placebo subjects. Tachycardia was identified as an adverse event for 3% (8/269) of these adult atomoxetine subjects compared with 0.8% (2/263) of placebo subjects.

Atomoxetine hydrochloride -treated adult subjects experienced mean increases in systolic (about 3 mm Hg) and diastolic (about 1 mm Hg) blood pressures compared with placebo. At the final study visit before drug discontinuation, 1.9% (5/258) of atomoxetine hydrochloride -treated adult subjects had systolic blood pressure measurements ≥ 150 mm Hg compared with 1.2% (3/256) of placebo subjects. At the final study visit before drug discontinuation, 0.8% (2/257) of atomoxetine hydrochloride -treated adult subjects had diastolic blood pressure measurements ≥ 100 mm Hg compared with 0.4% (1/257) of placebo subjects. No adult subject had a high systolic or diastolic blood pressure detected on more than one occasion.

Orthostatic hypotension has been reported in subjects taking atomoxetine hydrochloride. In short-term child- and adolescent-controlled trials, 5.2% (34/657) of atomoxetine hydrochloride -treated subjects experienced symptoms of postural hypotension compared with 2.0% (8/408) of placebo-treated subjects. TEVA-ATOMOXETINE should be used with caution in any condition that may predispose patients to hypotension.

Sudden Death and Pre-existing Structural Cardiac Abnormalities or Other Serious Heart Problems

Children and Adolescents - Sudden death has been reported in association with atomoxetine treatment at usual doses in children and adolescents with structural cardiac abnormalities or other serious heart problems. Although some serious heart problems alone carry an increased risk of sudden death, atomoxetine generally should not be used in children or adolescents with known serious structural cardiac abnormalities, cardiomyopathy, serious heart rhythm abnormalities, or other serious cardiac problems that may place them at increased vulnerability to the noradrenergic effects of atomoxetine.

Adults – Sudden deaths, stroke, and myocardial infarction have been reported in adults taking

atomoxetine at usual doses for ADHD. Although the role of atomoxetine in these adult cases is also unknown, adults have a greater likelihood than children of having serious structural cardiac abnormalities, cardiomyopathy, serious heart rhythm abnormalities, coronary artery disease, or other serious cardiac problems. Consideration should be given to not treating adults with clinically significant cardiac abnormalities.

General

Children: Theoretically there exists a pharmacological potential for all ADHD drugs to increase the risk of sudden/cardiac death. Although confirmation of an incremental risk for adverse cardiac events arising from treatment with ADHD medications is lacking, prescribers should consider this potential risk.

All drugs with sympathomimetic effects prescribed in the management of ADHD should be used with caution in patients who: *a)* are involved in strenuous exercise or activities, *b)* use stimulants, or *c)* have a family history of sudden/cardiac death. Prior to the initiation of treatment with sympathomimetic medications, a personal and family history (including assessment for a family history of sudden death or ventricular arrhythmia) and physical exam should be obtained to assess for the presence of cardiac disease. In patients with relevant risk factors and based on the clinician's judgment, further cardiovascular evaluation may be considered (e.g., electrocardiogram and echocardiogram). Patients who develop symptoms such as exertional chest pain, unexplained syncope, or other symptoms suggestive of cardiac disease during ADHD treatment should undergo a prompt cardiac evaluation. Patients who are considered to need extended treatment with atomoxetine should undergo periodic evaluation of their cardiovascular status (see DOSAGE AND ADMINISTRATION).

Genitourinary

Effects on urine outflow from the bladder — In adult ADHD controlled trials, the rates of urinary retention and urinary hesitation were increased among atomoxetine subjects compared with placebo subjects. A complaint of urinary retention or urinary hesitancy should be considered potentially related to atomoxetine.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis — Atomoxetine HCl was not carcinogenic in rats and mice when given in the diet for 2 years at time-weighted average doses up to 47 and 458 mg/kg/day, respectively. The highest dose used in rats is approximately 8 and 5 times the maximum human dose in children and adults, respectively, on a mg/m² basis. Plasma levels (AUC) of atomoxetine at this dose in rats are estimated to be 1.8 times (extensive metabolizers) or 0.2 times (poor metabolizers) those in humans receiving the maximum human dose. The highest dose used in mice is approximately 39 and 26 times the maximum human dose in children and adults, respectively, on a mg/m² basis.

Mutagenesis — Atomoxetine HCl was negative in a battery of genotoxicity studies that included a reverse point mutation assay (Ames Test), an in vitro mouse lymphoma assay, a chromosomal

aberration test in Chinese hamster ovary cells, an unscheduled DNA synthesis test in rat hepatocytes, and an in vivo micronucleus test in mice. However, there was a slight increase in the percentage of Chinese hamster ovary cells with diplochromosomes, suggesting endoreduplication (numerical aberration).

The metabolite N-desmethyatomoxetine HCl was negative in the Ames Test, mouse lymphoma assay, and unscheduled DNA synthesis test.

Impairment of fertility — Atomoxetine HCl did not impair fertility in rats when given in the diet at doses of up to 57 mg/kg/day, which is approximately 6 times the maximum human dose on a mg/m² basis.

Teratogenicity — Pregnant rabbits were treated with up to 100 mg/kg/day of atomoxetine by gavage throughout the period of organogenesis. At this dose, in 1 of 3 studies, a decrease in live fetuses and an increase in early resorptions was observed. Slight increases in the incidences of atypical origin of carotid artery and absent subclavian artery were observed. These findings were observed at doses that caused slight maternal toxicity. The no-effect dose for these findings was 30 mg/kg/day. The 100-mg/kg dose is approximately 23 times the maximum human dose on a mg/m² basis; plasma levels (AUC) of atomoxetine at this dose in rabbits are estimated to be 3.3 times (extensive metabolizers) or 0.4 times (poor metabolizers) those in humans receiving the maximum human dose.

Rats were treated with up to approximately 50 mg/kg/day of atomoxetine (approximately 6 times the maximum human dose on a mg/m² basis) in the diet from 2 weeks (females) or 10 weeks (males) prior to mating through the periods of organogenesis and lactation. In 1 of 2 studies, decreases in pup weight and pup survival were observed. The decreased pup survival was also seen at 25 mg/kg (but not at 13 mg/kg). In a study in which rats were treated with atomoxetine in the diet from 2 weeks (females) or 10 weeks (males) prior to mating throughout the period of organogenesis, a decrease in fetal weight (female only) and an increase in the incidence of incomplete ossification of the vertebral arch in fetuses were observed at 40 mg/kg/day (approximately 5 times the maximum human dose on a mg/m² basis) but not at 20 mg/kg/day. No adverse fetal effects were seen when pregnant rats were treated with up to 150 mg/kg/day (approximately 17 times the maximum human dose on a mg/m² basis) by gavage throughout the period of organogenesis.

Special Populations

Pregnant Women: No adequate and well-controlled studies have been conducted in pregnant women. TEVA-ATOMOXETINE should not be used during pregnancy unless the potential benefit justifies the potential risk to the fetus. The effect of atomoxetine hydrochloride on labour and delivery in humans is unknown.

The extent of exposure in pregnancy during clinical trials was very limited.

Nursing Mothers: — Atomoxetine and/or its metabolites were excreted in the milk of rats. It is not known if atomoxetine is excreted in human milk. Caution should be exercised if

atomoxetine hydrochloride is administered to a nursing woman.

Pediatrics (<6 years of age) — The safety and efficacy of atomoxetine hydrochloride in pediatric patients less than 6 years of age have not been established.

Pediatrics (6–18 years of age)

Risk of Suicide-Related Behaviours and Ideation in Children (*see also* WARNINGS AND PRECAUTIONS, POTENTIAL ASSOCIATION WITH THE OCCURRENCE OF BEHAVIOURAL AND EMOTIONAL CHANGES, INCLUDING SELF-HARM).

Pediatric Placebo-Controlled Clinical Trial Data: An increased risk over placebo, for suicide-related events in children and adolescents taking atomoxetine hydrochloride was identified in a combined analysis of 12 short-term (6–18 weeks) placebo-controlled trials (11 in ADHD and 1 in enuresis). Of 1357 patients who received atomoxetine hydrochloride, 5 (0.37%) had reports of **suicidal ideation** compared to 0% of 851 patients who received placebo. In addition, one **suicide attempt** (overdose) was identified which occurred in a atomoxetine hydrochloride patient. These 6 events occurred in atomoxetine hydrochloride patients 7 to 12 years of age who were male. There were no events in older adolescents, who comprised about 25 percent of the study population. Time to onset ranged from 9 to 32 days, and doses ranged from 0.48 to 1.40 mg/kg/day. A similar analysis in adult patients treated with atomoxetine hydrochloride for either ADHD or major depressive disorder (MDD) found no increased risk over placebo of suicidal ideation or behavior with the use of atomoxetine hydrochloride.

Not included in these numerators were 6 cases (3 in the atomoxetine hydrochloride arms and 3 in the placebo arms) of non-fatal potentially self-injurious actions where the intent is unknown, including burns and taking more than one dose of medication at a time.

Post Marketing Data: There have been very rare reports of suicidal ideation, suicidal attempts, suicidal depression and completed suicides in children and adolescents (*see* ADVERSE REACTIONS, Post-Market Adverse Drug Reactions, Tables 5 and 6).

Rigorous clinical monitoring for suicidal ideation or other indicators of potential for suicidal behaviour is advised in patients of all ages. This includes monitoring for agitation-type of emotional and behavioural changes, and clinical worsening.

Irritability and Mood Swings: Clinical Trial Data

Clinical trial data in children and adolescents show higher rates than placebo for irritability, mood swings, aggression, crying and tearfulness (*see* ADVERSE REACTIONS, Tables 1 and 2). The relationship, if any, between these events and suicide-related behaviours in children and adolescents with ADHD is unclear.

Aggressive Behaviour or Hostility

Aggressive behaviour or hostility has been observed in children and adolescents with ADHD,

and has been reported with some medications indicated for the treatment of ADHD. Although there is no conclusive evidence that atomoxetine causes aggressive behaviour or hostility, this was observed more frequently in clinical trials among children and adolescents treated with atomoxetine compared to placebo (overall risk ratio of 1.33 – not statistically significant). Patients beginning treatment for ADHD should be monitored for the appearance of, or worsening of, aggressive behaviour or hostility.

Caregivers/patients should be instructed to call their doctor as soon as possible should they notice an increase in aggression or hostility.

Animal Data: Growth and Neurobehavioural/Sexual Development

A study was conducted in young rats to evaluate the effects of atomoxetine on growth and neurobehavioral and sexual development. Rats were treated with 1, 10, or 50 mg/kg/day (approximately 0.2, 2, and 8 times, respectively, the maximum human dose on a mg/m² basis) of atomoxetine given by gavage from the early postnatal period (Day 10 of age) through adulthood. Slight delays in onset of vaginal patency (all doses) and preputial separation (10 and 50 mg/kg), slight decreases in epididymal weight and sperm number (10 and 50 mg/kg), and a slight decrease in corpora lutea (50 mg/kg) were seen, but there were no effects on fertility or reproductive performance. A slight delay in onset of incisor eruption was seen at 50 mg/kg. A slight increase in motor activity was seen on Day 15 (males at 10 and 50 mg/kg and females at 50 mg/kg) and on Day 30 (females at 50 mg/kg) but not on Day 60 of age. There were no effects on learning and memory tests. The significance of these findings to humans is unknown.

Geriatrics (> 65 years of age) — The safety and efficacy of atomoxetine hydrochloride in geriatric patients have not been established.

Monitoring and Laboratory Tests

Routine laboratory tests are not required.

CYP2D6 metabolism — Poor metabolizers (PMs) of CYP2D6 have a 10-fold higher AUC and a 5-fold higher peak concentration to a given dose of atomoxetine hydrochloride compared with extensive metabolizers (EMs). Approximately 7% of a Caucasian population are PMs. Laboratory tests are available to identify CYP2D6 PMs. The blood levels in PMs are similar to those attained by taking strong inhibitors of CYP2D6. The higher blood levels in PMs lead to a higher rate of some adverse effects of atomoxetine hydrochloride (*see* ADVERSE REACTIONS).

ADVERSE REACTIONS

Atomoxetine hydrochloride was administered to 3262 children or adolescent patients with ADHD and 471 adults with ADHD in clinical studies. During the ADHD clinical trials, 1409 patients (1236 pediatric and 173 adults) were treated for longer than 1 year and 1940 patients (1704 pediatric and 236 adults) were treated for over 6 months.

The data in the following tables and text cannot be used to predict the incidence of side effects in the course of usual medical practice where patient characteristics and other factors differ from those that prevailed in the clinical trials. Similarly, the cited frequencies cannot be compared with data obtained from other clinical investigations involving different treatments, uses, or investigators. The cited data provide the prescribing physician with some basis for estimating the relative contribution of drug and non-drug factors to the adverse event incidence in the population studied.

Child and Adolescent Clinical Trials

Reasons for discontinuation of treatment due to adverse events in child and adolescent clinical trials — In acute child and adolescent placebo-controlled trials, 4.1% (27/661) of atomoxetine subjects and 1.2% (5/410) placebo subjects discontinued for adverse events. For all studies, (including open-label and long-term studies), 5.8% of extensive metabolizer (EM) patients and 8.9% of poor metabolizer (PM) patients discontinued because of an adverse event. Among atomoxetine hydrochloride -treated patients, somnolence (0.8%, N=5); aggression (0.5%, N=3); irritability (0.5%, N=3); vomiting (0.5%, N=3) and abdominal pain (0.3%, N=2) were the reasons for discontinuation reported by more than 1 patient.

Commonly observed adverse events in acute child and adolescent, placebo-controlled trials — Commonly observed adverse events associated with the use of atomoxetine hydrochloride (incidence of 2% or greater) and not observed at an equivalent incidence among placebo-treated patients (atomoxetine hydrochloride incidence greater than placebo) are listed in Table 1 for all acute placebo-controlled trials. Results were similar in the BID trials and QD trials except as shown in Table 2, which shows both BID and QD results for selected adverse events. The most commonly observed adverse events in patients treated with atomoxetine hydrochloride (incidence of 5% or greater and at least twice the incidence in placebo patients, for either BID or QD dosing) were: appetite decreased, dizziness, dyspepsia, fatigue and/or lethargy, irritability, nausea, somnolence, and vomiting (*see* Table 2).

Table 1: Common Treatment-Emergent Adverse Events Reported in Atomoxetine Hydrochloride Placebo-Controlled Clinical Trials in Children and Adolescents with ADHD^a

Adverse Event	Percentage of Patients Reporting Events	
	atomoxetine hydrochloride (N=657)	Placebo (N=408)
Gastrointestinal Disorders		
Abdominal pain upper	18	13
Dyspepsia	5	1
Nausea	9	6

Adverse Event	Percentage of Patients Reporting Events	
	atomoxetine hydrochloride (N=657)	Placebo (N=408)
Vomiting	11	6
General Disorders		
Fatigue and/or Lethargy	8	4
Investigations		
Weight decreased	2	0
Metabolism and Nutritional Disorders		
Anorexia	2	<1
Appetite decreased	16	6
Nervous System Disorders		
Dizziness	5	2
Headache	21	20
Somnolence	10	4
Psychiatric Disorders		
Irritability	7	4
Mood swings	2	<1
Respiratory Disorders		
Rhinorrhea	4	2
Skin and Subcutaneous Tissue Disorders		
Rash	3	1

^a Events reported by at least 2% of patients treated with atomoxetine, and greater than placebo. The following events did not meet this criterion but were reported by more atomoxetine-treated patients than placebo-treated patients and are possibly related to atomoxetine treatment: aggression, blood pressure increased, early morning awakening, flushing, mydriasis, sinus tachycardia, crying, tearfulness, suicidal ideation. The following events were reported by at least 2% of patients treated with atomoxetine, and equal to or less than placebo: cough, diarrhea, insomnia, nasal congestion, nasopharyngitis, pharyngitis, pyrexia, upper respiratory tract infection.

Table 2: Common Treatment-Emergent Adverse Events Associated with the Use of atomoxetine hydrochloride in Acute (up to 9 weeks) Child and Adolescent Trials^a

Adverse Event	Percentage of Patients Reporting Events from BID Trials		Percentage of Patients Reporting Events from QD Trials	
	atomoxetine hydrochloride (N=340)	Placebo (N=207)	atomoxetine hydrochloride (N=317)	Placebo (N=201)
Gastrointestinal Disorders				
Abdominal pain	1	<1	3	<1
Abdominal pain upper	21	16	15	8
Constipation	3	1	<1	0
Dyspepsia	4	2	6	<1
Nausea	7	8	10	4
Vomiting	12	9	11	2
General Disorders				

Adverse Event	Percentage of Patients Reporting Events from BID Trials		Percentage of Patients Reporting Events from QD Trials	
	atomoxetine hydrochloride (N=340)	Placebo (N=207)	atomoxetine hydrochloride (N=317)	Placebo (N=201)
Fatigue and/or Lethargy	5	5	11	2
Pyrexia	5	7	6	4
Infections and Infestations				
Ear infection	3	1	1	<1
Influenza	3	1	1	<1
Pharyngitis streptococcal	2	<1	<1	<1
Investigations				
Weight decreased	3	0	2	0
Metabolism and Nutritional Disorders				
Anorexia	2	<1	3	<1
Appetite decreased	13	6	19	5
Nervous System Disorders				
Dizziness	6	3	4	<1
Headache	28	25	14	15
Sedation	1	1	3	1
Somnolence	7	5	14	3
Psychiatric Disorders				
Aggression	1	1	3	<1
Crying	2	1	1	0
Mood swings	2	0	3	1
Irritability	8	5	6	3
Respiratory Disorders				
Cough	11	7	6	9
Rhinorrhoea	4	3	3	1
Skin and Subcutaneous Tissue Disorders				
Rash	4	1	2	1

^a Events reported by at least 2% of patients treated with atomoxetine, and greater than placebo in either BID or QD trials.

The following adverse events occurred in at least 2% of PM patients and were either twice as frequent or statistically significantly more frequent in PM patients compared with EM patients: appetite decreased (24% of PMs, 17% of EMs); insomnia and middle insomnia (14% of PMs, 8% of EMs); sedation (4% of PMs, 2% of EMs); depression and/or depressed mood (5% of PMs, 3% of EMs); abrasion (4% of PMs, 2% of EMs); tremor (5% of PMs, 1% of EMs); early morning awakening (2% of PMs, 1% of EMs); enuresis (3% of PMs, 1% of EMs); pruritus (3% of PMs, 1% of EMs); mydriasis (2% of PMs, 1% of EMs); conjunctivitis (3% of PMs, 1% of EMs); syncope (2% of PMs, 1% of EMs); animal bite (2% of PMs, 1% of EMs).

Adult Clinical Trials

Reasons for discontinuation of treatment due to adverse events in acute adult placebo-controlled trials — In the acute adult placebo-controlled trials, 8.5% (23/270) atomoxetine subjects and 3.4% (9/266) placebo subjects discontinued for adverse events. Among atomoxetine hydrochloride -treated patients, insomnia (1.1%, N=3); chest pain (0.7%, N=2); palpitations (0.7%, N=2); and urinary retention (0.7%, N=2) were the reasons for discontinuation reported by more than 1 patient.

Commonly observed adverse events in acute adult placebo-controlled trials — Commonly observed adverse events associated with the use of atomoxetine hydrochloride (incidence of 2% or greater) and not observed at an equivalent incidence among placebo-treated patients (atomoxetine hydrochloride incidence greater than placebo) are listed in Table 3.

The most commonly observed adverse events in patients treated with atomoxetine hydrochloride (incidence of 5% or greater and at least twice the incidence in placebo patients) were: constipation, dry mouth, nausea, appetite decreased, dizziness, insomnia, decreased libido, ejaculatory problems, erectile disturbance, urinary hesitation and/or urinary retention and/or difficulty in micturition, and dysmenorrhea (*see* Table 3).

Table 3: Common Treatment-Emergent Adverse Events Associated with the Use of atomoxetine hydrochloride in Acute (up to 10 weeks) Adult Trials

Adverse Event ^a	Percentage of Patients Reporting Event	
	atomoxetine hydrochloride (N=269)	Placebo (N=263)
Cardiac Disorders		
Palpitations	4	1
Gastrointestinal Disorders		
Constipation	10	4
Dry mouth	21	6
Dyspepsia	6	4
Flatulence	2	1
Nausea	12	5
General Disorders		
Fatigue and/or Lethargy	7	4
Pyrexia	3	2
Chills	3	1
Infections		
Sinusitis	6	4
Investigations		
Weight decreased	2	1
Metabolism and Nutritional Disorders		
Appetite decreased	10	3
Musculoskeletal Disorders		
Myalgia	3	2

Adverse Event ^a	Percentage of Patients Reporting Event	
	atomoxetine hydrochloride (N=269)	Placebo (N=263)
Nervous System Disorders		
Dizziness	6	2
Headache	17	17
Insomnia and/or middle insomnia	17	8
Paraesthesia	4	2
Sinus headache	3	1
Psychiatric Disorders		
Abnormal dreams	4	3
Libido decreased	6	2
Sleep disorder	4	2
Renal and Urinary Disorders		
Urinary hesitation and/or urinary retention and dysuria	8	0
Reproductive System and Breast Disorders		
Dysmenorrhea ^c	7	3
Ejaculation failure ^b and/or ejaculation disorder ^s	5	2
Erectile dysfunction ^b	7	1
Impotence ^b	3	0
Menses delayed ^c	2	1
Menstrual disorder ^s	3	2
Menstruation irregular ^c	2	0
Orgasm abnormal	2	1
Prostatitis ^b	3	0
Skin and Subcutaneous Tissue Disorders		
Rash	2	1
Sweating increased	4	1
Vascular Disorders		
Hot flushes	3	1

^a Events reported by at least 2% of patients treated with atomoxetine, and greater than placebo. The following events did not meet this criterion but were reported by more atomoxetine-treated patients than placebo-treated patients and are possibly related to atomoxetine treatment: early morning awakening, peripheral coldness, tachycardia. The following events were reported by at least 2% of patients treated with atomoxetine, and equal to or less than placebo: abdominal pain upper, arthralgia, back pain, cough, diarrhea, influenza, irritability, nasopharyngitis, sore throat, upper respiratory tract infection, vomiting.

^b Based on total number of males (atomoxetine hydrochloride, N=174; placebo, N=172).

^c Based on total number of females (atomoxetine hydrochloride, N=95; placebo, N=91).

Male and female sexual dysfunction — Atomoxetine appears to impair sexual function in some patients. Changes in sexual desire, sexual performance, and sexual satisfaction are not well assessed in most clinical trials because they need special attention and because patients and physicians may be reluctant to discuss them. Accordingly, estimates of the incidence of untoward sexual experience and performance cited in product labelling are likely to

underestimate the actual incidence. The table below displays the incidence of sexual side effects reported by at least 2% of adult patients taking atomoxetine hydrochloride in placebo-controlled trials.

Table 4

	Atomoxetine hydrochloride	Placebo
Erectile dysfunction ^a	7%	1%
Libido decreased	6%	2%
Ejaculation failure ^a and/or ejaculation disorder ^a	5%	2%
Impotence ^a	3%	0%
Orgasm abnormal	2%	1%

^a

Males only.

There are no adequate and well-controlled studies examining sexual dysfunction with atomoxetine hydrochloride treatment. While it is difficult to know the precise risk of sexual dysfunction associated with the use of atomoxetine hydrochloride, physicians should routinely inquire about such possible side effects.

Post-Market Adverse Drug Reactions

During the first 5 years of post-market experience, it is estimated that over 5 million patients have been treated with atomoxetine hydrochloride, for 1,715,000 patient-years of therapy.

Tables 5 and 6 are based on post-market spontaneous adverse event reports. The percentages shown are calculated by dividing the number of adverse events reported to the company by the estimated number of patients exposed to the drug during the same time period. The causal relationship between atomoxetine hydrochloride and the emergence of these events has not been established.

Table 5: Atomoxetine hydrochloride Post-Market Spontaneous Adverse Event Reports in Children and Adolescents with ADHD.

Adverse Event	Frequency			
	≥1%	<1% and ≥0.1%	<0.1% and ≥0.01%	<0.01%
Cardiac Disorders				
Palpitations				X
Sinus tachycardia			X	
Electrocardiogram QT prolonged ^a				X
Eye Disorders				
Mydriasis			X	
Gastrointestinal Disorders				
Abdominal pain			X	
Dyspepsia				X
Hepatobiliary effects				X
Liver Function Tests abnormal				X

Adverse Event	Frequency			
	≥1%	<1% and ≥0.1%	<0.1% and ≥0.01%	<0.01%
Nausea			X	
Vomiting			X	
General Disorders				
Sudden death				X
Injury				
Overdose				X
Investigations				
Weight decreased			X	
Metabolism and Nutritional Disorders				
Anorexia			X	
Appetite decreased			X	
Nervous System Disorders				
Dizziness			X	
Seizure ^b			X	
Somnolence		X		
Syncope ^c				X
Tics				X
Psychiatric Disorders				
Aggression/Hostility			X	
Anxiety			X	
Depression and depressed mood			X	
Early morning awakening				X
Irritability			X	
Mood swings			X	
Suicidal ^d				X
Skin and Subcutaneous Tissue Disorders				
Pruritis				X
Rash			X	
Urogenital Disorders				
Painful or prolonged penile erection				X
Vascular Disorders				
Peripheral vascular instability, e.g., Raynaud's phenomenon				X

^a These spontaneously reported cases are not well documented, and the method of correction is unknown

^b Frequency of seizures reported = 0.01%

^c Includes reports of 'loss of consciousness'.

^d Includes reports of completed suicide, suicidal ideation, suicide attempt and suicidal depression.

Table 6: Atomoxetine hydrochloride Post-Market Spontaneous Adverse Event Reports in Adult Patients with ADHD.

Adverse Event	Frequency			
	≥1%	<1% and ≥0.1%	<0.1% and ≥0.01%	<0.01%
Cardiac Disorders				
Palpitations				X
Tachycardia			X	
Electrocardiogram QT prolonged ^a				X
Gastrointestinal Disorders				
Abdominal pain			X	
Constipation			X	
Dry mouth			X	
Dyspepsia				X
Flatulence				X
Hepatobiliary effects				X
Liver Function Tests abnormal				X
Nausea			X	
General Disorders				
Fatigue			X	
Lethargy				X
Chills			X	
Sudden death				X
Injury				
Overdose				X
Investigations				
Weight decreased			X	
Metabolism and Nutritional Disorders				
Appetite decreased			X	
Nervous System Disorders				
Dizziness			X	
Insomnia			X	
Middle insomnia				X
Seizure ^b				X
Sinus headache				X
Syncope ^c				X
Tics				X
Psychiatric Disorders				
Anxiety			X	
Depression and depressed mood			X	
Early morning awakening				X
Libido decreased				X
Sleep disorder				X
Suicidal ^d				X

Adverse Event	Frequency			
	≥1%	<1% and ≥0.1%	<0.1% and ≥0.01%	<0.01%
Renal and Urinary Disorders				
Difficulty in micturition			X	
Urinary hesitation			X	
Urinary retention			X	
Reproductive System and Breast Disorders				
Dysmenorrhea				X
Ejaculation disorder			X	
Ejaculation failure				X
Erectile dysfunction			X	
Male genital pain				X
Menstruation irregular				X
Orgasm abnormal				X
Prostatitis				X
Skin and Subcutaneous Tissue Disorders				
Dermatitis				X
Sweating increased			X	
Vascular Disorders				
Hot flushes			X	
Peripheral coldness				X
Peripheral vascular instability, e.g., Raynaud's phenomenon				X

^a These spontaneously reported cases are not well documented, and the method of correction is unknown.

^b Frequency of seizures reported = 0.0035%

^c Includes reports of 'loss of consciousness'.

^d Includes reports of completed suicide, suicidal ideation, suicide attempt, and suicidal depression.

In post-market experience, serious skin reactions were reported at a spontaneous reporting rate less than 0.001%.

DRUG ABUSE AND DEPENDENCE

Atomoxetine hydrochloride is not a controlled substance.

Atomoxetine hydrochloride is not a stimulant drug. In a randomized, double-blind, placebo-controlled, abuse-potential study in adults comparing effects of atomoxetine hydrochloride and placebo, atomoxetine hydrochloride was not associated with a pattern of response that suggested stimulant or euphoriant properties.

Clinical trials data in over 4000 children, adolescents, and adults with ADHD showed only isolated incidents of drug diversion or inappropriate self-administration associated with atomoxetine hydrochloride. There was no evidence of symptom rebound or adverse events suggesting a drug-discontinuation or withdrawal syndrome.

In preclinical studies, atomoxetine did not show a behavioural profile or stimulant properties associated with drugs that have abuse liability.

DRUG INTERACTIONS

Overview

Atomoxetine hydrochloride is primarily metabolized by the CYP2D6 pathway to 4-hydroxyatomoxetine. In extensive metabolizers (EMs), inhibitors of CYP2D6 (e.g., paroxetine, fluoxetine, quinidine) increase atomoxetine steady-state plasma concentrations to exposures similar to those observed in poor metabolizers (PMs). Dosage adjustment of atomoxetine hydrochloride may be necessary when coadministered with CYP2D6 inhibitors, e.g., paroxetine, fluoxetine, and quinidine (*see* DOSAGE AND ADMINISTRATION). In EM individuals treated with paroxetine or fluoxetine, the AUC of atomoxetine is approximately 6- to 8-fold and $C_{ss,max}$ is about 3- to 4-fold greater than atomoxetine alone.

In vitro studies suggest that coadministration of cytochrome P450 inhibitors to PMs will not further increase the plasma concentration of atomoxetine.

Atomoxetine did not cause clinically significant inhibition or induction of cytochrome P450 enzymes, including CYP1A2, CYP3A, CYP2D6, and CYP2C9.

Drugs that affect norepinephrine should be used cautiously when co-administered with atomoxetine hydrochloride because of the potential for additive or synergistic pharmacological effects.

Drug-Drug Interactions

Table 7: Established or Potential Drug-Drug Interactions with Atomoxetine hydrochloride

Drug	Ref	Effect	Clinical comment
MAO Inhibitors	T	There have been reports of serious, sometimes fatal, reactions when MAO Inhibitors are given concurrently or in close proximity to other drugs that affect brain monoamine concentrations.	See CONTRAINDICATIONS
Desipramine	CT	Coadministration of Atomoxetine hydrochloride with desipramine, a model compound for CYP2D6 metabolized drugs, did not alter the pharmacokinetics of desipramine.	Because desipramine has noradrenergic effects, it should not be used in combination with Atomoxetine hydrochloride.
Fluoxetine, Paroxetine	CT	Coadministration of selective inhibitors of CYP2D6 (e.g. fluoxetine, paroxetine) may increase atomoxetine steady-state	Slower titration of Atomoxetine hydrochloride may be necessary in those patients who are also taking fluoxetine, paroxetine or

Drug	Ref	Effect	Clinical comment
		plasma concentrations to exposures similar to those observed in CYP2D6 poor metabolizer patients.	other CYP2D6 inhibitor drugs (<i>see</i> Dosage and Administration, Special Populations).
Salbutamol or other β -adrenergic receptor agonists	CT	In adult volunteers, the effects on blood pressure and heart rate of a standard inhaled dose of salbutamol (200 mcg) are not clinically significant compared to IV administration, and not increased by coadministration of atomoxetine (80 mg QD for 5 days). Heart rate after multiple inhalations of salbutamol (800 mcg) was similar in the presence or absence of atomoxetine.	Atomoxetine may be used in combination with inhaled β_2 agonists, e.g., salbutamol, but should be used with caution in patients being treated with systemically administered (oral or intravenous) β_2 agonists, including atomoxetine.
Pressor Agents	T	Possible effects on blood pressure.	Atomoxetine hydrochloride should be used with caution in patients being treated with pressor agents.
Methylphenidate	CT	Co-administration of methylphenidate with Atomoxetine hydrochloride did not increase cardiovascular effects beyond those seen with methylphenidate administration alone.	
Midazolam	CT	Co-administration of Atomoxetine hydrochloride with midazolam, a model compound for CYP3A4 metabolized drugs, resulted in small increases in midazolam plasma concentrations.	No dosage adjustment required.
Drugs highly bound to plasma protein	In vitro	Atomoxetine did not affect the binding of <i>warfarin</i> , <i>acetylsalicylic acid</i> , <i>phenytoin</i> , or <i>diazepam</i> to human albumin in-vitro. Similarly, these compounds did not affect the binding of atomoxetine to human albumin.	No dosage adjustment required.
Drugs affecting gastric pH	CT	Drugs that elevate gastric pH (<i>magnesium hydroxide/aluminum hydroxide</i> , <i>omeprazole</i>) had no effect on Atomoxetine hydrochloride bioavailability.	No dosage adjustment required.

Legend: C = Case Study; CT = Clinical Trial; T = Theoretical

Drug-Food Interactions

TEVA-ATOMOXETINE may be taken with or without food.

Drug-Lifestyle Interactions

Alcohol: Consumption of ethanol with atomoxetine hydrochloride did not change the intoxicating effects of ethanol.

DOSAGE AND ADMINISTRATION

Dosing Considerations

TEVA-ATOMOXETINE should be administered starting at the lowest possible dose. Dosage should then be individually and slowly adjusted to the lowest effective dose, since individual patient response to TEVA-ATOMOXETINE varies widely.

TEVA-ATOMOXETINE should not be used in patients with symptomatic cardiovascular disease and should not generally be used in patients with known structural cardiac abnormalities (See CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS).

Children: Theoretically there exists a pharmacological potential for all ADHD drugs to increase the risk of sudden/cardiac death. Although confirmation of an incremental risk for adverse cardiac events arising from treatment with ADHD medications is lacking, prescribers should consider this potential risk.

All drugs with sympathomimetic effects prescribed in the management of ADHD should be used with caution in patients who: *a)* are involved in strenuous exercise or activities, *b)* use stimulants, or *c)* have a family history of sudden/cardiac death. Prior to the initiation of treatment with sympathomimetic medications, a personal and family history (including assessment for a family history of sudden death or ventricular arrhythmia) and physical exam should be obtained to assess for the presence of cardiac disease. In patients with relevant risk factors and based on the clinician's judgment, further cardiovascular evaluation may be considered (e.g., electrocardiogram and echocardiogram). Patients who develop symptoms such as exertional chest pain, unexplained syncope, or other symptoms suggestive of cardiac disease during ADHD treatment should undergo a prompt cardiac evaluation. Patients who are considered to need extended treatment with TEVA-ATOMOXETINE should undergo periodic evaluation of their cardiovascular status. (see WARNINGS AND PRECAUTIONS).

Rigorous clinical monitoring for suicidal ideation or other indicators of potential for suicidal behavior is advised in patients of all ages. This includes monitoring of agitation-type emotional and behavioral changes, and clinical worsening (see WARNING AND PRECAUTIONS, POTENTIAL ASSOCIATION WITH THE OCCURENCE OF BEHAVIOURAL AND EMOTIONAL CHANGES, INCLUDING SELF-HARM)

TEVA-ATOMOXETINE (atomoxetine hydrochloride) is intended for oral administration and may be taken with or without food, either as a single daily dose in the morning or as divided doses in the morning and late afternoon/early evening.

Initial improvement of ADHD symptoms is generally observed within 1 to 4 weeks of initiating therapy.

Atomoxetine hydrochloride did not worsen tics in pediatric patients, and may be used in patients with ADHD and comorbid motor tics or diagnosis of Tourette's Disorder. Atomoxetine hydrochloride did not worsen anxiety in either pediatric or adult patients, and may be used in patients with ADHD and comorbid anxiety disorders. See ACTION AND CLINICAL

PHARMACOLOGY, Special Populations and Conditions, Patients with Concomitant Illness.

If patients miss a dose, they should take it as soon as possible; however, they should not take more than the prescribed total daily amount of TEVA-ATOMOXETINE in any 24-hour period.

TEVA-ATOMOXETINE may be discontinued without tapering the dose.

Recommended Dose and Dosage Adjustment

Children (6 years and over) and Adolescents up to 70 kg Body Weight:

Do not exceed the recommended initial dose and subsequent dose escalations of TEVA-ATOMOXETINE. More rapid dose escalation may be associated with increased rates of somnolence and digestive system complaints. Do not exceed the recommended maximum total daily dose of 1.4 mg/kg or 100 mg, whichever is less. No additional benefit has been demonstrated at doses greater than 1.2mg/kg/day (see Clinical Trials). The safety of single doses over 1.8 mg/kg/day and total daily doses above 1.8 mg/kg have not been systematically evaluated, and therefore should not be administered because of potential side-effects (see ACTION AND CLINICAL PHARMACOLOGY- Cardiovascular Safety, OVERDOSAGE and DETAILED PHARMACOLOGY [Safety Pharmacology, Cardiac Function and Pharmacokinetics]).

TEVA-ATOMOXETINE should be initiated at a total daily dose of approximately 0.5 mg/kg/day (Step 1) for 7 to 14 days. Based on tolerability, the dose should be successively increased to approximately 0.8 mg/kg/day (Step 2) for 7 to 14 days, and then to approximately 1.2 mg/kg/day (Step 3). Table 8 below provides for each titration step the corresponding daily atomoxetine hydrochloride dose according to body weight. After a minimum of 30 days, the maintenance dose should be reassessed and adjusted according to clinical response.

The total daily dose in children and adolescents up to 70 kg should not exceed 1.4 mg/kg or 100 mg, whichever is less.

As the lowest available capsule strength is 10 mg, the child should weigh at least 20 kg at the time of initiation of therapy. Only whole capsules should be administered.

Table 8 Daily Atomoxetine Hydrochloride Dose for Each Titration Step as per Body Weight, in Children and Adolescents up to 70 kg Body Weight.

Body Weight	Step 1 (approx. 0.5 mg/kg/day)	Step 2 (approx. 0.8 mg/kg/day)	Step 3 (approx. 1.2 mg/kg/day)	Maximum Dose
20-29 kg	10 mg/day	18 mg/day	25 mg/day	1.4 mg/kg/day or 100 mg/day (whichever is less)
30-44 kg	18 mg/day	25 mg/day	40 mg/day	
45-64 kg	25 mg/day	40 mg/day	60 mg/day	
65-70 kg	40 mg/day	60 mg/day	80 mg/day	

Children and Adolescents over 70 kg Body Weight, and Adults

Do not exceed the recommended initial dose and subsequent dose escalations of TEVA-ATOMOXETINE. Do not exceed the recommended maximum total daily dose of 100 mg. The safety of single doses over 120 mg and total daily doses above 150 mg have not been systematically evaluated and therefore should not be administered because of potential side effects (see ACTION AND CLINICAL PHARMACOLOGY- Cardiovascular Safety, OVERDOSAGE and DETAILED PHARMACOLOGY [Safety Pharmacology, Cardiac Function and Pharmacokinetics]).

TEVA-ATOMOXETINE should be initiated at a total daily dose of 40 mg/day (Step 1) for 7 to 14 days. Based on tolerability, the dose should be successively increased to 60 mg/day (Step 2) for 7 to 14 days, and then to 80 mg/day (Step 3). After 2 to 4 additional weeks, the total daily dose may be increased to a maximum of 100 mg in patients who have not achieved an optimal response.

The maximum recommended total daily dose in children and adolescents over 70 kg and adults is 100 mg.

Dosage Adjustment for Special Populations

Hepatic Impaired

Atomoxetine clearance may be reduced in ADHD patients with hepatic impairment. For patients with moderate hepatic impairment (Child-Pugh Class B), initial and target doses should be reduced to 50% of the normal dose. For patients with severe hepatic impairment (Child-Pugh Class C), initial dose and target doses should be reduced to 25% of the normal dose.

Renal Impaired

Subjects with end stage renal disease had higher systemic exposure to atomoxetine than healthy subjects (about a 65% increase), but there was no difference when exposure was corrected for mg/kg dose. Atomoxetine can therefore be administered to ADHD patients with end stage renal disease or lesser degrees of renal insufficiency using the usual dosing regimen. Atomoxetine may exacerbate hypertension in patients with end stage renal disease.

Dosing adjustment for use with a strong CYP2D6 inhibitor

In children (≥ 6 years old) and adolescents up to 70 kg body weight administered strong CYP2D6 inhibitors, e.g., paroxetine, fluoxetine, and quinidine, TEVA-ATOMOXETINE should be initiated at 0.5 mg/kg/day and only increased to the next dose level if symptoms fail to improve after 14 days and the previous dose is well tolerated.

In children (≥ 6 years old) and adolescents over 70 kg body weight and adults administered strong CYP2D6 inhibitors, e.g., paroxetine, fluoxetine, and quinidine, TEVA-ATOMOXETINE should be initiated at 40 mg/day and only increased to the next dose level if symptoms fail to

improve after 14 days and the previous dose is well tolerated.

Maintenance/Extended Treatment

Pharmacological treatment of ADHD may be needed for extended periods. The efficacy of atomoxetine hydrochloride in maintaining symptom response during long-term treatment in children and adolescents was studied in an 18-month trial (3 months of acute open-label treatment followed by up to 15 months of placebo-controlled maintenance treatment). The results from this study suggest that atomoxetine may be beneficial in the long-term treatment of ADHD. Too few patients completed the study to permit an adequate assessment of the long-term safety profile of atomoxetine hydrochloride in this study. The long-term safety of atomoxetine hydrochloride has been demonstrated in double-blind and open-label clinical trials of at least 24 months. The physician who elects to use atomoxetine hydrochloride for extended periods should periodically reevaluate the long-term usefulness of the drug for the individual patient (see INDICATIONS AND CLINICAL USE).

OVERDOSAGE

There is limited clinical trial experience with atomoxetine overdose and no fatalities were observed in clinical studies.

During post-marketing, there have been reports of non-fatal, acute and chronic overdoses of atomoxetine. The most commonly reported symptoms accompanying acute and chronic overdoses were somnolence, agitation, hyperactivity, abnormal behaviour, and gastrointestinal symptoms. Signs and symptoms consistent with mild to moderate sympathetic nervous system activation (e.g., mydriasis, tachycardia, dry mouth) have also been observed. Seizures have been reported in several cases, including myoclonus of the extremities. Less commonly, there have been reports of QT prolongation and mental changes, including disorientation and hallucinations. All patients recovered from these events.

In the first 18 months of market availability in the US, among the mixed overdose reports involving atomoxetine where at least one other drug was taken in overdose amounts, there were 3 deaths reported, all in adults. The largest quantity of atomoxetine alone in a single overdose was 1400 mg, taken by a 17 year old patient. He experienced chest pain and drowsiness and was treated with activated charcoal about 2 hours after the ingestion and recovered fully after an overnight hospital stay for observation.

There were no deaths involving overdose of atomoxetine hydrochloride alone.

Management of Overdose: There is no established antidote for atomoxetine hydrochloride overdose. Treatment has been supportive, including establishing an airway when necessary, monitoring of cardiac and vital signs, along with appropriate symptomatic and supportive measures. Gastric lavage may be indicated if performed soon after ingestion. Activated charcoal may be useful in limiting absorption. Because atomoxetine is highly protein-bound, dialysis is not likely to be useful in the treatment of overdose.

For management of a suspected drug overdose, contact your regional Poison Control Centre.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Atomoxetine hydrochloride is a selective norepinephrine reuptake inhibitor. Its therapeutic effect in ADHD is thought to be related to its potent inhibition of the pre-synaptic norepinephrine transporter, with minimal affinity for other noradrenergic receptors or for other neurotransmitter transporters or receptors.

Pharmacodynamics

In ex vivo uptake and neurotransmitter depletion studies, atomoxetine was found to selectively inhibit the pre-synaptic norepinephrine transporter without directly affecting the serotonin or dopamine transporters. Atomoxetine has minimal affinity for other receptor systems.

Atomoxetine is primarily oxidized to 4-hydroxyatomoxetine, which is also a potent inhibitor of the pre-synaptic norepinephrine transporter.

Cardiovascular Safety

The safety and tolerability of gradually increasing multiple-dose regimes of atomoxetine 60 to 150 mg/day was studied in 16 healthy adults (10 EM subjects and 6 PM subjects). None of the mean or individual QT_c(F) intervals exceeded the upper limits of normal for each gender. The EM group had no statistically significant changes in mean QT_c(F) interval compared to the placebo treatment. No statistically significant changes in QT_c(F) were noted 1 hour post dose (during peak plasma concentrations) in the PM group. The PM group had a statistically significant increase in the mean QT_c(F) interval measured at time 0 (during trough plasma concentrations) on the last day of the 60- and 75-mg atomoxetine twice-daily dosing regimens compared to the placebo. The greatest mean prolongation was about 17 msec at the 60-mg BID dose level, with the mean interval length of 417.2 msec. At the 75-mg BID atomoxetine dose level, the greatest mean prolongation was 15 msec, and the mean interval length was 414.9 msec. The 60 mg BID and 75 mg BID doses correspond to 1.4-2.24 mg/kg/day and 1.75-2.8 mg/kg/day, respectively. Baseline ECGs obtained during screening of pediatric patients for atomoxetine clinical trials were reviewed for cases of QT_c prolongation. Using a correction method based on data from baseline ECGs, there were 32/3902 cases (0.8%) with QT_c(D)>450 msec and 5/3902 cases (0.1%) with QT_c(D)>500 msec. In a meta-analysis of ECG data from patients who received atomoxetine in pediatric clinical trials, no relationship was observed between changes from baseline to final QT_c(D) and prescribed atomoxetine dose, or between changes from baseline to QT_c(D) at time of expected peak exposure and prescribed atomoxetine dose.

Overall, the data do not suggest a meaningful relationship between atomoxetine plasma concentrations and the length of the QT interval corrected for heart rate in the recommended dosage range. However, since there is no requirement for a priori screening of ADHD patients for CYP2D6 metabolizer status before initiating treatment with atomoxetine, it is important that the lowest effective dose be used, so as to minimize potential cardiac side effects.

Pharmacokinetics

Atomoxetine is well-absorbed after oral administration and is minimally affected by food. It is eliminated primarily by oxidative metabolism through the cytochrome P450 2D6 (CYP2D6) enzymatic pathway and subsequent glucuronidation. Atomoxetine has a half-life of about 5 hours. A fraction of the population (about 7% of Caucasians and 2% of African Americans) are poor metabolizers (PMs) of CYP2D6 metabolized drugs. These individuals have reduced activity in this pathway resulting in 10-fold higher AUCs, 5-fold higher peak plasma concentrations, and slower elimination (plasma half-life of 21.6 hours) of atomoxetine compared with people with normal activity [extensive metabolizers (EMs)]. Drugs that inhibit CYP2D6, such as fluoxetine, paroxetine, and quinidine, cause similar increases in exposure.

The pharmacokinetics of atomoxetine have been evaluated in more than 400 children and adolescents in selected clinical trials using a population approach. Single-dose and steady-state individual pharmacokinetic data were also obtained in children, adolescents, and adults. When doses were normalized to a mg/kg basis, similar half-life, C_{max} , and AUC values were observed in children, adolescents, and adults. Clearance and volume of distribution after adjustment for body weight were also similar.

Atomoxetine pharmacokinetics are dose proportional within the therapeutic range; hence, administration of atomoxetine hydrochloride once- or twice-daily is expected to result in the same systemic exposure (AUC) over a 24-hour period. Results of efficacy analysis show that once-daily (QD) dosing with atomoxetine hydrochloride is efficacious in the treatment of ADHD.

Absorption: Atomoxetine is rapidly absorbed after oral administration, with absolute bioavailability of about 63% in extensive metabolizers (EMs) and 94% in poor metabolizers (PMs). Mean maximal plasma concentrations (C_{max}) are reached approximately 1 to 2 hours after dosing.

TEVA-ATOMOXETINE can be administered with or without food. In clinical trials with children and adolescents, administration of atomoxetine hydrochloride with food resulted in a 9% lower C_{max} . Administration of atomoxetine hydrochloride with a standard high-fat meal in adults did not affect the extent of oral absorption of atomoxetine (AUC), but did decrease the rate of absorption, resulting in a 37% lower C_{max} and delayed T_{max} by 3 hours.

Distribution: The steady-state volume of distribution after intravenous administration was approximately 0.85 L/kg indicating that atomoxetine distributes primarily into total body water. In children and adolescents, volume of distribution increased nearly proportionally to increases in body weight. Volume of distribution is similar across the patient weight range after normalizing for body weight.

At therapeutic concentrations, 98% of atomoxetine in plasma is bound to protein, primarily albumin.

Metabolism: Atomoxetine undergoes biotransformation primarily through the cytochrome P450 2D6 (CYP2D6) enzymatic pathway. People with reduced activity in the CYP2D6 pathway (PMs) have higher plasma concentrations of atomoxetine compared with people with normal activity (EMs). For PMs, AUC of atomoxetine at steady-state is approximately 10-fold higher and $C_{ss,max}$ is about 5-fold greater than for EMs.

Coadministration of atomoxetine hydrochloride with potent inhibitors of CYP2D6, such as fluoxetine, paroxetine, or quinidine, results in a substantial increase in atomoxetine plasma exposure, and dosing adjustment may be necessary (*see Drug-Drug Interactions*). In EM patients treated with potent CYP2D6 inhibitors such as fluoxetine, and paroxetine, the AUC of atomoxetine is approximately 6- to 8-fold and $C_{ss,max}$ is about 3- to 4-fold greater than with atomoxetine alone.

In vitro studies suggest that coadministration of cytochrome P450 inhibitors to PMs will not increase the plasma concentration of atomoxetine.

Atomoxetine did not inhibit or induce the CYP2D6 pathway.

The major oxidative metabolite formed regardless of CYP2D6 status is 4-hydroxy-atomoxetine, which is rapidly glucuronidated. 4-Hydroxyatomoxetine is equipotent to atomoxetine as an inhibitor of the norepinephrine transporter, but circulates in plasma at much lower concentrations (1% of atomoxetine concentration in EMs and 0.1% of atomoxetine concentration in PMs). 4-Hydroxyatomoxetine is primarily formed by CYP2D6. In individuals that lack CYP2D6 activity (poor metabolizers), 4-hydroxyatomoxetine is formed by several other cytochrome P450 enzymes, but at a slower rate. N-Desmethyl-atomoxetine is formed by CYP2C19 and other cytochrome P450 enzymes, but has much less pharmacological activity than atomoxetine, and plasma concentrations are lower (5% of atomoxetine concentration in EMs and 45% of atomoxetine concentration in PMs).

Elimination: The mean elimination half-life of atomoxetine after oral administration is 5.2 hours and 21.6 hours in EM and PM subjects, respectively. The elimination half-life of 4-hydroxyatomoxetine is similar to that of N-desmethylassatomoxetine (6 to 8 hours) in EM subjects, while the half-life of N-desmethylassatomoxetine is much longer in PM subjects (34 to 40 hours).

Atomoxetine is excreted primarily as 4-hydroxyatomoxetine-*O*-glucuronide, mainly in the urine (greater than 80% of the dose) and to a lesser extent in the feces (less than 17% of the dose). Only a small fraction (less than 3%) of the atomoxetine hydrochloride dose is excreted as unchanged atomoxetine, indicating extensive biotransformation.

Special Populations and Conditions

Pediatrics: The pharmacokinetics of atomoxetine in children and adolescents are similar to those in adults. The pharmacokinetics of atomoxetine have not been evaluated in children under 6 years of age.

Geriatrics: The pharmacokinetics of atomoxetine have not been systematically evaluated in the

geriatric population.

Gender: Gender did not influence atomoxetine disposition.

Race: Ethnic origin did not influence atomoxetine disposition.

Hepatic Insufficiency: Single doses of atomoxetine hydrochloride administered to EM subjects with moderate to severe hepatic insufficiency (Child-Pugh Class B and C) resulted in increased atomoxetine exposure, reduced atomoxetine clearance, and prolonged half-life of parent drug compared with healthy subjects. Dosage adjustment is recommended for patients with moderate or severe hepatic impairment (*see* DOSAGE AND ADMINISTRATION).

Renal Insufficiency: Single doses of atomoxetine hydrochloride administered to EM subjects with end stage renal disease resulted in higher atomoxetine exposure (AUC) than in healthy subjects (about a 65% increase), but there was no difference when exposure was corrected for mg/kg dose. Atomoxetine hydrochloride can therefore be administered to ADHD patients with end stage renal disease or lesser degrees of renal insufficiency using the normal dosing regimen.

Genetic Polymorphism: There are two major phenotypes associated with CYP2D6: extensive metabolizers that comprise >90% of the population, and poor metabolizers. Approximately 7% of the Caucasian population and 2% of Black population are poor metabolizers of CYP2D6.

Patients with Concomitant Illness

Pediatric Patients with ADHD and Comorbid Tics: Study LYAS enrolled patients meeting DSM-IV criteria for ADHD and comorbid Tourette's Disorder or chronic motor tics. Atomoxetine hydrochloride met the primary study objective of non-inferiority to placebo with respect to worsening of tics. Atomoxetine hydrochloride was also markedly superior to placebo in reducing symptoms of ADHD as assessed by the ADHDRS-IVParent: Inv Total score ($p=0.002$).

Patients with ADHD and Comorbid Anxiety Disorder

Study LYBP: Pediatric patients ($n = 176$, age 8-17), who met DSM-IV criteria for ADHD and at least one of the anxiety disorders of separation anxiety disorder, generalized anxiety disorder or social phobia were randomized 1:1 into a 12-week double-blind, placebo-controlled trial. The statistical analysis plan specified that patients responding (i.e., > 25% reduction in Pediatric Anxiety Rating Scale (PARS) total scores) during a blinded placebo lead-in period be excluded from the primary efficacy analyses. However, all of the enrolled patients with a baseline and at least one post-baseline score were included in the remaining efficacy analyses. Initial atomoxetine hydrochloride dose was 0.8 mg/kg/day, with increase to a target dose of 1.2 mg/kg/day (median dose 1.3 mg/kg/day \pm 0.3 mg/kg/day). Atomoxetine hydrochloride was markedly superior to placebo in reducing symptoms of ADHD as assessed by the ADHDRS-IV-Parent: Inv Total Score ($p<0.001$) and mean PARS total score improved significantly for atomoxetine relative to placebo ($p = 0.011$). Worsening of anxiety was defined post-hoc as a 25% increase in PARS, or patient-reported anxiety as an adverse event. Based on these criteria,

patients on atomoxetine hydrochloride did not have worsening of anxiety relative to the placebo group. Of the 158 patients who completed the double-blind placebo lead-in, 26 (16%) discontinued the study.

Study LYDQ: Adult patients (n=442, age 18-65), who met DSM-IV criteria for adult ADHD and social anxiety disorder (23% of whom also had Generalized Anxiety Disorder) were randomized into a 16-week double-blind, placebo-controlled trial. The statistical analysis plan specified that patients responding (i.e., >25% decrease in social anxiety symptoms as measured by the Liebowitz Social Anxiety Scale) during a blinded placebo lead-in period be excluded from the primary efficacy analyses. However, all of the enrolled patients with a baseline and at least one post-baseline score were included in the remaining efficacy analyses. Atomoxetine hydrochloride was initiated at 40 mg/day to a maximum dose of 100 mg/day (mean daily dose 83 mg/day \pm 19.5 mg/day). Atomoxetine hydrochloride was markedly superior to placebo in reducing symptoms of ADHD as assessed by the Conners' Adult ADHD Rating Scale ($p < 0.001$) and mean LSAS total score improved significantly for atomoxetine relative to placebo ($p \leq 0.01$). Worsening of anxiety was defined post-hoc as a 25% increase in LSAS, or patient-reported anxiety as an adverse event. Based on these criteria, patients on atomoxetine hydrochloride did not have worsening of anxiety relative to the placebo group. Of the 436 patients who completed the double-blind placebo lead-in, 172 (39%) discontinued the study.

STORAGE AND STABILITY

Store at controlled room temperature, 15 to 30°C.

TEVA-ATOMOXETINE capsules are not intended to be opened, they should be taken whole.

DOSAGE FORMS, COMPOSITION AND PACKAGING

TEVA-ATOMOXETINE (atomoxetine hydrochloride) capsules are supplied in 10, 18, 25, 40, 60 mg strengths, as atomoxetine base equivalent.

- 10 mg: Hard gelatin capsules with a white opaque cap and white opaque body, filled with a white to off-white powder with small agglomerates. Imprinting on body and cap: A10. They are supplied in HDPE bottles containing 100 and 500 capsules and in unit dose of 6 capsules.
- 18 mg: Hard gelatin capsules with a yellow opaque cap and white opaque body, filled with a white to off-white powder with small agglomerates. Imprinting on body and cap: A18. They are supplied in HDPE bottles containing 100 and 500 capsules and in unit dose of 10 capsules.
- 25 mg: Hard gelatin capsules with a blue opaque cap and white opaque body, filled with a white to off-white powder with small agglomerates. Imprinting on body and cap: A25. They are supplied in HDPE bottles containing 100 and 500 capsules and in unit dose of 10 capsules.

40 mg: Hard gelatin capsules with a blue opaque cap and blue opaque body, filled with a white to off-white powder with small agglomerates. Imprinting on body and cap: A40. They are supplied in HDPE bottles containing 100 and 500 capsules and in unit dose of 10 capsules.

60 mg: Hard gelatin capsules with a blue opaque cap and yellow opaque body, filled with a white to off-white powder with small agglomerates. Imprinting on body and cap: A60. They are supplied in HDPE bottles containing 100 and 500 capsules and in unit dose of 10 capsules.

Composition

In addition to Atomoxetine HCl, each capsule contains the inactive ingredients pregelatinized starch, sodium laurylsulfate and hard gelatin capsule contains gelatin, titanium dioxide, D&C Yellow 10 (18 mg capsules), FD&C Blue 1/Brilliant Blue FCF (25 mg and 40 mg capsules) and D&C yellow 10, FD&C Blue 1 and Brilliant Blue FCF (60 mg capsules).

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION

Drug Substance


Common Name: Atomoxetine Hydrochloride

Chemical Name: Benzenepropanamine N-methyl-gamma(2-methylphenoxy) hydrochloride,(-)

Molecular Formula: C₁₇H₂₁NO·HCl

Molecular Weight: 291.82

Structural Formula:



Physicochemical Properties: Atomoxetine hydrochloride is a white to off-white crystalline powder, which has a solubility of 27.8 mg/mL in water and 272.7 mg/mL in Methanol. Atomoxetine hydrochloride is the R(-) isomer, as determined by x-ray diffraction.

pH: 4.80 (1% aqueous solution)

pKa: 10.13

Melting Point: Onset temperature 168-169°C.

CLINICAL TRIALS

This is a three-way, blinded, randomized, single dose (1 x 10 mg), crossover bioequivalence study conducted in 19 healthy, non-smoking, male and female subjects 23 to 48 years of age to compare the bioavailability of atomoxetine from Teva-Atomoxetine (Teva Canada Limited), the Canadian reference product, Strattera (Eli Lilly Canada Inc.) and the UK reference product Strattera (Eli Lilly and Company Limited) under fasting condition. The results from the comparison of Teva-Atomoxetine and Strattera (Eli Lilly Canada Inc.) are presented in the table below.

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

Atomoxetine (1 x 10 mg) From measured data uncorrected for potency Geometric Mean Arithmetic Mean (CV %)				
Parameter	Teva-Atomoxetine	Strattera¹	% Ratio of Geometric Means	90% Confidence interval
AUC_T (ng·h/mL)	453.90 620.23 (121)	434.19 580.11 (118)	104.54	97.82 – 111.72
AUC_I (ng·h/mL)	468.58 677.34 (139)	450.58 630.35 (132)	103.96	97.42 – 110.93
C_{max} (ng/mL)	92.57 97.61 (33)	103.27 107.49 (31)	89.64	83.10 – 96.69
T_{max}[§] (h)	1.38 (73)	1.11 (112)		
T_{1/2}[§] (h)	3.71 (82)	3.69 (78)		

¹ Strattera (Eli Lilly Canada Inc.) was purchased in Canada

[§] Expressed as the arithmetic mean (%CV) only

This is a three-way, blinded, randomized, single dose (1 x 60 mg), crossover bioequivalence study conducted in 21 healthy, non-smoking, male and female subjects 24 to 54 years of age to compare the bioavailability of atomoxetine from Teva-Atomoxetine (Teva Canada Limited), the Canadian reference product, Strattera (Eli Lilly Canada Inc.) and the UK reference product Strattera (Eli Lilly and Company Limited) under fasting condition. The results from the comparison of Teva-Atomoxetine and Strattera (Eli Lilly Canada Inc.) are presented in the table below.

Atomoxetine (1 x 60 mg) From measured data uncorrected for potency Geometric Mean Arithmetic Mean (CV %)				
Parameter	Teva-Atomoxetine	Strattera ¹	% Ratio of Geometric Means	90% Confidence interval
AUC_T (ng·h/mL)	3858.05 4668.48 (68)	3908.45 4770.33 (78)	98.71	92.68 – 105.14
AUC_I (ng·h/mL)	3937.24 4978.15 (85)	3988.13 5134.23 (98)	98.72	92.75 – 105.09
C_{max} (ng/mL)	680.46 702.43 (22)	704.06 728.38 (31)	96.65	88.16 – 105.96
T_{max} [§] (h)	1.20 (95)	1.23 (92)		
T_{1/2} [§] (h)	4.48 (70)	4.50 (75)		

¹ Strattera (Eli Lilly Canada Inc.) was purchased in Canada

[§] Expressed as the arithmetic mean (%CV) only

The efficacy and safety of atomoxetine hydrochloride for the treatment of ADHD in children, adolescents and adults (Table 9) who met the *Diagnostic and Statistical Manual 4th edition* (DSM-IV) criteria were established in 6 acute, randomized, double-blind, placebo-controlled studies. Examination of population subsets (gender, age) did not reveal any differential responsiveness on the basis of these subgroupings.

Four additional double-blind, placebo-controlled studies were also conducted with children and adolescents in the school setting, and in adults (Table 9). Efficacy and safety were assessed in patients with comorbid tic disorder (pediatrics), and comorbid anxiety disorders (pediatrics and adults). As well, evening efficacy was assessed with once daily dosing (pediatrics and adults).

Long-term efficacy of atomoxetine hydrochloride was established in an 18 month, randomized, double-blind, placebo-controlled relapse-prevention study. Too few patients completed to allow an adequate assessment of long-term safety from this study.

The long-term safety of atomoxetine hydrochloride has been demonstrated in double-blind and open-label clinical trials involving 5382 children or adolescent patients and 525 adult patients with ADHD. During the ADHD clinical trials, 1625 children and adolescent patients were treated for longer than 1 year and 2529 children and adolescent patients were treated for over 6 months.

In both pediatric and adult studies, atomoxetine hydrochloride showed efficacy in patients who had been previously treated with stimulants, as well as in stimulant-naïve patients. atomoxetine hydrochloride has a different mechanism of action from stimulant therapies. Data on evening efficacy suggest that atomoxetine hydrochloride is not associated with the ‘on/off’ phenomena characteristic of stimulants.

Study Demographics and Trial Design

Table 9: Summary of Patient Demographics for Controlled Clinical Trials in ADHD Patients

Study No.	Trial design	Atomoxetine Dose/Duration	No. of Subjects	Mean Age (range)	Gender
<u>Randomized, Double-Blind Studies in Children and Adolescents</u>					
HFBD	Acute, placebo-controlled, parallel groups.	BID dosing, max 2.0 mg/kg/day; 9 weeks.	n = 147	9.7 years (7 – 13)	M = 119 F = 28
HFBK	Acute, placebo-controlled, parallel groups.	BID dosing, max 2.0 mg/kg/day; 9 weeks.	n = 144	9.9 years (7 – 13)	M = 117 F = 27
LYAC	Acute, dose-ranging, placebo-controlled, parallel groups.	Fixed BID doses 0.5, 1.2, and 1.8 mg/kg/day; 8 weeks.	n = 297	11.2 years (8 – 18)	M = 212 F = 85

Study No.	Trial design	Atomoxetine Dose/Duration	No. of Subjects	Mean Age (range)	Gender
LYAT	Once-daily dosing, acute, placebo-controlled, parallel groups.	QD dosing, max 1.5 mg/kg/day; 6 weeks.	n = 171	10.3 years (6 - 16)	M = 120 F = 51
LYBG	Once-daily dosing, acute, placebo-controlled, parallel groups, and assessment of evening efficacy.	QD dosing, max 1.8 mg/kg/day; 8 weeks.	n = 197	9.5 years (6 - 13)	M = 139 F = 58
LYAW	Once- daily dosing, acute, placebo-controlled, parallel groups, and assessment of efficacy in the school setting.	QD dosing, max 1.8 mg/kg/day; 7 weeks.	n = 153	9.9 years (7 - 13)	M = 123 F = 30
LYAS ¹	Acute, placebo-controlled, parallel groups, patients with ADHD and comorbid tic disorder.	BID dosing, max 1.5 mg/kg/day; up to 18 weeks	n = 148	11.2 years (7 - 17)	M = 131 F = 17
LYBI	Parallel-group comparison of atomoxetine, extended-release methylphenidate and placebo.	BID dosing max 1.8 mg/kg/day; approx. 6 weeks.	n = 516	10.2 years (6 - 17)	M = 383 F = 133
LYAF	Long-term relapse prevention, placebo-controlled, parallel groups.	BID dosing, max 1.8 mg/kg/day; 52 weeks double-blind.	n = 416	10.4 years (6 - 16)	M = 373 F = 43
LYBP ²	Placebo-controlled, parallel group study in patients with ADHD and comorbid anxiety.	BID dosing max 1.8 mg/kg/day; approx. 12 weeks.	n = 176	12.0 years (8 - 18)	M = 114 F = 62
<u>Randomized, Double-Blind Studies in Adults</u>					
LYAA	Acute, placebo-controlled, parallel groups.	BID dosing, max 120 mg/day 10 weeks	n = 280	40.3 years (18 - 67)	M = 178 F = 102
LYAO	Acute, placebo-controlled, parallel groups.	BID dosing max 120 mg/day 10 weeks	n = 256	42.1 years (19 - 77)	M = 170 F = 86
LYDQ ³	Placebo-controlled, parallel group study in adult patients with ADHD and comorbid social anxiety.	BID dosing max 100 mg/day; approx. 16 weeks.	n = 442	38.0 years (18 - 65)	M = 237 F = 205

^{1, 2 and 3}: See *Action and Clinical Pharmacology, Special Populations, Patients with concomitant illness.*

Children and Adolescents: Improvement on signs and symptoms of ADHD were evaluated by a comparison of mean change from baseline to endpoint for atomoxetine hydrochloride - and placebo-treated patients. The primary efficacy measure in all studies was the *ADHD Rating Scale-IV-Parent Version* (ADHDRS), with the exception of Study LYAW which used the *ADHD Rating Scale-IV-Teacher Version*. Both scales have been well-validated and contain 18 items corresponding exactly to the DSM-IV symptom criteria, thus providing strong content validity. Both scales were administered and scored by an investigator following interviews with a parent or teacher.

Doses ranged from 0.5 to 2.0 mg/kg/day. In clinical trials, the mean weight of children 6 to

12 years of age was 35 kg, and their mean final dose was 45 mg/day. Atomoxetine hydrochloride was dosed twice daily, morning and late afternoon/early evening in Studies HFBD, HFBK, and LYAC, or once daily in the morning in Studies LYAT, LYAW, and LYBG.

Adults: The primary outcome measure in the adult studies was the validated Conners' Adult ADHD Rating scale (CAARS), which contains 18 items that correspond exactly to the DSM-IV symptom criteria. This instrument is investigator-administered and scored, with ratings based on patient self-report. Adult patients with current diagnoses of depression and anxiety were excluded, which ensured that improvements related to atomoxetine hydrochloride were attributable to improvements in ADHD. Doses ranged from 30 to 60 mg twice daily, which correspond, on a weight-adjusted basis for a 70 kg adult, to the dosing used in children.

Study Results

Children and Adolescents

Acute, Double-Blind, Placebo-Controlled Studies: The effectiveness of atomoxetine hydrochloride in the treatment of ADHD was established in 6 randomized, double-blind, placebo-controlled studies of pediatric patients, ages 6 to 18 (Studies HFBD, HFBK, LYAC, LYAT, LYAW, LYBG). Approximately one-third of the patients met DSM-IV criteria for inattentive subtype and two-thirds met criteria for both inattentive and hyperactive/impulsive subtypes. In all studies, atomoxetine hydrochloride demonstrated a statistically significantly ($p < 0.001$) greater mean reduction in core ADHD symptoms compared with placebo on the primary efficacy measure (Figure 1).




Figure 1. Mean change from baseline to endpoint on the ADHDRS-IV-Parent:Inv Total score in the acute, double-blind, placebo-controlled pediatric studies.

The effect size in the atomoxetine hydrochloride treatment group was similar across studies, providing evidence of a consistent treatment effect regardless of dosing regimen (once or twice daily) or which observer (teacher or parent) was providing information for the primary outcome measure. The efficacy of atomoxetine hydrochloride in children and adolescents was demonstrated in both Hyperactive-Impulsive and Inattentive symptom domains, in ADHD patients with or without comorbidity (e.g., Oppositional Defiant Disorder, depression), and in patients with or without prior stimulant use.

Improvements in the primary outcome measure were consistently accompanied by improvements in broader measures of quality of life and functioning, including behaviour at school, self-esteem and family functioning.

The results of the fixed dose study (LYAC) provided evidence of dose response (onset of effect at 0.5 mg/kg/day, with maximal effect at 1.2 mg/kg/day). For the overall group, no further benefit was demonstrated at 1.8 mg/kg/day.

Study LYAT demonstrated the effectiveness of atomoxetine hydrochloride when administered once daily in the morning (Figure 2). Clinical improvement was typically seen within 1 week of starting treatment, and at 2 to 4 weeks, mean reductions in symptom severity scores were approaching maximal reductions. Study LYBG further demonstrated that the efficacy of atomoxetine hydrochloride persisted from morning until late evening following a once-daily morning dose.




Figure 2: Once-Daily Administration of atomoxetine hydrochloride.

Long-Term Placebo-Controlled Study: The effectiveness of atomoxetine in maintaining symptom response was investigated in Study LYAF, an 18-month study of children and

adolescents aged 6 to 15 who met DSM-IV criteria for ADHD. A total of 604 patients were enrolled in this study. Patients who responded to atomoxetine during approximately 3-months open-label treatment (n = 416) were randomized to double-blind, placebo-controlled continuation at their current doses of atomoxetine (n = 292) or on placebo (n = 124) for approximately 9 months to observe for relapse. Patients who were still on atomoxetine (n = 163) at the end of 12 months of treatment were further randomized to an additional 6 months double-blind, placebo-controlled continuation at their current doses of atomoxetine (n = 81) or on placebo (n = 82). The results from this study suggest that atomoxetine may be beneficial in the long-term treatment of ADHD. Too few patients completed to allow an adequate assessment of long-term safety from this study.

Children and adolescents enrolled in acute double-blind, placebo-controlled clinical trials, as well as those enrolled in open-label trials, were followed prospectively to assess safety, tolerability, and effects on growth. The long-term safety of atomoxetine hydrochloride has been demonstrated in double-blind and open-label clinical trials involving 5382 children or adolescent patients. During the ADHD clinical trials, 1625 children and adolescent patients were treated for longer than 1 year and 2529 children and adolescent patients were treated for over 6 months. In these trials, atomoxetine was safe and well tolerated, and there were minimal, if any, long-term effects on weight and height compared to normal growth curves.

Active Comparator Study: Study LYBI was a randomized, double-blind, parallel group, 6 week pediatric study to test the non-inferiority of atomoxetine to a standard extended-release stimulant comparator. The comparator was shown to be associated with superior response rates compared to atomoxetine ($p=0.016$). However, this study excluded patients who were stimulant nonresponders, so it is possible that the results were biased favouring the stimulant comparator. For patients not previously treated with a stimulant, there was no significant difference in response rates between stimulant comparator and atomoxetine ($p=0.423$). Both atomoxetine and the comparator were statistically superior to placebo. Both atomoxetine and comparator were safe and well-tolerated. Neither medication produced statistically significant QT prolongation compared to placebo.

Studies in Adult ADHD Patients

The results of 2 large placebo-controlled studies in adult patients with ADHD (N=536) have demonstrated the efficacy and safety of amotoxetine in this population. Amotoxetine was statistically significantly superior to placebo at reducing ADHD symptoms as measured on the CAARS Total ADHD symptom score (Figure 3, LYAA $p = 0.004$, LYAO $p = 0.002$). Efficacy was observed for overall ADHD symptoms as well as in the individual domains of Predominately Inattentive and Predominately Hyperactive-Impulsive symptoms.




Figure 3: CAARS Total Scores – Mean change from baseline to endpoint in randomized, double-blind, placebo-controlled adult studies

DETAILED PHARMACOLOGY

Efficacy Pharmacology

Atomoxetine hydrochloride was designed to be a potent and selective inhibitor of the norepinephrine transporter. In vitro, atomoxetine was found to have highly selective potency for the norepinephrine transporter over the dopamine and serotonin transporters, and was highly selective for inhibiting the uptake of radiolabelled norepinephrine into hypothalamic synaptosomes. Of the two primary phase I metabolites, 4-hydroxyatomoxetine was found to have similar in vitro potency for the norepinephrine transporter and inhibition of norepinephrine uptake compared to atomoxetine, while *N*-desmethyatomoxetine had lower affinity than atomoxetine for all three monoamine transporters. Further, atomoxetine and these metabolites were found to have relatively low affinities for neurotransmitter receptors and ion channels. In vivo evaluation of atomoxetine also demonstrated high potency and selectivity of atomoxetine for the norepinephrine transporter. In the prefrontal cortex, a site believed to be important for attention in humans, dose-dependent increases in the extracellular concentrations of norepinephrine and dopamine occurred when atomoxetine was given intraperitoneally or when atomoxetine was delivered directly through a microdialysis probe, suggesting that the increases in prefrontal cortex norepinephrine and dopamine were produced by local inhibition of the norepinephrine transporter. Atomoxetine did not produce a significant increase in the extracellular concentrations of dopamine in the nucleus accumbens, an area believed to mediate the rewarding effects of dopamine transporter inhibitors, and the striatum, an area believed to mediate the effects of dopamine transporter inhibitors on locomotor activity.

Safety Pharmacology

Atomoxetine was evaluated in animals or in vitro for potential effects on the central nervous system and behaviour, including abuse potential, and for effects on cardiovascular, respiratory, renal, gastrointestinal, and immune function.

In contrast to the high affinity of atomoxetine for the norepinephrine transporter site, atomoxetine, 4-hydroxyatomoxetine, and *N*-desmethylatomoxetine, have relatively low binding affinities for adrenergic, muscarinic, histaminergic, GABAergic, serotonergic, and dopaminergic receptors, as well as many other receptor sites and ion channels. Atomoxetine exhibited no agonist effects on isolated smooth and cardiac tissues and did not inhibit acetylcholine- or isoproterenol-induced contractions of these respective tissues. Therefore, atomoxetine would not be expected to interfere with the physiologic function mediated by these receptor systems or produce toxicity associated with muscarinic or adrenergic receptors. Importantly, clinical signs observed in rodents and dogs were consistent only with effects on the central nervous system (for example, reduced motor activity, leg weakness, jerky gait, tremors, and convulsions).

Atomoxetine is not associated with abuse potential, based upon its receptor binding profile and nonclinical behavioural pharmacology, as well as animal drug discrimination studies conducted with atomoxetine.

Atomoxetine produced no important changes in respiratory or renal function, gastrointestinal motility, or immune function in animals.

Cardiac Function

Atomoxetine was evaluated for effects on cardiac function in vitro and in dogs. In vitro blockade of the cloned human cardiac I_{Kr} (HERG) channel by atomoxetine occurred with an IC_{50} of 0.869 μ M. Studies with canine Purkinje tissue and in dogs given atomoxetine orally did not produce effects predictive or consistent with a compound-related QT interval prolongation. Concomitant inhibition of sodium and calcium inward currents in vitro helps explain the lack of in vivo findings. These observations, together with the clinical experience with atomoxetine, indicate that the in vitro HERG blockade observation is not likely predictive of an increased cardiovascular risk for atomoxetine.

Cardiovascular Safety Study:

The safety and tolerability of gradually increasing multiple-dose regimes of atomoxetine was studied during a 20 day period in a clinical pharmacology study involving 16 healthy adults (10 EM subjects and 6 PM subjects). Placebo or atomoxetine 60 to 150 mg/day, was given as twice daily doses of 30 mg, 45 mg, 60 mg, and 75 mg. None of the mean or individual QT_c(F) intervals (QT interval corrected by the Fridericia method) exceeded the upper limits of normal for each gender.

The EM group had no statistically significant changes in mean QT_c(F) interval following

any dose of atomoxetine compared to the placebo treatment. No statistically significant changes in QTc(F) were noted 1 hour post dose (during peak plasma concentrations) in the PM group. The PM group had a statistically significant increase in the mean QTc(F) interval measured at time 0 (during trough plasma concentrations) on the last day of the 60- and 75-mg atomoxetine twice-daily dosing regimens compared to the placebo. The greatest mean prolongation was about 17 msec at the 60-mg BID dose level, with the mean interval length of 417.2 msec. At the 75-mg BID atomoxetine dose level, the greatest mean prolongation was 15 msec, and the mean interval length was 414.9 msec. The 60 mg BID and 75 mg BID doses correspond to 1.4-2.24 mg/kg/day and 1.75-2.8 mg/kg/day, respectively.

Baseline ECGs obtained during screening of pediatric patients for atomoxetine clinical trials were reviewed for cases of QTc prolongation. Using Bazett's method for correcting QT (the usual method for automated QTc intervals), there were 131/3902 cases (3.4%) with QTc(B)>450 msec and 5/3902 cases (0.1%) with QTc(B)>500 msec. Using a correction method based on data from baseline ECGs (the method preferred by most regulatory agencies), there were 32/3902 cases (0.8%) with QTc(D)>450 msec and 5/3902 cases (0.1%) with QTc(D)>500 msec. In a meta-analysis of ECG data from patients who received atomoxetine in pediatric clinical trials, no relationship was observed between changes from baseline to final QTc(D) and prescribed atomoxetine dose (n=3127 patients), or between changes from baseline to QTc(D) at time of expected peak exposure and prescribed atomoxetine dose (n=2084 all patients, n=151 PM patients). For clinical trial patients where plasma concentrations were available, no relationship was observed between changes from baseline QTc(D) and atomoxetine plasma concentration during peak atomoxetine exposure (n=728 all patients, n=83 PM patients).

Overall, the data do not suggest a meaningful relationship between atomoxetine plasma concentrations and the length of the QT interval corrected for heart rate in the recommended dosage range. However, since there is no requirement for a priori screening of ADHD patients for CYP2D6 metabolizer status before initiating treatment with atomoxetine, it is important that the lowest effective dose be used, so as to minimize potential cardiac side effects.

Pharmacokinetics

Atomoxetine was well absorbed following oral administration to adult mice, rats, and dogs.

The primary oxidative (phase I) metabolite of atomoxetine was identified as 4-hydroxyatomoxetine, and its major ultimate metabolite is 4-hydroxyatomoxetine-*O*-glucuronide, in all species evaluated. The major routes of metabolism are aromatic ring-hydroxylation, benzylic/aliphatic hydroxylation, and *N*-demethylation. Subsequent *O*-glucuronidation and *O*-sulfation of the hydroxylated metabolites are the only phase II metabolic pathways to participate in the conjugation of the oxidized atomoxetine metabolites. Plasma concentrations of atomoxetine and its primary phase I metabolites,

N-desmethylatomoxetine and 4-hydroxyatomoxetine, increased with dose in each species. There was generally minimal accumulation with repeated dosing, which appeared to be due to the relatively rapid clearance of atomoxetine in each of the species evaluated. The half-life of atomoxetine following a single intravenous dose was 1.2 hr, 1.4 hr and 3.4 hr in the mouse, rat and dog, respectively. The most profound difference observed among species was the low bioavailability in rodents (approximately 4%) compared to the high bioavailability in dog (approximately 74%). These differences appear to be almost purely mediated by the efficient first-pass hepatic clearance of atomoxetine in rodents, resulting in poor systemic exposure to atomoxetine but high exposure to its metabolites. In the dog, atomoxetine does not undergo extensive first-pass metabolism.

Increases in hepatic cytochrome P450 activity were observed in mice and rats after large oral doses of atomoxetine. Slight increases in total cytochrome P450 content were observed at the highest dose tested (16 mg/kg/day) in male dogs. However, no evidence of such changes in cytochrome P450 activity have been observed in clinical studies.

Binding of atomoxetine and *N*-desmethylatomoxetine to plasma protein was high in each species evaluated including humans (82% to 99%), while the binding of 4-hydroxyatomoxetine to plasma protein was substantially lower (55% to 67%).

In the rat, atomoxetine and its metabolites were rapidly and widely distributed in tissues, with the highest levels being observed in the gastrointestinal tract and the liver. Milk excretion is not a major route of elimination for atomoxetine and/or its metabolites. Atomoxetine and/or its metabolites crossed the placenta resulting in relatively low fetal exposure compared to exposure in maternal tissues.

The major route of excretion of atomoxetine-derived radioequivalents was via the urine in each of the species evaluated. A low amount of fecal excretion appears to be due to biliary elimination. Very little atomoxetine was excreted intact, indicating that direct renal or biliary elimination of atomoxetine are minor routes of systemic clearance.

TOXICOLOGY

Acute Toxicity

The median lethal oral dose of atomoxetine hydrochloride in animals was estimated to be 25 mg/kg for cats, >37.5 mg/kg for dogs, and ≥190 mg/kg in rats and mice. Premonitory signs of toxicity following single oral doses of atomoxetine in animals included mydriasis and reduced pupillary light reflex, mucoid stools, salivation, vomiting, ataxia, tremors, myoclonic jerking, and convulsions.

Long-Term Toxicity

Toxicity studies of up to 1 year were conducted in adult rats and dogs to evaluate the potential chronic toxicity of atomoxetine. There was no major target organ toxicity

observed in dogs given oral doses up to 16 mg/kg/day or in rats given atomoxetine in the diet at time-weighted average doses up to 47 mg/kg/day. These doses are approximately 6-7 times the maximum recommended daily oral dose in children, and approximately 4-5 times the maximum recommended daily oral dose in adults, on a mg/m² basis. Mild hepatic effects, characterized by mottling and pallor of the liver, increased relative liver weights, hepatocellular vacuolation, and slightly increased serum ALT values, occurred in male rats given time-weighted average doses \geq 14 mg/kg/day. No hepatic effects were observed in dogs. Clinical signs of mydriasis, reduced pupillary light reflex, emesis, and tremors were observed in dogs, and these effects were minimal in adult dogs given \leq 8 mg/kg/day. Similar clinical signs were observed in young dogs (8 weeks of age) given atomoxetine orally for 1 month.

Toxicity studies were conducted to evaluate potential effects of atomoxetine on growth, neurobehavioural, and sexual development as well as reproductive performance in rats dosed from the early postnatal period to early adulthood. No important effects on bone growth, brain morphology, neurobehavioral or reproductive performance were observed in rats administered atomoxetine hydrochloride from 10 days of age through adulthood, at oral doses up to 50 mg/kg, approximately 7 times the maximum recommended daily oral dose in children on a mg/m² basis. There were no effects on the anatomy or morphology of either male or female reproductive organs. There was a minor, dose-related increase in time-to-onset of vaginal patency and preputial separation (1 to 3 days compared with controls), but all animals reached sexual maturity and no malformations of the vagina or prepuce/glans penis were observed. In male rats, spermatogenesis was normal. There were slight decreases in the number of sperm held in the epididymis but no effects on fertility. For both male and female rats, there were no effects on mating or fertility indices, no effects on embryo implantation or embryo viability, and no effects on learning and memory tests. The significance of these findings to humans is not known.

Carcinogenicity

Atomoxetine hydrochloride was not carcinogenic in rats and mice when given in the diet for 2 years at time-weighted average doses up to 47 and 458 mg/kg/day, respectively. The highest dose used in rats is approximately 8 and 5 times the maximum human dose in children and adults, respectively, on a mg/m² basis. Plasma levels (AUC) of atomoxetine at this dose in rats are estimated to be 1.8 times (extensive metabolizers) or 0.2 times (poor metabolizers) those in humans receiving the maximum human dose. The highest dose used in mice is approximately 39 and 26 times the maximum human dose in children and adults, respectively, on a mg/m² basis.

Mutagenicity

Atomoxetine hydrochloride was negative in a battery of genotoxicity studies that included a reverse point mutation assay (Ames Test), an in vitro mouse lymphoma assay, a chromosomal aberration test in Chinese hamster ovary cells, an unscheduled DNA synthesis test in rat hepatocytes and an in vivo micronucleus test in mice. However, there was a slight increase in the percentage of Chinese hamster ovary cells with

diplochromosomes, suggesting endoreduplication (numerical aberration).

The metabolite N-desmethyatomoxetine hydrochloride was negative in the Ames Test, mouse lymphoma assay, and unscheduled DNA synthesis test.

Teratogenicity

No evidence of drug-associated teratogenicity or retarded fetal development was produced in rabbits or rats administered atomoxetine hydrochloride throughout organogenesis at oral doses up to 100 mg/kg/day and 150 mg/kg/day (at least 20 times the maximum recommended daily oral dose in children and 13 times the maximum recommended daily oral dose in adults, on a mg/m² basis). In a rat fertility study, decreased pup weight and survival was observed, predominantly during the first week postpartum following maternal dietary atomoxetine time-weighted average doses of 23 mg/kg/day or higher. No adverse effects were observed in surviving pups.

Atomoxetine hydrochloride did not impair fertility when administered to rats from 10 days of age through adulthood at oral doses up to 50 mg/kg/day (up to 7 times the maximum recommended daily oral dose in children and 4 times the maximum recommended daily oral dose in adults, on a mg/m² basis). In addition, no evidence of impaired fertility was observed in either of two fertility studies in adult rats provided atomoxetine HCl in the diet at time-weighted average doses up to 57 mg/kg/day (up to 8 times the maximum recommended daily oral dose in children and 5 times the maximum recommended daily oral dose in adults, on a mg/m² basis).

Parturition in rats was not affected by atomoxetine.

REFERENCES

1. Adler L, Spencer T, Willims D, *et al.* Long-term, open label study of the safety and efficacy of atomoxetine in adults with attention-deficit/hyperactivity disorder: An interim Analysis. *J Clin Psychiatry* 2005; 66:294-299.
2. Adler L, Sutton V, Moore R, *et al.* Quality of life assessment in adult patients with attention-deficit/hyperactivity disorder treated with atomoxetine. *J Clin Psychopharmacol* 2006; 26:648-652.
3. Adler L, Liebowitz M, Kronenberger W, *et al.* Atomoxetine treatment in adults with Attention-Deficit/Hyperactivity Disorder and comorbid Social Anxiety Disorder. *Depression and Anxiety* 2009; 0:1–11.
4. Adler L, Spencer T, Brown T, *et al.* Once-daily atomoxetine for adult Attention-Deficit/Hyperactivity Disorder: A 6-month, double-blind trial. *J Clin Psychopharmacol* 2009; 29:44-50.
5. Allen AJ, Kurlan R, Gilbert D, *et al.* Atomoxetine treatment in children and adolescents with ADHD and comorbid tic disorders. *Neurology* 2005;65:1941–1949.
6. American Academy of Pediatrics (2001), Clinical Practice Guideline: Treatment of the School-Aged Child with Attention-Deficit/Hyperactivity Disorder. *Pediatrics* 108:1033-1044.
7. American Psychiatric Association (2000), Diagnostic and Statistical Manual of Mental Disorders: DSM-IV-TR. *American Psychiatric Association, Washington, DC.*
8. Buitelaar J, Michelson D, Danckaerts M, *et al.* A randomized, double-blind study of continuation treatment for Attention-Deficit/Hyperactivity Disorder after 1 year. *Biol Psychiatry* 2007; 61:694-699.
9. Canadian ADHD Resource Alliance (www.CADDRA.ca): Canadian ADHD Practice Guidelines: Medications for Attention Deficit/Hyperactivity Disorder, 2006 (1st Ed); Chapter 10 (App 17), 1-14.
10. Chalon S, Desager J, Desante K, *et al.* Effect of hepatic impairment on the pharmacokinetics of atomoxetine and its metabolites. *Clin Pharmacol Therap* 2003; 73:178-191.
11. Dell'Agnello G, Maschietto D, Bravaccio C, *et al.* Atomoxetine hydrochloride in the treatment of children and adolescents with Attention-Deficit/Hyperactivity Disorder and Comorbid Oppositional Defiant disorder: A placebo-controlled Italian study. *European Neuropsychopharmacol* (2009);19:822–834.

12. Faries D, Yalcin I, Harder D, *et al.* Validation of the ADHD Rating Scale as a clinician administered and scored instrument. *J Attn Disorders* 2001;5(2):107-115.
13. Geller D, Donnelly C, Lopez F, *et al.* Atomoxetine treatment for pediatric patients with attention-deficit/hyperactivity disorder with comorbid anxiety disorder. *J Amer Acad Child & Adolesc Psych* 2007; 46:1119-1127.
14. Hesdorffer DC, Ludvigsson P, Olafsson E, *et al.* ADHD as a risk factor for incident unprovoked seizures and epilepsy in children. *Arch Gen Psychiatry* 2004;61:731-736.
15. Heil S, Holmes H, Bickel W, *et al.* Comparison of the subjective, physiological, and psychomotor effects of atomoxetine and methylphenidate in light drug users. *Drug and Alcohol Dependence* 2002; 67:149-156.
16. Kelsey D, Sumner C, Casat C, Allen AJ, *et al.* Once-daily atomoxetine treatment for children with Attention-Deficit/Hyperactivity Disorder, including an assessment of evening and morning behavior: a double-blind, placebo-controlled trial. *Pediatrics* 2004; 114(1)e1-e8.
17. Kratochvil C, Heiligenstein J, Dittmann R, *et al.* Atomoxetine and methylphenidate treatment in children with ADHD: A prospective, randomized, open-label trial. *J Amer Acad Child Adolesc Psych* 2002;41:776-784.
18. Kratochvil C, Newcorn J, Arnold LE, *et al.* Atomoxetine alone or combined with fluoxetine for treating ADHD with comorbid depressive or anxiety symptoms. *J Am Acad Child Adolesc Psychiatry* 2005;44(9):915-924.
19. Kratochvil C, Wilens T, Greenhill L, *et al.* Effects of long-term atomoxetine treatment for young children with attention-deficit/hyperactivity disorder. *J Amer Acad Child & Adolesc Psych* 2006; 45:919-927.
20. Lowe S, Lim M, Smith B, *et al.* Haemodynamic effects of IV salbutamol and oral atomoxetine, alone and in combination. *J Clin Pharmacol* 2001;41(9):1025.
21. Maziade M, Rouleau N, Lee B, *et al.* Atomoxetine and neuropsychological function in children with Attention-Deficit/Hyperactivity Disorder: Results of a pilot study. *J Child Adolesc Psychopharmacol* 2009;19(6):709-718.
22. Michelson D, Faries D, Wernicke J. Atomoxetine in the Treatment of Children and Adolescents with ADHD: A Randomized, Placebo-Controlled, Dose-Response Study. *Pediatrics* 2001;108(5):1-9.
23. Michelson D, Allen A, Busner J, *et al.* Once-daily atomoxetine treatment for children and adolescents with attention deficit hyperactivity disorder: a randomized, placebo-controlled study. *Amer J Psychiatry* 2002;159:1896-1901.

24. Michelson D, Adler L, Spencer T, *et al.* Atomoxetine in adults with ADHD: Two randomized, placebo-controlled studies. *Biol Psychiatry* 2003;53:112–120.
25. Michelson D, Buitelaar J, Danckaerts M, *et al.* Relapse prevention in pediatric patients with ADHD treated with atomoxetine: A randomized, double-blind, placebo-controlled study. *J. Am. Acad. Child Adolesc. Psychiatry* 2004;43:896–904.
26. Montoya A, Hervas A, Cardo E, *et al.* Evaluation of atomoxetine for first-line treatment of newly diagnosed, treatment-naïve children and adolescents with attention deficit/hyperactivity disorder. *Curr Med Research & Opinion* 2009;25(11):2745–2754.
27. Newcorn J, Spencer T, Biederman J, *et al.* Atomoxetine treatment in children and adolescents with Attention-Deficit/Hyperactivity Disorder and comorbid Oppositional Defiant Disorder. *J Am Acad Child Adolesc Psychiatry* 2005;44(3):240–248.
28. Perwien A, Kratochvil C, Faries DE, *et al.* Atomoxetine treatment in children and adolescents with attention-deficit hyperactivity disorder: What are the long-term health-related quality-of-life outcomes? *J Child & Adolesc Psychopharmacol* 2006; 16:713-724.
29. Quintana H, Cherlin E, *et al.* Transitioning from methylphenidate or amphetamine to atomoxetine in children and adolescents with attention-deficit/hyperactivity disorder – a preliminary tolerability and efficacy study. *Clinical Therapeutics*, June 2007.
30. Sangal R, Owens J, Allen AJ, *et al.* Effects of atomoxetine and methylphenidate on sleep in children with ADHD. *Sleep* 2006; 29:1573-1585.
31. Sauer JM, Ring B, Witcher J. Clinical pharmacokinetics of atomoxetine. *Clin Pharmacokinet* 2005;44(6):571-590.
32. Simpson D, Plosker G. Atomoxetine: A Review of its Use in Adults with Attention Deficit Hyperactivity Disorder. *Drugs* 2004;64:205-222.
33. Smith B, Soon D, Teng C, *et. al.* Hemodynamic effects of atomoxetine are less pronounced than methylphenidate, and not additive in combination. *J Clin Pharmacol* 2001;41:1014-1033.
34. Spiller HA, Lintner CP, Winter ML: Atomoxetine ingestions in children: A report from Poison Centers. *Ann Pharmacother* 2005;39:1045-1048.
35. Svanborg P, Thernlund G, Gustafsson P, *et al.* Efficacy and safety of atomoxetine as add-on to psychoeducation in the treatment of attention deficit/hyperactivity disorder: A randomized, double-blind, placebo-controlled study in Swedish children and adolescents. *Eur Child Adolesc Psychiatry* 2009;18:240–249.

36. Svanborg P, Thernlund G, Gustafsson P, *et al.* Atomoxetine improves patient and family coping in attention deficit/hyperactivity disorder: A randomized, double-blind, placebo-controlled study in Swedish children and adolescents. *Eur Child Adolesc Psychiatry* 2009; *Published On-line May 23, 2009.*
37. Weiss M, Tannock R, Kratochvil C, *et al.* A randomized, placebo-controlled study of once-daily atomoxetine in the school setting in children with ADHD. *J Am Acad Child Adolesc Psychiatry* 2005;44(7):647–655.
38. Wernicke J, Adler L, Spencer T, *et al.* Changes in symptoms and adverse events after discontinuation of atomoxetine in children and adults with Attention Deficit/Hyperactivity Disorder: A Prospective, Placebo-Controlled Assessment. *J Clin Psychopharmacol* 2004; 24:30-35.
39. Wernicke J, Faries D, Girod D *et al.* Cardiovascular effects of atomoxetine in children, adolescents, and adults. *Drug Safety* 2003;26:729-40.
40. Wernicke JF, Holdridge KC, Jin L, *et al.* Seizure Risk in Patients with ADHD Treated with Atomoxetine. *Dev Med Child Neur* 2007; 49: 498-502
41. Wilens T. Drug therapy for adults with Attention-Deficit Hyperactivity Disorder. *Drugs* 2003;63:2395-2411.
42. Wilens T, Newcorn J, Kratochvil C, *et al.* Long-term atomoxetine treatment in adolescents with attention-deficit/hyperactivity disorder. *J Pediatrics* 2006; 149:112-119.
43. Pr[®]Strattera[®] 10, 18, 25, 40, 60, 80 and 100 mg Capsules Product Monograph, Eli Lilly Canada Inc., Canada. Date of Revision: Dec. 4, 2007
44. A single dose, comparative bioavailability study of three formulations of Atomoxetine Hydrochloride 10 mg Capsules under fasting conditions. Data is on file at Teva Canada Limited.
45. A single dose, comparative bioavailability study of three formulations of Atomoxetine Hydrochloride 60 mg Capsules under fasting conditions. Data is on file at Teva Canada Limited.

PART III: CONSUMER INFORMATION

Pr **TEVA-ATOMOXETINE**

(atomoxetine hydrochloride capsules)

This leaflet is part III of a three-part "Product Monograph" published when TEVA-ATOMOXETINE was approved for sale in Canada and is designed specifically for adults and parents of children/adolescents who will be prescribed this medication. This leaflet is a summary and will not tell you everything about TEVA-ATOMOXETINE. Contact your doctor or pharmacist if you have any questions about this drug.

The following have been reported with use of ATOMOXETINE HCl and also with stimulant medications:

1. Suicidal thoughts and actions in children and teenagers

Some children and teenagers may have a higher chance of having suicidal thoughts or actions. Tell your child or teenager's doctor if your child or teenager (or there is a family history of):

- has bipolar illness (manic-depressive illness)
- had suicide thoughts or actions before starting Teva-Atomoxetine.

The chance for suicidal thoughts and actions are higher:

- early during Teva-Atomoxetine treatment
- during dose adjustments.

Prevent suicidal thoughts and action in your child or teenager by:

- paying close attention to your child or teenager's moods, behaviors, thoughts, and feelings during Teva-Atomoxetine treatment
- keeping all follow-up visits with your child or teenager's doctor as scheduled.

Watch for the following signs in your child or teenager during Teva-Atomoxetine treatment:

- anxiety
- agitation
- panic attacks
- trouble sleeping
- irritability
- hostility

- aggressiveness
- impulsivity
- restlessness
- mania
- depression
- suicide thoughts.

Call your child or teenager's doctor right away if they have any of the above signs, especially if they are new, sudden, or severe.

Your child or teenager may need to be closely watched for suicidal thoughts and actions or need a change in medicine.

2. Severe liver damage

Call your doctor right away if you or your child have the following signs of liver problems:

- itching
- right upper belly pain
- dark urine
- yellow skin or eyes
- unexplained flu-like symptoms

3. Heart-related problems:

- **sudden death in patients who have heart problems or heart defects**
- **stroke and heart attack in adults**
- **increased blood pressure and heart rate**

Tell your doctor if you or your child have any heart problems, heart defects, high blood pressure, or a family history of these problems. Your doctor may wish to check you or your child carefully for heart problems before starting Teva-Atomoxetine.

Your doctor may wish to check you or your child's blood pressure and heart rate regularly during treatment with Teva-Atomoxetine.

Call your doctor right away if you or your child has any signs of heart problems such as chest pain, shortness of breath, or fainting while taking Teva-Atomoxetine.

4. New mental (psychiatric) problems in children and teenagers:

- new psychotic symptoms (such as hearing voices, believing things that are not true, are suspicious) or new manic symptoms.

Call your child or teenager's doctor right away about any new mental symptoms. Teva-Atomoxetine treatment may be stopped.

ABOUT THIS MEDICATION

What TEVA-ATOMOXETINE is used for:

TEVA-ATOMOXETINE is a medicine that is taken either once or twice a day for the treatment of Attention-Deficit/Hyperactivity Disorder (ADHD) in children 6 years of age and over, adolescents, and adults. TEVA-ATOMOXETINE should not be used in children under 6 years of age.

TEVA-ATOMOXETINE is a part of your overall treatment program for ADHD that may include other measures (psychological, educational, and social). Your doctor may also recommend other therapy.

TEVA-ATOMOXETINE works differently from other medicines used for the treatment of ADHD. Atomoxetine hydrochloride is not a stimulant and studies have shown that it has no potential for drug abuse or dependence.

What is ADHD:

ADHD has 3 main types of symptoms: inattention, hyperactivity, and impulsiveness. Symptoms of inattention include not paying attention, making careless mistakes, not listening, not finishing tasks, not following directions, and being easily distracted. Symptoms of hyperactivity and impulsiveness include fidgeting, talking excessively, running around at inappropriate times, and interrupting others. Some patients have more symptoms of hyperactivity and impulsiveness while others have more symptoms of inattentiveness. Some patients have all 3 types of symptoms.

Symptoms of ADHD in adults may include a lack of organization, problems starting tasks, impulsive actions, daydreaming, daytime drowsiness, slow processing of information, difficulty learning new things, irritability, lack of motivation, sensitivity to criticism, forgetfulness, low self-esteem, and excessive effort to maintain some organization. The symptoms shown by adults who primarily have attention problems but not hyperactivity have been commonly described as Attention-Deficit Disorder (ADD).

Many people have these symptoms from time to time. However, people with ADHD have these symptoms most of the time. Symptoms must be present for at least 6 months to be certain of the diagnosis. In addition, the symptoms cause problems in more than one area of life (home, school, work, or social situations).

When it should not be used:

Do not take TEVA-ATOMOXETINE if you:

- are taking, or have recently taken, an antidepressant medicine known as a monoamine oxidase inhibitor (MAOI). Some names of MAOI medicines are phenelzine sulfate (Nardil®) and tranylcypromine sulfate (Parnate®).
- have narrow angle glaucoma, an eye disease.
- are allergic to atomoxetine or any other ingredient of TEVA-ATOMOXETINE.
- have symptomatic cardiovascular disease.
- have moderate to severe high blood pressure.
- have advanced arteriosclerosis (hardened arteries).
- have uncontrolled hyperthyroidism (an overactive thyroid gland).

What the medicinal ingredient is:

Atomoxetine Hydrochloride

What the nonmedicinal ingredients are:

In addition to Atomoxetine HCl, each capsule contains the inactive ingredients pregelatinized starch, sodium laurylsulfate and hard gelatin capsule contains gelatin, titanium dioxide, D&C Yellow 10 (18 mg capsules), FD&C Blue 1/Brilliant Blue FCF (25 mg and 40 mg capsules) and D&C yellow 10, FD&C Blue 1 and Brilliant Blue FCF (60 mg capsules).

What dosage forms it comes in:

Each capsule of TEVA-ATOMOXETINE contains atomoxetine hydrochloride equivalent to 10 mg, 18 mg, 25 mg, 40 mg, or 60 mg of atomoxetine.

WARNINGS AND PRECAUTIONS

BEFORE you use TEVA-ATOMOXETINE, talk to your doctor or pharmacist if you:

- have structural heart abnormalities,
- have mental problems, including psychosis, mania, bipolar illness, or depression;
- have had seizures (convulsions, epilepsy) or abnormal EEGs (electroencephalograms);
- have or had any disorder of the blood vessels in the brain (e.g. aneurysm, stroke, vasculitis);
- have a family history of sudden death or death related to heart problems;
- do strenuous exercise;
- take other drugs for ADHD;
- have or had liver problems. You may need a lower dose;
- have mild high blood pressure. Atomoxetine hydrochloride can increase blood pressure;
- have problems with your heart or an irregular heartbeat. Atomoxetine hydrochloride can increase heart rate (pulse);
- have low blood pressure. Atomoxetine hydrochloride can cause dizziness or fainting in people with low blood pressure
- are nursing, pregnant, or thinking of becoming pregnant.

Other Important Safety Information:

Severe liver damage: Call your doctor right away if you or your child have the following signs of liver problems:

- dark urine
- yellow skin or eyes
- right upper belly pain
- itching
- unexplained flu-like symptoms

Sudden death has been reported in association with stimulant drugs for ADHD treatment in children with structural heart abnormalities. Although atomoxetine hydrochloride is not a stimulant drug, it generally should not be used in children, adolescents or adults with known structural heart abnormalities.

New or Worsened Emotional or Behavioural Problems: Particularly in the first few weeks or when doses are adjusted, a small number of patients taking drugs of this type may feel worse instead of better; for example, they may experience unusual feelings of agitation, hostility or anxiety, or have impulsive or disturbing thoughts such as thoughts of self-harm or harm to others. Should this happen to you, or to those in your care if you are a caregiver or guardian, consult your doctor immediately. Close observation by a doctor is necessary in this

situation.

During treatment with these types of medication it is important that you and your doctor have good ongoing communication about how you are feeling.

Do not drive a car or operate hazardous machinery until you know how TEVA-ATOMOXETINE affects you.

This medicine was prescribed for your use only. Do not let anyone else take your TEVA-ATOMOXETINE.

INTERACTIONS WITH TEVA-ATOMOXETINE

Tell your doctor about all the medicines you take or plan to take, including prescription and non-prescription medicines, dietary supplements, and herbal remedies. Your doctor will decide if you can take TEVA-ATOMOXETINE with your other medicines.

Certain medicines may change the way your body reacts to TEVA-ATOMOXETINE. These include medicines used to treat depression, like paroxetine hydrochloride (Paxil®) and fluoxetine hydrochloride (Prozac®), and certain other medicines (like quinidine).

Your doctor may need to change your dose of TEVA-ATOMOXETINE if you are taking it with these medicines.

You should not take TEVA-ATOMOXETINE if you are taking desipramine.

TEVA-ATOMOXETINE may change the way your body reacts to oral, intravenous, or nebulized salbutamol (or drugs with similar actions), but the effectiveness of these drugs will not be changed. Talk with your doctor before taking TEVA-ATOMOXETINE if you are taking salbutamol.

TEVA-ATOMOXETINE should be used with caution if you are being treated with drugs for high blood pressure. Talk with your doctor before taking TEVA-ATOMOXETINE if you are taking any drug for high blood pressure.

PROPER USE OF TEVA-ATOMOXETINE

Usual dose:

Take TEVA-ATOMOXETINE exactly as

directed by your doctor. It is very important that you do not take a larger dose of TEVA-ATOMOXETINE than prescribed by your doctor.

Your doctor may tell you to take TEVA-ATOMOXETINE once a day or twice a day (morning and late afternoon/early evening). To help you remember to take TEVA-ATOMOXETINE, you may want to take it at the same time every day.

Improvement of your ADHD symptoms is generally observed within 1 to 4 weeks of starting TEVA-ATOMOXETINE.

TEVA-ATOMOXETINE may be taken with or without food.

You should not open TEVA-ATOMOXETINE capsules, but if they are accidentally opened or broken, avoid contact with the powder and wash away any loose powder as soon as possible with water. If any of the powder gets in your eyes you should rinse them with water immediately and contact your doctor.

Overdose:

In the event of overdosage, contact your doctor, hospital emergency department or regional Poison Control Centre.

Missed Dose:

If you miss a dose, take it as soon as possible, but do not take more than your total daily dose in any 24-hour period.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

All prescription medicines may cause side effects in some patients. If you have some side-effects such as upset stomach, nausea, sleepiness or tiredness, your doctor may ask you to take TEVA-ATOMOXETINE twice a day with meals, or in the evening. Most side effects will disappear after the first few weeks.

Weight loss may occur after starting TEVA-ATOMOXETINE, especially in the first few weeks. Growth rates (weight and height) after 3 years of treatment are near normal. It is not known if growth will be slowed in children who use TEVA-ATOMOXETINE for a longer period

of time. Your doctor will watch your weight and height. If you are not growing or gaining weight as expected, your doctor may change your treatment of TEVA-ATOMOXETINE.

Atomoxetine hydrochloride can cause liver damage in rare cases. Call your doctor right away if you have dark urine, yellow skin/eyes, upper right-sided abdominal tenderness, or unexplained nausea, tiredness, itching or flu-like symptoms.

Stop taking Teva-Atomoxetine and call your doctor right away if you get swelling or hives. Teva-Atomoxetine can cause a serious allergic reaction in rare cases.

New or Worsened Emotional or Behavioural Problems: A small number of patients taking drugs of this type may feel worse instead of better; for example, they may experience new or worsened feelings of agitation, hostility or anxiety, or thoughts about suicide. Your doctor should be informed of such changes immediately. Close observation by a doctor is necessary in this situation. See also the WARNINGS AND PRECAUTIONS section.

The following common side effects were reported in clinical trials with ATOMOXETINE:

In teenagers and children over 6:

- upset stomach
- decreased appetite
- nausea or vomiting
- dizziness
- tiredness
- constipation
- low blood pressure

In Adults:

- constipation
- dry mouth
- nausea
- decreased appetite
- dizziness
- problems sleeping
- sexual side effects
- problems urinating
- menstrual cramps
- rapid or irregular heartbeat
- tiredness

SERIOUS SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Symptom / effect	Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
	Only if severe	In all cases	
Swelling or hives			√
Dark urine, yellow skin/eyes, upper right-sided abdominal tenderness, or flu-like symptoms		√	

This is not a complete list of side effects. For any unexpected effects while taking TEVA-ATOMOXETINE, contact your doctor or pharmacist.

HOW TO STORE IT

TEVA-ATOMOXETINE should be stored at room temperature (15 to 30°C).

Keep all medicines, including TEVA-ATOMOXETINE, out of the reach of children.

REPORTING SUSPECTED SIDE EFFECTS

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

1. Report online at
www.healthcanada.gc.ca/medeffect
2. Call toll-free at 1-866-234-2345
3. Complete a Canada Vigilance Reporting Form and:
 - Fax toll-free to 1-866-678-6789, or
 - Mail to: Canada Vigilance Program
Health Canada
Postal Locator 0701E
Ottawa, ON K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect™ Canada Web site at www.healthcanada.gc.ca/medeffect.

NOTE: Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals can be found by contacting Teva Canada Limited at:
1-800-268-4127 ext. 5005 (English);
1-877-777-9117 (French)
or druginfo@tevacanada.com

This leaflet was prepared by:
Teva Canada Limited
30 Novopharm Court
Toronto, Ontario
Canada, M1B 2K9

Last revised: May 11, 2012