

University of Toronto
Faculty of Arts and Science
Quiz 3
MAT2371Y - Advanced Calculus
Duration - 50 minutes
No Aids Permitted

Surname: _____

First Name: _____

Student Number: _____

Tutorial:

T0101	T5101	T5102
T4/T5	R4/R5	T5/R5
Chris	Anne	Ivan
SS1074	SS1070	BA1240

This exam contains 6 pages (including this cover page) and 3 problems. Check to see if any pages are missing and ensure that all required information at the top of this page has been filled in.

No aids are permitted on this examination. Examples of illegal aids include, but are not limited to textbooks, notes, calculators, or any electronic device.

Unless otherwise indicated, you are required to show your work on each problem on this exam. The following rules apply:

- **Organize your work**, in a reasonably neat and coherent way, in the space provided. Work scattered all over the page without a clear ordering will receive very little credit.
- **Mysterious or unsupported answers will not receive full credit.** A correct answer, unsupported by calculations, explanation, or algebraic work will receive no credit; an incorrect answer supported by substantially correct calculations and explanations might still receive partial credit.
- If you need more space, use the back of the pages; clearly indicate when you have done this.

Problem	Points	Score
1	7	
2	12	
3	11	
Total:	30	

1. (a) (7 points) Let C be the curve formed by intersecting the sphere, $S = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1\}$, with the plane $P = \{(x, y, z) \in \mathbb{R}^3 \mid x + z = 1\}$. So $C = S \cap P$. Find a map $\gamma : [0, 1] \rightarrow \mathbb{R}^3$ which parameterizes C .

-
2. (a) (2 points) Define what it means for $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$ to be differentiable at a point $a \in \mathbb{R}^n$
- (b) (5 points) Using any method you know, show that $f(x, y) = \sin(xy)$ is differentiable at any point (a, b)

- (c) (5 points) Let $\gamma : [0, 1] \rightarrow \mathbb{R}^3$ be the curve $\gamma(t) = (\cos t, \sin t, t)$, and let $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ be the function $f(x, y, z) = x^2 + y^2 + z^2$. Find $(f \circ \gamma)'(0)$

-
3. (a) (2 points) Give the statement of the 1-variable Taylor's theorem with Lagrange's remainder for a $C^{k+1}(U)$ function, where $U \subseteq \mathbb{R}$ is open.
- (b) (4 points) Use Lagrange's form of the remainder to estimate the error on the interval $h \in [0, 1]$ of the 3rd order Taylor expansion of $f(x) = e^x$ about the point $x = 0$.

- (c) (5 points) Find the 3rd order Taylor polynomial of $f(x, y) = \log(1 + x - y)$ based at the point $(0, 0)$