

Aware in Medicine

Making medical knowledge accessible for all.

A Guide to Understanding Mast Cell Activation Syndrome (MCAS)

When Your Body's Alarm System Won't Stop Ringing

Author: Bernardo Werneck da Costa

Date Published: 1.9.26

Reading Time: 7 minutes



Abstract

Mast Cell Activation Syndrome (MCAS) is an immune condition in which mast cells release excessive chemical mediators in response to ordinary, non-threatening stimuli, producing allergy-like symptoms across multiple organ systems. Because its presentation overlaps with many other conditions and no single test confirms it, MCAS is frequently missed or misdiagnosed, leaving patients to manage unpredictable flare-ups for years without a clear explanation. This article breaks down what is currently understood about the causes, symptoms, diagnostic criteria, and treatment options for MCAS, and offers guidance for patients navigating daily life with the condition, so that readers can better recognize the signs, engage in informed conversations with healthcare providers, and advocate for the care they need.

Introduction

Imagine your home's fire alarm going off at random times, triggered by useless things like a slight shift in temperature, a meal, or a stressful thought. For people living with Mast Cell Activation Syndrome (MCAS), this can feel like their daily reality. MCAS is a condition affecting the immune system in which certain white blood cells, called "mast cells," become dangerously overreactive. Instead of defending the body from actual threats, they mistake everyday stimuli, or outside inputs, as a danger. This causes them to flood the body with a massive stream of chemicals and hormones directly into the bloodstream. This release leads to a host of chronic and acute allergy-like symptoms that can affect the entire body simultaneously. Medical research shows that the extent to which individuals are affected can vary wildly. Some people may be bedridden, while others experience only a slight inconvenience. Due to this variation and the similarities MCAS has to other diseases and allergies, it can be misdiagnosed by doctors. People may suffer for years without answers. Breaking down the complexities of MCAS into simple terms is a crucial step toward helping patients recognize their symptoms, advocate for their health, and secure a proper diagnosis.

Causes and Risk Factors

To understand what drives Mast Cell Activation Syndrome, it is vital to examine how our immune systems are wired to protect us. Mast cells serve as a frontline form of defence, positioned to detect and react to harmful invaders such as bacteria or toxins. In a healthy individual, these cells only release chemical signals when a legitimate threat is present. However, in an individual suffering from MCAS, these guards may react too easily, meaning that even a harmless stimulus can set them off (Akin, 2017). Recent research suggests that specific cellular and genetic changes may contribute to some cases of MCAS rather than being the confirmed primary cause of the condition (Özdemir et al., 2024). A key example involves mutations within a certain receptor (such as c-KIT) responsible for controlling mast-cell growth and signalling. When this receptor mutates, it may cause mast cells to multiply abnormally and behave erratically. This can lead to the inappropriate release of chemicals and hormones even when the body is safe from harm (Molderings et al., 2011; Özdemir et al., 2024). Furthermore, studies suggest that MCAS does not always exist in isolation. Researchers have found some shared genetic patterns and similar symptoms between MCAS and other conditions. Most notably, MCAS has been associated with a condition

called Postural Orthostatic Tachycardia Syndrome (POTS), which affects an individual's heart rate and blood pressure. This may point to broader problems in how the immune and nervous systems are regulated, which could make certain individuals more susceptible to developing the syndrome. However, information about whether MCAS can directly cause other conditions remains speculative (Özdemir et al., 2024).

Signs & Symptoms

Because mast cells live inside almost every body tissue, especially those exposed to the outside, such as the skin, gut, and airways, a flare-up can strike anywhere at any time. Sometimes, an individual might experience symptoms across multiple body parts at once. This unpredictable nature is what makes the condition extremely disruptive to daily life. On the outside, a flare-up may look like a usual allergic reaction, causing sudden skin redness, hives, or swelling. Internally, it can manifest as severe stomach aches, nausea, diarrhoea, or abdominal distress (Castells et al., 2024). However, the impact of MCAS goes beyond typical physical allergy symptoms. Many people suffer from significant neurological and psychological symptoms, most often a profound impairment of cognitive abilities, such as brain fog (Castells et al., 2024). This means that, alongside the physical discomfort, people with MCAS may struggle with memory loss, concentration issues, and mental fatigue during a flare-up. These difficulties can often resemble ADHD-like symptoms (Castells et al., 2024).

Diagnosis

Getting an official diagnosis of MCAS is difficult because its symptoms mimic other illnesses and there is no single unique symptom. To make matters worse, routine blood work and medical scans are ineffective for identifying the illness on their own. For a patient to find answers, a doctor must first suspect MCAS and then work through a long and strict checklist of criteria.

According to medical consensus, a formal diagnosis of MCAS requires meeting three specific criteria (Valent et al., 2018):

1. Episodic Symptoms: Severe, recurring symptoms affecting two or more organ systems at the same time.
2. Tryptase Spike: A blood test must show a rise in serum tryptase (a specific enzyme released by mast cells) taken during an active flare-up. This spike is measured against the patient's baseline level, specifically calculated as a 20% increase plus 2 ng/mL over baseline.
3. Clinical Response: The patient must show clear, obvious improvement when administered medications that target these chemical mediators.

Note: Symptoms vary widely between individuals; readers should not self-diagnose based on this list. If someone suspects they have this condition, they should consult a healthcare provider.

Treatment & Management

Although there is currently no cure for MCAS, there are several forms of treatment that can help control symptoms and reduce the severity of flare-ups. Since the condition affects each individual differently, there is no single medication that works for everyone. Instead, treatment is often based on the specific chemicals being released by the mast cells and the symptoms they are causing. The most common treatments for MCAS are antihistamines. H1 antihistamines are generally used to reduce symptoms involving the skin, such as itching, hives, and swelling. H2 antihistamines can be used to control symptoms affecting the digestive system, such as stomach pain, nausea, or acid reflux. Furthermore, medications known as mast cell stabilizers can be used to prevent mast cells from releasing large amounts of chemicals into the body in the first place (Weiler et al., 2019). In more serious cases, stronger medications may be considered when the patient continues to experience severe or frequent reactions. If a flare-up develops into anaphylaxis—a severe, potentially life-threatening allergic reaction, epinephrine must be administered as quickly as possible. Although antihistamines may reduce certain symptoms, they cannot replace epinephrine during a serious emergency (Weiler et al., 2019).

Emphasize: Treatment plans are highly individualized; readers should always consult their personal physician before starting or modifying treatment.

Living with MCAS

Living with MCAS can be extremely difficult because flare-ups are often unpredictable and may be caused by normal parts of everyday life. Certain foods, medications, exercise, changes in temperature, and even emotional stress may cause the mast cells to react. However, triggers vary wildly between patients, meaning something that causes a severe reaction in one person may cause no reaction in another. Due to this, identifying personal triggers is an important part of managing the condition. Many individuals record their symptoms, meals, activities, and medications in order to determine whether a pattern exists. This information can then be shared with a doctor, helping both the patient and the medical professional create a more effective and personalized treatment plan. Furthermore, individuals at risk of experiencing anaphylaxis may be required to carry epinephrine and have an emergency plan prepared. Avoiding confirmed triggers and taking medication as prescribed can reduce

the number of flare-ups. However, because MCAS can change over time and affects every patient differently, consistent communication with a medical professional

remains vital to safely living with the condition (Weiler et al., 2019). **Conclusion**

Mast Cell Activation Syndrome is a complex condition where mast cells become dangerously overreactive and release large amounts of chemicals even when the body is not facing an actual threat. Since these cells exist throughout almost every body tissue, the symptoms can affect multiple organs simultaneously and vary significantly between individuals.

The condition is especially difficult to diagnose because its symptoms closely resemble those of allergies and several other illnesses. In order to receive an official diagnosis, an individual must experience repeated symptoms across multiple organs, show measurable evidence of mast cell activation, and improve after receiving medications that target the chemicals being released. Although there is currently no cure for MCAS, treatments such as antihistamines, mast cell stabilizers, and emergency epinephrine can help patients control their symptoms and prevent serious reactions. Research surrounding the condition is still developing, meaning its exact causes and connections with conditions such as POTS are not yet fully understood. Increasing awareness and breaking down the complexities of MCAS into understandable terms can help individuals recognize their symptoms, communicate with doctors, and advocate for the medical care they require.

References

All claims in this article are supported by evidence-based sources.

1. Akin, C. (2017). Mast cell activation syndromes. *Journal of Allergy and Clinical Immunology*, 140(2), 349–355.
<https://doi.org/10.1016/j.jaci.2017.06.007>
2. Castells, M., et al. (2024). Mast cell activation syndrome: Current understanding and research needs. *Journal of Allergy and Clinical Immunology*, 154(2), 255–263.
<https://doi.org/10.1016/j.jaci.2024.05.025>
3. Molderings, G. J., et al. (2011). Mast cell activation disease: A concise practical guide for diagnostic workup and therapeutic options. *Journal of Hematology & Oncology*, 4(10), 1–8.
<https://doi.org/10.1186/1756-8722-4-10>
4. Özdemir, Ö., et al. (2024). Mast cell activation syndrome: An up-to-date review of literature. *World Journal of Clinical Pediatrics*, 13(2), 92813, 1–12.
<https://doi.org/10.5409/wjcp.v13.i2.92813>

5. Valent, P., et al. (2018). Mast cell activation syndrome: Importance of consensus criteria and call for research. *Journal of Allergy and Clinical Immunology*, 142(3), 1008–1010. <https://doi.org/10.1016/j.jaci.2018.06.005>
6. Weiler, J. M., et al. (2019). Management of mast cell activation syndrome: A multidisciplinary approach. *Annals of Allergy, Asthma & Immunology*, 123(5), 429–438. <https://doi.org/10.1016/j.anai.2019.08.010>

Aware in Medicine (AiM)

Making medical knowledge accessible for all.

Aware in Medicine is a youth-led non-profit learning hub dedicated to improving health literacy by breaking down complex medical conditions into clear, research-backed explanations for patients, students, and families.

awareinmedicine.org | awareinmedicine@gmail.com

© 2026 Aware in Medicine. All rights reserved.