HIST293. AGE-ADJUSTED DEATH RATES FOR SELECTED CAUSES BY RACE AND SEX USING YEAR 2000 STANDARD POPULATION: DEATH REGISTRATION STATES, 1900-32 AND UNITED STATES, 1933-49

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the applicable revision of the international Classification of Diseases. The classification is revised periodically to incorporate and capture changes in medical knowledge. The revisions, is nown instances, do create breaks in the comparability of a cause of death the between two years. Refer to Comparability Study reports for further information.]

SUICIDE

ICD Revision	Years in use	ICD No
Sixth	1949-1949	E963, E970-E979
Fifth	1939-1948	163, 164
Fourth	1930-1938	163-171
Third	1921-1929	165-174
Second	1910-1920	155-163
First	1900-1909	155-163

	First	1900-1909 155-1	63						
Cause of death		All races			White			All other	
and year	Both	Male	Female	Both	Male	Female	Both	Male	Female
1949	13.2	21.4	5.5	14.0	22.5	5.9	5.2	8.9	1.7
1948	13.0	20.5	5.8	13.7	21.6	6.2	4.9	8.4	1.5
1947	13.4	21.1	6.1	14.2	22.3	6.5	4.8	8.1	1.6
1946	13.2	20.5	6.3	14.0	21.7	6.7	4.5	7.3	1.9
1945	12.7	19.7	6.3	13.5	20.9	6.8	4.2	7.1	1.5
1944	11.5	17.5	6.0	12.2	18.6	6.4	3.6	5.7	1.6
1943	12.0	18.3	6.0	12.7	19.4	6.4	3.5	5.5	1.5
1942	14.0	21.7	6.4	14.8	23.0	6.8	4.4	7.0	1.9
1941	15.0	23.3	6.9	15.9	24.6	7.4	4.9	7.9	1.9
1940	16.8	26.1	7.5	17.8	27.7	8.0	5.4	8.5	2.3
1939	16.7	26.1	7.3	17.8	27.8	7.8	5.1	8.0	2.2
1938	18.1	28.4	7.7	19.2	30.1	8.1	6.0	9.4	2.7
1937	17.9	27.9	7.9	19.0	29.6	8.4	5.8	9.0	2.5
1936	17.3	26.8	7.6	18.3	28.5	8.1	5.6	8.3	2.8
1935	17.5	27.3	7.5	18.5	28.9	8.0	6.0	9.1	2.7
1934	18.4	29.0	7.7	19.5	30.7	8.1	6.5	9.6	3.2
1933	20.1	32.3	7.7	21.3	34.2	8.2	6.6	10.5	2.6
1932	21.9	35.2	8.2	23.1	37.2	8.7	7.4	11.9	2.7
1931	21.4	34.1	8.2	22.6	36.2	8.7	6.6	10.3	2.7
1930	19.9	31.4	8.0	21.1	33.3	8.5	6.6	10.1	2.7
1929	18.0	28.0	7.7	19.0	29.5	8.2	6.4	10.2	2.5
1928	17.7	27.8	7.3	18.7	29.4	7.8	6.2	9.4	2.7
1927	17.2	26.8	7.3	18.1	28.2	7.7	5.7	8.6	2.4
1926 1925	16.6 15.8	25.5 24.4	7.4 6.9	17.4 16.6	26.8 25.7	7.8 7.3	5.6 5.2	8.1 8.0	2.9 2.1
									2.1
1924 1923	15.7 15.3	24.4 23.4	6.7 7.0	16.5 16.2	25.7 24.7	7.0 7.4	5.4 4.8	8.0 7.2	2.4
1923	15.6	23.4	6.9	16.5	25.5	7.4	4.8 5.4	7.2 8.3	2.0
1921	16.3	25.4	6.8	17.1	26.7	7.2	5.6	8.7	2.2
1921	13.4	19.8	6.9	14.1	20.8	7.2	4.6	6.7	2.1
1919	15.1	22.4	7.6	15.9	23.6	8.1	4.5	6.8	2.0
1918	15.7	23.6	7.5	16.2	24.5	7.8	6.9	10.4	3.0
1917	16.7	25.2	7.8				0.5		5.0
1916	17.8	27.6	7.5	18.4	28.3	7.9	6.0	9.5	2.0
1915	20.5	31.6	8.9	20.9	32.3	9.0	10.0	14.1	5.1
1914	20.1	30.9	8.8	20.5	31.5	8.9	10.4	15.8	4.2
1913	19.2	29.6	8.1						
1912	19.7	30.6	8.3						
1911	20.0	30.8	8.6						
1910	19.4	30.0	8.3						
1909	20.0	31.4	8.2						
1908	21.1	32.8	8.9						
1907	18.2	28.5	7.7						
1906	16.4	26.1	6.6						
1905	17.5	27.6	7.7						
1904	15.4	24.3	6.6						
1903	14.3	22.5	6.1						
1902	13.5	21.1	6.0						
1901	13.3	20.7	6.0						
1900	13.1	21.3	5.2						

⁻⁻⁻ Data not available. No tables exist by age, race, sex, and cause for 1900-13 and 1917.

Note: 1) See last page of table to see when each state was admitted to the death-registration area.

²⁾ No corrections were made for those deaths for which the age of the decedent was not stated or was given as unknown.

Source: CDC/National Center for Health Statistics, National Vital Statistics System, Mortality.

HIST293. AGE-ADJUSTED DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY RACE AND SEX: UNITED STATES, 1950-59

[Based on age-specific death rates per 100,000 population in specified group. Computed by the direct method, using as the standard population the age distribution of the total population of the United States as projected for 2000. Numbers after causes of death are category numbers of the seventh revision, International Classification of Diseases]

CAUSE OF DEATH CODE	AL	L RACES -	-		WHITE	ALL OTHER			
AND YEAR	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
SUICIDE (E963, E970-E979)									
1959	12.3	20.2	5.2	13.0	21.2	5.5	6.0	10.0	2.4
1958	12.4	20.4	5.3	13.1	21.4	5.6	5.6	9.2	2.2
1957	11.4	18.7	4.8	12.0	19.7	5.1	5.2	8.9	1.8
1956	11.6	19.0	4.9	12.3	20.0	5.2	4.7	7.7	2.0
1955	11.8	19.2	5.1	12.5	20.3	5.4	4.7	7.8	1.7
1954	11.7	19.4	4.6	12.3	20.4	4.9	5.0	8.4	1.8
1953	11.7	19.3	4.7	12.3	20.3	5.0	4.8	8.4	1.4
1952	11.6	18.9	4.9	12.3	20.0	5.2	4.5	7.7	1.5
1951	12.0	19.4	5.2	12.7	20.4	5.4	4.9	8.0	1.9
1950	13.2	21.2	5.6	13.9	22.3	6.0	5.2	8.6	1.9
HOMICIDE (E964, E980-E985)									
1959	5.0	7.7	2.5	2.7	3.8	1.5	25.4	41.8	10.4
1958	4.9	7.5	2.5	2.6	3.7	1.5	24.9	41.0	10.2
1957	4.8	7.4	2.3	2.4	3.5	1.4	25.5	42.5	9.9
1956	4.9	7.5	2.4	2.4	3.5	1.3	25.9	42.4	10.9
1955	4.8	7.5	2.3	2.5	3.6	1.3	25.2	41.7	10.1
1954	5.1	7.9	2.3	2.6	3.7	1.4	26.9	45.3	10.1
1953	5.1	7.9	2.3	2.6	3.7	1.4	27.0	45.5	10.0
1952	5.4	8.5	2.4	2.6	3.9	1.4	29.2	49.2	10.9
1951	5.0	7.8	2.4	2.5	3.7	1.4	26.8	44.4	10.6
1950	5.3	8.3	2.4	2.7	4.0	1.4	28.6	47.6	10.9

Source: CDC/NCHS, National Vital Statistics System, Mortality.

HIST293.. AGE-ADJUSTED DEATH RATES FOR 60 SELECTED CAUSES BY RACE AND SEX USING YEAR 2000 STANDARD PAGE 35 POPULATION: UNITED STATES, 1960-67

(RATES PER 100,000 U.S. STANDARD YEAR 2000 POPULATION. NUMBERS AFTER CAUSES OF DEATH ARE CATEGORY NUMBERS OF THE SEVENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1955)

CAUSE OF DEATH,	JSE OF DEATH, ALL RACES				- WHITE -		ALL OTHER			
AND YEAR	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	
SUICIDE (E963,	, E970-E979	9)								
1967	12.5	19.1	6.9	13.2	20.1	7.3	6.3	10.0	3.1	
1966	12.7	19.7	6.7	13.4	20.7	7.1	6.5	10.6	2.9	
1965	13.0	20.0	7.0	13.7	21.0	7.4	6.5	10.4	3.2	
1964	12.7	19.8	6.4	13.4	20.8	6.9	5.9	9.4	2.7	
1963	13.0	20.2	6.6	13.8	21.4	7.0	6.5	10.8	2.7	
1962	12.8	20.3	6.2	13.6	21.5	6.6	6.0	9.7	2.7	
1961	12.2	19.7	5.5	12.8	20.6	5.9	6.0	10.1	2.3	
1960	12.5	20.0	5.6	13.1	21.1	5.9	5.9	9.5	2.5	
HOMICIDE (E964	4, E980-E98	35)								
1967	7.5	11.9	3.4	3.9	5.9	2.1	35.3	60.5	13.3	
1966	6.5	10.2	3.1	3.4	5.0	1.9	31.4	53.3	12.0	
1965	6.1	9.6	2.8	3.2	4.8	1.7	29.3	49.5	11.3	
1964	5.7	8.8	2.7	3.0	4.3	1.7	27.3	46.3	10.3	
1963	5.4	8.4	2.6	2.9	4.3	1.6	26.3	44.2	10.2	
1962	5.4	8.2	2.6	2.9	4.2	1.7	26.0	43.6	10.1	
1961	5.2	7.9	2.6	2.8	4.0	1.7	24.7	41.1	9.9	
1960	5.2	7.9	2.6	2.7	3.9	1.5	25.4	41.4	10.9	

FOOTNOTES: * FIGURE DOES NOT MEET STANDARD OF RELIABILITY OR PRECISION, ESTIMATION IS BASED ON FEWER THAN 20 EVENTS IN THE NUMERATOR OR DENOMINATOR.

0.0 QUANTITY MORE THAN ZERO BUT LESS THAN 0.05.

... CATEGORY NOT APPLICABLE.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished table NEWSTAN 60-67S created on 00/02/10.

HIST293. AGE-ADJUSTED DEATH RATES FOR 69 SELECTED CAUSES BY RACE AND SEX USING YEAR 2000 STANDARD PAGE 40 POPULATION: UNITED STATES, 1968-78

(RATES PER 100,000 U.S. STANDARD YEAR 2000 POPULATION. NUMBERS AFTER CAUSES OF DEATH ARE CATEGORY NUMBERS OF THE EIGHTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, ADAPTED, 1965)

							ALL OTHER					
CAUSE OF DEATH,		- ALL RACE	S		- WHITE -			- TOTAL -			- BLACK -	
AND YEAR	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
ALL OTHER ACCI		300-E807,E	825-E949)									
1978	26.5	38.3	16.2	25.2	36.2	15.6	35.1	54.3	19.4	36.5	56.4	20.2
1977	27.2	39.2	16.8	25.9	37.1	16.1	36.2	55.1	20.3	37.9	57.8	21.3
1976	27.8	40.0	17.1	26.5	37.9	16.6	36.1	55.7	19.7	37.5	58.1	20.3
1975	30.0	43.1	18.4	28.6	40.7	17.8	39.4	60.3	21.6	40.8	62.7	22.5
1974	31.4	44.3	19.8	30.0	42.0	19.1	40.6	61.1	23.0	42.0	63.8	23.6
1973	33.1	46.7	20.9	31.6	44.1	20.3	43.4	66.3	23.9	44.3	67.8	24.6
1972	33.2	46.5	21.2	31.5	43.8	20.5	44.8	67.4	25.3	46.1	69.8	25.9
1971	33.7	46.9	21.7	31.9	44.0	20.9	46.1	69.4	26.2	47.2	71.1	26.9
1970	34.7	48.6	22.0	32.7	45.5	21.1	48.1	72.8	26.5	50.1	76.4	27.5
1969	36.4	50.4	23.7	34.7	47.7	22.9	48.5	71.6	28.3	49.1	72.3	29.0
1968	36.9	50.0	24.8	35.0	46.8	24.0	50.4	75.4	28.6	51.2	77.0	29.2
SUICIDE (E950-E9	1501											
1978	12.9	20.2	6.6	13.6	21.2	7.1	7.3	12.1	3.3	7.1	12.0	3.0
1977	13.7	21.4	7.1	14.5	22.5	7.1	7.6	12.1	3.6	7.1	11.8	3.2
1976	13.7	20.4	7.1	13.9	21.3	7.6	7.6	12.5	3.4	7.1	12.1	3.1
1975	13.6	20.4	7.3	14.3	21.3	7.8	7.4	12.0	3.5	6.8	11.5	2.9
1974	13.0	20.9	7.1	13.9	21.9	7.6	7.4		3.2	6.8	11.5	2.9
								12.0				
1973	13.1	20.1	7.1	13.8	21.1	7.6	7.2	11.6	3.4	6.6	11.2	2.8
1972	13.3	20.3	7.5	14.0	21.2	7.9	7.4	11.9	3.6	6.8	11.1	3.1
1971	13.1	19.7	7.6	13.9	20.7	8.1	6.9	10.5	3.8	6.2	9.7	3.3
1970	13.1	19.8	7.4	13.8	20.8	7.9	6.7	10.5	3.4	6.2	10.0	2.9
1969	12.7	19.4	7.1	13.4	20.3	7.6	6.6	10.7	3.2	6.0	9.8	2.8
1968	12.4	19.1	6.6	13.1	20.1	7.1	5.9	9.6	2.9	5.4	9.0	2.4

FOOTNOTES: * FIGURE DOES NOT MEET STANDARD OF RELIABILITY OR PRECISION, ESTIMATION IS BASED ON FEWER THAN 20 EVENTS IN THE NUMERATOR OR DENOMINATOR.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished table NEWSTAN 68-78S created on 00/01/27.

^{0.0} QUANTITY MORE THAN ZERO BUT LESS THAN 0.05.

^{...} CATEGORY NOT APPLICABLE.

HIST293. AGE-ADJUSTED DEATH RATES FOR 72 SELECTED CAUSES BY RACE AND SEX USING YEAR 2000 STANDARD PAGE 83

POPULATION: UNITED STATES, 1979-98

(RATES PER 100,000 U.S. STANDARD YEAR 2000 POPULATION. NUMBERS AFTER CAUSES OF DEATH ARE CATEGORY NUMBERS OF THE NINTH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES, 1975)

									ALL OTH	ER	-	
CAUSE OF DEATH,	A	LL RACES			- WHITE			- TOTAL			- BLACK	
AND YEAR	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE	BOTH	MALE	FEMALE
SUICIDE (E950-E9	59)											
1998	11.3	19.2	4.3	12.2	20.6	4.7	6.3	11.0	2.3	5.8	10.6	1.8
1997	11.4	19.4	4.4	12.3	20.6	4.8	6.7	11.7	2.4	6.3	11.4	2.0
1996	11.7	20.0	4.3	12.5	21.3	4.7	6.8	11.7	2.6	6.6	11.9	2.0
1995	12.0	20.6	4.4	12.8	21.9	4.7	7.1	12.3	2.6	6.9	12.5	2.1
1994	12.1	20.7	4.5	12.8	21.9	4.8	7.3	12.8	2.6	7.0	12.9	2.1
1993	12.2	20.9	4.6	13.0	22.1	5.0	7.3	12.6	2.7	7.1	13.0	2.2
1992	12.1	20.6	4.7	12.9	21.9	5.0	7.0	12.1	2.6	6.9	12.6	2.1
1991	12.3	21.2	4.7	13.2	22.5	5.1	7.1	12.6	2.4	7.0	13.0	2.0
1990	12.5	21.5	4.8	13.4	22.8	5.2	7.2	12.5	2.7	7.1	12.8	2.4
1989	12.3	21.0	4.9	13.1	22.3	5.2	7.2	12.4	2.8	7.2	13.0	2.5
1988	12.5	21.2	5.1	13.3	22.5	5.5	6.9	11.9	2.7	6.9	12.2	2.5
1987	12.8	21.7	5.3	13.7	23.0	5.7	7.1	12.7	2.6	6.9	12.8	2.1
1986	13.0	21.9	5.5	13.9	23.2	6.0	7.0	12.1	2.8	6.8	12.2	2.4
1985	12.5	21.1	5.2	13.4	22.4	5.7	6.8	11.7	2.7	6.6	11.8	2.3
1984	12.6	20.9	5.6	13.5	22.1	6.0	6.8	11.6	2.8	6.6	11.7	2.3
1983	12.4	20.4	5.5	13.2	21.6	6.0	6.5	11.2	2.7	6.2	11.1	2.2
1982	12.5	20.4	5.8	13.3	21.6	6.2	6.5	11.1	2.7	6.3	11.2	2.3
1981	12.3	19.8	6.0	13.1	20.9	6.4	6.9	11.5	3.1	6.5	11.4	2.5
1980	12.2	19.9	5.7	13.0	20.9	6.1	6.8	11.6	2.8	6.5	11.4	2.4
1979	12.6	19.9	6.3	13.2	20.7	6.7	7.7	13.0	3.3	7.3	12.7	2.9

FOOTNOTES: * FIGURE DOES NOT MEET STANDARD OF RELIABILITY OR PRECISION, ESTIMATION IS BASED ON FEWER THAN 20 EVENTS IN THE NUMERATOR OR DENOMINATOR.

- 0.0 QUANTITY MORE THAN ZERO BUT LESS THAN 0.05.
- ... CATEGORY NOT APPLICABLE.
- 1/ INCLUDED IN "ALL OTHER INFECTIOUS AND PARASITIC DISEASES" CATEGORY SHOWN ABOVE.
- 2/ INCLUDED IN "ALL OTHER DISEASES (RESIDUAL)" CATEGORY SHOWN ABOVE.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished table NEWSTAN 79-98S created on 00/03/02.