[Rates per 100,000 population for specified group.]

ICD Revision	Years in use	ICD Nos
Fifth	1939	163, 164
Fourth	1930-1938	163-171
Third	1921-1929	165-174
Second	1910-1920	155-163
First	1900-1909	155-163

ALL	KAU	_£5,	RO	ΙH	2F	KE:

ALL RACES	, BOTH SEXES	S										
ears/	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	10.2				6.8	11.2	15.8	19.9	23.5	26.1	29.2	
1901	10.4				6.8	11.4	15.4	20.5	25.6	25.6	29.2	
1902	10.3				7.4	10.3	16.1	20.0	24.3	25.5	30.3	
1903	11.3				6.6	13.3	18.9	22.9	23.0	27.9	24.4	
1904	12.2				8.5	14.0	17.2	25.3	27.9	27.7	32.2	
1905	13.5				9.4	14.9	19.6	26.5	30.3	33.0	37.9	57.1
1906	12.8				8.0	14.1	19.8	26.5	30.5	31.4	31.0	
1907	14.5				10.8	17.0	21.1	28.4	33.6	35.5	32.7	
1908	16.8				11.3	19.4	25.2	36.0	40.2	35.7	34.3	33.2
1909	16.0			0.3	10.5	18.1	23.8	33.7	37.5	35.9	34.0	30.5
1910	15.3			0.3	10.7	17.7	22.4	29.9	37.0	35.3	38.7	33.4
1911	16.0			0.2	10.9	18.8	24.5	32.1	37.3	35.6	34.8	27.8
1912	15.6			0.3	9.8	18.4	23.6	31.1	38.7	34.2	38.1	28.9
1913	15.4			0.2	10.4	18.9	22.9	30.1	36.7	35.1	31.6	27.1
1914	16.1			0.3	10.8	18.9	23.6	32.5	40.0	35.7	32.7	31.3
1915	16.2			0.3	9.8	18.6	23.9	32.2	41.3	39.7	35.1	36.8
1916	13.7			0.2	8.7	15.5	20.1	26.9	35.9	36.3	34.4	29.9
1917	13.0			0.3	8.3	16.7	18.9	25.0	30.1	34.7	32.5	27.0
1918	12.3			0.2	7.3	15.8	19.2	22.0	29.0	30.0	30.4	31.5
1919	11.5			0.2	6.6	13.7	18.1	21.6	27.7	29.4	32.2	39.5
1920	10.2			0.2	6.2	11.5	15.7	20.4	24.2	27.4	28.2	32.5
1921	12.4			0.3	6.9	13.3	19.0	26.2	31.3	32.5	32.3	35.0
1922	11.7			0.2	6.4	12.1	18.0	24.8	29.7	33.8	32.8	35.1
1923	11.5			0.2	5.7	11.9	17.8	24.0	29.5	31.2	35.0	35.4
1924	11.9			0.2	6.3	11.9	18.6	24.9	31.3	33.4	30.8	31.0
1925	12.0			0.2	5.9	12.2	18.0	25.2	30.3	35.8	31.7	32.4
1926	12.6			0.2	6.5	12.2	19.2	26.4	31.9	36.6	36.1	32.1
1927	13.2			0.3	6.4	13.1	19.6	27.7	33.7	37.4	35.9	34.0
1928	13.5			0.2	6.7	13.4	19.4	29.0	34.3	40.4	39.4	29.9
1929	13.9			0.2	7.1	13.6	20.3	28.7	35.0	41.0	40.8	25.3
1930	15.6			0.2	7.5	14.9	22.7	33.0	41.2	42.8	40.1	29.2
1931	16.8			0.2	7.2	15.1	22.9	36.0	46.9	48.6	42.1	33.9
1932	17.4			0.2	7.3	15.5	22.8	36.4	49.1	51.3	45.5	27.5
1933	15.9			0.2	7.2	13.9	21.3	33.2	43.0	46.1	45.1	31.7
1934	14.9			0.2	7.4	14.5	19.9	29.6	37.2	40.9	41.2	25.6
1935	14.3			0.2	7.1	14.3	19.0	28.6	34.9	35.8	38.7	27.3
1936	14.3			0.3	6.8	14.5	19.7	27.6	34.1	34.7	38.0	26.8
1937	15.0			0.3	6.9	14.4	20.7	30.1	35.5	34.5	38.6	27.7
1938	15.3			0.2	6.7	14.5	21.0	30.7	37.5	34.4	36.1	27.7
1939	14.1			0.3	5.9	12.9	18.9	28.2	34.4	34.0	36.0	24.1

^{...} Category not applicable
* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.
Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (#6/91917).
Source: CD/CN/CN, Sutational VMS alstatics System, Mortality, Unpublished tables and National Office of VMS alstatics, Valual Statistics System, Mortality, Unpublished tables and National Office of VMS alstatics, Valual Statistics System, Mortality, Unpublished tables and National Office of VMS alstatiscs, Vital Statistics Special Reports, Volume 43, 1956.

[Rates per 100,000 population for specified group.]

SUICIDE

ALL RACES,												
Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years		85 years and over
1900	15.7				7.7	15.2	24.5	32.9	40.0	44.6	56.4	
1901	15.5				8.0	15.6	20.7	32.1	43.9	42.5	58.3	
1902	15.4				8.4	13.9	24.7	30.4	39.4	42.1	59.1	
1903	17.2				8.1	18.0	28.7	37.2	36.9	47.7	43.8	
1904	18.5				9.7	18.9	26.3	40.1	45.8	48.9	57.6	
1905	20.2				10.6	20.1	29.5	41.3	51.4	58.1	67.0	108.8
1906	19.6				9.8	20.0	30.1	41.3	50.0	55.0	57.6	
1907	21.9				13.5	24.0	30.8	44.1	54.4	61.5	59.4	
1908	25.5				13.8	27.0	37.9	56.2	66.8	59.1	59.8	66.8
1909	24.2				12.3	25.0	36.7	52.3	60.9	62.2	62.4	64.5
1910	23.0				12.9	24.9	32.6	46.4	61.2	58.2	68.9	61.7
1911	23.9				13.0	26.1	36.8	49.4	60.7	58.6	61.9	54.5
1912	23.4				11.5	25.7	34.9	48.1	62.2	58.8	68.3	57.5
1913	23.2				12.0	26.7	33.9	47.4	59.9	59.7	55.8	44.6
1914	24.1				12.5	27.0	34.8	50.4	62.9	59.5	58.2	62.3
1915	24.3				11.5	26.1	35.0	49.2	66.5	65.4	61.8	75.6
1916	20.7				10.4	21.7	30.1	40.1	57.3	61.0	63.2	64.1
1917	19.2			0.3	9.9	23.4	26.8	36.5	47.8	57.0	58.1	48.3
1918	18.2				9.2	23.4	28.0	31.6	43.7	48.5	52.1	60.7
1919	16.5			0.3	7.5	19.3	25.7	29.8	41.2	46.9	56.5	80.0
1920	14.5			0.2	7.1	15.1	21.5	28.7	36.6	44.6	49.9	63.5
1921	18.9			0.4	8.3	18.7	28.5	39.9	49.6	54.4	57.9	73.0
1922	17.6			0.2	7.1	17.0	26.2	37.4	46.6	56.7	57.6	70.9
1923	17.0			0.3	6.0	16.1	26.0	36.3	45.4	52.7	61.6	67.8
1924	18.1			0.2	7.4	16.4	27.5	38.0	49.7	56.1	56.0	61.6
1925	18.0			0.3	7.2	16.3	26.1	37.7	48.3	60.4	59.0	63.9
1926	18.8			0.3	7.4	16.1	28.0	39.6	50.3	61.9	64.4	62.6
1927	19.9			0.3	7.9	17.9	28.6	42.4	53.7	63.6	64.9	67.6
1928	20.7			0.2	7.8	18.4	28.6	45.1	55.4	68.4	73.1	57.1
1929	21.1			0.3	8.6	18.4	30.0	43.9	54.8	69.4	73.5	51.2
1930	24.1			0.3	9.1	20.6	34.2	51.9	66.6	72.3	72.6	58.3
1931	26.2			0.3	8.7	20.9	34.4	56.9	77.0	84.3	76.1	70.4
1932	27.4			0.3	9.0	21.8	34.7	58.0	81.4	87.6	84.9	53.4
1933	24.9			0.2	8.8	19.4	31.6	52.8	70.2	79.9	83.8	64.5
1934	22.8			0.3	9.2	20.3	29.2	45.0	60.1	70.6	75.1	51.3
1935	21.7			0.3	9.0	19.7	27.5	44.1	55.5	60.1	71.9	57.0
1936	21.7			0.4	8.8	19.7	29.0	41.9	54.4	58.2	69.3	53.9
1937	22.8			0.5	9.0	20.4	30.4	45.5	56.1	58.1	70.3	57.1
1938	23.5			0.3	9.0	20.9	31.1	47.0	60.2	58.0	66.0	54.6
1939	21.7			0.4	8.0	18.8	27.5	42.1	54.8	56.3	67.3	47.7

... Category not applicable

* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 3910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: COZ/NOR, Sattonian VMS attainskies System, Mortality, Unpublished adabes and National Office of VMS attainskies, Valual Sattainskies, System, Mortality, Unpublished adabes and National Office of VMS attainskies, Valual Sattainskies, Special Reports, Volume 43, 1956.

[Rates per 100,000 population for specified group.]

SUICIDE

ALL RACES,												
ears/	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900	4.7				6.0	7.1	6.5	6.3	7.3	8.6		
1901	5.3				5.6	7.1	9.7	8.5	7.7	9.5		
1902	5.2				6.3	6.6	6.9	9.0	9.6	9.9		
1903	5.4				5.1	8.5	8.4	7.8	9.3	9.2		
1904	6.0				7.3	8.9	7.5	9.7	10.2	7.7		
1905	6.8				8.3	9.6	9.2	10.8	9.4	9.3		
1906	5.8				6.2	7.8	8.4	10.3	10.4	8.4		
1907	6.9				8.0	9.5	10.4	11.0	12.1	10.2		
1908	7.8				8.8	11.1	11.1	13.5	12.1	12.4	10.9	
1909	7.3				8.8	10.6	9.5	13.0	12.7	9.8	8.0	•
1910	7.2				8.4	9.6	11.0	11.4	11.1	12.5	11.1	•
1911	7.5				8.8	10.8	10.8	12.7	12.0	12.3	9.8	
1912	7.3				8.1	10.4	11.0	12.0	13.2	9.5	10.4	
1913	7.1				8.7	10.3	10.7	10.5	11.6	10.3	9.6	
1914	7.7				9.2	10.2	11.2	12.2	15.0	11.6	9.3	
1915	7.6				8.0	10.4	11.5	13.0	13.7	13.6	10.8	
1916	6.4				6.9	8.9	9.2	11.9	12.5	11.1	8.2	•
1917	6.6				6.7	9.7	10.3	11.9	10.6	11.8	9.3	•
1918	6.2				5.7	8.7	9.6	11.0	12.9	11.1	10.6	•
1919	6.3				5.7	8.2	9.8	12.2	12.7	11.4	10.0	•
1920	5.7				5.3	7.7	9.3	11.1	10.6	9.5	8.3	
1921	5.7				5.5	7.8	8.7	10.8	11.3	9.8	8.8	
1922	5.7				5.7	7.2	9.0	10.8	11.4	10.1	10.1	
1923	5.8				5.4	7.7	8.9	10.6	12.2	9.0	10.5	
1924	5.6				5.2	7.4	8.9	10.5	11.4	9.7	7.5	•
1925	5.8				4.7	8.0	9.3	11.6	10.7	10.3	6.5	•
1926	6.2				5.7	8.2	9.7	12.0	11.8	10.3	10.0	•
1927	6.2			0.2	5.1	8.4	10.1	11.7	11.8	10.4	8.9	•
1928	6.2				5.7	8.4	9.6	11.6	11.3	11.4	7.9	•
1929	6.6				5.7	8.7	10.1	12.0	13.4	11.7	10.2	•
1930	6.9				6.1	9.1	10.7	12.3	13.8	12.2	9.5	•
1931	7.1				5.6	9.4	10.9	13.2	14.5	11.8	10.2	•
1932	7.1				5.7	9.3	10.5	13.0	14.3	14.0	8.5	•
1933	6.8				5.7	8.4	10.5	12.0	13.9	11.4	8.7	•
1934	6.8				5.5	8.7	10.4	12.9	12.7	10.4	9.6	•
1935	6.8				5.2	9.1	10.2	12.0	13.1	11.2	7.7	•
1936	6.8			0.2	4.8	9.3	10.1	12.4	12.6	11.0	9.1	•
1937	7.0				4.9	8.5	10.7	13.7	13.6	10.8	9.3	•
1938	6.9				4.4	8.2	10.8	13.3	13.4	10.8	8.5	•
1939	6.5				3.8	7.2	10.1	13.4	12.9	11.7	7.4	•

... Category not applicable

* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 3910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: COZ/NOR, Sattonian VMS attainskies System, Mortality, Unpublished adabes and National Office of VMS attainskies, Valual Sattainskies, System, Mortality, Unpublished adabes and National Office of VMS attainskies, Valual Sattainskies, Special Reports, Volume 43, 1956.

[Rates per 100,000 population for specified group.]

WHITE, BO												
Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900												
1901												
1902												
1903												
1904												
1905												
1906												
1907												
1908												
1909												
1910	15.4											
1911	16.1											
1912	15.8											
1913	15.6											
1914	16.4	-		0.3	11.0	19.0	24.0	33.5	40.8	36.3	33.4	32.8
1915	16.5			0.3	9.9	18.6	24.3	33.0	42.2	40.6	35.8	38.6
1916	14.3			0.2	9.0	15.9	20.9	27.9	37.3	37.6	35.5	32.0
1917	13.6											
1918	12.8			0.3	7.5	16.4	19.9	23.0	30.0	30.7	31.3	34.2
1919	12.2			0.2	6.9	14.2	19.1	22.9	29.0	30.9	33.9	43.5
1920	10.8	-		0.2	6.5	12.0	16.7	21.5	25.2	28.5	29.4	35.7
1921	13.1	-		0.3	7.1	13.8	20.0	27.8	32.8	33.7	33.8	37.6
1922	12.5			0.2	6.7	12.7	18.9	26.4	31.3	35.5	34.2	38.9
1923	12.2			0.2	6.0	12.5	18.7	25.4	31.1	32.7	36.8	38.4
1924	12.7			0.2	6.6	12.4	19.5	26.4	32.9	34.9	32.3	33.5
1925	12.7			0.2	6.2	12.7	19.0	26.8	31.9	37.2	33.1	35.0
1926	13.4			0.2	6.9	12.7	20.3	27.8	33.4	38.1	37.8	34.8
1927	14.0			0.3	6.8	13.6	20.7	29.4	35.5	39.0	37.8	37.5
1928	14.5			0.2	7.1	14.0	20.5	31.0	36.1	42.4	41.4	32.3
1929	14.9			0.2	7.5	14.2	21.6	30.5	36.9	42.9	42.6	27.2
1930	16.8			0.2	8.0	15.7	24.2	35.3	43.4	44.5	42.0	31.5
1931	18.0			0.2	7.6	15.8	24.4	38.6	49.3	51.0	44.2	36.3
1932	18.7			0.2	7.6	16.4	24.2	38.9	51.5	53.9	47.4	30.1
1933	17.2			0.2	7.5	14.6	22.5	35.4	45.5	48.5	47.4	34.9
1934	16.1			0.2	7.6	15.2	21.3	31.6	39.2	43.1	42.9	27.4
1935	15.4			0.2	7.4	15.0	20.1	30.6	36.7	37.9	40.6	30.0
1936	15.4			0.3	7.1	15.4	21.0	29.6	35.9	36.7	39.8	29.8
1937	16.1			0.3	7.3	15.2	22.1	32.1	37.4	36.5	40.5	30.5
1938	16.4			0.2	7.0	15.3	22.5	32.7	39.4	36.6	37.5	30.2
1939	15.3			0.3	6.2	13.7	20.2	30.0	36.4	36.2	37.5	26.8

⁻⁻⁻ Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.
--- Category not applicable
--- Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military death; in registration states, occurring after the declaration of war (Af/91917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

[Rates per 100,000 population for specified group.]

WHITE, MA												
Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900												
1901												
1902												
1903												
1904												
1905												
1906												
1907												
1908												
1909												
1910	23.2											
1911	24.1											
1912	23.8											
1913	23.6											
1914	24.5		***	:	12.6	27.0	35.3	52.1	64.1	60.6	59.4	65.2
1915	24.8			:	11.7	26.2	35.8	50.6	68.2	66.8	63.1	79.1
1916	21.5			•	10.8	22.1	31.1	41.8	59.6	63.2	65.4	50.5
1917	20.1											
1918	19.0	-			9.5 7.7	24.1	29.0	33.3	45.2	49.8	53.9	65.7
1919	17.5	-		0.3		19.9	27.1	31.7	43.3	49.4	59.7	88.1
1920	15.4				7.5	15.7	22.7	30.2	38.3	46.6	52.2	69.6
1921	20.0	-		0.4	8.5	19.4	29.9	42.5	52.2	56.6	60.7	78.2
1922 1923	18.7 18.1	-		0.3	7.4 6.3	17.8 16.9	27.4 27.3	40.0	49.3	59.7	60.1 65.2	78.0 74.1
1923	19.2	-	***	0.3	7.7	17.1	27.3	38.4 40.5	48.0 52.3	55.3 59.0	58.9	74.1 67.3
1924	19.2		***	0.3	7.7	17.1	27.3	40.3	51.1	63.0	61.5	68.4
1925	20.0	-	***	0.3	7.4	16.8	27.3	40.3	53.0	64.7	67.7	67.1
1926	21.2			0.3	8.3	18.4	30.0	45.2	56.8	66.3	68.5	74.1
1928	22.1			0.3	8.1	19.3	30.1	48.2	58.6	72.1	77.0	61.1
1929	22.5			0.2	8.9	19.2	31.8	46.7	58.1	72.6	76.8	54.8
1930	25.9			0.3	9.6	21.6	36.2	55.7	70.6	75.3	76.1	63.8
1931	28.2			0.3	9.2	21.8	36.4	61.2	81.4	88.6	80.1	75.2
1932	29.4			0.3	9.4	23.0	36.5	62.1	85.9	92.2	88.5	58.3
1933	26.8			0.2	9.1	20.4	33.2	56.3	74.7	84.1	88.2	70.9
1934	24.6			0.3	9.6	21.3	31.0	48.1	63.6	74.8	78.1	56.2
1935	23.3			0.3	9.5	20.7	29.0	47.0	58.5	63.6	75.8	62.6
1936	23.3			0.4	9.1	21.0	30.7	44.9	57.3	61.6	73.0	59.9
1937	24.5	-		0.4	9.5	21.5	32.2	48.5	59.2	61.6	73.8	62.6
1938	25.3			0.3	9.4	22.2	33.1	50.1	63.6	61.9	68.5	59.9
1939	23.4			0.5	8.5	19.8	29.3	44.8	58.2	60.2	70.4	52.9
-555	23.4			0.5	0.5	13.0	25.5	44.0	30.2	00.2	70.4	52.5

⁻⁻⁻ Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.
--- Category not applicable
--- Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

Note: Values shown here differ from those published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military death; in registration states, occurring after the declaration of war (Af/91917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

[Rates per 100,000 population for specified group.]

WHITE, FEI												
ears/	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900												
1901												
1902												
1903												
1904												
1905												
1906												
1907												
1908												
1909												
1910	7.2											
1911	7.6											
1912	7.4											
1913	7.2											
1914	7.8	-			9.3	10.3	11.5	12.6	15.5	11.7	9.5	
1915	7.7	-		:	8.1	10.4	11.6	13.3	14.0	14.0	10.8	•
1916	6.8	-			7.2	9.3	9.7	12.4	12.9	11.5	8.5	•
1917	6.9											
1918	6.5			:	5.8	9.1	10.1	11.5	13.4	11.2	10.9	
1919	6.7				6.1	8.5	10.5	12.9	13.4	11.9	10.6	•
1920	6.1	-			5.5	8.2	10.0	11.7	11.1	9.9	8.6	•
1921	6.1	-			5.7	8.0	9.2	11.5	11.8	10.2	9.3	•
1922	6.1	-		•	6.0	7.6	9.6	11.4	11.9	10.7	10.7	•
1923	6.2		***	•	5.7	8.1	9.4	11.3	12.9	9.5	10.9	:
1924	5.9			:	5.4	7.6	9.5	11.1	11.9	10.1	8.0	
1925	6.2			:	5.0	8.4	9.9	12.3	11.3	10.7	6.9	
1926	6.6				6.1	8.6	10.2	12.6	12.3	10.7	10.3	
1927	6.6			0.2	5.3	8.8	10.6	12.4	12.4	10.9	9.4	
1928	6.6			:	6.1	8.8	10.3	12.5	11.9	11.8	8.2	
1929	7.1			:	6.0	9.3	10.8	12.9	14.1	12.3	10.7	
1930	7.4				6.4	9.8	11.4	13.2	14.5	12.7	10.0	
1931	7.6				5.9	9.9	11.8	14.2	15.2	12.3	10.5	
1932 1933	7.7 7.3				5.9 5.9	9.9	11.1	13.9	15.0	14.7	9.0	:
				:		9.0	11.3	12.9	14.6	12.1	9.1	:
1934	7.3			:	5.7	9.2	11.1	13.8	13.3	10.9	9.9	:
1935	7.2				5.4	9.4	10.9	13.0	13.8	11.9	7.9	
1936	7.3		***	0.2	5.0	9.8	10.9	13.3	13.4	11.6	9.0	:
1937	7.6			:	5.1	9.0	11.6	14.6	14.4	11.5	9.8	•
1938	7.4			:	4.6	8.5	11.7	14.3	14.0	11.4	9.0	•
1939	7.1				3.9	7.6	11.0	14.3	13.7	12.4	7.5	•

⁻⁻⁻ Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

.-- Category not applicable

Note: Values shown here differ from those upublished in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military death; in registration states or curring after the declaration of war (Af-/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

[Rates per 100,000 population for specified group.]

SUICIDE

ALL OTHER, BOTH SEXES

	, BUTH SEXE											
Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900												
1901												
1902												
1903												
1904												
1905												
1906												
1907												
1908												
1909												
1910	11.8											
1911	12.7											
1912	9.7											
1913	8.9											
1914	9.8				8.1	18.3	15.5	9.1	17.4			
1915	9.3				6.3	17.4	14.8	13.1				
1916	5.0				4.3	9.5	9.2	8.5				
1917	4.9											
1918	5.5				5.2	9.3	9.9	6.1	10.3	16.5		
1919	4.0				3.8	8.0	7.2	5.8				
1920	3.6	-			3.7	5.6	5.4	7.6	6.3			
1921	4.6	-			5.0	7.7	8.0	6.7	6.7			
1922	4.2	-			3.8	6.1	8.9	7.6	6.7			
1923	3.7	-			3.0	5.5	7.8	8.0	6.4			
1924	4.3	-			3.6	7.2	8.4	6.9	8.8			
1925	4.0				3.5	6.7	7.7	6.0	5.8	10.1		
1926	4.4				3.3	7.3	7.2	8.6	9.3			
1927	4.7				3.5	8.6	8.6	8.2	8.2			
1928	4.8	-			4.0	7.9	8.6	8.7	10.5			
1929	4.9	-			4.4	7.8	7.7	10.1	9.3	10.2		
1930	5.0	-			4.0	7.7	8.5	9.0	10.6	13.8		
1931	5.2				4.0	8.8	8.5	8.7	13.2	9.0		
1932	5.5				4.8	8.1	10.3	10.1	15.7	9.3		
1933	4.9				4.9	7.4	9.8	10.9	9.0	9.8		
1934	4.7				5.2	8.4	7.7	8.8	10.5	6.9		
1935	5.1				4.3	8.4	8.4	8.3	10.7	6.2		
1936	4.7	-			4.2	7.0	7.9	6.9	10.8	6.9		
1937	4.9				3.9	7.6	8.3	8.7	10.0	7.5		
1938	4.9				4.1	7.5	7.8	8.7	11.2	6.1		
1939	4.2	-			3.3	6.5	7.0	8.6	7.3	6.0		

⁻⁻⁻ Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

.-- Category not applicable

Note: Values shown here differ from those upublished in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1917-1918 because this includes military death; in registration states or curring after the declaration of war (Af-/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

[Rates per 100,000 population for specified group.]

ALL OTHER	R, MALE											
Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
1900												
1901												
1902												
1903												
1904												
1905												
1906												
1907												
1908												
1909												
1910	16.9											
1911	19.3											
1912	13.8											
1913	12.7											
1914	14.6				8.4	26.9	23.6	13.9	29.7			
1915	13.3					24.0	20.2	18.7				
1916	7.9				4.8	15.5	15.4	12.6				
1917	7.2											
1918	8.4	-			6.7	14.8	15.7	8.6	16.1			
1919	6.0	-			5.2	11.8	10.9	8.8				
1920	5.2				3.7	8.5	8.3	10.7				
1921	6.9				6.3	10.8	12.8	10.3				
1922	6.3	-			4.6	9.2	14.1	10.7	9.9			
1923	5.6	-			3.5	7.5	11.9	13.2	10.4			
1924	6.3	-			4.1	9.6	13.6	10.2	13.8			
1925	6.1				4.8	9.7	12.0	8.2	9.1	•		
1926	6.2				3.8	9.7	10.3	12.1	14.0			
1927	7.0				4.3	12.8	12.8	12.4	12.6			
1928	7.2	-			5.2	10.8	14.0	14.2	15.8			
1929	7.6				5.7	11.6	11.9	16.6	13.8	17.5		
1930	7.6				4.4	11.9	13.4	14.3	16.5	23.0		
1931	8.0				4.6	12.7	14.2	14.6	20.0	14.9		
1932	8.6				5.9	11.9	16.1	16.8	24.9	16.3		
1933	7.7				5.9	11.2	16.3	17.5	13.8	17.4		
1934	6.9				6.3	11.9	11.3	13.8	16.7			
1935	7.4				5.1	10.7	13.0	14.8	17.8	10.8		
1936	7.0				5.5	9.2	13.2	10.6	17.8	11.2		
1937	7.4				4.4	10.7	13.7	13.2	15.9	13.0		
1938	7.4				5.5	9.8	12.7	14.0	17.2	9.0		
1939	6.5				3.9	10.1	11.8	13.1	12.0	9.3		

⁻⁻⁻ Data not available. No original tables for the time period 1900-13 and 1917 included all of the following variables at the same time: age, race, sex, and cause.

* Figure does not meet standard of reliability or precision, estimation is based on fewer than 20 events in the numerator or denominator.

^{*} Figure does not meet standarfor feationity or precious, issumation to base on new user or precious and the standard or precious published in the annual volumes of 1910-1915 because North Carolina is not included in these tables. The values are greater than those shown in the annual volumes of 1910-1915 because this includes military deaths (in registration states) occurring after the declaration of war (4/6/1917).

Source: CDC/NCHS, National Vital Statistics System, Mortality, Unpublished tables and National Office of Vital Statistics, Vital Statistics Special Reports, Volume 43, 1956.

[Rates per 100,000 population for specified group.]

SUICIDE

ALL OTHER											
Years	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	85 years and over
1900											
1901											
1902											
1903											
1904											
1905											
1906											
1907											
1908											
1909											
1910	6.0										
1911	5.3										
1912	5.2										
1913	4.7										
1914	4.6				7.8	8.9				•	
1915	5.1					10.4				•	
1916	2.0										
1917	2.5										
1918	2.5				3.8	4.1				•	
1919	2.0					4.6				•	
1920	2.0				3.7						
1921	2.3				3.7	4.9					
1922	2.1				3.1	3.3					
1923	1.8				2.5	3.6	3.6				
1924	2.3				3.2	4.9					
1925	2.0				2.3	3.9					
1926	2.6				2.8	5.1	3.9				
1927	2.4				2.8	4.8	4.3				
1928	2.4				2.9	5.3	3.2				
1929	2.3				3.3	4.2	3.6				
1930	2.4				3.6	3.8	3.7				
1931	2.5				3.5	5.2	3.0				
1932	2.5				3.7	4.5	4.7				
1933	2.2				4.0	3.9	3.5	3.4			
1934	2.6				4.3	5.1	4.2				
1935	2.7				3.6	6.3	4.0				
1936	2.4				3.0	4.9	2.7				
1937	2.5				3.4	4.7	3.0	3.8			
1938	2.5				2.9	5.3	3.1				
1939	2.0				2.7	3.2	2.4	3.7			

Both sexes	40	60	64
Male	28	41	45
Fomalo	12	10	10