




sparc

Deriving Proton Pitch Angle Distributions Using Sentinel 6 – MF NGRM Observations

esa




E. Christodoulou⁽¹⁾, H. Evans⁽²⁾, C. Katsavrias⁽¹⁾, M. Vuolo⁽²⁾, G. Santin⁽²⁾, I. Sandberg⁽³⁾, I. A. Daglis⁽¹⁾, and P. Nieminen⁽²⁾

⁽¹⁾ National and Kapodistrian University of Athens, Athens, Greece; ⁽²⁾ ESTEC, European Space Agency, Noordwijk, The Netherlands; ⁽³⁾ Space Applications and Research Consultancy, Athens, Greece


Motivation and Data

The inner radiation belt exhibits strongly anisotropic pitch angle distributions (PADs), which are a particularly powerful diagnostic for understanding the physical processes governing particle behaviour in the inner magnetosphere.

Sentinel 6 (S6) is an Earth observation mission in near-polar, LEO orbit, that carries a **Next Generation Radiation Monitor (NGRM)**, which measures electrons and protons across multiple energy ranges [1-2]. However, the detector is mounted in a fixed orientation, (the detector bore site is aligned along the spacecraft flight direction), which fixes the pitch angle of the instrument. The orbit has a 9 day repeat cycle, which provides the minimum temporal binning limit.




- We utilise the **Level 1 Version 1 Proton Differential Flux Dataset** (Produced by SPARC [3], publicly available through SWE HAPI [4]).
- We focus on **9 MeV protons** (SDSS-2).
- Most measurements cluster within two narrow pitch-angle ranges, approximately $10^\circ - 75^\circ$ and $100^\circ - 170^\circ$ (See Figure 3).
- Northbound and southbound passes sample different pitch-angle intervals (See Figure 4).



Figures 3: Example of the PAs sampled during (a) northbound and (b) southbound passages of a quiet month.

- Pitch angle is computed as $PA = \cos^{-1} \hat{B} \hat{X}_{sc}$, where $\hat{B}$ is the magnetic field vector calculated from the IGRF magnetic field model with the Olson and Pfitzer (1977) quiet time external model, and $\hat{X}_{sc}$ is the spacecraft +X.
- Key assumption:** northbound and southbound passes sample the same underlying population in the same geographic region over the subsampled periods.



Figures 4: PAD of the measured proton fluxes at 9 MeV, where colour indicates the different L-shells.

GRAS/GEANT4 Simulations and Fitting Process




Figure 5: Heat map of the channel's response as a function of incident angle and proton energy + Schematic of how θ is measured




Figure 6: Histogram of the angular detector response as a function of $\cos \theta$. Colour corresponds to different energy bins.




Figure 7: Green curve: the best-fit PAD, blue curve: computed total efficiency. Dashed red line indicates 90° .

- Data split into 9-day segments (Jan 2024 – Jan 2026) and $4^\circ \times 4^\circ$ lat–lon spatial bins; passes are separated into northbound ($PA > 90^\circ$) and southbound ($PA < 90^\circ$) samples.
- PAD modelled as $f(\alpha) \propto \sin^n \alpha$; detector efficiency for a given n: $E_{model}(N) = \frac{\int_0^{180^\circ} R(\alpha) \sin^N \alpha da}{\int_0^{180^\circ} \sin^N \alpha da}$
- For each bin: compute the observed north/south flux ratio and compare to the modelled efficiency ratio across a grid of n (1-12, step 0.25). **Best-fit n:** $R_{observed} \approx R_{modelled}(n)$.

Results




Figure 8: Flux-map with colour showing S6-MF/NGRM Level 1 Version 1 differential fluxes in log axis, for 15-24 March 2024. The enhancement of the SAA (corresponding to the outer edge of the inner radiation belt at low latitudes) is visible.




Figure 11: Fitted PAD exponent n in B-L space for the time periods (right) 15--24 February 2024, (left) 15--24 March 2024.




Figure 12: Derived omnidirectional proton flux at 9 MeV, for (top) 15-24 February 2024, (bottom) 15-24 March 2024.

- Flux enhancement of **at least an order of magnitude** following the May 2024 storm.
- Localized "hotspots" near the west coast of South America and central Africa are **fitting artifacts, not physical** - there, the detector look angle is nearly parallel to the field lines, therefore the fitting process leads to overestimations of the flux.

References:
 [1] Desorgher et al.(2013), "The Next Generation Radiation Monitor—NGRM", IEEE/NSS/MIC.
 [2] Sandberg et al.(2022), "First Results and Analysis From ESA Next Generation Radiation Monitor Unit Onboard EDRS-C", IEEE/TNS, 69(7)
 [3] Sandberg (2023), S6-MF/NGRM Level 1 Data Description, Tech. Rep., ESA.
 [4] ESA, Next Generation Radiation Monitor (NGRM), <https://swe.ssa.esa.int/en/ngrm>
 [5] Santin G. et al.(2005), "GRAS: a general-purpose 3-D Modular Simulation tool for space environment effects analysis", IEEE/TNS, 52(6).




Figure 9: Fitted PAD exponent n as a function of time, with error bars, for January - August 2024, grouped by L-shell.

- Heart of inner belt** ($L = 1.4 - 1.8$): $n \approx 5.5 \pm 1$, close to isotropy and largely insensitive to geomagnetic activity.
- Outer edge of inner belt** ($L = 1.8 - 2.2$): $n \approx 11 \pm 0.5$ in quiet time, dropping by ~ 2 units during the May 2024 superstorm.
- Approaching slot region** ($L = 2.2 - 2.6$): drops from $n \approx 10$ to $n \approx 6$ during the storm.
- Post-storm recovery of n is **non-monotonic** at $L = 2.2 - 2.6$, repeatedly interrupted by subsequent injections.

$$\Phi_{\text{omni},X} = \frac{F_X}{E_X(n)}, \text{ for } X \in \{N, S\},$$

where n is the best-fit n.




Figure 13: Equatorial Pitch-Angle Distributions of (left panel) measured and (right panel) retrieved omnidirectional flux for the time periods (top) 15-24 February 2024, (bottom) 15-24 March 2024. Colour indicates L-shell.

- Two humps arising from the northbound/southbound passes become **much more aligned in magnitude** after retrieval, as expected from the method.
- Quiet-time shape: smooth rise from low flux near 10° to a broad peak. Storm-time: around an order-of-magnitude enhancement with two peaks near 25° and 30° .

Conclusions:

- The anisotropy index n is fairly stable over time in the heart of the inner belt, but shows strong dependence on geomagnetic activity near the outer edge.
- Raw directional fluxes underestimate omnidirectional flux by **at least half an order of magnitude** - confirming the need for high-fidelity response modelling for accurate LEO radiation interpretation.
- Future work:** extend to additional NGRM energy channels; apply the same framework to Sentinel-6B and MetOp-SG (also carrying NGRM).

Acknowledgments: E. Christodoulou wishes to express her profound gratitude to the Space Environment and Effects Section of the European Space Research and Technology Centre (ESTEC) for the internship during which this work was conducted.