Functional Equations

June 29, 2006

- 1. Find all functions $f: \mathbb{R} \to \mathbb{R}$ such that $f(x) + f(\frac{1}{1-x}) = x$ for all $x \in \mathbb{R}, x \neq 1$.
- 2. Find all functions $f: \mathbb{R} \to \mathbb{R}$ such that f(x+y) = f(x) + f(y) and f(xy) = f(x)f(y) for all $x, y \in \mathbb{R}$.
- 3. Find all functions $f: \mathbb{R} \to [0, \infty)$ such that for all $x, y \in \mathbb{R}$,

$$f(x^2 + y^2) = f(x^2 - y^2) + f(2xy).$$

- 4. Find all functions $f: \mathbb{R} \to \mathbb{R}$ such that the equality $f(f(x) + y) = f(x^2 y) + 4f(x)y$ holds for all pairs of real numbers x, y.
- 5. Let $f:[0,1]\to\mathbb{R}$ be a function such that
 - (a) f(1) = 1,
 - (b) $f(x) \ge 0$ for all $x \in [0, 1]$,
 - (c) if x, y and x + y all lie in [0, 1], then $f(x + y) \ge f(x) + f(y)$.

Prove that $f(x) \leq 2x$ for all $x \in [0, 1]$.

6. Let $f: \mathbb{R} \to \mathbb{R}$ be a function such that f(0) = 1/2 and

$$f(x+y) = f(x)f(5-y) + f(y)f(5-x)$$

for all $x, y \in \mathbb{R}$. Prove that f is constant.

- 7. Find all functions $f: \mathbb{R} \to \mathbb{R}$ which satisfy $f(xf(x) + f(y)) = f(x)^2 + y$ for all x, y.
- 8. (IMO 1992/2) Find all functions $f: \mathbb{R} \to \mathbb{R}$ such that $f(x^2 + f(y)) = y + f(x)^2$ for all x, y.
- 9. Let n > 2 be an integer and let $f: \mathbb{R}^2 \to \mathbb{R}$ be a function scuh that for any regular n-gon $A_1 A_2 \cdots A_n$,

$$f(A_1) + f(A_2) + \dots + f(A_n) = 0.$$

Prove that f is the zero function.

10. Find all polynomials p(x) such that for all x,

$$(x-16)p(2x) = 16(x-1)p(x).$$

11. Find all functions $f: \mathbb{R} \to [0, \infty)$ such that for all $x, y \in \mathbb{R}$,

$$f(x^2 + y^2) = f(x^2 - y^2) + f(2xy).$$

- 12. Find all functions $f: \mathbb{R} \to \mathbb{R}$ such that the equality $f(f(x) + y) = f(x^2 y) + 4f(x)y$ holds for all pairs of real numbers x, y.
- 13. Let $f: \mathbb{R} \to \mathbb{R}$ be a function such that for all $x, y \in \mathbb{R}$,

$$f(x^3 + y^3) = (x + y)(f(x)^2 - f(x)f(y) + f(y)^2).$$

Prove that for all $x \in \mathbb{R}$, f(1996x) = 1996f(x).

14. Find all functions $f: \mathbb{R} \to \mathbb{R}$ such that for all real numbers x, y, z, t,

$$(f(x) + f(z))(f(y) + f(t)) = f(xy - zt) + f(xt + yz).$$

- 15. Let $f: \mathbb{N} \to \mathbb{N}$ be a function satisfying
 - (a) For every $n \in \mathbb{N}$, f(n+f(n)) = f(n).
 - (b) For some $n_0 \in \mathbb{N}$, $f(n_0) = 1$.

Show that f(n) = 1 for all $n \in \mathbb{N}$.

- 16. Find all functions $f: \mathbb{Z} \to \mathbb{Z}$ which satisfy f(m+f(n)) = f(m) + n for all $m, n \in \mathbb{Z}$.
- 17. Let S denote the set of nonnegative integers. Find all function $f: S \to S$ such that

$$f(m+f(n)) = f(f(m)) + f(n)$$
 for all $m, n \in S$.

18. Let \mathbb{Q}^+ denote the set of positive rational numbers. Find all functions $f: \mathbb{Q}^+ \to \mathbb{Q}^+$ such that for all $x \in \mathbb{Q}^+$

$$f(x+1) = f(x) + 1$$
 and $f(x^2) = f(x)^2$.

- 19. For which integers k does there exist a function $f: \mathbb{N} \to \mathbb{Z}$ such that
 - (a) f(1995) = 1996, and
 - (b) $f(xy) = f(x) + f(y) + kf(\gcd(x, y))$ for all $x, y \in \mathbb{N}$?
- 20. Let S denote the set of nonnegative integers. Find a bijective function $f: S \to S$ such that for all m, $n \in S$,

$$f(3mn + m + n) = 4f(m)f(n) + f(m) + f(n).$$

21. Determine all functions $f: \mathbb{N} \to \mathbb{N}$ such that for all $n \in \mathbb{N}$,

$$f(n) + f(n+1) = f(n+2)f(n+3) - 1996.$$

22. Consider all functions $f: \mathbb{N} \to \mathbb{N}$ such that $f(t^2 f(s)) = s f(t)^2$ for all $s, t \in \mathbb{N}$. Determine the least possible value of f(1998).