Journal of Back and Musculoskeletal Rehabilitation -1 (2019) 1–7 DOI 10.3233/BMR-181303

IOS Press

Association of COPD with osteoporosis in male smokers: A case control study in a tertiary medical college hospital in Bangladesh

Mohammad Zabed Jillul Bari^a, Ismail Patwary^b, Delwar Hussain^c, SAHM Mesbahul Islam^c and Johannes J. Rasker^{d,*}

Abstract.

OBJECTIVES: Chronic obstructive pulmonary disease (COPD) in y increase the risk of osteoporosis and resulting fractures can contribute to disability and mortality of patients. We intended to evaluate the frequency of osteoporosis in male smokers with and without COPD and study whether any correlation existed between osteoporosis and COPD.

MATERIALS AND METHODS: This case-control crucy was carried out in the Department of Medicine, Sylhet M.A.G. Osmani Medical College Hospital, Sylhet, Banglade h 'et veen July 2013 and June 2015. Seventy four male smokers with COPD and 66 age-matched male smokers without COPD were enrolled. All individuals underwent Bone Mass Densitometry (BMD) by Dual-Energy X-Ray Absorptiometry (DEXA).

RESULTS: COPD and non-COPD groups district differ regarding age and smoking pack-years. Osteoporosis at femoral neck (48.6% versus 16.7%; p < 0.001) and lumbal spine (68.9% versus 37.9%; p < 0.01) was significantly higher in COPD compared to controls. Osteopenia did not differ significantly. Patients with COPD were 4.5 times more likely to develop osteoporosis than controls after adjusting age, smoking-back years and BMI (adjusted OR = 4.5; 95% CI = 1.8–11.5).

CONCLUSIONS: Osteoporosis is more frequent in male smokers with COPD compared to smokers without COPD. COPD is a risk factor of osteoporosis independent of age, smoking and BMI.

Keywords: Osteoporosis, male smokers, COPD, developing country, bangladesh

1. Introduction

Chronic obstructive pulmonary disease (COPD) is a major cause of chronic morbidity and mortality world-

wide. The Global Initiative for Chronic Obstructive

*Corresponding author: Johannes J. Rasker, Faculty of Behavioral Management and Social Sciences, Department Psychology Health and Technology, PO Box 217 7500AE Enschede, The Netherlands. Tel.: +31 623 628 967; E-mail: J.J.Rasker@utwente.nl.

Lung Disease (GOLD) has defined COPD as a preventable and treatable disease that is primarily characterized by progressive airflow limitation. This airflow limitation is not fully reversible and is associated with an abnormal inflammatory response of the lung to noxious particles or gases, most often cigarette smoke [1].

In addition to progressive loss of lung function, there is an increasing awareness of the development of extrapulmonary co-morbidities, posing additional problems in the management of COPD [2]. There is a grow-

^aDepartment of Medicine, Sylhet MAG Osmani Medical College, Sylhet, Bangladesh

^bDepartment of Medicine, Sylhet Women's Medical College, Sylhet, Bangladesh 🥏

CDepartment of Respiratory Medicine, Sylhet MAG Osmani Medical College, Sylhet, Bangladesh

 $^{^{}m d}$ University of Twente, Enschede, The Netherlands

16

17

18

20

21

22

23

24

25

26

27

28

30

31

32

33

34

35

36

37

38

40

41

42

43

44

46

47

48

50

51

Variables	COPD ($n = 74$)	Control $(n = 66)$	p value
Age			
41–50 years	6 (8.1)	3 (4.5)	p > 0.05
51–60 years	25 (33.8)	33 (50.0)	
61–70 years	37 (50.0)	26 (39.4)	
71–80 years	6 (8.1)	4 (6.1)	
Mean \pm SD	62.57 ± 8.02	60.65 ± 7.12	$^{\dagger}p = 0.139$
Smoking (pack-years)	37.50 ± 16.77	33.39 ± 13.82	$^{\dagger}p = 0.118$
BMI (Kg/M ²)	18.22 ± 2.58	21.33 ± 4.43	$^{\dagger}p < 0.001$

^{*}Chi-square (Yates' corrected) test and †unpaired t-test were applied to analyse the data. Figure in the parenthesis indicates the corresponding percentage

ing evidence that osteoporosis is one of the systemic effects associated with COPD [3]. A low bone mineral density (BMD), leading to osteoporosis is common in COPD, studies reporting osteoporosis in 24 to 60% of patients with COPD [4–9], and osteopenia in 35% to 72% [10] Osteoporosis and its related fractures are common in patients with COPD and may have significant impact on quality of life and even respiratory function [11].

Several studies showed that the prevalence of osteoporosis is two-to five-fold higher in COPD than in agematched subjects without airflow obstruction [6,7]. Graat-Verboom et al. [12] found an overall prevalence of osteoporosis of 31.7% in COPD versus 5.8% in healthy subjects, p < 0.001). Naghshin et al. [13] found that the frequency of osteoporosis in male smokers with COPD is much higher that in male smoker controls. Indeed, COPD patients ir 12.5 times more likely to develop osteoporosis Turniermore, in a screening tool for males at risk for esteoporosis, the presence of COPD is one of the parameters increasing this risk almost four times [14] COPD patients have a higher prevalence of escoporosis than healthy elderly subjects [15]. However, Karadag et al. [16] and Sim et al. [17] did not find a significant difference in prevalence of osteoporosis between COPD patients and healthy subjects.

This problem has not yet been studied in Bangladesh, a developing country with a high percentage (31.1%) of male smokers [18], and COPD (13.5%) [19]. Moreover, in osteoporosis, subsequent vertebral fractures or hip fractures may further compromise lung function and quality of life and increase the mortality of COPD [20,21]. Hence, this study is undertaken to compare the frequency and association of osteoporosis between male smokers with and without COPD in a developing country, Bangladesh. Knowledge of exact data would make it possible to improve

the quality of life in this risk group, by appropriate preventive strategies that could a oid or reduce the consequences of osteoporosis,

55

57

59

60

63

64

65

68

69

70

72

73

77

80

83

2. Materials and methods

2.1. Study participants

This case-control study was carried out in the Department of Medicine, Sylhet M.A.G. Osmani Medical College Hospital, Sylhet, Bangladesh between July, 2013 and June, 2015. All data were collected prospectively using pre-fabricated data sheets.

All male smokers with at least 10 pack years aged over 40 years and diagnosed as cases of COPD fulfilling the selection criteria were enrolled. Controls were volunteers selected from accompanying persons of the patients or other patients or hospital staff members, who were age matched male smokers without COPD/chronic respiratory diseases (e.g. interstitial lung diseases). Exclusion criteria for patients and controls were: bronchial asthma, chronic heart failure, liver cirrhosis, thyroid dysfunction and rheumatologic disorders, malignancies, chronic renal disease, as well as patients taking systemic corticosteroids for more than 3 months during the last year, bisphosphonates, ergocalciferol, levothyroxin, lithium, calcium and/or vitamin D preparations.

Considering an anticipated effect size of 20% and a number of predictors of 4 with 5% significance level and 80% power, the minimum sample size was 65. In this study 74 COPD patients and 66 non-COPD subjects were enrolled.

2.2. Diagnosis of COPD and osteoporosis

To diagnose COPD, all subjects underwent spirometry (Forced expiratory volume in 1 second [FEV₁],

89

90

91

92

93

97

98

100

101

103

104

105

106

107

108

109

112

113

114

115

116

117

119

120

121

122

123

124

125

126

127

130

131

133

134

135

136

137

138

139

141

142

143

145

146

147

149

150

151

152

153

154

155

156

157

158

160

161

162

163

164

165

Table 2
Lung function measured by spirometry in the two groups

Spirometry	COPD $(n = 74)$	Control $(n = 66)$	P value
FEV ₁ (% predicted)	33.93 ± 12.67	93.12 ± 11.63	*p < 0.001
FVC (% predicted)	52.49 ± 13.87	88.32 ± 11.88	*p < 0.001
FEV ₁ /FVC (%)	46.83 ± 9.24	81.65 ± 5.99	*p < 0.001

^{*}Unpaired t-test was applied to analyse the data. Data were presented as mean \pm standard deviation.

Forced Vital Capacity [FVC], and FEV1/FVC ratio) using RMS Helios 702 spirometer (Recorders and Medicare systems private limited, MEDSPIROR, India). The diagnosis of COPD was made based on clinical history (cough and sputum on most days for at least 3 consecutive months for at least 2 successive years along with breathlessness with history of exposure to risk factors especially tobacco smoking) and confirmed by pulmonary function testing. Staging of COPD was done as per GOLD criteria: stage-I, FEV₁/FVC < 0.70, FEV₁ $\ge 80\%$ predicted; stage-II, $FEV_1/FVC < 0.70$, FEV_1 50–79% predicted; stage-III, $FEV_1/FVC < 0.70$, FEV_1 30–49% predicted and stage-IV, $FEV_1/FVC < 0.70$, $FEV_1 < 30\%$ predicted or FEV₁ < 50% predicted if respiratory failure present [22]. Spirometry was also performed in controls, and all the observations were compared with those of the COPD patients (Table 2).

To measure osteoporosis, bone mineral density of the cases and controls was determined using whole body densitometer, DEXA (Dual Energy X-Ray Absorptiometry) scan (GE Healthcare Lunar progley vance, scanner serial no. PA + 302343, sof ware version \pm ENCORE 2008 version 12.2, Ger var y). BMD, bone mineral content (BMC), and area vere measured at the femoral neck and at the lumber spine (vertebrae L1-L4) [23]. All parameters were expressed in standard globally accepted terms: bMD (g/cm²), BMC (g), and area (cm²). A patient's PMD was given a T-score, which was derived by comparing it to an average score for a healthy 30-year-old male [8,9]. The differences between the "normal young" score and the patient's score were referred to as standard deviation (SD). Tscore values below-2.5 SD were defined as osteoporosis, between-1.0 and-2.5 SD as osteopenia and-1 SD as normal bone density [9]. Osteoporosis score was measured both in cases and controls and compared.

Data regarding the use of corticosteroids orally and as inhalers are collected from history taking and previous clinical records.

2.3. Statistical analyses

The statistical analyses were performed using SPSS (Statistical Package for Social Science) version 21 for

Windows. Descriptive statistics and frequency distributions were generated for the data. A chi-square test was used to show a relationship between categorical variables and a unpaired t-test was used to show a relationship between numerical variables. Statistical significance was set at p < 0.05 for all tests.

2.4. Ethical disclosure

Approval of the stacy protocol was obtained from the Institutional Educal Committee of Sylhet M.A.G Osmani Medical College, Sylhet, Bangladesh and informed written consent was obtained from the patients or attendants (where appropriate) after full explanation of the details of the disease process and purpose of the study

3. Results

The mean age of the COPD group was 62.57 ± 8.02 years and of controls 60.65 ± 7.12 years; (p = 0.139 NS). The mean smoking pack-years also did not differ significantly (p = 0.118 NS) (Table 1). In this study, FEV₁ (% predicted), FVC (% predicted) and FEV₁/FVC (%) were significantly lower in the COPD group than that of the control group (p < 0.001 each) (Table 2). GOLD stage-III was the most frequent stage of COPD and constituted 55.4% of the cases, followed by stage-IV (35.1%) and stage-II (9.5%). In the COPD group oral corticosteroids were used by 9 (12.2%) patients, inhaled steroids by 47 (63.5%) and 18 (24.3%) patients did not use steroids.

Using femoral neck densitometry normal bone mineral density was found significantly fewer in the COPD group than in the controls (8.1% versus 27.3%; p < 0.05). Osteoporosis was found significantly more in the COPD group than in the controls (48.6% versus 16.7%; p < 0.001); whereas osteopenia did not differ significantly between the COPD and control groups (43.2% versus 56.1%; p > 0.05) (Table 3).

With lumber spinal densitometry normal bone mineral density (5.4% versus 18.2%; p < 0.05) and os-

Table 3

Comparison of bone densit	v and octeonoro	cic in emokere wi	th COPD and	l emoking controle
Comparison of bone densit	y and osteoporo	SIS III SIIIOKEIS WI	iii COFD aiic	i sinoking controls

•	•			C
Variables	COPD $(n = 74)$	Control $(n = 66)$	Statistical values	p value
T score				
Femoralneck	-2.31 ± 0.92	-1.65 ± 0.93	t = -4.187	*p < 0.001
Lumbarspine	-3.01 ± 1.26	-1.92 ± 1.13	t = -5.352	*p < 0.001
Osteoporosis				
Femoralneck	36 (48.6)	11 (16.7)	Z = 4.308	$^{\dagger}p < 0.001$
Lumbarspine	51 (68.9%)	25 (37.9)	Z = 3.856	$^{\dagger}p < 0.01$
Osteopenia				
Femoralneck	32 (43.2)	37 (56.1)	Z = -1.537	$^{\dagger}p > 0.05$
Lumbarspine	19 (25.7)	29 (43.9)	Z = -2.291	$^{\dagger} p < 0.05$
Normal				
Femoralneck	6 (8.1)	18 (27.3)	Z = -3.031	$^{\dagger}p < 0.05$
Lumbarspine	4 (5.4)	12 (18.2)	Z = -2.358	$^{\dagger}p < 0.05$

^{*}Unpaired t-test and $^{\dagger}Z$ test for proportion were applied to analyse the data.

Table 4

Risk factors of osteoporosis in patients with COPD by multiple logistic regression analyses

Variables	Crude OR	Adjusted OR	
	(95% CI)	(95% CI)	
Age			
41–50 years	Reference	Reference	
51–60 years	4.67 (0.67-32.38)	0.83 (0.26-4.30)	
61–70 years	5.19 (1.20-22.38)	1.102 (0.22-5.56)	
71–80 years	5.40 (1.26–23.17)	0.15 (0.02-1.36)	
Smoking-pack years			
≤ 20 pack/year	Reference	Reference	
21-40 pack/year	4.25 (0.82-22.13)	0.634 (0.19-2.06)	
41-60 pack/year	2.74 (0.68-11.05)	0.54 (0.13–2.18)	
61–80 pack/year	1.71 (0.37–7.85)	0.312 (0.05-2.02)	
BMI in kg/M ²		XV	
$< 18.5 \text{ Kg/M}^2$	Reference	Reference	
18.5-22.9 Kg/M ²	2.12 (0.96-4.69)	$0.85 (0.14 \pm 0.05)$	
23.0-24.9 Kg/M ²	1.04 (0.30-3.65)	0.15 (0.15–8.95)	
25.0-29.9 Kg/M ²	4.09 (0.83-20.03)	0.73 (0.09-5.85)	
COPD			
No	Reference	Reference	
Yes	4.737 (2.146–10 455)	4.549 (1.793–11.537)	

COPD: Chronic obstructive puln many disease; OR = Odds ratio, CI = Confident Interval.

teopenia (25.7% versus 43.9%; p < 0.05) was found significantly fewer in the COPD group than in controls; while osteoporosis was found significantly more in the COPD group than in controls (68.9% versus 37.9%; p < 0.01) (Table 3).

Multivariate analysis showed that the presence of COPD significantly correlates with osteoporosis (adjusted OR = 4.55; 95% CI = 1.79-11.54), but age, smoking-pack years and BMI did not correlate with osteoporosis (Table 4).

Use of oral corticosteroids (adjusted OR = 1.77; 95% CI = 0.29-10.71) and inhalation steroids (adjusted OR = 1.74; 95% CI = 0.51-5.91) compared to no use of steroids was not associated with osteoporosis.

Association between osteope to is and steroid use in COPD patients (n = 74). Crude OR and adjusted OR for age, smoking pack years and BMI

Steroiduse	C.vde OR (95% CI)	Adjusted OR (95% CI)
Oral	1.96 (0.39-9.93)	1.77 (0.29-10.71)
Inhalat on	1.95 (0.64–5.89)	1.74 (0.51-5.91)
No use of oral	Reference	Reference
or ir nala ional		
rei lus		

OR Odds ratio, CI = Confident interval.

in patients with COPD when adjusted for age, smoking pack year and BMI (Table 5).

4. Discussion

Osteoporosis is a major problem in men with chronic ailments. In men with COPD, osteoporosis may be particularly disabling because vertebral fractures reduce vital capacity, which further compromises ventilation [24]. Evidence suggests that the prevalence of osteoporosis in patients with COPD is high and potentially important [25,26]. When studying the relationship between osteoporosis and COPD, one has to realize that the two diseases have a number of risk factors in common including: smoking, older age, long-term treatment with corticosteroids, and low body mass index.

In the current study the mean BMI was significantly lower in the COPD group than in the smoking controls (p < 0.001) (Table 1), but with multivariate multiple logistic regression analyses BMI did not correlate with osteoporosis (Table 4). In an Indian uncontrolled study among 102 COPD patients, after using multivariate logistic regression analysis, BMI was not found to be a significant risk factor for osteoporosis

205

207

208

211

212

213

215

216

219

220

221

222

223

226

227

228

229

230

233

234

235

237

238

241

242

244

245

248

249

250

251

252

255

256

259

262

263

266

267

270

271

274

275

277

278

281

284

285

286

288

289

291

292

293

295

296

297

299

300

303

304

in COPD patients [26]. A significantly lower BMI in the COPD group than controls was reported in several studies [17,26–28].

In the present study T-scores of the femoral neck were significantly lower in COPD patients than in controls (p < 0.001). These results were in agreement with other studies [13,17]. In contrast, Karadag et al. [16] found T-scores of the femoral neck were significantly lower in COPD patients but difference was not significant (p = 0.9).

This study revealed that osteoporosis in the femoral neck was in 36 (48.6%) COPD compared to 11 (16.7%) in controls (p < 0.001). Several studies were comparable with our findings [17,30-32]. Karadag et al. [16] disagreed with this finding, as they reported that the frequency of osteoporosis was higher in COPD but that the difference was not significant (p = 0.68). They performed their study in COPD patients who were clinically stable and perfectly treated; this choice may have influenced their findings. Graat-Verboom et al. [33] assessed risk factors for developing osteoporosis in clinically stable COPD outpatients at baseline and after 3 years. The prevalence of osteoporosis in COPD patients increased from 47% to 61% in 3 years mostly due to an increase of vertebral fractures. Lower baseline T-scores at the trochanter independently increased the risk for the development of osteoporosis

In our study significantly more osteoporosis vas found in the lumbar spine and femoral neck in the smoking COPD group than in smoking con rols (p < 0.01). This result was comparable with a study by Naghshin et al. [13].

In our study we showed by rachivariate analysis that the presence of COPD significantly correlates with osteoporosis (adjusted OR - 150; 95% CI = 1.79-11.54), but age, smoking pack years and BMI did not predict osteoporosis. This result was supported by the study by Naghshin et al. [13], who stated that the risk of osteoporosis in patients with COPD was almost 12.5 fold compared to control group (OR: 12.46, CI 95% = 3.9–39.85). In Korea osteoporosis was found in 191 (17.7%) of 1,081 COPD patients. In multivariate analyses, older age (odds ratio [OR] = 1.10, P < 0.001) was a risk factor for osteoporosis. Patients with male sex (OR = 0.06, P < 0.001), high house income (OR = 0.75, P = 0.045), and high BMI (OR = 0.74, P <0.001) were less likely to have osteoporosis. In addition, osteoporosis was associated with poor HRQOL $(\beta = -0.21, P = 0.023)$ [34].

In the current study use of oral corticosteroids compared to no use of steroids was not associated with os-

teoporosis in patients with COPD when adjusted for age, smoking pack year and BMI (adjusted OR = 1.77; 95% CI = 0.29–10.71). Inhalation steroids compared to no use of steroids was also not associated with osteoporosis in patients with COPD (adjusted OR = 1.74; 95% CI = 0.51–5.91). Some studies showed that one of the most obvious causes of osteoporosis in COPD patients is treatment with glucocorticoids, both as systemic therapy and as inhaled glucocorticoids [1,9,15,35,36], whereas others reported little or no effect of glucocorticoids on osteoporosis [23,37]. So glucocorticoid use does not fully account for the low bone mineral density (BMD) and high prevalence of osteoporosis in COPD patients [12]. Furthermore the classic explanation of ost opporosis in COPD as a result of accelerated lec'h e in bone mineral density among users of inhaled corticosteroids is not supported by clinical that [35,36,38,39]. Moreover in a randomized controlled trial, Mathioudakis et al. [40] have shown that long-term use of low-dose inhaled corticosteroids protects the COPD patients from developing oscoporosis. This is secondary to the decrease of in lan mation in the lungs, which further decreases the systemic spill-over. The long-term treatment with phaled corticosteroids had also no effect on fracture risk in patients with COPD [41], and at conventional doses [35]. One year of inhaled corticosteroid treatment was shown to exert no effects on bone mineral density [42], while a treatment of 3 years with inhaled triamcinolone was found to reduce bone mineral density [43]. In this study use of oral steroid in COPD patients was less than 3 months and this may explain differences with some of the above mentioned studies.

In an Indian controlled study among 30 COPD patients, the risk of osteoporosis and osteopenia was found to increase with the increase of COPD severity [44]. This fits in with our finding that COPD is an independent risk factor for osteoporosis.

Our study has some limitations. First, this was a single-centre study. A second limitation is that we did not include X-Ray studies of the vertebra. Third, we did not assess the physical activity of COPD and control subjects as this might have influenced osteoporosis; none of our patients or controls were bed-ridden. Fourth, we did not record the life time cumulative doses of inhaled or systemic corticosteroids.

The strength of the study is the fact that it is the first study in Bangladesh and one of the first large controlled studies in a developing country in Asia studying the frequency of osteoporosis in smokers with and without COPD.

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

372

373

374

375

377

378

380

381

382

383

384

385

386

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

404

405

406

407

408

409

410

412

413

414

415

5. Conclusions

306

307

308

310

311

312

315

316

317

318

319

320

323

324

326

327

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

The frequency of osteoporosis is higher in male smokers with COPD compared to those without COPD. Thus COPD appears to be a risk factor for osteoporosis independent of smoking. During rehabilitation of COPD patients back problems due to osteoporotic deformation of the spine and or fractures should be a point of special attentions Physicians should be aware of this complication and BMD should be measured in every male smoker with COPD; prevention of osteoporosis should be part of the medical care for COPD patients.

Acknowledgments

The authors thank Dr. Md. Tabibul Islam, MD (Dermatology and Venereology), Assistant Professor, Department of Dermatology and Venereology, Sylhet MAG Osmani Medical College, Sylhet, Bangladesh for his assistance in the analysis of the data in this study.

Conflict of interest

The authors have no conflict of interest to report

References

- Lehouck A, Boonen S, Decramer M Ja ssens WCO. PD, bone metabolism, and osteoporosis Ch. st. 2011; 139: 648-57
- [2] Sin DD, Anthonisen NR, Soriar o JF, Agusti AG. Mortality in COPD: Role of comorbidities. Fur Respir J. 2006; 28: 1245-57
- [3] Nuti R, Siviero P, Magg S, Guglielmi G, Caffarelli C, Crepaldi G, Gonelli S. Vertebral fractures in patients with chronic obstructive pulmonary disease: The EOLO study. Osteoporos Int. 2009; 20: 989-98.
- [4] Vrieze A, de Greef MHG, Wijkstra PJ, Wempe JB. Low bone mineral density in COPD patients related to worse lung function, low weight and decreased fat-free mass. Osteoporos Int. 2007; 18: 1197-202.
- [5] Biskobing DM. COPD and osteoporosis. Chest. 2002; 121: 609-20.
- [6] Bolton CE, Ionescu AA, Shiels KM, Pettit RJ, Edwards PH, Stone MD, Nixon LS, Evans WD, Griffiths TL, Shale DJ. Associated loss of fat-free mass and bone mineral density in chronic obstructive pulmonary disease. Am J Respir Crit Care Med. 2004; 170: 1286-93.
- [7] Sabit R, Bolton CE, Edwards PH, Pettit RJ, Evans WD, McEniery CM, Wilkinson IB, Cockcroft JR, Shale DJ. Arterial stiffness and osteoporosis in chronic obstructive pulmonary disease. Am J Respir Crit Care Med. 2007; 175: 1259-65.

- [8] Kanis JA, Melton III LJ, Christiansen C, Johnston CC, Khaltaev N. The diagnosis of osteoporosis. J Bone Miner Res. 1994; 9: 1137-41.
- [9] Jorgensen NR, Schwarz P, Holme I, Henriksen BM, Petersen LJ, Backer V. The prevalence of osteoporosis in patients with chronic obstructive pulmonary disease: A cross sectional study. Respir Med. 2007; 101: 177-85.
- [10] Gregg EW, Cauley JA, Seeley DG, Ensred KE, Bauer DC. Physical activity and osteoporotic fracture risk in older women. Ann Intern Med. 1998: 29: 81-88.
- [11] Inoue D, Watanabe R, Okazaki RCO. PD and osteoporosis: Links, risks, and treatment challenges. Int J Chron Obstruct Pulmon Dis. 2016; 11: 637-48.
- [12] Graat-Verboom L, Wouters EFM, Smeenk FWJM, van den Borne BE, Lunde R, Spruit MA. et al. Current status of research on osteoporosis in COPD: A systematic review. Eur Respir J. 2009; 34: 209-18.
- [13] Naghshin R, Javadzadeh A, Mousavi SAJ, Adeli SH, Heydari A. Comparison of the steo rorsis between male smokers with and without chron robstructive pulmonary disease. Tanaffos. 2004; 3: 13-3.
- Tanaffos. 2004; 3: 13-3.

 [14] Shepherd AJ, Cas AP Carlson CA, Ray L. Development and internal validation of the male osteoporosis risk estimation score. Ann Tar. M.ed. 2007; 5: 540-6.
- [15] Kjensli A, F. & JA, Ryg M, Blenk T, Armbrecht G, Diep LM, Ellingsen T High prevalence of vertebral deformities in COPD parints: Relation to disease severity. Eur Respir J. 2009; 33: 10.3-24.
- [16] Yar dag F, Cildag O, Yurekli Y, Curgey O. Should COPD paients be routinely evaluated for bone mineral density? J Bone Miner Metab. 2003; 21: 242-6.
- [17] Sim YS, Lee JH, Ryu YJ, Chun EM, Chang JH. Prevalence and risk factors of osteoporosis in patients with chronic obstructive pulmonary disease. Tuberc Respir Dis. 2009; 66: 186-91.
- [18] Ng M, Freeman MK, Fleming TD, Robinson M, Dwyer-Lindgren L, Thomson B, Wollum A, Sanman E, Wulf S, Lopez AD, Murray CJ, Gakidou E. Smoking prevalence and cigarette consumption in 187 countries, 1980-2012. JAMA. 2014; 311: 183-92.
- [19] Alam DS, Chowdhury MA, Siddiquee AT, Ahmed S, Clemens JD. Prevalence and determinants of chronic obstructive pulmonary disease (COPD) in bangladesh. COPD. 2015; 12: 658-67.
- [20] Duckers JM, Evans BAJ, Fraser WD, Stone MD, Bolton CE, Shale DJ. Low bone mineral density in men with chronic obstructive pulmonary disease. Respir Res. 2011; 12: 101.
- [21] Onescu AA, Schoon E. Osteoporosis in chronic obstructive pulmonary disease. Eur Respir J. 2003; 22(Suppl): 64S-75.
- [22] Ralston SH, McInnes IB. Rheumatology and bone disease. in: Davidson's Principles and Practice of Medicine. Walker BR, Colledge NR, Ralston SH, Penman ID, Editors. 22Nd ed. Edinburgh: Churchill Livingstone, Elsevier. 2014; 1057-136.
- [23] Oh JY, Lee YS, Min KH, Lee SY, Shim JJ, Kang KH, Hur GY. Osteoporosis in patients with asthma-chronic obstructive pulmonary disease overlap syndrome. Tuberc Respir Dis. 2018; 81: 73-79.
- [24] Sin DD, Man JP, Man SF. The risk of osteoporosis in caucasian men and women with obstructive airways disease. Am J Med. 2003; 114: 10-4.
- [25] de Vries F, van Staa TP, Bracke MS, Cooper C, Leufkens HG, Lammers JW. Severity of obstructive airway disease and risk of osteoporotic fracture. Eur Respir J. 2005; 25: 879-84.

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443 444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

485

486

487

488

489

490

491

493

494

495

496

497

M.Z.J. Bari et al. / Association of COPD with osteoporosis in male smokers

- [26] Hattiholi J, Gaude GS. Prevalence and correlates of osteoporosis in chronic obstructive pulmonary disease patients in india. Lung India. 2014; 31: 221-7.
- [27] El-Gazzar AG, Abdalla ME, Almahdy MA. Study of osteoporosis in chronic obstructive pulmonary disease. Egypt J Chest Dis Tuberc. 2013; 62: 91-5.
- [28] Iqbal F, Michaelson J, Thaler L, Rubin J, Roman J, Nanes MS. Declining bone mass in men with chronic pulmonary disease. Chest. 2004; 128: 1616-24.
- [29] Abbasi M, Zohal M, Atapour B, Yazdi Z. Prevalence of osteoporosis and its risk factors in men with COPD in qazvin. Int J Chronic Dis. 2016; 4038530.
- [30] Katsura H, Kida K. A comparison of bone mineral density in elderly female patients with COPD and bronchial asthma. Chest. 2002; 122: 1949-55.
- [31] Lin CH, Chen KH, Chen CM, Chang CH, Huang TJ, Lin CH. Risk factors for osteoporosis in male patients with chronic obstructive pulmonary disease in taiwan. Peer J. 2018 Jan 12; 6: e4232.
- [32] Nayyar N, Sood RG, Sarkar M, Tomar A, Thakur V, Bhoil R. Prevalence of osteoporosis and osteopenia in stable patients of chronic obstructive pulmonary disease in sub-himalayan region of himachal pradesh, india. J Family Med Prim Care. 2017; 6: 595-9.
- [33] Graat-Verboom L, Smeenk FW, van den Borne BE, Spruit MA, Jansen FH, van Enschot JW, Wouters EF. Progression of osteoporosis in patients with COPD: A 3-year follow up study. Respir Med. 2012; 106: 861-70.
- [34] Lee SH, Kwon H-Y. Prevalence of osteoporosis in korean patients with chronic obstructive pulmonary disease and their health-related quality of life according to the korea national health and nutrition examination survey 2008-2011. J Bone Metab. 2017; 24: 241-8.
- [35] Jones A, Fay JK, Burr M, Stone M, Hood K, Roberts G. n-haled corticosteroid effects on bone metabolism in ascittate and mild chronic obstructive pulmonary disease. Co are ne Database Syst Rev. 2002; (1): CD003537.
- [36] Ferguson GT, Calverley PM, Anderson JA, Jenki, s CR, Jones PW, Willits LR, Yates JC, Vestbo J, Celli E Prevalence and progression of osteoporosis in patients with COPD: Results from the towards a revolution in COPL health study. Chest. 2009; 136: 1456-65.

- [37] Medici TC, Grebski E, Hacki M, Ruegsegger P, Maden C, Efthimiou J. Effect of one year treatment with inhaled fluticasone propionate or beclomethasone dipropionate on bone density and bone metabolism: A randomised parallel group study in adult asthmatic subjects. Thorax. 2000; 55: 375-82.
- [38] Graat-Verboom L, Smeenk FW, van den Borne BE, Spruit MA, Donkers-van Rossum AB, Aarts RP, Wouters EF. Risk factors for osteoporosis in caucasian patients with moderate chronic obstructive pulmonary disease: A case control study. Bone. 2012; 50: 1234-9.
- [39] Anar C, YükselYavuz M, Güldaval F, Varol Y, Kalenci D. Assessment of osteoporosis using the FRAX method and the importance of vitamin D levels in COPD patients. Multidiscip Respir Med. 2018 Jan 6; 13: 1.
- [40] Mathioudakis AG, Amanetopoulou SG, Gialmanidis IP, Chatzimavridou-Grigoriadou V, Siasos G, Evangellopulou E, Mathioudakis GA. Impact of long-term treatment with lowdose inhaled corticosteroids on the bone mineral density of chronic obstructive pulmon, ry c sease patients: Aggravating or beneficial? Respirology. 20:3; 18: 147-53.
- [41] Johnell O, Pauwels R, Le Glahl CG, Laitinen LA, Postma DS, Pride NB, Ohlsso SV Lone mineral density in patients with chronic obstructive rulmonary disease treated with budes-onide turbular r. Lur Respir J. 2002; 19: 1058-63.
- [42] Halpern M. Schmier JK, Van Kerkhove MD, Watkins M, Kelberg C. Impact of long-term inhaled corticosteroid therap, on bone mineral density: Results of a meta-analysis. Ann Anngy Asthma Immunol. 2004; 92: 201-67.
- [43] Ca Ilon PD, Connett JE, Wise RA, Tashkin DP, Madhok T, Skeans M, Carpenter PC, Bailey WC, Buist AS, Eichenhorn M, Kanner RE, Weinmann G, et al. Lung health study research group, loss of bone density with inhaled triamcinolone in lung health study II. Am J Respir Crit Care Med. 2004; 170: 1302-9.
- [44] Daga MK, Gupta BK, Mawari G, Hira HS, Sekhri T, Bharali D. Assessment of osteoporosis and osteopenia and its corelation with disease severity in patients of chronic obstructive pulmonary disease with respect to associated risk factors: A case control study in india. J Pulm Respir Med. 2018; 8: 456.