

Nutritional status of children with neurodevelopmental disorders: A cross-sectional study at a tertiary-level hospital in northern Bangladesh.

Rabeya Khatun¹, Md. Kaoser Bin Siddique², Mst. Reshma Khatun³, Maskura Benzir⁴, Md. Rafiqul Islam⁵, Olav Muurlink⁶, Sohel Ahmed⁷

Authors details:

1. Rabeya Khatun, email: rabeya1389@gmail.com, ORCID ID: <https://orcid.org/0000-0001-9715-9621>. Assistant Professor, Department of Pediatrics, TMSS Medical College & Rafatullah Community Hospital, Bogura, Bangladesh.
2. Md. Kaoser Bin Siddique, email: kaoserbd@gmail.com, ORCID ID: <https://orcid.org/0000-0002-0336-2694>. Coordinator of Research & Training, Research, Planning & Development, Health Sector, TMSS, Rangpur Road, Thengamara, Bogura, Bangladesh.
3. Mst. Reshma Khatun, email: reshma@manarat.ac.bd, ORCID ID: <https://orcid.org/0000-0001-6770-3585>. Lecturer, Department of Pharmacy, Manarat International University, Dhaka, Bangladesh.
4. Maskura Benzir, email: mkbenzir@gmail.com, ORCID ID: <https://orcid.org/0000-0001-9533-9821>. Department of Anatomy, TMSS Medical College, Bogura-5800, Bangladesh.
5. Md. Rafiqul Islam, email: drmrslamped@yahoo.com, ORCID ID: <https://orcid.org/0000-0003-3056-7940>. Professor & Head of Department of Pediatrics, TMSS Medical College & Rafatullah Community Hospital, Bogura, Bangladesh.
6. Olav Muurlink, email: o.muurlink@cqu.edu.au, ORCID ID: 0000-0002-8251-9521. Associate Professor, Sustainable Innovation, School of Business and Law, Central Queensland University, Brisbane, Australia.
7. Sohel Ahmed, email: ptsohel@gmail.com, ORCID ID: <https://orcid.org/0000-0001-8706-9841>. Consultant & Founder at Ahmed Physiotherapy & Research Center, Kalabagan, Dhaka-1205, Bangladesh.

Address of Correspondence

Md. Kaoser Bin Siddique, Email: kaoserbd@gmail.com, Orcid ID: <https://orcid.org/0000-0002-0336-2694>. Coordinator, Research & Training, Health Sector, TMSS, Rangpur Road, Thengamara, Bogura, Bangladesh.

Word count: Abstract- 236; Full-text-2983; Tables-06; References-32

Abstract

Background: Malnutrition in children with neurodevelopmental disorders (NDDs) is a significant global public health issue. It has become increasingly clear that proper nutritional assessment with management or advice will support children and families to reach their best potential outcomes.

Objectives: The objective of this study to assess nutritional status and the sociodemographic profile of children with neurodevelopmental disorders in Bangladesh.

Methods: A cross-sectional study was conducted from December to April 2020 among the population of children with NDDs who presented to the pediatric department of the TMSS Medical College and Rafatullah Community Hospital in Bogura during this period. Socio-demographic data along with anthropometric measurements of the children were taken. Assessment of nutritional status were made using metrics such as z-scores for weight-for-age (WAZ), height-for-age (HAZ), and body mass index-for-age (BAZ). Using SPSS Version 25, descriptive statistics (number and percentage) and analytical statistics (*chi*-square and logistic regression) were included.

Results: The result shows that around 58.6% of children displayed malnutrition, with 47.8% showing undernutrition (WHZ / BAZ $-1\text{ SD} \leq -3\text{ SD}$), and 10.8% had overnutrition (BAZ $> 2\text{SD}$). Significant negative associations were found between malnutrition and parental education level, in particular the mother's, father's employment status, urban residency, and monthly family income. Children diagnosed with cerebral palsy exhibited twice the likelihood to be malnourished (AOR 2.39, 95% CI 0.83-6.87). Furthermore, residing in rural regions was associated with an increased risk of experiencing malnutrition, as indicated by an adjusted odds ratio of 1.60 (95% CI 0.12-3.09).

Conclusions: While the results are cross-sectional, over half of children with NDDs were found to be malnourished, suggesting that children with NDD in Bangladesh are vulnerable to developing any form of malnutrition. Therefore, regular assessments and timely nutritional support may improve their situation.

Keywords: Children, nutritional status, malnutrition, neurodevelopmental disorders, Bangladesh

[Text]

Introduction

Malnutrition in children with neurodevelopmental disorders (NDDs) is a global public health issue. Children with NDDs are typically very prone to malnutrition, either under- or over-nutrition (1)(+) both of which are associated with negative health consequences. Adequate nutrition is independently associated with basic health outcomes, including survival and quality of life. (1.2)(+2) According to the Diagnostic and Statistical Manual of Mental Disorders (Fifth Edition) neurodevelopmental disorders are described as "a group of conditions with onset in the developmental period. The disorders typically manifest early in development, often before the child enters [formal schooling], and are characterized by developmental deficits that produce impairments of personal, social, academic, or occupational functioning". (3)(+) Disability is neither exclusively biological nor social in origin or manifestation; rather, it results from the combination of personal and environmental variables. (4)(+) Globally 15.6% of people have a disability, and an alarming 80% of those living with a disability present in low- and middle-income countries (LMICs). (5)(+) The prevalence of disability in Bangladesh is 15.6% (6)(+) but the prevalence of malnutrition among children in this category remains unclear.

Malnutrition remains one of the leading causes of childhood mortality in LMICs, and complications emerge when malnutrition is coupled with a disability. (7)(+) Malnutrition accounts for over half of infant and child deaths worldwide each year, and 3.1 million children with disabilities die from malnutrition. (7)(+) Children with disabilities are three times more likely to be underweight than typically developed children. (8)(+) Even in developing countries, the actual prevalence of malnutrition and its consequences vary significantly. In Kenya, disabled children are 2.2 and 1.7 times more likely to be underweight and stunted, respectively, than their age- and gender-matched healthy peers (9)(+) By contrast, in Bangladesh, a previous study suggested these children are 6.6 times higher odds of being severely underweight and 11.8 times higher odds of being severely stunted. (10)(+) These results suggest that sociocultural differences may be amplifying malnutrition.

An influential framework described by Groce et al. shows that malnutrition may occur in children with NDDs when there is an increased need for nutrients, greater nutrient loss, and decreased nutrient intake. (11)(+) Malnutrition is also associated with a child's growth and development; it reduces peripheral blood circulation, which causes delayed wound healing and increases spasticity and irritability. (12)(+) As for the third aspect of the Groce model, decreased intake, malnutrition resulting from inadequate parental care affects 70% of neurologically challenged children residing in residential settings, placing them at risk for stunted growth, inadequate nutrition, and delayed physical and mental development. (12)(+2)

There have been limited studies examining the interaction between malnutrition and disability in Bangladesh. A study focusing on children with permanent disabilities paired with age-matched controls found that severe acute malnutrition was 4.0 times as likely among children with disability, and the prevalence of underweight and stunting in children with CP was 71.2% and 75.0%, respectively, and in children with epilepsy it was 75.9% and 79.4%, respectively. (10)(+) A study examining malnutrition in Bangladeshi children with cerebral palsy found that more than two-thirds of the sample of 726 were underweight, with moderate to severe undernutrition significantly associated with sociodemographic factors such as age, monthly family income, and physiological factors including gross motor function and neurological type. (13)(+)

Formatted: Indent: First line: 0"

Formatted: Font: Times New Roman, Font color: Auto

Formatted: Pattern: Clear

The current study adds to this small body of work, offering the strength of a hospital-based study with physician-conducted diagnoses, and using exclusion criteria that allows the study to examine the role played by associated factors other than the disability in contributing to disability. The study thus illuminates the role of socioeconomic factors in driving malnutrition in developing world children with a disability, so that a proper nutritional assessment, support management can be designed for children and their caregivers.

Methods

Study Design, ~~and~~ setting and sample selection

This descriptive cross-sectional study was conducted from December 2020 to April 2022 at TMSS Medical College Hospital and Rafatullah Community Hospital, Bogura, at the Department of ~~Paediatrics~~Pediatrics and Child Development Center. The facilities include outdoor/outpatient, clinical wards, and rehabilitation services. Patients are usually assessed in the outdoor/outpatient department, and referred either to the hospital or the rehabilitation centre. In this case, patients who met the sample selection criteria (see below) and were referred to the rehabilitation centre (not the hospital) were taken as the study population. Parents and legal guardians of children with NDDs were interviewed through a pretested structured questionnaire, and anthropometric measurements were taken of each child to assess nutritional status.

Sample selection criteria

- Inclusion Criteria:** Only children with NDD diagnoses independently confirmed by a physician, within the age range of 6 to 144 months, accompanied by parents who were able and willing to participate and provide written consent were included in the study.
- Exclusion criteria:** Children with known chronic ailments affecting nutrition like congenital heart disease, chronic renal disease, human immunodeficiency virus (HIV) infection, and other congenital anomalies that would independently affect their food intake, e.g., cleft lip and cleft palate, were excluded from the study.

According to the Neuro Developmental Disorder (NDD) Protection Trust Act 2013, NDD comprises four conditions: Autism Spectrum Disorder (ASD), Cerebral Palsy (CP), Down Syndrome (DS) and 4. Intellectual Disability (ID). However, this study used six additional conditions under NDD: CP with epilepsy; epilepsy; DS with epilepsy; Attention-deficit/hyperactivity disorder (ADHD); speech delay and developmental delay. Note that none of the sample exhibited DS with epilepsy.

Sample size estimation

The sample size has been calculated by applying the single population proportion formula, under the assumptions of a 95% level of significance, a 5% margin of error, and a response distribution of 50% were used to calculate the sample size. The following calculation, where n = sample size, p = prevalence, and d = margin of error, was used to determine the necessary sample size:

$$N = z^2 p (1-p) / d^2, n = (1.96)^2 \times (0.5) (0.5) / (0.03)^2, = 384$$

However, the challenges posed by the COVID-19 pandemic, including lockdown measures and time limitations, made it arduous to gather data and identify children with neurodevelopmental disorders (NDD) as responders. Consequently, the research team

Formatted: Default Paragraph Font, Font:

Formatted: Default Paragraph Font, Font:

Formatted: pf0

Formatted: Default Paragraph Font, Font:

Formatted: Default Paragraph Font, Font:

Formatted: Default Paragraph Font, Font:

Formatted: Default Paragraph Font, Font:

Formatted: Default Paragraph Font, Font:

Formatted: Default Paragraph Font, Font:

Formatted: Default Paragraph Font, Font:

Formatted: Default Paragraph Font, Font:

Formatted: Default Paragraph Font, Font:

Formatted: Default Paragraph Font, Font:

Formatted: Font: (Default) Times New Roman, 12 pt, Font color: Auto, English (United States), Pattern: Clear

Formatted: Font: Not Bold, Not Italic

had difficulties in locating 384 participants within the designated timeframe of our research. Consequently, there was a need to increase the margin of error from 5% to 8%, necessitating a final sample size of 150. In light of an anticipated 5% rate of incomplete responses, the final minimum needed sample size for this research has been determined to be 157.

Survey tool development

A questionnaire was developed based on the objectives of this study. The Gross Motor Function Classification System (GMFCS) developed by CanChild in Canada was used as a basis for the questionnaire. ~~Questionnaire-~~The questionnaire was translated into Bangla language for ease of comprehension in a population that includes the full spectrum of education levels.

[Survey Questionnaire as supplementary file]

- a. **Parents' socio-demographic variables:** Demographic characteristics such as level of education, occupation, area of residency (rural/urban), family size, and monthly family income were included.
- b. **Demographic variable of children:** Demographic variables including age, sex, and clinical diagnosis, weight, height and BMI were included.

Outcome variables

The criteria developed by WHO (2009) for the classification of acute malnutrition in children was employed. ~~(14)(14)~~ The WHO recommends using the formula $z\text{-score} = (X - m) / SD$, where X is the observed value (height, weight, or BMI), m is the mean value, and SD is the value of the standard deviation of the distribution that corresponds to the reference population.

Weight for height, BMI for age, and height for age comparisons were used to establish wasting, which means acute malnutrition, and stunting, that is, chronic malnutrition, respectively. ~~(15)(15)~~ According to the Z score classification, weight-for-height z-score $WHZ/BAZ \leq -3$ SD indicates acute severe malnutrition; -2.99 to -1 SD Mild to moderate malnutrition and overnutrition were indicated by >2 SD.

Data collection and management

Clearance was obtained from hospital authorities, and informed written consent was obtained from each parent or guardian at the time of the interview. Detailed information was collected using a structured, pretested standard questionnaire by the face-to-face interview method. Children with a diagnosis of NDD made by a medical doctor and confirmed by patient records attending the ~~paediatrics~~paediatrics department and Child Development Center (Shishu Bikash Kendro) were consecutively recruited for the study. ~~Relevant data were collected from hospital records.~~ Children previously identified as having a disability underwent a detailed clinical assessment to confirm the diagnosis. The measures included in the instrument were presented orally to parents or guardians in the Bangla language. The anthropometry measurement were conducted by clinicians or nurses with the weight and height of each study participant measured according to the standard guidelines of the World Health Organisation. ~~(16)(16)~~ Each child's weight was measured to the nearest 0.1kg using a baby scale for children under one year and a digital bathroom scale for the rest of the children. Children who could not stand were measured in their parent's or guardian's arms. The difference between the combined weight of the bearer of the child and child and that of the bearer alone

Formatted: Indent: First line: 0"

Formatted: Font: Not Bold

Formatted: Font: Not Bold

was recorded as the child's weight. Height was measured using a stadiometer for children older than 2 years who were able to stand flat-footed and straight. The recombinant length of the participants less than 2 years of age and those who were 2 years of age or older and were unable to stand upright was measured using an infantometer with a fixed head and a movable footboard. For children with physical deformities who were unable to stand independently, skeletal contracture segmental measurements are used to estimate stature. The segmental measurements are upper-arm length and tibial length. These approaches are considered valid for measuring the stature of children with cerebral palsy. In this study, tibial length was preferred. We utilized the equation for estimating stature in CP by Stevenson et al.. (17)(17)

All z scores were calculated using WHO Anthro and WHO Anthro Plus software. Then nutritional status was defined by WHO standards based on the z scores. Height for age and weight for height was used for stunting and wasting respectively. (16)(16)

Formatted: Indent: First line: 0"

Survey validation and administration

The draft of the survey questionnaire was not subjected to content validation, as we used a WHO-validated tool. (18)(18) The final draft of the questionnaire was piloted initially on five ~~young patients who would have been qualified for the full study, representative populations~~ to check the response time; it took around 20 minutes to complete the questionnaire. Children with neurodevelopmental disorders (NDDs) who were regularly attending the out-patient department (OPD), and in-patient department (IPD) of the paediatrics department, and child development centre of TMSS Medical College and Rafatullah Community Hospital, Bogura. The patient's medical records and clinical examination were used to verify their diagnosis.

Statistical analysis

Data was first logically sequenced using Microsoft Excel for Windows. Data analysis was carried out using the software SPSS 20.0 (IBM Corporation, New York, USA) for Windows. The continuous variables are presented as the mean and a 95% confidential interval. Categorical variables are presented as numbers and percentages. The Pearson chi-square test or Fisher's exact test was used to find out the association between variables. Binary ~~Logistic~~ logistic regression was used to assess the predicted risk factors for nutritional status, using odds ratio (OR) and a 95% confidence interval (CI). The level of significance was set at <0.05.

Results

Demographics of the child with NDDs and their parents

A total of 157 children (57.3% male) with neurodevelopmental disorders took part in this study. The mean age of the participants was 63.68 months, ranging ~~in age~~ from six to 144 months. Cerebral palsy was most common; 27.4% of children were diagnosed with cerebral palsy, with a further 24.8% of children diagnosed as suffering from cerebral palsy with epilepsy. In this study, about 28% of parents had no formal education, 33.4% of parents' occupation was listed as a business, most of the participants ~~about~~ (68.8%) were from rural areas, and 42.7% of participants' gross monthly family income was <10000 Bangladeshi ~~Takataka~~¹. Nearly half of the

¹ By comparison, media reports notes that the Bangladesh labour ministry in 2023 proposed a minimum wage of 12,500 Bangladesh taka per month. This suggests that a significantly higher

participants about 47.1% of participants had 4-5 household members. The details are presented in Table 1.

Nutritional status of the children

Of the total sample, ~~58.6~~47.8% exhibited ~~wasting with~~ 8.9% ~~of the total sample of children exhibited~~ severe wasting ($BAZ \leq -3$ SD) and ~~38.8% of the balance children~~ had mild to moderate wasting ($BAZ -2.99$ to -1 SD). On the other hand, 10.8% of children were overweight ($BAZ > 2$ SD). ~~On the other hand, 41.4% of children were in good health ($BAZ 0.99$ to 1.99 SD).~~ 59.57.3% of children in the sample were stunted and the remaining 42.7% of had age-appropriate height ($HAZ -0.99$ to 1.99 SD). The nutritional status of sample is presented in Table 2 with the disease-specific nutritional status of the child presented in Table 3.

Association between the nutritional status of child and socio-demographic of their parents

There was a significant association between the nutritional status of children and their parents' education level, in particular that of the mother (father, $P = 0.019$ and mother, $p = 0.003$) ~~and as well as with the father's occupation ($p = 0.022$).~~ Findings also reveal a highly significant association between the nutritional status of children and the family's monthly income ($P = < 0.001$). ~~Children from families in rural areas ($p = 0.042$) and children from families with income below 10,000 taka ($p < 0.001$) were significantly more likely to be malnourished.~~ Details are presented in Table 4.

Result of regression analysis

Prior to conducting binary logistic regression, we assessed the presence of multicollinearity among the independent variables through the utilization of the Variance Inflation Factor (VIF). The VIF values of all variables were less than 3 suggesting there was no issue with multicollinearity. In the unadjusted model, Age; Father's Education; Father's Occupation were significantly associated with malnutrition among the NDD children whereas the adjusted model revealed only significant association between child's age and malnutrition. Children diagnosed with Cerebral Palsy exhibit twice as great a likelihood to be susceptible to malnourishment (AOR 2.39, 95% CI 0.83-6.87, $P < 0.001$). Furthermore, residing in rural regions is associated with an increased risk of experiencing malnutrition, as indicated by an adjusted odds ratio of 1.60 (95% CI 0.12-3.09). Children aged in categories greater than 25 months (25-48, 49-72, 73-96, 97-120, and > 120 months) all had a greater chance of being malnourished with odds ratios of (AOR 4.70, 95% CI 1.55 - 14.27, $p = 0.006$), (AOR 4.74, 95% CI 1.37 - 16.32, $p = 0.014$), (AOR 5.61, 95% CI 1.29 - 24.29, $p = 0.014$), (AOR 4.65, 95% CI 0.99 - 21.66, $p = 0.050$), and (AOR 4.20, 95% CI 1.26 - 13.97, $p = 0.019$) respectively than children aged at or below 24 months. Details are presented in Table 5.

[Table 1-5 insert here]

~~proportion of the families in this sample were living below the poverty line than the World Bank estimate at the time (24.3%).~~

Discussion

Malnutrition is considered an important risk factor that can have a negative impact on the prognosis of patients with chronic neurological disease. (19)(19) If malnutrition is not treated in time, it may impair the immune system (20)(20) and aggravate neuromuscular disabilities. (21)(21) Therefore, early identification of malnutrition in children with NDDs is important to ensure timely intervention, and to improve outcomes. Research confirms that children with disabilities are more likely to suffer from malnutrition, much more likely to be underweight or stunted. (8,10,17)(8,10,17) The causal relationship between NDDs and malnutrition is bidirectional. Crepin *et al.*'s research is an example of studies examining the biochemical link between malnutrition and epilepsy, for example, finding that protein-energy malnutrition or a lack of a few specific micronutrients in early childhood can affect the normal functioning of the Central Nervous System (CNS) and may be linked to the development of epilepsy in children. (22)(22)

This present study takes a different approach, aiming to explore malnutrition in a developing world hospital setting, where socioeconomic and cultural factors may significantly determine availability and provision of nutrition, working with a sample where disability in itself did not prevent adequate maintenance of food intake. By focusing on children who did not exhibit symptoms that should impede nourishment, the study allows a focus on the socioeconomic factors that may contribute to malnourishment. The result of this study showed that 58.6% of children in this category exhibited malnutrition, with a ratio of undernutrition to overnutrition of 4.4 to 1. There were significant negative associations between malnutrition and parental education level, in particular mother's education level, fathers' employment, urban residency, and monthly family income.

Collectively, the results suggest that malnourishment is likely mediated and moderated by socioeconomic factors. It is more pronounced for example in families with lower levels of education (particularly on the mother's side) suggesting a role for health awareness education in improving outcomes. It is more pronounced in families living in rural areas, where access to health care is limited. Education and family location are thus likely mediating factors. –The clearest likely moderating factor is household income. As the modal child in this study belonged to rural-based family with a monthly income below 10,000 BDT, access to adequate nourishment is a household problem. A high proportion of the household budget needs to be allocated to food, and food inflation in recent years has had a disproportionate effect on malnutrition in developing countries.(23)(SOURCE). Nevertheless, poverty and lack of access to healthcare and public health advice and even higher quality education resources may have been confounded in the study, as all are associated with rural locations in Bangladesh. —A longitudinal study of greater power may be able to disentangle the causal paths.

The-Interestingly, the current study of 157 children with NDDs showed an uneven gender split: - there was a higher prevalence of NDDs in boys compared to girls, with a ratio of 1.13:1.00, -with gender differences in cerebral palsy (24)(23) being frequently observed. Few studies, particularly in LMICs, explore the role of gender in malnutrition among children with disabilities. (8)(8)

Commented [A1]: Headey D, Ruel M. Food inflation and child undernutrition in low and middle income countries. *Nature Communications*. 2023 Sep 16;14(1):5761.

Commented [A2R1]: Reference Added

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font: Not Bold

In our study, children with NDDs accounted for around 47.8% of those who had muscle wasting, while children with CP (34.7%), CP combined with epilepsy (32%), speech delay (10.7%), and epilepsy (8%) had higher rates of undernutrition. The statistics also revealed that 57.3% of the respondents showed stunting. The children's dietary status and their clinical diagnoses were also statistically significantly associated. f

Formatted: Normal, Space Before: 0 pt

In our study, malnutrition was found to be prominent in children with autism spectrum disorder (ASD) (23.5%) and ADHD, which in Bangladeshi law is a sub-category of ASD (17.6%) common in a range of conditions in which it is not inevitable. Even in the case of cerebral palsy (where in excess of 60% of the current sample exhibited malnutrition), clinicians now accept that under- or over-nutrition can be addressed. It was discovered that children with ASD and attention deficit hyperactivity disorder (ADHD) often expressed concern about their dietary habits. Children with nervous system or brain problems such as those with cerebral palsy, Prader-Willi syndrome, Down syndrome, spina bifida, Rett syndrome, cognitive disabilities, developmental delays, ASD, and ADHD, experience obesity and overweight more severely. (25)(24) The prevalence of obesity varies between 9.7% (26-28)(25-27) and 35.7% (25)(24) although in the current sample was much lower. Since it restricts their capacity to move independently, impairs their ability to breathe, and increases their risk of developing cardiovascular issues, obesity has always been a serious concern for children with NDD. For example, overweight and obesity problems obesity in children with Down syndrome are is harmful because it predicts they increase the risk of adult obesity and the subsequent noncommunicable diseases consequences like diabetes and hypertension. (20,25)(20,24) Usually, individuals with ASD suffer more from feeding problems and obesity compared to their peers as they have a specific preference towards sweet, cereal-based products and fats and oil, milk and milk-based food and high protein and high fat foods, etc. They also have a routine in their diet, which likely expands the tendency to have an unbalanced diet. (29)(28)

Formatted: Indent: First line: 0"

As most of the modal child in this study children belonged to rural based family with an income areas (68.8%) and their family income (42.7%) was below 10,000 BDT, this may reflect that the people in rural remote sites are poor poverty and lack of access to healthcare and public health advice may have been confounded in the study. They have no proper knowledge or awareness about feeding their children, who lived in rural (77.3%) areas compared to children living in urban areas. There was a similar study by Jahan et al., where underweight and/or stunting among children with disabilities were significantly associated with parental educational qualification, socioeconomic status, and mainstream school attendance. (10)(10) Children with NDD may later have impaired motor function, bone health, social involvement, and total healthcare usage as a result of their low nutritional condition and poor development. (30)(29)

Strength and limitations

While relatively novel in the developing world context in that it used hospital clinicians and a hospital setting to conduct the key components of this study, the sample size was relatively small, and cannot be extrapolated onto a national scale. The study is also correlational, hence it cannot convincingly whether the child's nourishment or disability status partly caused aspects of the family's socioeconomic status; in other words, the burden of looking after a disabled child has been associated with impact on the family budget in previous studies. (30)(29) [SOURCE] There were also limitations in the measurement approach; simpler measures were chosen for analysing malnutrition

Formatted: Indent: First line: 0"

and stunting, measures that are common to such studies but still not gold standard approaches. We incorporated BMI for age (BAZ) for 0–19 years to assess wasting or undernutrition in our research sample because WHO AnthroPlus only calculates WAZ for children aged 0-121 months; we were unable to determine the existence of general malnutrition (WAZ SD). Although peripheral blood film (PBF) and a complete blood count (CBC) may be used to identify iron deficiency, these invasive techniques are not usually used for this condition. (31)(30)

This study does, however, offer plausible evidence that socioeconomic status does impact on malnourishment outcomes in disabled children in Bangladesh. By only including children who have no physiological reason to be malnourished we were able to isolate how socioeconomic factors impact on malnourishment. It is possible that dietary factors and oromotor dysfunction are responsible for causing malnutrition in these children however.

Conclusion: The study results provide evidence of the impact of socio-demographic factors on the malnourishment of disabled Bangladeshi children. Parental education level, father's occupation, and monthly family income all appear to be significantly related to the nutritional status of children with NDD, and cases of malnourishment appear more pronounced in children from rural areas.

Recommendations

1. Children with NDD should be regularly screened for malnutrition in Bangladesh. This should happen more frequently in cases where physical factors impede feeding, or promote choking, regurgitation, and spilling of food.
2. Mass screening programmes should particularly in remote regions should form the basis of improving the nutritional status of these children. Expanding access to affordable health care for families with disabled children and offering nutritional advice and support will help mitigate malnourishment prevalence.
3. Public health education programs focused on parents, offering advice on care and nutrition for their children may exert a positive influence on disabled children's health and survival in Bangladesh.

Abbreviations

ASD	Autism Spectrum Disorder
ADHD	Attention-Deficit / Hyperactivity Disorder
BAZ	BMI for Age Z Score
BMI	Body Mass Index
CBC	Complete Blood Count
CNS	Central Nervous System
CP	Cerebral Palsy
HIV	Human Immunodeficiency Virus
IPD	Indoor Patient Department
MAM	Moderate Acute Malnutrition
NDD	Neuro Developmental Disorder
OPD	Outdoor Patient Department
PBF	Peripheral Blood Film
SAM	Severe Acute Malnutrition
TMSS	Thengamara Mohila Sabuj Sangha
WAZ	Weight for Age Z-Score
WHZ	Weight-for-Height Z-Score
WHO	World Health Organization

Declarations:

Author Contributions: Conceptualization, R.K.; methodology, R.K.; M.K.B.S.; and S.A software, R.K.; and M.R.K, validation and investigation, R.K.; M.R.K.; M.K.B.S.; M.B; & M.R.I., formal analysis, R.K.; MKBS.; and M.R.K, writing—original draft preparation, RK, M.R.K.; M.B and MKBS.; writing—review and editing, O.M revised whole manuscript and finalized; S.A., M.K.B.S. and R.K; All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Ethics approval and consent to participate

This study strictly followed the ethical guidelines by Bangladesh medical research council 2014 and the Helsinki Declaration 2013 (revised) involving human participants in research. The design of the study, data collection procedure, presentation of the data and citation comply with the standard Committee on Publication Ethics (COPE) guideline.

Institutional review board statement: Ethical clearance has been obtained from the institutional ethical board of Pundra University of Bangladesh (Reference PUB/PH/2021-49).

Informed consent statement: Written consent was obtained from parents or legal guardians for participants under 16 years of age. The parents or legal guardians were

informed about the nature and purpose of the study before participating, and after that, written informed consent was obtained from them.

Consent to publication

Not applicable

Availability of data and materials: All the raw de-identified data will be made available upon reasonable request from the corresponding author.

Competing interests: The authors declare no competing interests.

Acknowledgements

The authors would like to thank all the faculties of Public Health Department of Pundra University of Science Technology and all the staff of Pediatric Department of TMSS Medical College for their support in conducting this study. The authors would also like to acknowledge Rtn. Dr. Md. Matiur Rahman, Prof. Dr. Raghieb Ahsan, Dr. Mohammad Mahbub Ur Rahim, Md. Rahidul Islam and colleagues of TMC & RCH and Grand Health Sector, TMSS, for their cordial support. The authors are grateful to all the participants who provided valuable data for this study.

References

1. Canadian Paediatric Society NCommittee. Undernutrition in children with a neurodevelopmental disability. Can Med Assoc J. 1994;151:753-9.
2. Samson-Fang L, Fung E, Stallings VA, Conaway M, Worley G, Rosenbaum P, et al. Relationship of nutritional status to health and societal participation in children with cerebral palsy. J Pediatr. 2002 Nov;141(5):637-43.
3. American Psychiatric Association D 5 TForce. Diagnostic and statistical manual of mental disorders: DSM-5™ (5th ed.). 5th ed. American Psychiatric Publishing, Inc.; 2013.
4. Wong ME. United Nations Convention on the Rights of Persons with Disabilities in Singapore: Considering Some Post-Ratification Implications. In Special Needs In Singapore: Trends And Issues. 2021;11-30.
5. World Health Organization. World Report on Disability. 2011;
6. Bangladesh Bureau of Statistics (BBS). National Survey on Persons with Disabilities (NSPD) conducted in 2019. June 2022. 2022 Jun [cited 2024 Jan 9]; Available from: https://bbs.portal.gov.bd/sites/default/files/files/bbs.portal.gov.bd/page/b343a8b4_956b_45ca_872f_4cf9b2f1a6e0/2022-06-13-15-24-ca6f018ab83c88a4db8ff51386439794.pdf
7. Groce N CEKMCL. Stronger together : nutrition-disability links and synergies|Briefing note. Nutrition working group. global partnership for children with disabilities New York: Unicef [Internet]. 2013 [cited 2024 Jan 9]; Available from: [https://sites.unicef.org/disabilities/files/Stronger-Together Nutritio...](https://sites.unicef.org/disabilities/files/Stronger-Together_Nutritio...)
8. Hume-Nixon M, Kuper H. The association between malnutrition and childhood disability in low- and middle- income countries: systematic

- [review and meta-analysis of observational studies. Tropical Medicine & International Health. 2018 Nov 10;23\(11\):1158–75.](#)
9. [Kuper H, Nyapera V, Evans J, Munyendo D, Zuurmond M, Frison S, et al. Malnutrition and Childhood Disability in Turkana, Kenya: Results from a Case-Control Study. PLoS One. 2015 Dec 21;10\(12\):e0144926.](#)
 10. [Jahan I, Karim T, Al Imam MH, Das MC, Ali KM, Muhit M, et al. Childhood Disability and Nutrition: Findings from a Population-Based Case Control Study in Rural Bangladesh. Nutrients. 2019 Nov 11;11\(11\):2728.](#)
 11. [Groce N, Challenger E, Berman-Bieler R, Farkas A, Yilmaz N, Schultink W, et al. Malnutrition and disability: unexplored opportunities for collaboration. Paediatr Int Child Health. 2014 Nov 13;34\(4\):308–14.](#)
 12. [Pancheva R, Toneva A, Nikolova S, Huysentruyt K, Dimitrov L, Yoncheva N, et al. Malnutrition in Institutionalized and Non-institutionalized Neurologically Impaired Children in Bulgaria. Ann Nutr Metab. 2023 Nov 9;](#)
 13. [Jahan I, Muhit M, Karim T, Smithers-Sheedy H, Novak I, Jones C, et al. What makes children with cerebral palsy vulnerable to malnutrition? Findings from the Bangladesh cerebral palsy register \(BCPR\). Disabil Rehabil. 2019 Sep 11;41\(19\):2247–54.](#)
 14. [World Health Organization. WHO Child Growth Standards and the Identification of Severe Acute Malnutrition in Infants and Children: A Joint Statement by the World Health Organization and the United Nations Children’s Fund. 2009.](#)
 15. [Schaffrath Rosario A, Schienkiewitz A, Neuhauser H. German height references for children aged 0 to under 18 years compared to WHO and CDC growth charts. Ann Hum Biol. 2011 Mar 13;38\(2\):121–30.](#)
 16. [Stevenson RD. Use of segmental measures to estimate stature in children with cerebral palsy. Arch Pediatr Adolesc Med. 1995 Jun;149\(6\):658–62.](#)

17. [Mwangome MK, Berkley JA. The reliability of weight-for-length/height <scp>Z</scp> scores in children. Matern Child Nutr. 2014 Oct;10\(4\):474–80.](#)
18. [Chandra RK, Kumari S. Nutrition and Immunity: An Overview. J Nutr. 1994 Aug;124:1433S-1435S.](#)
19. [Caglià P, Luca S, Gandolfo L, Amodeo C. Enteral nutrition in patients with chronic neurological diseases. Minerva Gastroenterol Dietol. 2000 Dec;46\(4\):199–206.](#)
20. [ENGSDNER G, HABTE D, SJÖGREN I, VAHLQUIST B. BRAIN GROWTH IN CHILDREN WITH KWASHIORKOR A Study using Head Circumference Measurement, Transillumination und Ultrasonic Echo Ventriculography. Acta Paediatr. 1974 Sep 21;63\(5\):687–94.](#)
21. [Mutch L, Alberman E, Hagberg B, Kodama K, Perat MV. Cerebral Palsy Epidemiology: Where are We Now and Where are We Going? Dev Med Child Neurol. 1992 Jun 12;34\(6\):547–51.](#)
22. [Reinehr T, Dobe M, Winkel K, Schaefer A, Hoffmann D. Obesity in Disabled Children and Adolescents. Dtsch Arztebl Int. 2010 Apr 16;](#)
23. [Headey D, Ruel M. Food inflation and child undernutrition in low and middle income countries. Nat Commun. 2023 Sep 16;14\(1\):5761.](#)
24. [Romeo DM, Venezia I, Pede E, Brogna C. Cerebral palsy and sex differences in children: A narrative review of the literature. J Neurosci Res. 2023 May 9;101\(5\):783–95.](#)
25. [Holenweg-Gross C, Newman CJ, Faouzi M, Poirot-Hodgkinson I, Bérard C, Roulet-Perez E. Undernutrition in children with profound intellectual and multiple disabilities \(<scp>PIMD</scp> \): its prevalence and influence on quality of life. Child Care Health Dev. 2014 Jul;40\(4\):525–32.](#)
26. [Jahan I, Sultana R, Afroz M, Muhit M, Badawi N, Khandaker G. Dietary Intake, Feeding Pattern, and Nutritional Status of Children with Cerebral Palsy in Rural Bangladesh. Nutrients. 2023 Sep 29;15\(19\):4209.](#)

27. [Manikandan B, Gloria J. K, Samuel R, Russell PSS. Feeding Difficulties Among Children With Special Needs: A Cross-Sectional Study From India. OTJR: Occupational Therapy Journal of Research. 2023 Oct 14;43\(4\):592-9.](#)
28. [Adams MS, Khan NZ, Begum SA, Wirz SL, Hesketh T, Pring TR. Feeding difficulties in children with cerebral palsy: low-cost caregiver training in Dhaka, Bangladesh. Child Care Health Dev. 2012 Nov 15;38\(6\):878-88.](#)
29. [Sánchez-Lastres J, Eirís-Puñal J, Otero-Cepeda JL, Pavón-Belinchón P, Castro-Gago M. Nutritional status of mentally retarded children in north-west Spain. I. Anthropometric indicators. Acta Paediatr. 2003 Jun;92\(6\):747-53.](#)
30. [Rothwell DW, Lach LM, Kohen DE, Findlay LC, Arim RG. Income trajectories of families raising a child with a neurodisability. Disabil Rehabil. 2022 May 8;44\(10\):1923-32.](#)
31. [Lark RK, Williams CL, Stadler D, Simpson SL, Henderson RC, Samson-Fang L, et al. Serum prealbumin and albumin concentrations do not reflect nutritional state in children with cerebral palsy. J Pediatr. 2005 Nov;147\(5\):695-7.](#)
- ~~1. Canadian Paediatric Society NCommittee. Undernutrition in children with a neurodevelopmental disability. Can Med Assoc J. 1994;151:753-9.~~
- ~~2. Samson-Fang L, Fung E, Stallings VA, Conaway M, Worley G, Rosenbaum P, et al. Relationship of nutritional status to health and societal participation in children with cerebral palsy. J Pediatr. 2002 Nov;141(5):637-43.~~
- ~~3. American Psychiatric Association D 5 TForce. Diagnostic and statistical manual of mental disorders: DSM 5™ (5th ed.). 5th ed. American Psychiatric Publishing, Inc.; 2013.~~
- ~~4. Wong ME. United Nations Convention on the Rights of Persons with Disabilities in Singapore: Considering Some Post-Ratification Implications. In Special Needs In Singapore: Trends And Issues. 2021;11-30.~~
- ~~5. World Health Organization. World Report on Disability. 2011;~~

6. ~~Bangladesh Bureau of Statistics (BBS). National Survey on Persons with Disabilities (NSPD) conducted in 2019. June 2022. 2022 Jun [cited 2024 Jan 9]; Available from: https://bbs.portal.gov.bd/sites/default/files/files/bbs.portal.gov.bd/page/b343a8b4_956b_45ea_872f_4cf9b2f1a6e0/2022-06-13-15-24-ca6f018ab83c88a4db8ff51386439794.pdf~~
7. ~~Groce N, CEKMCL. Stronger together: nutrition disability links and synergies|Briefing note. Nutrition working group, global partnership for children with disabilities New York: Unicef [Internet]. 2013 [cited 2024 Jan 9]; Available from: https://sites.unicef.org/disabilities/files/Stronger-Together_Nutritio...~~
8. ~~Hume Nixon M, Kuper H. The association between malnutrition and childhood disability in low and middle income countries: systematic review and meta-analysis of observational studies. Tropical Medicine & International Health. 2018 Nov 10;23(11):1158–75.~~
9. ~~Kuper H, Nyapera V, Evans J, Munyendo D, Zuurmond M, Frison S, et al. Malnutrition and Childhood Disability in Turkana, Kenya: Results from a Case Control Study. PLoS One. 2015 Dec 21;10(12):e0144926.~~
10. ~~Jahan I, Karim T, Al Imam MH, Das MC, Ali KM, Muhit M, et al. Childhood Disability and Nutrition: Findings from a Population-Based Case Control Study in Rural Bangladesh. Nutrients. 2019 Nov 11;11(11):2728.~~
11. ~~Groce N, Challenger E, Berman-Bieler R, Farkas A, Yilmaz N, Schultink W, et al. Malnutrition and disability: unexplored opportunities for collaboration. Paediatr Int Child Health. 2014 Nov 13;34(4):308–14.~~
12. ~~Pancheva R, Toneva A, Nikolova S, Huysentruyt K, Dimitrov L, Yoncheva N, et al. Malnutrition in Institutionalized and Non institutionalized Neurologically Impaired Children in Bulgaria. Ann Nutr Metab. 2023 Nov 9;~~
13. ~~Jahan I, Muhit M, Karim T, Smithers-Sheedy H, Novak I, Jones C, et al. What makes children with cerebral palsy vulnerable to malnutrition? Findings from the Bangladesh cerebral palsy register (BCPR). Disabil Rehabil. 2019 Sep 11;41(19):2247–54.~~

14. World Health Organization. WHO Child Growth Standards and the Identification of Severe Acute Malnutrition in Infants and Children: A Joint Statement by the World Health Organization and the United Nations Children's Fund. 2009.
15. Schaffrath Rosario A, Schienkiewitz A, Neuhauser H. German height references for children aged 0 to under 18 years compared to WHO and CDC growth charts. *Ann Hum Biol.* 2011 Mar 13;38(2):121–30.
16. Stevenson RD. Use of segmental measures to estimate stature in children with cerebral palsy. *Arch Pediatr Adolesc Med.* 1995 Jun;149(6):658–62.
17. Mwangome MK, Berkley JA. The reliability of weight-for-length/height <sep>Z</sep> scores in children. *Matern Child Nutr.* 2014 Oct;10(4):474–80.
18. Chandra RK, Kumari S. Nutrition and Immunity: An Overview. *J Nutr.* 1994 Aug;124:1433S–1435S.
19. Caglià P, Luca S, Gandolfo L, Amodeo C. Enteral nutrition in patients with chronic neurological diseases. *Minerva Gastroenterol Dietol.* 2000 Dec;46(4):199–206.
20. ENGSNER G, HABTE D, SJÖGREN I, VAHLQUIST B. BRAIN GROWTH IN CHILDREN WITH KWASHIORKOR *A Study using Head Circumference Measurement, Transillumination und Ultrasonic Echo Ventriculography.* *Acta Paediatr.* 1974 Sep 21;63(5):687–94.
21. Mutch L, Alberman E, Hagberg B, Kodama K, Perat MV. Cerebral Palsy Epidemiology: Where are We Now and Where are We Going? *Dev Med Child Neurol.* 1992 Jun 12;34(6):547–51.
22. Reinehr T, Dobe M, Winkel K, Schaefer A, Hoffmann D. Obesity in Disabled Children and Adolescents. *Dtsch Arztebl Int.* 2010 Apr 16;
23. Romeo DM, Venezia I, Pede E, Brogna C. Cerebral palsy and sex differences in children: A narrative review of the literature. *J Neurosci Res.* 2023 May 9;101(5):783–95.

24. — Holenweg-Gross C, Newman CJ, Faouzi M, Poirot-Hodgkinson I, Bérard C, Roulet Perez E. Undernutrition in children with profound intellectual and multiple disabilities (<scp>PIMD</scp>): its prevalence and influence on quality of life. *Child Care Health Dev.* 2014 Jul;40(4):525–32.
25. — Jahan I, Sultana R, Afroz M, Muhit M, Badawi N, Khandaker G. Dietary Intake, Feeding Pattern, and Nutritional Status of Children with Cerebral Palsy in Rural Bangladesh. *Nutrients.* 2023 Sep 29;15(19):4209.
26. — Manikandan B, Gloria J. K, Samuel R, Russell PSS. Feeding Difficulties Among Children With Special Needs: A Cross-Sectional Study From India. *OTJR: Occupational Therapy Journal of Research.* 2023 Oct 14;43(4):592–9.
27. — Adams MS, Khan NZ, Begum SA, Wirz SL, Hesketh T, Pring TR. Feeding difficulties in children with cerebral palsy: low cost caregiver training in Dhaka, Bangladesh. *Child Care Health Dev.* 2012 Nov 15;38(6):878–88.
28. — Sánchez Lastres J, Eirís Puñal J, Otero Cepeda JL, Pavón Belinchón P, Castro Gago M. Nutritional status of mentally retarded children in north-west Spain. I. Anthropometric indicators. *Acta Paediatr.* 2003 Jun;92(6):747–53.
29. — Rothwell DW, Lach LM, Kohen DE, Findlay LC, Arim RG. Income trajectories of families raising a child with a neurodisability. *Disabil Rehabil.* 2022 May 8;44(10):1923–32.
30. — Lark RK, Williams CL, Stadler D, Simpson SL, Henderson RC, Samson-Fang L, et al. Serum prealbumin and albumin concentrations do not reflect nutritional state in children with cerebral palsy. *J Pediatr.* 2005 Nov;147(5):695–7.

-

Table 1: Bio-demographic status of children with NDD and their parents (n=157)

Variable	Number	Percentage
Age in months		
<24	33	21
25-48	42	26.8
49-72	27	17.2
73-96	15	9.6
97-120	24	8.9
>120	26	16.6
Total	157	100
Mean age	63.68	
Range	6-144	
SD	42.95	
Sex		
Male	90	57.3
Female	67	42.7
Total	157	100
Clinical diagnosis		
Cerebral palsy	43	27.4
Cerebral palsy with Epilepsy epilepsy	39	24.8
Epilepsy	10	6.4
Down syndrome	11	07
Autism spectrum disorder	21	13.4
ADHD	05	3.2
Speech delay	15	9.6
Developmental Delay delay	10	6.4
Intellectual Disability disability	03	1.9
Total	157	100
Residence		
Urban	37	23.6
Semi-urban	12	7.6
Rural	108	68.8
Total	157	100
Monthly Family income (BDT/USD)		
<10000 (<101 USD)	67	42.7
11000-25000 (102-240 USD)	46	29.3
26000-35000 (241-342USD)	16	10.2
≥35000 (≥342USD)	28	17.8
Total	157	100
Family type		
Nuclear	109	69.4
Extended	48	30.6
Household size		
Small Family 1-3 Members	28	17.8
Medium Family 4-5 Members	78	47.1
Large Family ≥6 Members	55	35
Total	157	100
	Father	Mother

	Number	Percentage	Number	Percentage
Education				
No formal Education	31	19.7	13	8.3
Up to Primary	26	16.6	24	15.3
Below SSC	30	19.1	48	30.6
SSC	22	14	32	20.4
HSC	11	7	14	8.9
Bachelors Honors	16	10.2	12	7.6
Masters and above	21	13.4	14	8.9
Total	157	100	157	100
Occupation				
Govt. Service	7	4.5	7	4.5
Other Service	20	12.7	5	3.2
Business	51	32.5	3	1.9
Day labor	14	8.9	01	0.6
Farmer	35	22.3	-	-
Housewife	-	-	135	86
Unemployed	01	0.6	-	-
Others	29	18.5	06	3.8
Total	157	100	157	100

Table 2: Nutritional status of the study population (n=157)

Nutritional indices reported as Z score		
Categories	Weight for height (WHZ) / BMI for age Z score (BAZ)	Children's height/length for age Z score (HAZ)
Z Score	f (%)	f (%)
≤-3 SD	14 (8.9)	29 (18.5)
-2.99 to -1 SD	61(38.8)	61 (38.8)
-0.99 to 1.99 SD	65(41.4)	67(42.7)
>2 SD	17(10.8)	-
Total	157(100%)	157(100%)

Table 3: Nutritional status and children's clinical diagnosis (n=157)

variables	Nutritional status		x2	P value	Nutritional status	x2	P value
	Wasting	Over weight			Stunting		
▲Cerebral palsy (CP)	26 (34.7%)	02 (11.8%)			29 (32.2%)		
▲Cerebral palsy with Epilepsy	24 (32%)	02 (11.8%)			28 (31.1%)		
▲Epilepsy	06 (8%)	0			06 (6.7%)		
▲Down syndrome	02 (2.7%)	02 (11.8%)		0.001	09 (10%)		0.001
▲Autism Spectrum Disorder (ASD)	04 (5.3%)	04 (23.5%)	38.052		06 (6.7%)	25.533	
▲ADHD	01 (1.3%)	03 (17.6%)			0		
▲Speech delay	08 (10.7%)	01 (5.9%)			05 (5.6%)		
▲Developmental Delay	02 (2.7%)	02 (11.8%)			05 (5.6%)		
▲Intellectual Disability	02 (2.7%)	01 (5.9%)			02 (2.2%)		
Total	75 (100%)	17 (100%)			90 (100%)		

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

[illegible]2

Family type	Extended	18 (24%)	08 (47.1%)		
Household size	Small Family 1-3 Members	12 (16%)	03 (17.6%)	3.7768	0.438
	Medium Family 4-5 Members	41 (54.7%)	06 (35.3%)		
	Large Family ≥6 Members	22 (29.3%)	08 (47.1%)		
	Total	75 (100%)	17 (100%)		

- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt
- Formatted: Font: 10 pt

**Table 5: Factors associated with malnutrition among the 6 to 144-month-olds
NDD children (Binary logistic regression analysis)**

Characteristics	Unadjusted OR (95% CI)	p- value s	Adjusted OR* (95% CI)	p- value s
Age (month)				
<24 (ref)	1			
25-48	3.15 (1.22-8.14)	0.018 *	4.70 (1.55-14.27)	0.006 *
49-72	2.97 (1.03-8.54)	0.043 *	4.74 (1.37-16.32)	0.014 *
73-96	4.81 (1.25-18.50)	0.022 *	5.61 (1.29-24.29)	0.021 *
97-120	2.33 (0.65-8.34)	0.192	4.65 (0.99-21.66)	0.050 *
>121	3.30 (1.13-9.69)	0.029 *	4.20 (1.26-13.97)	0.019 *
Father education				
Masters and above (ref)	1			
Honors Bachelors(Graduation)	1.71 (0.46-6.37)	0.421	2.00 (0.42-9.50)	0.383
HSC	1.11 (0.25- 4.82)	0.888	0.61 (0.09-4.18)	0.620
SSC	4.53 (1.21- 16.96)	0.025 *	3.32 (0.51-21.64)	0.210
Below SSC	1.33 (0.43- 4.09)	0.615	0.73 (0.13-3.97)	0.713
Up to Primary	2.52 (0.77- 8.22)	0.126	1.26 (0.20-7.92)	0.802
No formal Education	2.42 (0.78- 7.54)	0.126	1.53 (0.26-8.95)	0.637
Father occupation				
Service (ref)	1			
Business	2.91 (1.11- 7.62)	0.030 *	2.34 (0.67-8.12)	0.181
Day labor	2.01 (0.77- 5.20)	0.151	0.91 (0.19-4.37)	0.910
Others	2.38 (0.81-6.96)	0.113	1.18 (0.25-5.57)	0.834
Household Income				
>35000 Taka (ref)	1			
26000-35000 Taka	0.87 (0.25- 2.96)	0.819	0.72 (0.16-3.27)	0.716
11000-25000 Taka	1.03 (0.40- 2.65)	0.948	0.63 (0.13-3.07)	0.570
100-10000Taka (1-101USD)	1.66 (0.67- 4.07)	0.269	0.90 (0.16-5.12)	0.910
Place of residence				
Urban (ref)	1			
Semi-urban	1.05 (0.29- 3.88)	0.93	0.63 (0.12-3.30)	0.584

Rural	1.79 (0.84- 3.81)	0.13	1.60 (0.46-5.52)	0.455
Clinical diagnosis				
Autism Spectrum Disorders (ref)	1			
Down Syndrome	0.54 (0.14- 2.15)	0.385	0.61 (0.12-3.09)	0.555
Cerebral Palsy	1.68 (0.81- 3.48)	0.164	2.39 (0.83-6.87)	0.106

*Sig (p<0.05)