Nanoparticle trafficking in the tumor immune microenvironment is driven by coupling of cell-intrinsic and extrinsic immune mechanisms

Stephanie Monson, Sahil Kulkarni, Kshitiz Parihar, Jake Brenner, Jacob Myerson, Ravi Radhakrishnan
The University of Pennsylvania

Contact + more info!

Motivation

Two Clinical Challenges

THERAPEUTICS

Only ~1% of engineered therapeutic nanoparticles (NP) reach their target in solid tumors, in part due to immune recognition and clearance.

DIAGNOSTICS

Tumors exploit extracellular vesicles (EVs) to modulate the tumor immune microenvironment (TIME), making it difficult to predict patient prognoses




Figure created with BioRender

Hypothesis Nanoparticle trafficking in the TIME is governed by coupling between cell-extrinsic immune-recognition processes and cell-intrinsic biophysical mechanisms of cellular uptake, shaping tissue-level behavior.


Impact Improved therapeutic design principles and non-invasive diagnostics

Objective Determine how extrinsic factors influence immune recognition and uptake of nanoparticles

Methods

We develop spatially explicit biophysical models of relevant extrinsic and intrinsic phenomena involved in immune recognition and uptake

Complement opsonization




Agent-based simulation model
- Complement tickover, autocatalysis, diffusion, and decay

For more details see [4,5].

Control kinetic and transport parameters and NP geometry

Measure dynamic complement activity and final opsonized fraction

Phagocytic recognition




Monte Carlo simulation model
- Deformable membrane surface and asymmetrically opsonized NP
- Receptor-ligand binding
- Clathrin coat formation

For more details see [3,6].

Control membrane biophysical properties by changing energy function parameters

Measure receptor engagement, and membrane deformation

Legend



Results


1. Nanoparticle recognition follows a reversible switch-like percolation as a function of extrinsic factors

Extrinsic factors influencing opsonization

- C3b activity in TIME: C3b amplification, deactivation, and diffusion
- NP ligand spacing/density


Clinical Implication

1. Small shifts in immune microenvironment may trigger dramatic shifts in complement response to therapies
2. These findings motivate engineering therapies informed by patient-specific biomarkers




2a. Four distinct regimes of spatial immune receptor engagement based on NP

Spatial receptor engagement and membrane deformation changes with increasing complement opsonization



2b. Immune engagement and deformation further depends on coupling of extrinsic + intrinsic factors



Receptor engagement and membrane deformation changes in response to **intrinsic** cytoskeleton remodeling (excess membrane area)

Clinical Implication

1. Heterogeneous membrane remodeling in the TIME influences NP clearance and EV-mediate signal transduction
2. Membrane biomarker profiling could help customize NP treatments and contextualize EV surface profiles

Conclusions and Impact

CONCLUSIONS

The coupling of cell-extrinsic (e.g complement activity and nanoparticle surface chemistry) and cell-intrinsic (e.g. excess membrane area from cytoskeletal and lipid remodeling) factors drive immune receptor engagement and membrane remodeling

SCIENTIFIC IMPACT


Our work delineates key physical mechanisms governing NP recognition and uptake under heterogeneous immune and biophysical conditions

CLINICAL IMPACT

These findings motivate integrating immune, membrane, and single EV profiling biomarkers to inform patient-specific diagnostics and therapy optimization to tackle inter- and intra- patient heterogeneity

These physics may govern heterogeneous signaling responsible for
1.) malignant and metastatic transformation via exosome trafficking and
2.) reduced therapeutic efficacy in tumors


Future Work



Parihar et al. 2025

Study tumor immune evasion by modeling EVs with surface PDL1 engaging with T-cell receptors

Visualize interface with STORM microscopy



Study effect of NP and EV uptake on tumor heterogeneity using tissue-level simulation

Acknowledgements

We thank funding and computing support in part from NIH, Physical Sciences-Oncology Network, Penn PARCC, CIPD Restore Grant and NSF ACCESS.

National Institutes of Health

References

1. Parihar, K., Liu, DA., Hassan, G. *et al. Nat. Biomed. Eng* **9**, 1202–1221 (2025). https://doi.org/10.1038/s41551-025-01446-0
2. Wilhelm, S., Tavares, A., Dai, Q. *et al. Nat Rev Mater* **1**, 16014 (2016). https://doi.org/10.1038/natrevmats.2016.14
3. Ramakrishnan N et al. 2016 R. Soc. open sci. 3: 160260. http://dx.doi.org/10.1098/rsos.160260S
4. Wang et al., 2025, Cell 188, 4058–4073. https://doi.org/10.1016/j.cell.2025.05.026
5. **Monson, S**, Kulkarni, S, Meyerson, J, Brenner, J, Radhakrishnan, R. bioRxiv 2026.08.14.744667; doi: https://doi.org/10.64898/2026.08.14.744667
6. **Monson, S**, Parihar, K. *manuscript in preparation*.