

## Value

## CISC 7404 - Decision Making

Steven Morad

University of Macau

Review	2
Policy-Conditioned Returns	3
Value Functions	17
Exercise	24
TD Value Functions	26
Q Functions	38
Q Learning	49
Homework	55

## Review

Trajectory optimization is model-based algorithm

Trajectory optimization is model-based algorithm

Guaranteed optimal policy, given infinite compute

Trajectory optimization is model-based algorithm

Guaranteed optimal policy, given infinite compute

We must make approximations to implement trajectory optimization

Trajectory optimization is model-based algorithm

Guaranteed optimal policy, given infinite compute

We must make approximations to implement trajectory optimization

These approximations break optimality guarantees

Trajectory optimization is model-based algorithm

Guaranteed optimal policy, given infinite compute

We must make approximations to implement trajectory optimization

These approximations break optimality guarantees

Today, we will look at new algorithms based on the notion of value

Trajectory optimization is model-based algorithm

Guaranteed optimal policy, given infinite compute

We must make approximations to implement trajectory optimization

These approximations break optimality guarantees

Today, we will look at new algorithms based on the notion of **value** 

Uses fewer approximations but can achieve optimal policy

Trajectory optimization is model-based algorithm

Guaranteed optimal policy, given infinite compute

We must make approximations to implement trajectory optimization

These approximations break optimality guarantees

Today, we will look at new algorithms based on the notion of **value** 

Uses fewer approximations but can achieve optimal policy

Can model infinitely long returns

Trajectory optimization is model-based algorithm

Guaranteed optimal policy, given infinite compute

We must make approximations to implement trajectory optimization

These approximations break optimality guarantees

Today, we will look at new algorithms based on the notion of **value** 

Uses fewer approximations but can achieve optimal policy

Can model infinitely long returns

Expensive to train, but very cheap to use

Recall the return from trajectory optimization

Recall the return from trajectory optimization

$$[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E} [\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots]$$

Recall the return from trajectory optimization

$$[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, \textcolor{red}{a_0, a_1, \ldots}] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E} [\mathcal{R}(s_{t+1}) \mid s_0, \textcolor{red}{a_0, a_1, \ldots}]$$

This is an action-conditioned discounted return

Recall the return from trajectory optimization

$$[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, \textcolor{red}{a_0, a_1, \ldots}] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E} [\mathcal{R}(s_{t+1}) \mid s_0, \textcolor{red}{a_0, a_1, \ldots}]$$

This is an action-conditioned discounted return

Conditioned/dependent on a sequence of actions

Recall the return from trajectory optimization

$$[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, \textcolor{red}{a_0, a_1, \ldots}] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E} [\mathcal{R}(s_{t+1}) \mid s_0, \textcolor{red}{a_0, a_1, \ldots}]$$

This is an action-conditioned discounted return

Conditioned/dependent on a sequence of actions

There is no structure to the actions

Recall the return from trajectory optimization

$$[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, \textcolor{red}{a_0, a_1, \ldots}] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, \textcolor{red}{a_0, a_1, \ldots}]$$

This is an action-conditioned discounted return

Conditioned/dependent on a sequence of actions

There is no structure to the actions

Random

Recall the return from trajectory optimization

$$[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, \textcolor{red}{a_0, a_1, \ldots}] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E} [\mathcal{R}(s_{t+1}) \mid s_0, \textcolor{red}{a_0, a_1, \ldots}]$$

This is an action-conditioned discounted return

Conditioned/dependent on a sequence of actions

There is no structure to the actions

- Random
- Picked by humans

Recall the return from trajectory optimization

$$[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, \textcolor{red}{a_0, a_1, \ldots}] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E} [\mathcal{R}(s_{t+1}) \mid s_0, \textcolor{red}{a_0, a_1, \ldots}]$$

This is an action-conditioned discounted return

Conditioned/dependent on a sequence of actions

There is no structure to the actions

- Random
- Picked by humans
- Maximize  $\mathcal{G}$

$$\mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots]$$

$$\mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots]$$

Last time, we introduced the policy

$$\mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots]$$

Last time, we introduced the policy

$$\pi: S \times \Theta \mapsto \Delta A$$

$$\mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots]$$

Last time, we introduced the policy

$$\pi: S \times \Theta \mapsto \Delta A$$

Example policy, greedy policy

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots]$$

Last time, we introduced the policy

$$\pi: S \times \Theta \mapsto \Delta A$$

Example policy, greedy policy

$$\pi(a_t \mid s_t; \theta_\pi) = \begin{cases} 1 \text{ if } a_t = \arg\max_{a_t \in A} \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] \\ 0 \text{ otherwise} \end{cases}$$

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots]$$

Last time, we introduced the policy

$$\pi: S \times \Theta \mapsto \Delta A$$

Example policy, greedy policy

$$\pi(a_t \mid s_t; \theta_\pi) = \begin{cases} 1 \text{ if } a_t = \arg\max_{a_t \in A} \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] \\ 0 \text{ otherwise} \end{cases}$$

$$[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots]$$

$$[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E} [\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots]$$

$$\pi: S \times \Theta \mapsto \Delta A$$

$$[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E} [\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots]$$

$$\pi: S \times \Theta \mapsto \Delta A$$

Conditioning the return on actions is annoying

$$[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E} [\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots]$$

$$\pi: S \times \Theta \mapsto \Delta A$$

Conditioning the return on actions is annoying

Must compute infinitely many actions and outcomes for the return

$$[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E} [\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots]$$

$$\pi: S \times \Theta \mapsto \Delta A$$

Conditioning the return on actions is annoying

Must compute infinitely many actions and outcomes for the return

What if we condition on a policy, instead of specific actions?

$$\mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots]$$

$$a_0 \sim \pi(\cdot \mid s_0; \theta_\pi), \quad a_1 \sim \pi(\cdot \mid s_1; \theta_\pi), \quad a_2 \sim \pi(\cdot \mid s_2; \theta_\pi), \quad \dots$$

$$\mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots]$$

$$a_0 \sim \pi(\cdot \mid s_0; \theta_\pi), \quad a_1 \sim \pi(\cdot \mid s_1; \theta_\pi), \quad a_2 \sim \pi(\cdot \mid s_2; \theta_\pi), \quad \dots$$

Condition on a function parameterized by  $\theta_{\pi}$  instead of many actions

$$\mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots]$$

$$a_0 \sim \pi(\cdot \mid s_0; \theta_{\pi}), \quad a_1 \sim \pi(\cdot \mid s_1; \theta_{\pi}), \quad a_2 \sim \pi(\cdot \mid s_2; \theta_{\pi}), \quad \dots$$

Condition on a function parameterized by  $\theta_{\pi}$  instead of many actions

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \boldsymbol{\theta}_{\pi}] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, \boldsymbol{\theta}_{\pi}]$$

$$\mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots]$$

$$a_0 \sim \pi(\cdot \mid s_0; \theta_\pi), \quad a_1 \sim \pi(\cdot \mid s_1; \theta_\pi), \quad a_2 \sim \pi(\cdot \mid s_2; \theta_\pi), \quad \dots$$

Condition on a function parameterized by  $\theta_{\pi}$  instead of many actions

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \boldsymbol{\theta}_{\pi}] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, \boldsymbol{\theta}_{\pi}]$$

The function outputs a distribution over the action space  $\pi(a \mid s; \theta_{\pi})$ 

$$\mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots]$$

$$\begin{split} \mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \ldots] &= \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots] \\ \mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \theta_{\pi}] &= \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, \theta_{\pi}] \end{split}$$

$$\begin{split} \mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \ldots] &= \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots] \\ \mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \theta_{\pi}] &= \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, \theta_{\pi}] \end{split}$$

Now conditioned on the policy

$$\begin{split} \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] &= \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots] \\ \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0; \theta_{\pi}] &= \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, \theta_{\pi}] \end{split}$$

Now conditioned on the policy

But remember,  $\mathcal{R}(s_{t+1})$  hides much of the magic

$$\begin{split} \mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \ldots] &= \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \ldots] \\ \mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \theta_{\pi}] &= \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, \theta_{\pi}] \end{split}$$

Now conditioned on the policy

But remember,  $\mathcal{R}(s_{t+1})$  hides much of the magic

How does  $\mathbb{E}[\mathcal{R}(s_{t+1})]$  change when we condition on  $\theta_{\pi}$ ?

$$\mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathcal{R}(s_{t+1}) \sum_{s_{t+1} \in S} \Pr(s_{t+1} \mid s_0, a_0, \ldots, a_t)$$

$$\mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathcal{R}(s_{t+1}) \sum_{s_{t+1} \in S} \Pr(s_{t+1} \mid s_0, a_0, \ldots, a_t)$$

**Question:** What changes when we condition on  $\theta_{\pi}$ ?

$$\mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathcal{R}(s_{t+1}) \sum_{s_{t+1} \in S} \Pr(s_{t+1} \mid s_0, a_0, \ldots, a_t)$$

**Question:** What changes when we condition on  $\theta_{\pi}$ ?

$$\Pr(s_{t+1} \mid s_0; \theta_{\pi})$$

$$\mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathcal{R}(s_{t+1}) \sum_{s_{t+1} \in S} \Pr(s_{t+1} \mid s_0, a_0, \ldots, a_t)$$

**Question:** What changes when we condition on  $\theta_{\pi}$ ?

$$\Pr(s_{t+1} \mid s_0; \theta_{\pi})$$

Maybe we can use  $\Pr(s_{t+1} \mid s_t, a_t)$  to figure this out

$$\mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathcal{R}(s_{t+1}) \sum_{s_{t+1} \in S} \Pr(s_{t+1} \mid s_0, a_0, \ldots, a_t)$$

**Question:** What changes when we condition on  $\theta_{\pi}$ ?

$$\Pr(s_{t+1} \mid s_0; \theta_{\pi})$$

Maybe we can use  $\Pr(s_{t+1} \mid s_t, a_t)$  to figure this out

**Question:** What was  $Pr(s_{t+1} \mid s_t, a_t)$ ?

$$\mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0, a_0, a_1, \ldots] = \sum_{t=0}^{\infty} \gamma^t \mathcal{R}(s_{t+1}) \sum_{s_{t+1} \in S} \Pr(s_{t+1} \mid s_0, a_0, \ldots, a_t)$$

**Question:** What changes when we condition on  $\theta_{\pi}$ ?

$$\Pr(s_{t+1} \mid s_0; \theta_{\pi})$$

Maybe we can use  $\Pr(s_{t+1} \mid s_t, a_t)$  to figure this out

**Question:** What was  $Pr(s_{t+1} \mid s_t, a_t)$ ?

**Answer:** State transition function

$$\operatorname{Tr}(s_{t+1} \mid s_t, a_t)$$

$$\operatorname{Tr}(s_{t+1} \mid s_t, a_t)$$

$$\operatorname{Tr}(s_{t+1} \mid s_t, a_t)$$

**Issue:** State transition function needs an action  $\boldsymbol{a}_t$ 

$$\operatorname{Tr}(s_{t+1} \mid s_t, a_t)$$

**Issue:** State transition function needs an action  $a_t$ 

Policy  $\pi$  outputs a distribution over the action space

$$\operatorname{Tr}(s_{t+1} \mid s_t, a_t)$$

**Issue:** State transition function needs an action  $a_t$ 

Policy  $\pi$  outputs a distribution over the action space

**Question:** What is  $\Pr(s_{t+1} \mid s_t, \theta_{\pi})$ ?

$$\operatorname{Tr}(s_{t+1} \mid s_t, a_t)$$

**Issue:** State transition function needs an action  $a_t$ 

Policy  $\pi$  outputs a distribution over the action space

**Question:** What is  $\Pr(s_{t+1} \mid s_t, \theta_{\pi})$ ? Hint: Consider all possible actions

$$\operatorname{Tr}(s_{t+1} \mid s_t, a_t)$$

**Issue:** State transition function needs an action  $a_t$ 

Policy  $\pi$  outputs a distribution over the action space

**Question:** What is  $\Pr(s_{t+1} \mid s_t, \theta_{\pi})$ ? Hint: Consider all possible actions

$$\Pr(s_{t+1} \mid s_t; \theta_{\pi}) = \sum_{a_t \in A} \operatorname{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_{\pi})$$

$$\operatorname{Tr}(s_{t+1} \mid s_t, a_t)$$

**Issue:** State transition function needs an action  $a_t$ 

Policy  $\pi$  outputs a distribution over the action space

**Question:** What is  $\Pr(s_{t+1} \mid s_t, \theta_{\pi})$ ? Hint: Consider all possible actions

$$\Pr(s_{t+1} \mid s_t; \theta_{\pi}) = \sum_{a_t \in A} \operatorname{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_{\pi})$$

Combine the policy distribution with next state distribution

$$\Pr(s_{t+1} \mid s_t; \theta_{\pi}) = \sum_{a_t \in A} \operatorname{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_{\pi})$$

$$\Pr(s_{t+1} \mid s_t; \theta_{\pi}) = \sum_{a_t \in A} \operatorname{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_{\pi})$$

Write out the first few timesteps

$$\Pr(s_{t+1} \mid s_t; \theta_{\pi}) = \sum_{a_t \in A} \operatorname{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_{\pi})$$

Write out the first few timesteps

$$\Pr(s_1 \mid s_0; \theta_{\pi}) = \sum_{a_0 \in A} \operatorname{Tr}(s_1 \mid s_0, a_0) \cdot \pi(a_0 \mid s_0; \theta_{\pi})$$

$$\Pr(s_{t+1} \mid s_t; \theta_{\pi}) = \sum_{a_t \in A} \operatorname{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_{\pi})$$

Write out the first few timesteps

$$\begin{split} \Pr(s_1 \mid s_0; \theta_\pi) &= \sum_{a_0 \in A} \mathrm{Tr}(s_1 \mid s_0, a_0) \cdot \pi(a_0 \mid s_0; \theta_\pi) \\ \Pr(s_2 \mid s_0; \theta_\pi) &= \sum_{s_1 \in S} \sum_{a_1 \in A} \mathrm{Tr}(s_2 \mid s_1, a_1) \cdot \pi(a_1 \mid s_1; \theta_\pi) \\ &\cdot \sum_{a_0 \in A} \mathrm{Tr}(s_1 \mid s_0, a_0) \cdot \pi(a_0 \mid s_0; \theta_\pi) \end{split}$$

$$\begin{split} \Pr(s_1 \mid s_0; \theta_\pi) &= \sum_{a_0 \in A} \mathrm{Tr}(s_1 \mid s_0, a_0) \cdot \pi(a_0 \mid s_0; \theta_\pi) \\ \Pr(s_2 \mid s_0; \theta_\pi) &= \sum_{s_1 \in S} \sum_{a_1 \in A} \mathrm{Tr}(s_2 \mid s_1, a_1) \cdot \pi(a_1 \mid s_1; \theta_\pi) \\ &\cdot \sum_{a_0 \in A} \mathrm{Tr}(s_1 \mid s_0, a_0) \cdot \pi(a_0 \mid s_0; \theta_\pi) \end{split}$$

$$\begin{split} \Pr(s_1 \mid s_0; \theta_\pi) &= \sum_{a_0 \in A} \mathrm{Tr}(s_1 \mid s_0, a_0) \cdot \pi(a_0 \mid s_0; \theta_\pi) \\ \Pr(s_2 \mid s_0; \theta_\pi) &= \sum_{s_1 \in S} \sum_{a_1 \in A} \mathrm{Tr}(s_2 \mid s_1, a_1) \cdot \pi(a_1 \mid s_1; \theta_\pi) \\ &\cdot \sum_{a_0 \in A} \mathrm{Tr}(s_1 \mid s_0, a_0) \cdot \pi(a_0 \mid s_0; \theta_\pi) \end{split}$$

Derive a general form for  $\Pr(s_{n+1} \mid s_0; \theta_{\pi})$ 

$$\begin{split} \Pr(s_1 \mid s_0; \theta_\pi) &= \sum_{a_0 \in A} \operatorname{Tr}(s_1 \mid s_0, a_0) \cdot \pi(a_0 \mid s_0; \theta_\pi) \\ \Pr(s_2 \mid s_0; \theta_\pi) &= \sum_{s_1 \in S} \sum_{a_1 \in A} \operatorname{Tr}(s_2 \mid s_1, a_1) \cdot \pi(a_1 \mid s_1; \theta_\pi) \\ &\cdot \sum_{a_0 \in A} \operatorname{Tr}(s_1 \mid s_0, a_0) \cdot \pi(a_0 \mid s_0; \theta_\pi) \end{split}$$

Derive a general form for  $\Pr(s_{n+1} \mid s_0; \theta_{\pi})$ 

$$\Pr(s_{n+1} \mid s_0; \theta_{\pi}) = \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \left( \sum_{a_t \in A} \operatorname{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta) \right)$$

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \left( \sum_{a_t \in A} \operatorname{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta) \right)$$

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \left( \sum_{a_t \in A} \operatorname{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta) \right)$$

Plug back into our expected reward

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \left( \sum_{a_t \in A} \operatorname{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta) \right)$$

Plug back into our expected reward

$$\mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_{\pi}] = \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_{\pi})$$

$$\mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_{\pi}] = \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_{\pi})$$

$$\mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_{\pi}] = \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_{\pi})$$

$$\mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_{\pi}] = \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_{\pi})$$

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \theta_{\pi}] =$$

$$\mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_{\pi}] = \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_{\pi})$$

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \boldsymbol{\theta}_{\pi}] = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \boldsymbol{\theta}_{\pi})$$

$$\mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_{\pi}] = \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_{\pi})$$

$$\begin{split} \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0; \theta_\pi] &= \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ &+ \gamma \sum_{s_2 \in S} \mathcal{R}(s_2) \cdot \Pr(s_2 \mid s_0; \theta_\pi) \end{split}$$

$$\mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_{\pi}] = \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_{\pi})$$

$$\begin{split} \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0; \theta_\pi] &= \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ &+ \gamma \sum_{s_2 \in S} \mathcal{R}(s_2) \cdot \Pr(s_2 \mid s_0; \theta_\pi) \\ &+ \gamma^2 \sum_{s_3 \in S} \mathcal{R}(s_3) \cdot \Pr(s_3 \mid s_0; \theta_\pi) \end{split}$$

$$\mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_{\pi}] = \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_{\pi})$$

$$\begin{split} \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0; \theta_\pi] &= \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ &+ \gamma \sum_{s_2 \in S} \mathcal{R}(s_2) \cdot \Pr(s_2 \mid s_0; \theta_\pi) \\ &+ \gamma^2 \sum_{s_3 \in S} \mathcal{R}(s_3) \cdot \Pr(s_3 \mid s_0; \theta_\pi) \end{split}$$

**Definition:** General form of policy-conditioned discounted return

**Definition:** General form of policy-conditioned discounted return

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \boldsymbol{\theta}_{\pi}] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_n, a_n)$$

## **Policy-Conditioned Returns**

**Definition:** General form of policy-conditioned discounted return

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \boldsymbol{\theta}_{\pi}] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_n, a_n)$$

Where the state distribution is

## **Policy-Conditioned Returns**

**Definition:** General form of policy-conditioned discounted return

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \boldsymbol{\theta}_{\pi}] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_n, a_n)$$

Where the state distribution is

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \left( \sum_{a_t \in A} \operatorname{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta) \right)$$

$$\Pr(s_{n+1} \mid s_0; \theta_{\pi}) = \sum_{a_0, \dots, a_n \in A} \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \operatorname{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_{\pi})$$

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{a_0, \dots, a_n \in A} \sum_{s_1, \dots, s_n \in S} \prod_{t=0} \operatorname{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \boldsymbol{\theta}_{\pi}] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \boldsymbol{\theta}_{\pi})$$

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{a_0, \dots, a_n \in A} \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \operatorname{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \boldsymbol{\theta}_{\pi}] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \boldsymbol{\theta}_{\pi})$$

These two equations form the basis of all reinforcement learning

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{a_0, \dots, a_n \in A} \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \operatorname{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \boldsymbol{\theta}_{\pi}] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{m+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \boldsymbol{\theta}_{\pi})$$

These two equations form the basis of all reinforcement learning

**Goal:** find the  $\theta_{\pi}$  (policy parameters) to maximize the expected return

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{a_0, \dots, a_n \in A} \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \operatorname{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \boldsymbol{\theta}_{\pi}] = \sum_{n=0}^{\infty} \gamma^n \sum_{s=1}^{\infty} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \boldsymbol{\theta}_{\pi})$$

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{a_0, \dots, a_n \in A} \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \operatorname{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \boldsymbol{\theta}_{\pi}] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \boldsymbol{\theta}_{\pi})$$

We have another name for  $\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \theta_{\pi}]$ 

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{a_0, \dots, a_n \in A} \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \operatorname{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \boldsymbol{\theta}_{\pi}] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \boldsymbol{\theta}_{\pi})$$

We have another name for  $\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \theta_{\pi}]$ 

We call it the **value function**  $V: S \times \Theta \mapsto \mathbb{R}$ 

$$\Pr(s_{n+1} \mid s_0; \theta_{\pi}) = \sum_{a_0, \dots, a_n \in A} \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \operatorname{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_{\pi})$$

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \boldsymbol{\theta}_{\pi}] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{m+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \boldsymbol{\theta}_{\pi})$$

We have another name for  $\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \theta_{\pi}]$ 

We call it the **value function**  $V: S \times \Theta \mapsto \mathbb{R}$ 

$$V(s_0,\theta_\pi) = \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0;\theta_\pi] = \sum_{n=0}^\infty \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0;\theta_\pi)$$

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s=1 \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0; \theta_\pi] = \sum_{n=0}^\infty \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0; \theta_\pi] = \sum_{n=0}^\infty \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0; \theta_\pi] = \sum_{n=0}^\infty \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

$$s = 240 \text{ km/h}$$

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0; \theta_\pi] = \sum_{n=0}^\infty \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

$$s=240~\mathrm{km/h}$$
  $\theta_{\pi}=\mathrm{Race~car~driver}$ 

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0; \theta_\pi] = \sum_{n=0}^\infty \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

$$s = 240 \text{ km/h}$$

$$\theta_{\pi} = \text{Race car driver} \qquad V(s, \theta_{\pi}) = \text{good}$$

$$V(s, \theta_{\pi}) = \text{good}$$

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0; \theta_\pi] = \sum_{n=0}^\infty \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

$$s = 240 \text{ km/h}$$

$$\theta_{\pi} = \text{Race car driver} \qquad V(s, \theta_{\pi}) = \text{good}$$

$$V(s, \theta_{\pi}) = \text{good}$$

$$s = 240 \text{ km/h}$$

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0; \theta_\pi] = \sum_{n=0}^\infty \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

$$s = 240 \text{ km/h}$$

$$\theta_{\pi} = \text{Race car driver} \qquad V(s, \theta_{\pi}) = \text{good}$$

$$V(s, \theta_{\pi}) = \text{good}$$

$$s = 240 \text{ km/h}$$

$$\theta_{\pi} = \text{Grandpa}$$

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0; \theta_\pi] = \sum_{n=0}^\infty \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

$$s = 240 \text{ km/h}$$

$$\theta_{\pi} = \text{Race car driver} \qquad V(s, \theta_{\pi}) = \text{good}$$

$$V(s, \theta_{\pi}) = \text{good}$$

$$s = 240 \text{ km/h}$$

$$\theta_{\pi} = \text{Grandpa}$$

$$V(s, \theta_{\pi}) = \text{not good}$$

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0; \theta_\pi] = \sum_{n=0}^\infty \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

Valuable states lead to good returns **under the current policy** 

$$s = 240 \text{ km/h}$$

$$\theta_{\pi} = \text{Race car driver} \qquad V(s, \theta_{\pi}) = \text{good}$$

$$V(s, \theta_{\pi}) = \text{good}$$

$$s = 240 \text{ km/h}$$

$$\theta_{\pi} = \text{Grandpa}$$

$$V(s, \theta_{\pi}) = \text{not good}$$

We use the value function to direct the policy to good states

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

Valuable states lead to good returns **under the current policy** 

$$s = 240 \text{ km/h}$$

$$\theta_{\pi} = \text{Race car driver} \qquad V(s, \theta_{\pi}) = \text{good}$$

$$V(s, \theta_{\pi}) = \text{good}$$

$$s = 240 \text{ km/h}$$

$$\theta_{\pi} = \text{Grandpa}$$

$$V(s, \theta_{\pi}) = \text{not good}$$

We use the value function to direct the policy to good states

It is a critical part of decision making

With the value function, we can use any state as a starting state

With the value function, we can use any state as a starting state

The state does not need to be the start of a trajectory

With the value function, we can use any state as a starting state

The state does not need to be the start of a trajectory

**Example:** Consider the sequence of states

$$s_a, s_b, s_c$$

With the value function, we can use any state as a starting state

The state does not need to be the start of a trajectory

**Example:** Consider the sequence of states

$$s_a, s_b, s_c$$

We can compute

$$V(s_a, \theta_\pi), V(s_b, \theta_\pi), V(s_c, \theta_\pi)$$

To find the value of any state

**Question:** Why does Prof. Steven keep showing stupid equations? How is the value function useful?

**Question:** Why does Prof. Steven keep showing stupid equations? How is the value function useful?

We can use the value of a state to make decisions

**Question:** Why does Prof. Steven keep showing stupid equations? How is the value function useful?

We can use the value of a state to make decisions

 $s_a = \text{Live in Macau}, s_b = \text{Live in Beijing}$ 

**Question:** Why does Prof. Steven keep showing stupid equations? How is the value function useful?

We can use the value of a state to make decisions

$$s_a = \text{Live in Macau}, s_b = \text{Live in Beijing}$$

Given all your preferences  $(\mathcal{R})$  and thoughts  $(\theta_{\pi})$ , we can determine which life is better for you

**Question:** Why does Prof. Steven keep showing stupid equations? How is the value function useful?

We can use the value of a state to make decisions

$$s_a = \text{Live in Macau}, s_b = \text{Live in Beijing}$$

Given all your preferences  $(\mathcal{R})$  and thoughts  $(\theta_{\pi})$ , we can determine which life is better for you

 $V(s, \theta_{\pi})$  considers your future friends, income, wife/husband, etc

**Question:** Why does Prof. Steven keep showing stupid equations? How is the value function useful?

We can use the value of a state to make decisions

$$s_a = \text{Live in Macau}, s_b = \text{Live in Beijing}$$

Given all your preferences  $(\mathcal{R})$  and thoughts  $(\theta_{\pi})$ , we can determine which life is better for you

 $V(s,\theta_{\pi})$  considers your future friends, income, wife/husband, etc

**Question:** Why does Prof. Steven keep showing stupid equations? How is the value function useful?

We can use the value of a state to make decisions

$$s_a = \text{Live in Macau}, s_b = \text{Live in Beijing}$$

Given all your preferences  $(\mathcal{R})$  and thoughts  $(\theta_{\pi})$ , we can determine which life is better for you

 $V(s,\theta_{\pi})$  considers your future friends, income, wife/husband, etc

$$V(s_a, \theta_\pi) = 1032$$

**Question:** Why does Prof. Steven keep showing stupid equations? How is the value function useful?

We can use the value of a state to make decisions

$$s_a = \text{Live in Macau}, s_b = \text{Live in Beijing}$$

Given all your preferences  $(\mathcal{R})$  and thoughts  $(\theta_{\pi})$ , we can determine which life is better for you

 $V(s,\theta_{\pi})$  considers your future friends, income, wife/husband, etc

$$V(s_a,\theta_\pi)=1032 \qquad \qquad V(s_b,\theta_\pi)=945$$

**Question:** Why does Prof. Steven keep showing stupid equations? How is the value function useful?

We can use the value of a state to make decisions

$$s_a = \text{Live in Macau}, s_b = \text{Live in Beijing}$$

Given all your preferences  $(\mathcal{R})$  and thoughts  $(\theta_{\pi})$ , we can determine which life is better for you

 $V(s,\theta_{\pi})$  considers your future friends, income, wife/husband, etc

$$V(s_a,\theta_\pi)=1032 \qquad \qquad V(s_b,\theta_\pi)=945$$

 $s_a = \text{Live in Macau}, s_b = \text{Live in Beijing}$ 

$$s_a = \text{Live in Macau}, s_b = \text{Live in Beijing}$$

$$V(s_a,\theta_\pi)=1032 \qquad \qquad V(s_b,\theta_\pi)=945$$

$$s_a = \text{Live in Macau}, s_b = \text{Live in Beijing}$$

$$V(s_a,\theta_\pi)=1032 \qquad \qquad V(s_b,\theta_\pi)=945$$

This value leads us to the right decisions

$$s_a = \text{Live in Macau}, s_b = \text{Live in Beijing}$$

$$V(s_a,\theta_\pi)=1032 \qquad \qquad V(s_b,\theta_\pi)=945$$

This value leads us to the right decisions

Some optimal decisions are hard for humans to make

$$s_a = \text{Live in Macau}, s_b = \text{Live in Beijing}$$

$$V(s_a,\theta_\pi)=1032 \qquad \qquad V(s_b,\theta_\pi)=945$$

This value leads us to the right decisions

Some optimal decisions are hard for humans to make

With value, we can be sure we make the right decision

• Think of two places you want to live after graduation  $s_a, s_b$ 

- Think of two places you want to live after graduation  $s_a, s_b$
- Consider your behavior  $(\theta_{\pi})$  and what is important to you  $(\mathcal{R})$

- Think of two places you want to live after graduation  $s_a, s_b$
- Consider your behavior  $(\theta_{\pi})$  and what is important to you  $(\mathcal{R})$
- Top 3 life goals as states  $s_x, s_y, s_z \in G$  (e.g., friends, money, hobby, etc)

- Think of two places you want to live after graduation  $s_a, s_b$
- Consider your behavior  $(\theta_{\pi})$  and what is important to you  $(\mathcal{R})$
- Top 3 life goals as states  $s_x, s_y, s_z \in G$  (e.g., friends, money, hobby, etc)
- Assign a reward  $\mathcal R$  for each goal, and choose discount factor  $\gamma$

- Think of two places you want to live after graduation  $s_a, s_b$
- Consider your behavior  $(\theta_{\pi})$  and what is important to you  $(\mathcal{R})$
- Top 3 life goals as states  $s_x, s_y, s_z \in G$  (e.g., friends, money, hobby, etc)
- Assign a reward  $\mathcal R$  for each goal, and choose discount factor  $\gamma$

For each location  $s_0 \in \{s_a, s_b\}$ :

- Think of two places you want to live after graduation  $s_a, s_b$
- Consider your behavior  $(\theta_{\pi})$  and what is important to you  $(\mathcal{R})$
- Top 3 life goals as states  $s_x, s_y, s_z \in G$  (e.g., friends, money, hobby, etc)
- Assign a reward  $\mathcal R$  for each goal, and choose discount factor  $\gamma$

For each location  $s_0 \in \{s_a, s_b\}$ :

- Estimate probability of reaching each goal  $\Pr(s_g \mid s_0); s_g \in \{s_x, s_y, s_z\}$
- Estimate time to accomplish each goal t = ...

$$V(s_0, \theta_\pi) = \sum_{s_g \in \{s_x, s_y, s_z\}} \gamma^t \mathcal{R}\big(s_g\big) \cdot \Pr\big(s_g \mid s_0; \theta_\pi\big)$$

**Note:** We define the value function in many different ways

**Note:** We define the value function in many different ways

It always approximates the expected discounted return from  $s_0$ 

**Note:** We define the value function in many different ways

It always approximates the expected discounted return from  $s_0$ 

We call the following equation the **Monte Carlo** value function

$$V(s_0, \theta_\pi) = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

**Note:** We define the value function in many different ways

It always approximates the expected discounted return from  $s_0$ 

We call the following equation the **Monte Carlo** value function

$$V(s_0, \theta_\pi) = \sum_{n=0}^\infty \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Difficult to compute the Monte Carlo value function

Note: We define the value function in many different ways

It always approximates the expected discounted return from  $s_0$ 

We call the following equation the **Monte Carlo** value function

$$V(s_0, \theta_\pi) = \sum_{n=0}^\infty \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Difficult to compute the Monte Carlo value function

Must either have a terminal states

**Note:** We define the value function in many different ways

It always approximates the expected discounted return from  $s_0$ 

We call the following equation the **Monte Carlo** value function

$$V(s_0, \theta_\pi) = \sum_{n=0}^\infty \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Difficult to compute the Monte Carlo value function

- Must either have a terminal states
- Or build the infinite decision tree

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0; \theta_\pi] = \sum_{t=0}^\infty \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0; \theta_\pi] = \sum_{t=0}^\infty \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

Infinite sums are icky

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0; \theta_\pi] = \sum_{t=0}^\infty \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

Infinite sums are icky

They make everything difficult and intractable

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0; \theta_\pi] = \sum_{t=0}^\infty \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

Infinite sums are icky

They make everything difficult and intractable

Let us try to delete the infinite sum

$$V(s_0, \theta_\pi) = \sum_{t=0}^\infty \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

$$V(s_0, \theta_\pi) = \sum_{t=0}^\infty \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

Factor out initial timestep t = 0 out of the outer sum

$$V(s_0, \theta_\pi) = \sum_{t=0}^\infty \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

Factor out initial timestep t = 0 out of the outer sum

$$V(s_0,\theta_\pi) = \gamma^0 \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0;\theta_\pi)$$

$$+ \sum_{t=1}^{\infty} \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_{\pi})$$

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \sum_{t=1}^{\infty} \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_{\pi})$$

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \sum_{t=1}^{\infty} \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_{\pi})$$

Rewrite sum starting from t = 0

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \sum_{t=1}^{\infty} \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_{\pi})$$

Rewrite sum starting from t = 0

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \sum_{t=0}^{\infty} \gamma^{t+1} \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \cdot \Pr(s_{t+2} \mid s_0; \theta_{\pi})$$

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \sum_{t=0}^{\infty} \gamma^{t+1} \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \cdot \Pr(s_{t+2} \mid s_0; \theta_{\pi})$$

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \sum_{t=0}^{\infty} \gamma^{t+1} \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \cdot \Pr(s_{t+2} \mid s_0; \theta_{\pi})$$

Factor out  $\gamma$ 

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \sum_{t=0}^{\infty} \gamma^{t+1} \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \cdot \Pr(s_{t+2} \mid s_0; \theta_{\pi})$$

Factor out  $\gamma$ 

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+\gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \cdot \Pr(s_{t+2} \mid s_0; \theta_{\pi})$$

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \cdot \Pr(s_{t+2} \mid s_0; \theta_{\pi})$$

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \cdot \Pr(s_{t+2} \mid s_0; \theta_{\pi})$$

Split Pr using Markov property

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \cdot \Pr(s_{t+2} \mid s_0; \theta_{\pi})$$

Split Pr using Markov property

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \gamma \sum_{t=0}^{\infty} \gamma^{t} \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \sum_{s_{1}} \Pr(s_{t+2} \mid s_{t+1}; \theta_{\pi}) \Pr(s_{1} \mid s_{0}; \theta_{\pi})$$

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \gamma \sum_{t=0}^{\infty} \gamma^{t} \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \sum_{s_{1}} \Pr(s_{t+2} \mid s_{t+1}; \theta_{\pi}) \Pr(s_{1} \mid s_{0}; \theta_{\pi})$$

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \sum_{s_1} \Pr(s_{t+2} \mid s_{t+1}; \theta_{\pi}) \Pr(s_1 \mid s_0; \theta_{\pi})$$

Move sum and Pr outside

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \sum_{s_1} \Pr(s_{t+2} \mid s_{t+1}; \theta_{\pi}) \Pr(s_1 \mid s_0; \theta_{\pi})$$

Move sum and Pr outside

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \sum_{s_1} \Pr(s_1 \mid s_0; \theta_\pi) \gamma \sum_{t=0}^\infty \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi)$$

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \sum_{s_1} \Pr(s_1 \mid s_0; \theta_\pi) \gamma \ \sum_{t=0}^\infty \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi)$$

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \sum_{s_1} \Pr(s_1 \mid s_0; \theta_\pi) \gamma \left[ \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi) \right]$$

**Question:** What is this term?

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \sum_{s_1} \Pr(s_1 \mid s_0; \theta_\pi) \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi)$$

**Question:** What is this term?

$$V(s_0, \theta_\pi) = \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \sum_{s_1} \Pr(s_1 \mid s_0; \theta_\pi) \gamma \left[ \sum_{t=0}^\infty \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi) \right]$$

**Question:** What is this term?

$$V(s_0, \theta_\pi) = \sum_{t=0}^\infty \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

$$V(s_1, \theta_\pi) = \sum_{t=0}^\infty \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \cdot \Pr(s_{t+2} \mid s_1; \theta_\pi)$$

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \sum_{s_1} \Pr(s_1 \mid s_0; \theta_\pi) \gamma \sum_{t=0}^\infty \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi)$$

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \sum_{s_1} \Pr(s_1 \mid s_0; \theta_\pi) \gamma \sum_{t=0}^\infty \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi)$$

$$V(s_0, \theta_\pi) = \left(\sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)\right) + \gamma V(s_1, \theta_\pi)$$

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \sum_{s_1} \Pr(s_1 \mid s_0; \theta_\pi) \gamma \sum_{t=0}^\infty \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi)$$

$$V(s_0, \theta_\pi) = \left(\sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)\right) + \gamma V(s_1, \theta_\pi)$$

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

$$+ \sum_{s_1} \Pr(s_1 \mid s_0; \theta_\pi) \gamma \sum_{t=0}^\infty \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi)$$

$$V(s_0, \theta_\pi) = \left(\sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)\right) + \gamma V(s_1, \theta_\pi)$$

$$V(s_0, \theta_{\pi}) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_{\pi}] + \gamma V(s_1, \theta_{\pi})$$

This is a huge finding!

$$V(s_0,\theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0;\theta_\pi] + \gamma V(s_1,\theta_\pi)$$

$$V(s_0,\theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0;\theta_\pi] + \gamma V(s_1,\theta_\pi)$$

Value function has a recursive definition

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

Value function has a recursive definition

Represent policy-conditioned discounted return without an infinite sum

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

Value function has a recursive definition

Represent policy-conditioned discounted return without an infinite sum

We call this the **Temporal Difference** (TD) value function

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

Value function has a recursive definition

Represent policy-conditioned discounted return without an infinite sum

We call this the **Temporal Difference** (TD) value function

Compute the return with a single transition  $s_0 \rightarrow s_1$ 

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

Value function has a recursive definition

Represent policy-conditioned discounted return without an infinite sum

We call this the **Temporal Difference** (TD) value function

Compute the return with a single transition  $s_0 \rightarrow s_1$ 

Evaluate infinite-depth decision tree with a single function call

To summarize, we can represent the value function in two ways:

To summarize, we can represent the value function in two ways:

The Monte Carlo value function

To summarize, we can represent the value function in two ways:

The Monte Carlo value function

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

To summarize, we can represent the value function in two ways:

The Monte Carlo value function

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

The Temporal Difference value function

To summarize, we can represent the value function in two ways:

The Monte Carlo value function

$$V(s_0,\theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0;\theta_\pi] + \gamma V(s_1,\theta_\pi)$$

The Temporal Difference value function

$$V(s_0, \theta_\pi) = \sum_{t=0}^\infty \gamma^t \mathbb{E} \big[ \mathcal{R} \big( s_{t+1} \big) \mid s_0, \theta_\pi \big]$$

We saw two forms of the value function

We saw two forms of the value function

The value function relies on a policy

We saw two forms of the value function

The value function relies on a policy

But it does not tell us the policy

We saw two forms of the value function

The value function relies on a policy

But it does not tell us the policy

How can we use the value function to find an optimal policy?

Consider the value function

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\pmb{\tau}) \mid s_0; \theta_\pi] = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

With trajectory optimization we conditioned on actions

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \ldots]$$

We conditioned the value function on policy parameters

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \theta_{\pi}]$$

What if we wanted a mix of both?

$$\mathbb{E}[\mathcal{G}(\boldsymbol{ au}) \mid s_0, a_0; \theta_{\pi}]$$

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0; \theta_{\pi}]$$

We call this the **Q** function

$$Q(s, a, \theta_{\pi}) = \mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0; \theta_{\pi}]$$

We can derive the Q function from the value function

$$V(s_0, \theta_{\pi}) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_{\pi}] + \gamma V(s_1, \theta_{\pi})$$

First, introduce the action  $a_0$ 

$$V(s_0, a_0, \theta_{\pi}) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_{\pi}] + \gamma V(s_1, \theta_{\pi})$$

$$V(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

Condition the initial reward on the action

$$V(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

Call it the Q function

$$Q(s_0, a_0, \theta_{\pi}) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_{\pi})$$

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

The Q function tells us:

- The value of an action  $a_0$
- In state  $s_0$
- If we follow  $\pi(a_t \mid s_t; \theta_{\pi})$  afterwards

**Question:** How can we use the Q function for decision making?

Hint: We can evaluate Q for every possible action

$$\operatorname*{arg\ max}_{a_0 \in A} Q(s_0, a_0, \theta_\pi) = \operatorname*{arg\ max}_{a_0 \in A} \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$


$$\underset{a_0 \in A}{\arg \max} \, Q(s_0, a_0, \theta_\pi) = \underset{a_0 \in A}{\arg \max} \, \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

This is a very powerful equation

- Compute  $Q(s_0, a_0)$  for all  $a_0$
- Pick the  $a_0$  that maximizes  $Q(s_0,a_0)$


This  $a_0$  is **guaranteed** to be the optimal action for the **infinite** future

We collapsed the infinite decision tree into a single level



$$t = 2$$

•



$$\operatorname*{arg\ max}_{a_0 \in A} Q(s_0, a_0, \theta_\pi) = \operatorname*{arg\ max}_{a_0 \in A} (\mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi))$$

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

We will use either Q or V in every other algorithm in the course!

It is the core of decision making

Q learning is a model-free algorithm first invented in the 1980s

It is still used heavily today

In fact, I am using it in our research right now

We now have all the information we need to implement Q learning

Our Q function relies on the value function for some  $\theta_{\pi}$ 

Right now, it is not clear what the policy is

So how can we use the Q function without knowing the policy?

Start with the Q function

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

We want to take the action that maximizes Q

$$\underset{a_0 \in A}{\arg \max} \, Q(s_0, a_0, \theta_\pi) = \underset{a_0 \in A}{\arg \max} (\mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi))$$

Recall the definition of value function

$$V(s_0, \theta_\pi) = \sum_{t=0}^\infty \gamma^t \mathbb{E} \big[ \mathcal{R} \big( s_{t+1} \big) \mid s_0; \theta_\pi \big]$$

$$V(s_0, \theta_\pi) = \sum_{t=0}^\infty \gamma^t \mathbb{E} \big[ \mathcal{R}(s_{t+1}) \mid s_0; \theta_\pi \big]$$

What should our policy be?

Well we know the following is optimal

$$\underset{a_0 \in A}{\arg \max} \, Q(s_0, a_0, \theta_\pi) = \underset{a_0 \in A}{\arg \max} (\mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi))$$

What if we say we follow a policy that maximizes Q?

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 \text{ if } a_0 = \arg\max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 \text{ otherwise} \end{cases}$$

What is the value function for this policy?

$$V(s_0,\theta_\pi) = \max_{a \in A} Q(s_0,a,\theta_\pi)$$

So we can rewrite the Q function without V

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \operatorname*{arg\ max}_{a \in A} Q(s_1, a, \theta_\pi)$$

## Homework

### Homework

Due in 2 weeks (Weds 12 March, 23:59)

Download and submit .py and .ipynb files

Uses turnitin for checking

https://colab.research.google.com/drive/1xtBxAaVc3ax6\_j59RC3

NLQQPFcIEoau-?usp=sharing