



# Value

CISC 7404 - Decision Making

Steven Morad

University of Macau

Review .....	2
Policy-Conditioned Returns .....	3
Value Functions .....	17
Exercise .....	24
TD Value Functions .....	26
Q Functions .....	38
Q Learning .....	48
Homework .....	67

# Review

---

# Policy-Conditioned Returns

---

# Policy-Conditioned Returns

Trajectory optimization is model-based algorithm

# Policy-Conditioned Returns

Trajectory optimization is model-based algorithm

Guaranteed optimal policy, given infinite compute

# Policy-Conditioned Returns

Trajectory optimization is model-based algorithm

Guaranteed optimal policy, given infinite compute

We must make approximations to implement trajectory optimization

# Policy-Conditioned Returns

Trajectory optimization is model-based algorithm

Guaranteed optimal policy, given infinite compute

We must make approximations to implement trajectory optimization

These approximations break optimality guarantees



# Policy-Conditioned Returns

Trajectory optimization is model-based algorithm

Guaranteed optimal policy, given infinite compute

We must make approximations to implement trajectory optimization

These approximations break optimality guarantees

Today, we will look at new algorithms based on the notion of **value**

# Policy-Conditioned Returns

Trajectory optimization is model-based algorithm

Guaranteed optimal policy, given infinite compute

We must make approximations to implement trajectory optimization

These approximations break optimality guarantees

Today, we will look at new algorithms based on the notion of **value**

Uses fewer approximations but can achieve optimal policy

# Policy-Conditioned Returns

Trajectory optimization is model-based algorithm

Guaranteed optimal policy, given infinite compute

We must make approximations to implement trajectory optimization

These approximations break optimality guarantees

Today, we will look at new algorithms based on the notion of **value**

Uses fewer approximations but can achieve optimal policy

Can model infinitely long returns

# Policy-Conditioned Returns

Trajectory optimization is model-based algorithm

Guaranteed optimal policy, given infinite compute

We must make approximations to implement trajectory optimization

These approximations break optimality guarantees

Today, we will look at new algorithms based on the notion of **value**

Uses fewer approximations but can achieve optimal policy

Can model infinitely long returns

Expensive to train, but very cheap to use

# Policy-Conditioned Returns

Recall the return from trajectory optimization

# Policy-Conditioned Returns

Recall the return from trajectory optimization

$$[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

# Policy-Conditioned Returns

Recall the return from trajectory optimization

$$[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

This is an **action-conditioned** discounted return

# Policy-Conditioned Returns

Recall the return from trajectory optimization

$$[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

This is an **action-conditioned** discounted return

Conditioned/dependent on a sequence of actions



# Policy-Conditioned Returns

Recall the return from trajectory optimization

$$[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

This is an **action-conditioned** discounted return

Conditioned/dependent on a sequence of actions

There is no structure to the actions

# Policy-Conditioned Returns

Recall the return from trajectory optimization

$$[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

This is an **action-conditioned** discounted return

Conditioned/dependent on a sequence of actions

There is no structure to the actions

- Random

# Policy-Conditioned Returns

Recall the return from trajectory optimization

$$[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

This is an **action-conditioned** discounted return

Conditioned/dependent on a sequence of actions

There is no structure to the actions

- Random
- Picked by humans

# Policy-Conditioned Returns

Recall the return from trajectory optimization

$$[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

This is an **action-conditioned** discounted return

Conditioned/dependent on a sequence of actions

There is no structure to the actions

- Random
- Picked by humans
- Maximize  $\mathcal{G}$

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

Last time, we introduced the policy

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

Last time, we introduced the policy

$$\pi : S \times \Theta \mapsto \Delta A$$

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

Last time, we introduced the policy

$$\pi : S \times \Theta \mapsto \Delta A$$

Example policy, greedy policy



# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

Last time, we introduced the policy

$$\pi : S \times \Theta \mapsto \Delta A$$

Example policy, greedy policy

$$\pi(a_t \mid s_t; \theta_\pi) = \begin{cases} 1 & \text{if } a_t = \arg \max_{a_t \in A} \mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] \\ 0 & \text{otherwise} \end{cases}$$

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

Last time, we introduced the policy

$$\pi : S \times \Theta \mapsto \Delta A$$

Example policy, greedy policy

$$\pi(a_t \mid s_t; \theta_\pi) = \begin{cases} 1 & \text{if } a_t = \arg \max_{a_t \in A} \mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] \\ 0 & \text{otherwise} \end{cases}$$

# Policy-Conditioned Returns

$$[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

# Policy-Conditioned Returns

$$[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

$$\pi : S \times \Theta \mapsto \Delta A$$

# Policy-Conditioned Returns

$$[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

$$\pi : S \times \Theta \mapsto \Delta A$$

Conditioning the return on actions is annoying

# Policy-Conditioned Returns

$$[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

$$\pi : S \times \Theta \mapsto \Delta A$$

Conditioning the return on actions is annoying

Must compute infinitely many actions and outcomes for the return

# Policy-Conditioned Returns

$$[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

$$\pi : S \times \Theta \mapsto \Delta A$$

Conditioning the return on actions is annoying

Must compute infinitely many actions and outcomes for the return

What if we condition on a policy, instead of specific actions?

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

$$a_0 \sim \pi(\cdot \mid s_0; \theta_{\pi}), \quad a_1 \sim \pi(\cdot \mid s_1; \theta_{\pi}), \quad a_2 \sim \pi(\cdot \mid s_2; \theta_{\pi}), \quad \dots$$



# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

$$a_0 \sim \pi(\cdot \mid s_0; \theta_{\pi}), \quad a_1 \sim \pi(\cdot \mid s_1; \theta_{\pi}), \quad a_2 \sim \pi(\cdot \mid s_2; \theta_{\pi}), \quad \dots$$

Condition on a function parameterized by  $\theta_{\pi}$  instead of many actions

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

$$a_0 \sim \pi(\cdot \mid s_0; \theta_{\pi}), \quad a_1 \sim \pi(\cdot \mid s_1; \theta_{\pi}), \quad a_2 \sim \pi(\cdot \mid s_2; \theta_{\pi}), \quad \dots$$

Condition on a function parameterized by  $\theta_{\pi}$  instead of many actions

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \theta_{\pi}] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, \theta_{\pi}]$$

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

$$a_0 \sim \pi(\cdot \mid s_0; \theta_{\pi}), \quad a_1 \sim \pi(\cdot \mid s_1; \theta_{\pi}), \quad a_2 \sim \pi(\cdot \mid s_2; \theta_{\pi}), \quad \dots$$

Condition on a function parameterized by  $\theta_{\pi}$  instead of many actions

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \theta_{\pi}] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, \theta_{\pi}]$$

The function outputs a distribution over the action space  $\pi(a \mid s; \theta_{\pi})$

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \theta_{\pi}] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, \theta_{\pi}]$$

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \theta_{\pi}] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, \theta_{\pi}]$$

Now conditioned on the policy

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \theta_{\pi}] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, \theta_{\pi}]$$

Now conditioned on the policy

But remember,  $\mathcal{R}(s_{t+1})$  hides much of the magic

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, a_0, a_1, \dots]$$

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \theta_{\pi}] = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, \theta_{\pi}]$$

Now conditioned on the policy

But remember,  $\mathcal{R}(s_{t+1})$  hides much of the magic

How does  $\mathbb{E}[\mathcal{R}(s_{t+1})]$  change when we condition on  $\theta_{\pi}$ ?



# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathcal{R}(s_{t+1}) \sum_{s_{t+1} \in S} \Pr(s_{t+1} \mid s_0, a_0, \dots, a_t)$$

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathcal{R}(s_{t+1}) \sum_{s_{t+1} \in S} \Pr(s_{t+1} \mid s_0, a_0, \dots, a_t)$$

**Question:** What changes when we condition on  $\theta_\pi$ ?

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathcal{R}(s_{t+1}) \sum_{s_{t+1} \in S} \Pr(s_{t+1} \mid s_0, a_0, \dots, a_t)$$

**Question:** What changes when we condition on  $\theta_\pi$ ?

$$\Pr(s_{t+1} \mid s_0; \theta_\pi)$$

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathcal{R}(s_{t+1}) \sum_{s_{t+1} \in S} \Pr(s_{t+1} \mid s_0, a_0, \dots, a_t)$$

**Question:** What changes when we condition on  $\theta_\pi$ ?

$$\Pr(s_{t+1} \mid s_0; \theta_\pi)$$

Maybe we can use  $\Pr(s_{t+1} \mid s_t, a_t)$  to figure this out

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathcal{R}(s_{t+1}) \sum_{s_{t+1} \in S} \Pr(s_{t+1} \mid s_0, a_0, \dots, a_t)$$

**Question:** What changes when we condition on  $\theta_\pi$ ?

$$\Pr(s_{t+1} \mid s_0; \theta_\pi)$$

Maybe we can use  $\Pr(s_{t+1} \mid s_t, a_t)$  to figure this out

**Question:** What was  $\Pr(s_{t+1} \mid s_t, a_t)$ ?

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0, a_0, a_1, \dots] = \sum_{t=0}^{\infty} \gamma^t \mathcal{R}(s_{t+1}) \sum_{s_{t+1} \in S} \Pr(s_{t+1} \mid s_0, a_0, \dots, a_t)$$

**Question:** What changes when we condition on  $\theta_\pi$ ?

$$\Pr(s_{t+1} \mid s_0; \theta_\pi)$$

Maybe we can use  $\Pr(s_{t+1} \mid s_t, a_t)$  to figure this out

**Question:** What was  $\Pr(s_{t+1} \mid s_t, a_t)$ ?

**Answer:** State transition function

$$\Pr(s_{t+1} \mid s_t, a_t)$$

# Policy-Conditioned Returns

$$\text{Tr}(s_{t+1} \mid s_t, a_t)$$

# Policy-Conditioned Returns

$$\text{Tr}(s_{t+1} \mid s_t, a_t)$$

**Issue:** State transition function needs an action  $a_t$



# Policy-Conditioned Returns

$$\text{Tr}(s_{t+1} \mid s_t, a_t)$$

**Issue:** State transition function needs an action  $a_t$

Policy  $\pi$  outputs a distribution over the action space

# Policy-Conditioned Returns

$$\text{Tr}(s_{t+1} \mid s_t, a_t)$$

**Issue:** State transition function needs an action  $a_t$

Policy  $\pi$  outputs a distribution over the action space

**Question:** What is  $\text{Pr}(s_{t+1} \mid s_t, \theta_\pi)$ ?

# Policy-Conditioned Returns

$$\text{Tr}(s_{t+1} \mid s_t, a_t)$$

**Issue:** State transition function needs an action  $a_t$

Policy  $\pi$  outputs a distribution over the action space

**Question:** What is  $\text{Pr}(s_{t+1} \mid s_t, \theta_\pi)$ ? Hint: Consider all possible actions

# Policy-Conditioned Returns

$$\text{Tr}(s_{t+1} \mid s_t, a_t)$$

**Issue:** State transition function needs an action  $a_t$

Policy  $\pi$  outputs a distribution over the action space

**Question:** What is  $\text{Pr}(s_{t+1} \mid s_t, \theta_\pi)$ ? Hint: Consider all possible actions

$$\text{Pr}(s_{t+1} \mid s_t; \theta_\pi) = \sum_{a_t \in A} \text{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

# Policy-Conditioned Returns

$$\text{Tr}(s_{t+1} \mid s_t, a_t)$$

**Issue:** State transition function needs an action  $a_t$

Policy  $\pi$  outputs a distribution over the action space

**Question:** What is  $\text{Pr}(s_{t+1} \mid s_t, \theta_\pi)$ ? Hint: Consider all possible actions

$$\text{Pr}(s_{t+1} \mid s_t; \theta_\pi) = \sum_{a_t \in A} \text{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

Combine the policy distribution with next state distribution

# Policy-Conditioned Returns

$$\Pr(s_{t+1} \mid s_t; \theta_\pi) = \sum_{a_t \in A} \text{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

# Policy-Conditioned Returns

$$\Pr(s_{t+1} \mid s_t; \theta_\pi) = \sum_{a_t \in A} \text{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

Write out the first few timesteps

# Policy-Conditioned Returns

$$\Pr(s_{t+1} \mid s_t; \theta_\pi) = \sum_{a_t \in A} \text{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

Write out the first few timesteps

$$\Pr(s_1 \mid s_0; \theta_\pi) = \sum_{a_0 \in A} \text{Tr}(s_1 \mid s_0, a_0) \cdot \pi(a_0 \mid s_0; \theta_\pi)$$



# Policy-Conditioned Returns

$$\Pr(s_{t+1} \mid s_t; \theta_\pi) = \sum_{a_t \in A} \text{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

Write out the first few timesteps

$$\Pr(s_1 \mid s_0; \theta_\pi) = \sum_{a_0 \in A} \text{Tr}(s_1 \mid s_0, a_0) \cdot \pi(a_0 \mid s_0; \theta_\pi)$$

$$\begin{aligned} \Pr(s_2 \mid s_0; \theta_\pi) &= \sum_{s_1 \in S} \sum_{a_1 \in A} \text{Tr}(s_2 \mid s_1, a_1) \cdot \pi(a_1 \mid s_1; \theta_\pi) \\ &\quad \cdot \sum_{a_0 \in A} \text{Tr}(s_1 \mid s_0, a_0) \cdot \pi(a_0 \mid s_0; \theta_\pi) \end{aligned}$$

# Policy-Conditioned Returns

$$\Pr(s_1 \mid s_0; \theta_\pi) = \sum_{a_0 \in A} \text{Tr}(s_1 \mid s_0, a_0) \cdot \pi(a_0 \mid s_0; \theta_\pi)$$

$$\begin{aligned} \Pr(s_2 \mid s_0; \theta_\pi) &= \sum_{s_1 \in S} \sum_{a_1 \in A} \text{Tr}(s_2 \mid s_1, a_1) \cdot \pi(a_1 \mid s_1; \theta_\pi) \\ &\quad \cdot \sum_{a_0 \in A} \text{Tr}(s_1 \mid s_0, a_0) \cdot \pi(a_0 \mid s_0; \theta_\pi) \end{aligned}$$

# Policy-Conditioned Returns

$$\Pr(s_1 \mid s_0; \theta_\pi) = \sum_{a_0 \in A} \text{Tr}(s_1 \mid s_0, a_0) \cdot \pi(a_0 \mid s_0; \theta_\pi)$$

$$\begin{aligned} \Pr(s_2 \mid s_0; \theta_\pi) &= \sum_{s_1 \in S} \sum_{a_1 \in A} \text{Tr}(s_2 \mid s_1, a_1) \cdot \pi(a_1 \mid s_1; \theta_\pi) \\ &\quad \cdot \sum_{a_0 \in A} \text{Tr}(s_1 \mid s_0, a_0) \cdot \pi(a_0 \mid s_0; \theta_\pi) \end{aligned}$$

Derive a general form for  $\Pr(s_{n+1} \mid s_0; \theta_\pi)$

# Policy-Conditioned Returns

$$\Pr(s_1 \mid s_0; \theta_\pi) = \sum_{a_0 \in A} \text{Tr}(s_1 \mid s_0, a_0) \cdot \pi(a_0 \mid s_0; \theta_\pi)$$

$$\begin{aligned} \Pr(s_2 \mid s_0; \theta_\pi) &= \sum_{s_1 \in S} \sum_{a_1 \in A} \text{Tr}(s_2 \mid s_1, a_1) \cdot \pi(a_1 \mid s_1; \theta_\pi) \\ &\quad \cdot \sum_{a_0 \in A} \text{Tr}(s_1 \mid s_0, a_0) \cdot \pi(a_0 \mid s_0; \theta_\pi) \end{aligned}$$

Derive a general form for  $\Pr(s_{n+1} \mid s_0; \theta_\pi)$

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \left( \sum_{a_t \in A} \text{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta) \right)$$

# Policy-Conditioned Returns

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \left( \sum_{a_t \in A} \text{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta) \right)$$

# Policy-Conditioned Returns

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \left( \sum_{a_t \in A} \text{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta) \right)$$

Plug back into our expected reward

# Policy-Conditioned Returns

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \left( \sum_{a_t \in A} \text{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta) \right)$$

Plug back into our expected reward

$$\mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_\pi] = \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_\pi] = \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$



# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_\pi] = \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

Discounted return is discounted sum of rewards

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_\pi] = \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

Discounted return is discounted sum of rewards

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] =$$

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_\pi] = \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

Discounted return is discounted sum of rewards

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi)$$

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_\pi] = \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

Discounted return is discounted sum of rewards

$$\begin{aligned} \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] &= \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ &\quad + \gamma \sum_{s_2 \in S} \mathcal{R}(s_2) \cdot \Pr(s_2 \mid s_0; \theta_\pi) \end{aligned}$$

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_\pi] = \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

Discounted return is discounted sum of rewards

$$\begin{aligned} \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] &= \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ &\quad + \gamma \sum_{s_2 \in S} \mathcal{R}(s_2) \cdot \Pr(s_2 \mid s_0; \theta_\pi) \\ &\quad + \gamma^2 \sum_{s_3 \in S} \mathcal{R}(s_3) \cdot \Pr(s_3 \mid s_0; \theta_\pi) \end{aligned}$$

# Policy-Conditioned Returns

$$\mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_\pi] = \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

Discounted return is discounted sum of rewards

$$\begin{aligned} \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] &= \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ &\quad + \gamma \sum_{s_2 \in S} \mathcal{R}(s_2) \cdot \Pr(s_2 \mid s_0; \theta_\pi) \\ &\quad + \gamma^2 \sum_{s_3 \in S} \mathcal{R}(s_3) \cdot \Pr(s_3 \mid s_0; \theta_\pi) \\ &\quad \dots \end{aligned}$$

# Policy-Conditioned Returns

**Definition:** General form of policy-conditioned discounted return

# Policy-Conditioned Returns

**Definition:** General form of policy-conditioned discounted return

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_n, a_n)$$



# Policy-Conditioned Returns

**Definition:** General form of policy-conditioned discounted return

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_n, a_n)$$

Where the state distribution is

# Policy-Conditioned Returns

**Definition:** General form of policy-conditioned discounted return

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_n, a_n)$$

Where the state distribution is

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \left( \sum_{a_t \in A} \Pr(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta) \right)$$

# Value Functions

---

# Value Functions

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{a_0, \dots, a_n \in A} \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \text{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

# Value Functions

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{a_0, \dots, a_n \in A} \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \text{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

# Value Functions

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{a_0, \dots, a_n \in A} \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \text{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

These two equations form the basis of all reinforcement learning

# Value Functions

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{a_0, \dots, a_n \in A} \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \text{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

These two equations form the basis of all reinforcement learning

**Goal:** find the  $\theta_\pi$  (policy parameters) to maximize the expected return

# Value Functions

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{a_0, \dots, a_n \in A} \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \text{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$



# Value Functions

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{a_0, \dots, a_n \in A} \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \text{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

We have another name for  $\mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi]$

# Value Functions

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{a_0, \dots, a_n \in A} \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \text{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

We have another name for  $\mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi]$

We call it the **value function**  $V : S \times \Theta \mapsto \mathbb{R}$

# Value Functions

$$\Pr(s_{n+1} \mid s_0; \theta_\pi) = \sum_{a_0, \dots, a_n \in A} \sum_{s_1, \dots, s_n \in S} \prod_{t=0}^n \text{Tr}(s_{t+1} \mid s_t, a_t) \cdot \pi(a_t \mid s_t; \theta_\pi)$$

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

We have another name for  $\mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi]$

We call it the **value function**  $V : S \times \Theta \mapsto \mathbb{R}$

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

# Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

# Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

# Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

Valuable states lead to good returns **under the current policy**

# Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

Valuable states lead to good returns **under the current policy**

$s = 240$  km/h

# Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

Valuable states lead to good returns **under the current policy**

$s = 240$  km/h

$\theta_\pi =$  Race car driver



# Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

Valuable states lead to good returns **under the current policy**

$s = 240 \text{ km/h}$                        $\theta_\pi = \text{Race car driver}$                        $V(s, \theta_\pi) = \text{good}$

# Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

Valuable states lead to good returns **under the current policy**

$s = 240 \text{ km/h}$                        $\theta_\pi = \text{Race car driver}$                        $V(s, \theta_\pi) = \text{good}$

$s = 240 \text{ km/h}$

# Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

Valuable states lead to good returns **under the current policy**

$s = 240 \text{ km/h}$                        $\theta_\pi = \text{Race car driver}$                        $V(s, \theta_\pi) = \text{good}$

$s = 240 \text{ km/h}$                        $\theta_\pi = \text{Grandpa}$

# Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

Valuable states lead to good returns **under the current policy**

$s = 240 \text{ km/h}$	$\theta_\pi = \text{Race car driver}$	$V(s, \theta_\pi) = \text{good}$
------------------------	---------------------------------------	----------------------------------

$s = 240 \text{ km/h}$	$\theta_\pi = \text{Grandpa}$	$V(s, \theta_\pi) = \text{not good}$
------------------------	-------------------------------	--------------------------------------

# Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

Valuable states lead to good returns **under the current policy**

$s = 240 \text{ km/h}$                        $\theta_\pi = \text{Race car driver}$                        $V(s, \theta_\pi) = \text{good}$

$s = 240 \text{ km/h}$                        $\theta_\pi = \text{Grandpa}$                        $V(s, \theta_\pi) = \text{not good}$

We use the value function to direct the policy to good states

# Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Value function takes any state  $s_0$ , and tells us how valuable  $s_0$  is

Valuable states lead to good returns **under the current policy**

$s = 240$  km/h                       $\theta_\pi =$  Race car driver                       $V(s, \theta_\pi) =$  good

$s = 240$  km/h                       $\theta_\pi =$  Grandpa                       $V(s, \theta_\pi) =$  not good

We use the value function to direct the policy to good states

It is a critical part of decision making

# Value Functions

With the value function, we can use any state as a starting state

# Value Functions

With the value function, we can use any state as a starting state

The state does not need to be the start of a trajectory



# Value Functions

With the value function, we can use any state as a starting state

The state does not need to be the start of a trajectory

**Example:** Consider the sequence of states

$$s_a, s_b, s_c$$

# Value Functions

With the value function, we can use any state as a starting state

The state does not need to be the start of a trajectory

**Example:** Consider the sequence of states

$$s_a, s_b, s_c$$

We can compute

$$V(s_a, \theta_\pi), V(s_b, \theta_\pi), V(s_c, \theta_\pi)$$

To find the value of any state

# Value Functions

**Question:** Why does Prof. Steven keep showing stupid equations? How is the value function useful?

# Value Functions

**Question:** Why does Prof. Steven keep showing stupid equations? How is the value function useful?

We can use the value of a state to make decisions

# Value Functions

**Question:** Why does Prof. Steven keep showing stupid equations? How is the value function useful?

We can use the value of a state to make decisions

$$s_a = \text{Live in Macau}, s_b = \text{Live in Beijing}$$

# Value Functions

**Question:** Why does Prof. Steven keep showing stupid equations? How is the value function useful?

We can use the value of a state to make decisions

$$s_a = \text{Live in Macau}, s_b = \text{Live in Beijing}$$

Given all your preferences ( $\mathcal{R}$ ) and thoughts ( $\theta_\pi$ ), we can determine which life is better for you

# Value Functions

**Question:** Why does Prof. Steven keep showing stupid equations? How is the value function useful?

We can use the value of a state to make decisions

$$s_a = \text{Live in Macau}, s_b = \text{Live in Beijing}$$

Given all your preferences ( $\mathcal{R}$ ) and thoughts ( $\theta_\pi$ ), we can determine which life is better for you

$V(s, \theta_\pi)$  considers your future friends, income, wife/husband, etc

# Value Functions

**Question:** Why does Prof. Steven keep showing stupid equations? How is the value function useful?

We can use the value of a state to make decisions

$$s_a = \text{Live in Macau}, s_b = \text{Live in Beijing}$$

Given all your preferences ( $\mathcal{R}$ ) and thoughts ( $\theta_\pi$ ), we can determine which life is better for you

$V(s, \theta_\pi)$  considers your future friends, income, wife/husband, etc

Combines all this info into one value, a single number of “goodness”



# Value Functions

**Question:** Why does Prof. Steven keep showing stupid equations? How is the value function useful?

We can use the value of a state to make decisions

$$s_a = \text{Live in Macau}, s_b = \text{Live in Beijing}$$

Given all your preferences ( $\mathcal{R}$ ) and thoughts ( $\theta_\pi$ ), we can determine which life is better for you

$V(s, \theta_\pi)$  considers your future friends, income, wife/husband, etc

Combines all this info into one value, a single number of “goodness”

$$V(s_a, \theta_\pi) = 1032$$

# Value Functions

**Question:** Why does Prof. Steven keep showing stupid equations? How is the value function useful?

We can use the value of a state to make decisions

$$s_a = \text{Live in Macau}, s_b = \text{Live in Beijing}$$

Given all your preferences ( $\mathcal{R}$ ) and thoughts ( $\theta_\pi$ ), we can determine which life is better for you

$V(s, \theta_\pi)$  considers your future friends, income, wife/husband, etc

Combines all this info into one value, a single number of “goodness”

$$V(s_a, \theta_\pi) = 1032$$

$$V(s_b, \theta_\pi) = 945$$

# Value Functions

**Question:** Why does Prof. Steven keep showing stupid equations? How is the value function useful?

We can use the value of a state to make decisions

$$s_a = \text{Live in Macau}, s_b = \text{Live in Beijing}$$

Given all your preferences ( $\mathcal{R}$ ) and thoughts ( $\theta_\pi$ ), we can determine which life is better for you

$V(s, \theta_\pi)$  considers your future friends, income, wife/husband, etc

Combines all this info into one value, a single number of “goodness”

$$V(s_a, \theta_\pi) = 1032$$

$$V(s_b, \theta_\pi) = 945$$

# Value Functions

$s_a$  = Live in Macau,  $s_b$  = Live in Beijing

# Value Functions

$s_a$  = Live in Macau,  $s_b$  = Live in Beijing

$$V(s_a, \theta_\pi) = 1032$$

$$V(s_b, \theta_\pi) = 945$$

# Value Functions

$s_a$  = Live in Macau,  $s_b$  = Live in Beijing

$$V(s_a, \theta_\pi) = 1032$$

$$V(s_b, \theta_\pi) = 945$$

This value leads us to the right decisions

# Value Functions

$s_a$  = Live in Macau,  $s_b$  = Live in Beijing

$$V(s_a, \theta_\pi) = 1032$$

$$V(s_b, \theta_\pi) = 945$$

This value leads us to the right decisions

Some optimal decisions are hard for humans to make

# Value Functions

$s_a$  = Live in Macau,  $s_b$  = Live in Beijing

$$V(s_a, \theta_\pi) = 1032$$

$$V(s_b, \theta_\pi) = 945$$

This value leads us to the right decisions

Some optimal decisions are hard for humans to make

With value, we can be sure we make the right decision



# Exercise

---

# Exercise

- Think of two places you want to live after graduation  $s_a, s_b$

# Exercise

- Think of two places you want to live after graduation  $s_a, s_b$
- Consider your behavior ( $\theta_\pi$ ) and what is important to you ( $\mathcal{R}$ )

# Exercise

- Think of two places you want to live after graduation  $s_a, s_b$
- Consider your behavior ( $\theta_\pi$ ) and what is important to you ( $\mathcal{R}$ )
- 3 life goals as states  $s_x, s_y, s_z \in G$  (e.g., friends, money, hobby, etc)

# Exercise

- Think of two places you want to live after graduation  $s_a, s_b$
- Consider your behavior ( $\theta_\pi$ ) and what is important to you ( $\mathcal{R}$ )
- 3 life goals as states  $s_x, s_y, s_z \in G$  (e.g., friends, money, hobby, etc)
- Assign a reward  $\mathcal{R}$  for each goal, and choose discount factor  $\gamma$

# Exercise

- Think of two places you want to live after graduation  $s_a, s_b$
- Consider your behavior ( $\theta_\pi$ ) and what is important to you ( $\mathcal{R}$ )
- 3 life goals as states  $s_x, s_y, s_z \in G$  (e.g., friends, money, hobby, etc)
- Assign a reward  $\mathcal{R}$  for each goal, and choose discount factor  $\gamma$

For each location  $s_0 \in \{s_a, s_b\}$ :

# Exercise

- Think of two places you want to live after graduation  $s_a, s_b$
- Consider your behavior ( $\theta_\pi$ ) and what is important to you ( $\mathcal{R}$ )
- 3 life goals as states  $s_x, s_y, s_z \in G$  (e.g., friends, money, hobby, etc)
- Assign a reward  $\mathcal{R}$  for each goal, and choose discount factor  $\gamma$

For each location  $s_0 \in \{s_a, s_b\}$ :

- Estimate probability of reaching each goal  $\Pr(s_g \mid s_0); s_g \in \{s_x, s_y, s_z\}$

# Exercise

- Think of two places you want to live after graduation  $s_a, s_b$
- Consider your behavior ( $\theta_\pi$ ) and what is important to you ( $\mathcal{R}$ )
- 3 life goals as states  $s_x, s_y, s_z \in G$  (e.g., friends, money, hobby, etc)
- Assign a reward  $\mathcal{R}$  for each goal, and choose discount factor  $\gamma$

For each location  $s_0 \in \{s_a, s_b\}$ :

- Estimate probability of reaching each goal  $\Pr(s_g \mid s_0); s_g \in \{s_x, s_y, s_z\}$
- Estimate time to accomplish each goal  $t = \dots$



# Exercise

- Think of two places you want to live after graduation  $s_a, s_b$
- Consider your behavior ( $\theta_\pi$ ) and what is important to you ( $\mathcal{R}$ )
- 3 life goals as states  $s_x, s_y, s_z \in G$  (e.g., friends, money, hobby, etc)
- Assign a reward  $\mathcal{R}$  for each goal, and choose discount factor  $\gamma$

For each location  $s_0 \in \{s_a, s_b\}$ :

- Estimate probability of reaching each goal  $\Pr(s_g \mid s_0); s_g \in \{s_x, s_y, s_z\}$
- Estimate time to accomplish each goal  $t = \dots$

$$V(s_0, \theta_\pi) = \sum_{s_g \in \{s_x, s_y, s_z\}} \gamma^{t_g} \mathcal{R}(s_g) \cdot \Pr(s_g \mid s_0; \theta_\pi)$$

# TD Value Functions

---

# TD Value Functions

**Note:** We define the value function in many different ways

# TD Value Functions

**Note:** We define the value function in many different ways

It always approximates the expected discounted return from  $s_0$

# TD Value Functions

**Note:** We define the value function in many different ways

It always approximates the expected discounted return from  $s_0$

We call the following equation the **Monte Carlo** value function

$$V(s_0, \theta_\pi) = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

# TD Value Functions

**Note:** We define the value function in many different ways

It always approximates the expected discounted return from  $s_0$

We call the following equation the **Monte Carlo** value function

$$V(s_0, \theta_\pi) = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Difficult to compute the Monte Carlo value function

# TD Value Functions

**Note:** We define the value function in many different ways

It always approximates the expected discounted return from  $s_0$

We call the following equation the **Monte Carlo** value function

$$V(s_0, \theta_\pi) = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Difficult to compute the Monte Carlo value function

- Must either have a terminal states

# TD Value Functions

**Note:** We define the value function in many different ways

It always approximates the expected discounted return from  $s_0$

We call the following equation the **Monte Carlo** value function

$$V(s_0, \theta_\pi) = \sum_{n=0}^{\infty} \gamma^n \sum_{s_{n+1} \in S} \mathcal{R}(s_{n+1}) \cdot \Pr(s_{n+1} \mid s_0; \theta_\pi)$$

Difficult to compute the Monte Carlo value function

- Must either have a terminal states
- Or build the infinite decision tree



# TD Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

# TD Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

Infinite sums make things difficult and intractable

# TD Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

Infinite sums make things difficult and intractable

Let us try to delete the infinite sum

# TD Value Functions

$$V(s_0, \theta_\pi) = \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

# TD Value Functions

$$V(s_0, \theta_\pi) = \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

Factor out initial timestep  $t = 0$  out of the outer sum

# TD Value Functions

$$V(s_0, \theta_\pi) = \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

Factor out initial timestep  $t = 0$  out of the outer sum

$$\begin{aligned} V(s_0, \theta_\pi) &= \gamma^0 \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ &+ \sum_{t=1}^{\infty} \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi) \end{aligned}$$

# TD Value Functions

$$\begin{aligned} V(s_0, \theta_\pi) &= \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ &+ \sum_{t=1}^{\infty} \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi) \end{aligned}$$

# TD Value Functions

$$\begin{aligned} V(s_0, \theta_\pi) &= \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ &+ \sum_{t=1}^{\infty} \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi) \end{aligned}$$

Rewrite sum starting from  $t = 0$



# TD Value Functions

$$\begin{aligned} V(s_0, \theta_\pi) &= \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ &+ \sum_{t=1}^{\infty} \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi) \end{aligned}$$

Rewrite sum starting from  $t = 0$

$$\begin{aligned} V(s_0, \theta_\pi) &= \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ &+ \sum_{t=0}^{\infty} \gamma^{t+1} \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \cdot \Pr(s_{t+2} \mid s_0; \theta_\pi) \end{aligned}$$

# TD Value Functions

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ + \sum_{t=0}^{\infty} \gamma^{t+1} \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \cdot \Pr(s_{t+2} \mid s_0; \theta_\pi)$$

# TD Value Functions

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ + \sum_{t=0}^{\infty} \gamma^{t+1} \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \cdot \Pr(s_{t+2} \mid s_0; \theta_\pi)$$

Factor out  $\gamma$

# TD Value Functions

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ + \sum_{t=0}^{\infty} \gamma^{t+1} \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \cdot \Pr(s_{t+2} \mid s_0; \theta_\pi)$$

Factor out  $\gamma$

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ + \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \cdot \Pr(s_{t+2} \mid s_0; \theta_\pi)$$

# TD Value Functions

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ + \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \cdot \Pr(s_{t+2} \mid s_0; \theta_\pi)$$

# TD Value Functions

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ + \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \cdot \Pr(s_{t+2} \mid s_0; \theta_\pi)$$

Split Pr using Markov property

# TD Value Functions

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ + \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \cdot \Pr(s_{t+2} \mid s_0; \theta_\pi)$$

Split Pr using Markov property

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ + \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \sum_{s_1} \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi) \Pr(s_1 \mid s_0; \theta_\pi)$$

# TD Value Functions

$$\begin{aligned} V(s_0, \theta_\pi) &= \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ &+ \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \sum_{s_1} \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi) \Pr(s_1 \mid s_0; \theta_\pi) \end{aligned}$$



# TD Value Functions

$$\begin{aligned} V(s_0, \theta_\pi) = & \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ & + \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \sum_{s_1} \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi) \Pr(s_1 \mid s_0; \theta_\pi) \end{aligned}$$

Move sum and Pr outside

# TD Value Functions

$$\begin{aligned} V(s_0, \theta_\pi) &= \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ &+ \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \sum_{s_1} \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi) \Pr(s_1 \mid s_0; \theta_\pi) \end{aligned}$$

Move sum and Pr outside

$$\begin{aligned} V(s_0, \theta_\pi) &= \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ &+ \sum_{s_1} \Pr(s_1 \mid s_0; \theta_\pi) \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi) \end{aligned}$$

# TD Value Functions

$$\begin{aligned} V(s_0, \theta_\pi) &= \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ &+ \sum_{s_1} \Pr(s_1 \mid s_0; \theta_\pi) \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi) \end{aligned}$$

# TD Value Functions

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ + \sum_{s_1} \Pr(s_1 \mid s_0; \theta_\pi) \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi)$$

**Question:** What is this term?

# TD Value Functions

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ + \sum_{s_1} \Pr(s_1 \mid s_0; \theta_\pi) \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi)$$

**Question:** What is this term?

$$V(s_0, \theta_\pi) = \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

# TD Value Functions

$$V(s_0, \theta_\pi) = \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ + \sum_{s_1} \Pr(s_1 \mid s_0; \theta_\pi) \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi)$$

**Question:** What is this term?

$$V(s_0, \theta_\pi) = \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+1} \in S} \mathcal{R}(s_{t+1}) \cdot \Pr(s_{t+1} \mid s_0; \theta_\pi)$$

$$V(s_1, \theta_\pi) = \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \cdot \Pr(s_{t+2} \mid s_1; \theta_\pi)$$

# TD Value Functions

$$\begin{aligned} V(s_0, \theta_\pi) &= \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ &+ \sum_{s_1} \Pr(s_1 \mid s_0; \theta_\pi) \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi) \end{aligned}$$

# TD Value Functions

$$\begin{aligned} V(s_0, \theta_\pi) &= \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ &+ \sum_{s_1} \Pr(s_1 \mid s_0; \theta_\pi) \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi) \\ V(s_0, \theta_\pi) &= \left( \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \right) + \gamma V(s_1, \theta_\pi) \end{aligned}$$



# TD Value Functions

$$\begin{aligned} V(s_0, \theta_\pi) &= \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ &+ \sum_{s_1} \Pr(s_1 \mid s_0; \theta_\pi) \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi) \end{aligned}$$

$$V(s_0, \theta_\pi) = \left( \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \right) + \gamma V(s_1, \theta_\pi)$$

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

# TD Value Functions

$$\begin{aligned} V(s_0, \theta_\pi) &= \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \\ &+ \sum_{s_1} \Pr(s_1 \mid s_0; \theta_\pi) \gamma \sum_{t=0}^{\infty} \gamma^t \sum_{s_{t+2} \in S} \mathcal{R}(s_{t+2}) \Pr(s_{t+2} \mid s_{t+1}; \theta_\pi) \end{aligned}$$

$$V(s_0, \theta_\pi) = \left( \sum_{s_1 \in S} \mathcal{R}(s_1) \cdot \Pr(s_1 \mid s_0; \theta_\pi) \right) + \gamma V(s_1, \theta_\pi)$$

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

This is a huge finding!

# TD Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

# TD Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

Value function has a recursive definition

# TD Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

Value function has a recursive definition

Represent policy-conditioned discounted return without an infinite sum

# TD Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

Value function has a recursive definition

Represent policy-conditioned discounted return without an infinite sum

We call this the **Temporal Difference** (TD) value function

# TD Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

Value function has a recursive definition

Represent policy-conditioned discounted return without an infinite sum

We call this the **Temporal Difference** (TD) value function

Compute the return with a single transition  $s_0 \rightarrow s_1$

# TD Value Functions

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

Value function has a recursive definition

Represent policy-conditioned discounted return without an infinite sum

We call this the **Temporal Difference** (TD) value function

Compute the return with a single transition  $s_0 \rightarrow s_1$

Evaluate infinite-depth decision tree with a single function call



# TD Value Functions

To summarize, we can represent the value function in two ways:

# TD Value Functions

To summarize, we can represent the value function in two ways:

The Monte Carlo value function

# TD Value Functions

To summarize, we can represent the value function in two ways:

The Monte Carlo value function

$$V(s_0, \theta_\pi) = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, \theta_\pi]$$

# TD Value Functions

To summarize, we can represent the value function in two ways:

The Monte Carlo value function

$$V(s_0, \theta_\pi) = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, \theta_\pi]$$

The Temporal Difference value function

# TD Value Functions

To summarize, we can represent the value function in two ways:

The Monte Carlo value function

$$V(s_0, \theta_\pi) = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0, \theta_\pi]$$

The Temporal Difference value function

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

They produce the same result, but with different computation

# Q Functions

---

# Q Functions

We saw two forms of the value function

# Q Functions

We saw two forms of the value function

The value function relies on a policy



# Q Functions

We saw two forms of the value function

The value function relies on a policy

But our goal was to find a policy, so how does value help?

# Q Functions

We saw two forms of the value function

The value function relies on a policy

But our goal was to find a policy, so how does value help?

There is a special connection between an optimal policy and the value function

# Q Functions

We saw two forms of the value function

The value function relies on a policy

But our goal was to find a policy, so how does value help?

There is a special connection between an optimal policy and the value function

We can use the value function to find an optimal policy

# Q Functions

Consider the Temporal Difference value function

# Q Functions

Consider the Temporal Difference value function

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

# Q Functions

Consider the Temporal Difference value function

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

We conditioned the value function on policy parameters

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi]$$

# Q Functions

Consider the Temporal Difference value function

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

We conditioned the value function on policy parameters

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi]$$

With trajectory optimization we conditioned on actions

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0, a_0, a_1, \dots]$$

# Q Functions

Consider the Temporal Difference value function

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi] = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

We conditioned the value function on policy parameters

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0; \theta_\pi]$$

With trajectory optimization we conditioned on actions

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0, a_0, a_1, \dots]$$

What if we wanted a mix of both?

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0, a_0; \theta_\pi]$$



# Q Functions

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0; \boldsymbol{\theta}_\pi]$$

# Q Functions

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0; \boldsymbol{\theta}_\pi]$$

Means:

# Q Functions

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0, a_0; \theta_\pi]$$

Means:

- Take a specific action  $a_0$  (trajectory optimization)

# Q Functions

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0; \theta_\pi]$$

Means:

- Take a specific action  $a_0$  (trajectory optimization)
- Follow  $\pi(a \mid s; \theta_\pi)$  for all future actions  $a_1, a_2, \dots$  (value function)

# Q Functions

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0; \boldsymbol{\theta}_\pi]$$

# Q Functions

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0, a_0; \theta_\pi]$$

We call this the **Q function**

# Q Functions

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0; \theta_\pi]$$

We call this the **Q function**

$$Q(s, a, \theta_\pi) = \mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0; \theta_\pi]$$

# Q Functions

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0; \theta_\pi]$$

We call this the **Q function**

$$Q(s, a, \theta_\pi) = \mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0; \theta_\pi]$$

We can derive the Q function from the value function



# Q Functions

$$\mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0; \theta_\pi]$$

We call this the **Q function**

$$Q(s, a, \theta_\pi) = \mathbb{E}[\mathcal{G}(\boldsymbol{\tau}) \mid s_0, a_0; \theta_\pi]$$

We can derive the Q function from the value function

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

# Q Functions

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0, a_0; \theta_\pi]$$

We call this the **Q function**

$$Q(s, a, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0, a_0; \theta_\pi]$$

We can derive the Q function from the value function

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

First, introduce the action  $a_0$

# Q Functions

$$\mathbb{E}[\mathcal{G}(\tau) \mid s_0, a_0; \theta_\pi]$$

We call this the **Q function**

$$Q(s, a, \theta_\pi) = \mathbb{E}[\mathcal{G}(\tau) \mid s_0, a_0; \theta_\pi]$$

We can derive the Q function from the value function

$$V(s_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

First, introduce the action  $a_0$

$$V(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

# Q Functions

$$V(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

# Q Functions

$$V(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

Condition the initial reward on the action

# Q Functions

$$V(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

Condition the initial reward on the action

$$V(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

# Q Functions

$$V(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

Condition the initial reward on the action

$$V(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

Call it the Q function

# Q Functions

$$V(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0; \theta_\pi] + \gamma V(s_1, \theta_\pi)$$

Condition the initial reward on the action

$$V(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

Call it the Q function

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$



# Q Functions

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

# Q Functions

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

The Q function tells us:

# Q Functions

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

The Q function tells us:

- The value of an action  $a_0$

# Q Functions

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

The Q function tells us:

- The value of an action  $a_0$
- In state  $s_0$

# Q Functions

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

The Q function tells us:

- The value of an action  $a_0$
- In state  $s_0$
- If we follow  $\pi(a_t \mid s_t; \theta_\pi)$  afterwards

# Q Functions

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

The Q function tells us:

- The value of an action  $a_0$
- In state  $s_0$
- If we follow  $\pi(a_t \mid s_t; \theta_\pi)$  afterwards

**Question:** How can we use the Q function for decision making?

# Q Functions

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

The Q function tells us:

- The value of an action  $a_0$
- In state  $s_0$
- If we follow  $\pi(a_t \mid s_t; \theta_\pi)$  afterwards

**Question:** How can we use the Q function for decision making?

Hint: We can evaluate  $Q$  for every possible action

# Q Functions

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

The Q function tells us:

- The value of an action  $a_0$
- In state  $s_0$
- If we follow  $\pi(a_t \mid s_t; \theta_\pi)$  afterwards

**Question:** How can we use the Q function for decision making?

Hint: We can evaluate  $Q$  for every possible action

$$\arg \max_{a_0 \in A} Q(s_0, a_0, \theta_\pi) = \arg \max_{a_0 \in A} (\mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi))$$



# Q Functions

$$\arg \max_{a_0 \in A} Q(s_0, a_0, \theta_\pi) = \arg \max_{a_0 \in A} (\mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi))$$

# Q Functions

$$\arg \max_{a_0 \in A} Q(s_0, a_0, \theta_\pi) = \arg \max_{a_0 \in A} (\mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi))$$

This is a very powerful equation

# Q Functions

$$\arg \max_{a_0 \in A} Q(s_0, a_0, \theta_\pi) = \arg \max_{a_0 \in A} (\mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi))$$

This is a very powerful equation

- Compute  $Q(s_0, a_0)$  for all  $a_0$

# Q Functions

$$\arg \max_{a_0 \in A} Q(s_0, a_0, \theta_\pi) = \arg \max_{a_0 \in A} (\mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi))$$

This is a very powerful equation

- Compute  $Q(s_0, a_0)$  for all  $a_0$
- Pick the  $a_0$  that maximizes  $Q(s_0, a_0)$

# Q Functions

$$\arg \max_{a_0 \in A} Q(s_0, a_0, \theta_\pi) = \arg \max_{a_0 \in A} (\mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi))$$

This is a very powerful equation

- Compute  $Q(s_0, a_0)$  for all  $a_0$
- Pick the  $a_0$  that maximizes  $Q(s_0, a_0)$

This  $a_0$  is **guaranteed** to be the optimal action

# Q Functions

$$\arg \max_{a_0 \in A} Q(s_0, a_0, \theta_\pi) = \arg \max_{a_0 \in A} (\mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi))$$

This is a very powerful equation

- Compute  $Q(s_0, a_0)$  for all  $a_0$
- Pick the  $a_0$  that maximizes  $Q(s_0, a_0)$

This  $a_0$  is **guaranteed** to be the optimal action

This considers the effect of  $a_0$  on the **infinite** future

# Q Functions

$$\arg \max_{a_0 \in A} Q(s_0, a_0, \theta_\pi) = \arg \max_{a_0 \in A} (\mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi))$$

This is a very powerful equation


- Compute  $Q(s_0, a_0)$  for all  $a_0$
- Pick the  $a_0$  that maximizes  $Q(s_0, a_0)$

This  $a_0$  is **guaranteed** to be the optimal action

This considers the effect of  $a_0$  on the **infinite** future

We collapsed the infinite decision tree into a single level


# Q Functions





# Q Functions

$t = 0$



# Q Learning

---

# Q Learning

Q learning is a **model-free** algorithm first discovered in the 1980s

# Q Learning

Q learning is a **model-free** algorithm first discovered in the 1980s

**Model-based**

# Q Learning

Q learning is a **model-free** algorithm first discovered in the 1980s

## Model-based

We know  $\text{Tr}(s_{t+1} \mid s_t, a_t)$

# Q Learning

Q learning is a **model-free** algorithm first discovered in the 1980s

## Model-based

We know  $\text{Tr}(s_{t+1} \mid s_t, a_t)$

Cheap to train, expensive to use

# Q Learning

Q learning is a **model-free** algorithm first discovered in the 1980s

## Model-based

We know  $\text{Tr}(s_{t+1} \mid s_t, a_t)$

Cheap to train, expensive to use

Closer to traditional control theory

# Q Learning

Q learning is a **model-free** algorithm first discovered in the 1980s

## Model-based

## Model-free

We know  $\text{Tr}(s_{t+1} \mid s_t, a_t)$

Cheap to train, expensive to use

Closer to traditional control theory



# Q Learning

Q learning is a **model-free** algorithm first discovered in the 1980s

## Model-based

We know  $\text{Tr}(s_{t+1} \mid s_t, a_t)$

Cheap to train, expensive to use

Closer to traditional control theory

## Model-free

We do not know  $\text{Tr}(s_{t+1} \mid s_t, a_t)$

# Q Learning

Q learning is a **model-free** algorithm first discovered in the 1980s

## Model-based

We know  $\text{Tr}(s_{t+1} \mid s_t, a_t)$

Cheap to train, expensive to use

Closer to traditional control theory

## Model-free

We do not know  $\text{Tr}(s_{t+1} \mid s_t, a_t)$

Expensive to train, cheap to use

# Q Learning

Q learning is a **model-free** algorithm first discovered in the 1980s

## Model-based

We know  $\text{Tr}(s_{t+1} \mid s_t, a_t)$

Cheap to train, expensive to use

Closer to traditional control theory

## Model-free

We do not know  $\text{Tr}(s_{t+1} \mid s_t, a_t)$

Expensive to train, cheap to use

Closer to deep learning

# Q Learning

Q learning is a **model-free** algorithm first discovered in the 1980s

## Model-based

We know  $\text{Tr}(s_{t+1} \mid s_t, a_t)$

Cheap to train, expensive to use

Closer to traditional control theory

## Model-free

We do not know  $\text{Tr}(s_{t+1} \mid s_t, a_t)$

Expensive to train, cheap to use

Closer to deep learning

# Q Learning

Q learning is still popular today

---

<sup>1</sup>*Simplifying Deep Temporal Difference Learning*. ICLR. 2024.

# Q Learning

Q learning is still popular today

Works well with deep neural networks

---

<sup>1</sup>*Simplifying Deep Temporal Difference Learning*. ICLR. 2024.

# Q Learning

Q learning is still popular today

Works well with deep neural networks

Researchers are still improving it<sup>1</sup>

---

<sup>1</sup>*Simplifying Deep Temporal Difference Learning*. ICLR. 2024.

# Q Learning

Q learning is still popular today

Works well with deep neural networks

Researchers are still improving it<sup>1</sup>

In fact, our lab is using it in our research right now

---

<sup>1</sup>*Simplifying Deep Temporal Difference Learning*. ICLR. 2024.



# Q Learning

Q learning is still popular today

Works well with deep neural networks

Researchers are still improving it<sup>1</sup>

In fact, our lab is using it in our research right now

We now have all the information we need to implement Q learning

---

<sup>1</sup>*Simplifying Deep Temporal Difference Learning*. ICLR. 2024.

# Q Learning

Our Q function relies on the value function for some  $\theta_\pi$

# Q Learning

Our  $Q$  function relies on the value function for some  $\theta_\pi$

Right now, it is not clear what the policy is

# Q Learning

Our  $Q$  function relies on the value function for some  $\theta_\pi$

Right now, it is not clear what the policy is

So how can we use the  $Q$  function without knowing the policy?

# Q Learning

Our Q function relies on the value function for some  $\theta_\pi$

Right now, it is not clear what the policy is

So how can we use the Q function without knowing the policy?

Let us find out

# Q Learning

Start with the  $Q$  function

# Q Learning

Start with the Q function

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

# Q Learning

Start with the Q function

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

We want to take the action that maximizes Q



# Q Learning

Start with the Q function

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

We want to take the action that maximizes Q

$$\arg \max_{a_0 \in A} Q(s_0, a_0, \theta_\pi) = \arg \max_{a_0 \in A} (\mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi))$$

# Q Learning

Start with the Q function

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

We want to take the action that maximizes Q

$$\arg \max_{a_0 \in A} Q(s_0, a_0, \theta_\pi) = \arg \max_{a_0 \in A} (\mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi))$$

Recall the definition of value function

# Q Learning

Start with the Q function

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

We want to take the action that maximizes Q

$$\arg \max_{a_0 \in A} Q(s_0, a_0, \theta_\pi) = \arg \max_{a_0 \in A} (\mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi))$$

Recall the definition of value function

$$V(s_0, \theta_\pi) = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_\pi]$$

# Q Learning

$$\arg \max_{a_0 \in A} Q(s_0, a_0, \theta_\pi) = \arg \max_{a_0 \in A} (\mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi))$$

$$V(s_0, \theta_\pi) = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_\pi]$$

# Q Learning

$$\arg \max_{a_0 \in A} Q(s_0, a_0, \theta_\pi) = \arg \max_{a_0 \in A} (\mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi))$$

$$V(s_0, \theta_\pi) = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_\pi]$$

**Question:** What should our policy be?

# Q Learning

$$\arg \max_{a_0 \in A} Q(s_0, a_0, \theta_\pi) = \arg \max_{a_0 \in A} (\mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi))$$

$$V(s_0, \theta_\pi) = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_\pi]$$

**Question:** What should our policy be?

Hint: The first equation provides the optimal action  $a_0$

# Q Learning

$$\arg \max_{a_0 \in A} Q(s_0, a_0, \theta_\pi) = \arg \max_{a_0 \in A} (\mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi))$$

$$V(s_0, \theta_\pi) = \sum_{t=0}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_0; \theta_\pi]$$

**Question:** What should our policy be?

Hint: The first equation provides the optimal action  $a_0$

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$

# Q Learning

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$



# Q Learning

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$

The policy uses the Q function

# Q Learning

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$

The policy uses the Q function

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

# Q Learning

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$

The policy uses the Q function

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

The Q function uses the policy (using the value function)

# Q Learning

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

# Q Learning


$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

Now that we know the policy, we can simplify the value function

# Q Learning

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$


$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$


Now that we know the policy, we can simplify the value function

# Q Learning

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$

$\arg \max_{a \in A}$


$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

Now that we know the policy, we can simplify the value function


**Question:** What is the value function for an optimal policy?

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

# Q Learning

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$

$\arg \max_{a \in A}$


$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma V(s_1, \theta_\pi)$$

Now that we know the policy, we can simplify the value function

**Question:** What is the value function for an optimal policy?

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$



# Q Learning

**Definition:** In Temporal Difference Q learning, we learn  $Q$  using

# Q Learning

**Definition:** In Temporal Difference Q learning, we learn  $Q$  using

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

# Q Learning

**Definition:** In Temporal Difference Q learning, we learn  $Q$  using

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

**Definition:** In Monte Carlo Q learning, we learn  $Q$  using

# Q Learning

**Definition:** In Temporal Difference Q learning, we learn  $Q$  using

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

**Definition:** In Monte Carlo Q learning, we learn  $Q$  using

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi]$$


# Q Learning

**Definition:** In Temporal Difference Q learning, we learn  $Q$  using

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

**Definition:** In Monte Carlo Q learning, we learn  $Q$  using

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi]$$

 Return following  $\pi$

# Q Learning

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi]$$

# Q Learning

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi]$$

If we want to learn the left hand side, we must know the right hand side

# Q Learning

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

$$Q(s_0, a_0, \theta_\pi) = \mathbb{E}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi]$$

If we want to learn the left hand side, we must know the right hand side

**Question:** How do we find these terms?



# Q Learning

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi]$$

**Question:** How do we find these terms?

**Answer:** Empirical expectation from episode data  $(s_t, a_t, \mathcal{R}(s_{t+1}))$

# Q Learning

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \mathbb{E}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi]$$

**Question:** How do we find these terms?

**Answer:** Empirical expectation from episode data  $(s_t, a_t, \mathcal{R}(s_{t+1}))$

$$E = \begin{bmatrix} s_0 & s_1 & s_2 & \dots \\ a_0 & a_1 & a_2 & \dots \\ r_0 & r_1 & r_2 & \dots \end{bmatrix}^\top$$

# Q Learning

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \hat{\mathbb{E}}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi]$$

# Q Learning

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \hat{\mathbb{E}}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi]$$

**Question:** How to find these terms?

# Q Learning

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \hat{\mathbb{E}}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi]$$

**Question:** How to find these terms?

$$\mathbf{E} = \begin{bmatrix} s_0 & s_1 & s_2 & \dots \\ a_0 & a_1 & a_2 & \dots \\ r_0 & r_1 & r_2 & \dots \end{bmatrix}^\top$$

# Q Learning

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \hat{\mathbb{E}}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi]$$

**Question:** How to find these terms?

$$\mathbf{E} = \begin{bmatrix} s_0 & s_1 & s_2 & \dots \\ a_0 & a_1 & a_2 & \dots \\ r_0 & r_1 & r_2 & \dots \end{bmatrix}^\top$$

**TD:** (Careful with terminal states)

$$\neg d \cdot \gamma \max_{a \in A} Q(s_{t+1}, a, \theta_\pi)$$

# Q Learning

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \hat{\mathbb{E}}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi]$$

**Question:** How to find these terms?

$$\mathbf{E} = \begin{bmatrix} s_0 & s_1 & s_2 & \dots \\ a_0 & a_1 & a_2 & \dots \\ r_0 & r_1 & r_2 & \dots \end{bmatrix}^\top$$

**TD:** (Careful with terminal states)

$$\neg d \cdot \gamma \max_{a \in A} Q(s_{t+1}, a, \theta_\pi)$$

**MC:**  $\gamma r_{t+1} + \gamma^2 r_{t+2} + \dots$

# Q Learning

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \hat{\mathbb{E}}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi]$$

**Question:** How to find these terms?

$$\mathbf{E} = \begin{bmatrix} s_0 & s_1 & s_2 & \dots \\ a_0 & a_1 & a_2 & \dots \\ r_0 & r_1 & r_2 & \dots \end{bmatrix}^\top$$

**TD:** (Careful with terminal states)  $\neg d \cdot \gamma \max_{a \in A} Q(s_{t+1}, a, \theta_\pi)$       **MC:**  $\gamma r_{t+1} + \gamma^2 r_{t+2} + \dots$

We know the right hand side, use it to learn the left hand side



# Q Learning

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \hat{\mathbb{E}}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi]$$

# Q Learning

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \hat{\mathbb{E}}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi]$$

Assume  $Q(s, a, \theta_\pi)$  has error  $\eta$  with right hand side

# Q Learning

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \hat{\mathbb{E}}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi]$$

Assume  $Q(s, a, \theta_\pi)$  has error  $\eta$  with right hand side

Use the error to update the Q function

# Q Learning

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \gamma \max_{a \in A} Q(s_1, a, \theta_\pi)$$

$$Q(s_0, a_0, \theta_\pi) = \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \hat{\mathbb{E}}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi]$$

Assume  $Q(s, a, \theta_\pi)$  has error  $\eta$  with right hand side

Use the error to update the Q function

$$Q_{i+1}(s, a, \theta_\pi) = Q_i(s, a, \theta_\pi) - \eta$$

Improve convergence with a learning rate  $\alpha$

$$Q_{i+1}(s, a, \theta_\pi) = \alpha(Q_i(s, a, \theta_\pi) - \eta)$$

# Q Learning

Monte Carlo update:

# Q Learning

Monte Carlo update:

$$Q_{i+1}(s_0, a_0, \theta_\pi) = \alpha \left( \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \hat{\mathbb{E}}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi] - Q_i(s_0, a_0, \theta_\pi) \right)$$

# Q Learning

**Monte Carlo update:**

$$Q_{i+1}(s_0, a_0, \theta_\pi) = \alpha \left( \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \hat{\mathbb{E}}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi] - Q_i(s_0, a_0, \theta_\pi) \right)$$

**Temporal Difference update:**

# Q Learning

**Monte Carlo update:**

$$Q_{i+1}(s_0, a_0, \theta_\pi) = \alpha \left( \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + \sum_{t=1}^{\infty} \gamma^t \hat{\mathbb{E}}[\mathcal{R}(s_{t+1}) \mid s_1; \theta_\pi] - Q_i(s_0, a_0, \theta_\pi) \right)$$

**Temporal Difference update:**

$$Q_{i+1}(s_0, a_0, \theta_\pi) = \alpha \left( \hat{\mathbb{E}}[\mathcal{R}(s_1) \mid s_0, a_0] + d\gamma \max_{a \in A} Q_i(s_1, a, \theta_\pi) - Q_i(s_0, a_0, \theta_\pi) \right)$$

Updates guarantee convergence to the true Q function ( $\lim_{i \rightarrow \infty} \eta = 0$ )



# Q Learning

Last thing, we must collect episodes to train Q!

# Q Learning

Last thing, we must collect episodes to train Q!

Can run policy in environment to create episodes

# Q Learning

Last thing, we must collect episodes to train Q!

Can run policy in environment to create episodes

```
states, next_states, rewards, terminateds = [], [], [], []
state = environment.reset()
while not terminated:
    action = policy.sample(state)
    next_state, reward, terminated = environment.step(action)

    states.append(state), next_states.append(next_state), ...
    state = next_state

episode = (states, next_states, rewards, terminateds)
```

# Q Learning

What policy do we sample actions from?

# Q Learning

What policy do we sample actions from?

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$

# Q Learning

What policy do we sample actions from?

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$

**Question:** Any issues?

# Q Learning

What policy do we sample actions from?

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$

**Question:** Any issues?

**Answer:** Always sample the same action

# Q Learning

What policy do we sample actions from?

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$

**Question:** Any issues?

**Answer:** Always sample the same action

If Q function is wrong, always sample bad actions



# Q Learning

What policy do we sample actions from?

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$

**Question:** Any issues?

**Answer:** Always sample the same action

If Q function is wrong, always sample bad actions

Without correct actions, Q function will not improve!

# Q Learning

What policy do we sample actions from?

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$

**Question:** Any issues?

**Answer:** Always sample the same action

If Q function is wrong, always sample bad actions

Without correct actions, Q function will not improve!

**Question:** What can we do?

# Q Learning

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$

# Q Learning

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$

Epsilon greedy policy!

# Q Learning

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$

Epsilon greedy policy!

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} (1 - \varepsilon) & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ \frac{\varepsilon}{|A|} & \text{for } a \in A \end{cases}$$

# Q Learning

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$

Epsilon greedy policy!

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} (1 - \varepsilon) & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ \frac{\varepsilon}{|A|} & \text{for } a \in A \end{cases}$$

Sample random action with probability  $\varepsilon$

# Q Learning

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} 1 & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ 0 & \text{otherwise} \end{cases}$$

Epsilon greedy policy!

$$\pi(a_0 \mid s_0; \theta_\pi) = \begin{cases} (1 - \varepsilon) & \text{if } a_0 = \arg \max_{a \in A} Q(s_0, a, \theta_\pi) \\ \frac{\varepsilon}{|A|} & \text{for } a \in A \end{cases}$$

Sample random action with probability  $\varepsilon$

In the limit, sample all possible actions in all states

# Q Learning

So far:



# Q Learning

So far:

- Defined training objective (TD and MC updates)

# Q Learning

So far:

- Defined training objective (TD and MC updates)
- Defined dataset (episodes)

# Q Learning

So far:

- Defined training objective (TD and MC updates)
- Defined dataset (episodes)
- Need to define model (Q function)!

# Q Learning

So far:

- Defined training objective (TD and MC updates)
- Defined dataset (episodes)
- Need to define model (Q function)!

Next time, we will use deep neural networks

# Q Learning

So far:

- Defined training objective (TD and MC updates)
- Defined dataset (episodes)
- Need to define model (Q function)!

Next time, we will use deep neural networks

Today and for homework, use a simple matrix

# Q Learning

Each state is a row, each action is a column in a matrix

# Q Learning

Each state is a row, each action is a column in a matrix

$$\begin{bmatrix} Q(s_a, a_a) & Q(s_b, a_b) & \dots \\ Q(s_b, a_a) & Q(s_b, a_b) & \dots \\ \vdots & \vdots & \vdots \end{bmatrix}$$

# Q Learning

Each state is a row, each action is a column in a matrix

$$\begin{bmatrix} Q(s_a, a_a) & Q(s_b, a_b) & \dots \\ Q(s_b, a_a) & Q(s_b, a_b) & \dots \\ \vdots & \vdots & \vdots \end{bmatrix}$$

$Q_{i,j}$  gives Q value for state  $s = i$  and action  $a = j$



# Homework

---

# Homework

You have everything you need to solve homework

# Homework

You have everything you need to solve homework

Due in 2 weeks (Weds 12 March, 23:59)

# Homework

You have everything you need to solve homework

Due in 2 weeks (Weds 12 March, 23:59)

Download and submit `.py` and `.ipynb` files

# Homework

You have everything you need to solve homework

Due in 2 weeks (Weds 12 March, 23:59)

Download and submit .py and .ipynb files

Uses turnitin for checking

# Homework

You have everything you need to solve homework

Due in 2 weeks (Weds 12 March, 23:59)

Download and submit .py and .ipynb files

Uses turnitin for checking

[https://colab.research.google.com/drive/1xtBxAaVc3ax6\\_j59RC3NLQQPFcIEoau-?usp=sharing](https://colab.research.google.com/drive/1xtBxAaVc3ax6_j59RC3NLQQPFcIEoau-?usp=sharing)