

Elastic Bounds of Pearson’s r : Theoretical and Empirical Insights for Ordinal Variables

Cees van der Eijk Scott Moser

21, March 2026

Table of contents

1	Introduction and Conceptual Setting	3
1.1	On ordinal-categorical variables (orcats), their level of measurement and their common use in quantitative analyses	4
1.2	Bounds of correlations between ordinal categorical variables (orcats)	5
1.3	Background and Definitions	7
2	Theoretical Framework	7
2.1	Optimal Coupling and Correlation Bounds	8
2.1.1	Illustration	13
2.2	Examples	14
2.2.1	Notation (permutation extremes, “Lane A”)	15
2.2.2	A) Same number of categories, but $ r_{\min} \neq r_{\max}$ (concrete numbers)	15
2.2.3	B) Different numbers of categories, yet $ r_{\min} = r_{\max}$ (concrete numbers)	16
2.2.4	C) Spearman invariance (same data as in B)	16
2.3	what changes and what doesn’t (Pearson vs Spearman)	17
2.4	Population Bounds	17
2.4.1	Lemma (signs of extreme correlations under monotone scoring)	17
2.5	Asymetry	19
2.5.1	1) setup (permutation extremes for a single sample)	20
2.5.2	2) pair-specific necessary & sufficient condition	20
2.5.3	3) universal condition (independent of X)	21
2.6	Estiimation and Joint Uncertainty	22
3	Measurement Issues and Metrics of Constraint	22
3.1	Possible “Metrics of Constraint”	24
3.1.1	Overlap-Based Measures of Asymmetry	24
4	Implications, Issues, and Questions	25

4.1	Interpretation of ‘Effect Size’	25
4.2	Comparison Across Studies / Measures	25
4.3	Hypothesis Testing	25
4.4	Rescaling	25
4.5	Variance “Explained”	25
5	Monte Carlo Simulations	26
5.1	Effects of Marginal Shapes	27
5.2	Symmetry Breaking in Correlation Bounds	27
6	Illustration with ‘Typical’ Data: BES	27
7	Discussion	28
7.1	Comparing ‘Typical’ Data (BES) with Simulated Data	28
7.2	Relation to Literature	28
7.3	Recommendations	28
8	Conclusion	29
8.1	Extensions and Open Questions	29
8.1.1	Correlation Matricies	30
9	References	33
A	Appendix A: Mathematical Proofs	35
A.1	Proof A.1	35
A.2	Proof A.2	35
B	Appendix B: Software Tools and Code Listings	35
C	Appendix C: Additional Tables and Simulations	35
D	Appendix D: Old Outline	35

Abstract

The use of product-moment correlations to assess relationships between ordered-categorical variables is widespread and generally tolerated in spite of the non-metric character of such variables. Potential problems of this practice have been identified in the literature. This article focusses on one of these, namely that correlation coefficients are poorly comparable across pairs of ordered-categorical variables because their ranges are constrained in non-uniform ways. This not only affects the validity of correlation coefficients for descriptive purposes, but also as bases for subsequent analyses, as in, e.g., principal component analysis, factor analysis, and structural equation modelling. This article focusses on this problem of ‘constrained’ correlations. It explains the problem and explores conditions under which it is most severe. It provides illustrations based on both fabricated and actual empirical data, the latter from a mass-survey that ubiquitously contains many ordered-categorical variables that are commonly analysed with product-moment correlations. The article is accompanied by a software tool in R for analysing the attainable upper- and lower bounds of correlations in actual data.

Pearson’s correlation coefficient r , when applied to ordinal data, is subject to elasticity due to constraints from marginal distributions. This paper develops theoretical bounds for r , based on Boole–Fréchet–Hoeffding inequalities and rearrangement inequalities, and derives analytic expressions for their limits. We investigate when symmetry in the bounds holds or breaks, and explore practical implications for correlation-based methods like PCA and SEM. An accompanying R package allows users to compute bounds and perform hypothesis tests given fixed marginals.

1 Introduction and Conceptual Setting

In this article we elaborate why correlations involving ordered-categorical variables are generally ‘constrained’, which means that their attainable maximum and minimum values are rarely +1 and -1, as is often thought, but values that have to be calculated for each pair of variables. We assess the magnitude of these constraints in fictitious as well as empirical data and analyse under which conditions they are most severe. The article is accompanied by a software tool in R: that allows applied researchers to easily assess the extent of the problem in their own data.

The use of product-moment correlations (also known as Pearson correlations) to assess linear relationships between ordered-categorical variables—henceforth referred to as *orcats*—is widespread and widely condoned in empirical studies, in spite of such variables not being metric while the product-moment correlation assumes that they are. We do not focus on this problem from an obsessive desire for methodological purity, were it only because many statistical procedures are quite robust for violations of their assumptions [fn: examples?]. Instead, we draw attention to problematic consequences of this violation of assumptions that, as far as we know, have been rarely recognised in the applied literature although they are well-known in the theoretical statistical literature. This is the problem that correlation coefficients involving *orcats* are constrained in their range which makes them mutually incomparable, leading to problems in hypothesis testing and in analytical procedures that use correlations as their input.

Despite long-standing cautions about applying Pearson’s correlation coefficient r to ordinal variables ((Labovitz, 1970); (Gaito, 1980)), it remains routine in survey-based social science. In political science and sociology alike, researchers frequently correlate 4-, 5-, or 7-point items as if they were interval-scaled, usually without examining the implications of the marginals. Examples include correlations among attitude items in large-scale survey batteries ((Alwin & Krosnick, 1991)), ideology and issue positions on mixed-resolution scales ((Brewer & Gross, 2005); (Kinder & Kam, 2009)), and summary indices that are subsequently modeled using correlation-based techniques. Reviews document that such practices are pervasive in applied survey analysis ((Kreuter & Valliant, 2007)). Methodological discussions sometimes defend pragmatic use of parametric tools with ordinal data ((Jacoby, 1999)), but even optimistic views do not address the *feasible range* of r imposed by fixed marginals—an omission with direct consequences for substantive interpretation.

Alternatives to Pearson’s r —notably Spearman’s ρ , Kendall’s τ , and the polychoric correlation—offer different trade-offs ((Spearman, 1904); (Kendall, 1938); (Olsson, 1979)). Rank-based measures ignore the marginals; polychorics assume latent bivariate normality. In contrast, we pursue bounds on r that *explicitly* depend on the observed marginals, drawing on rearrangement inequalities and the Fréchet–Hoeffding framework ((Fréchet, 1951); (Hoeffding, 1940); (Rüschendorf, 1981)). This perspective clarifies when the attainable range of r is narrow, when symmetry fails ($r_{\min} \neq -r_{\max}$), and when the lower bound can be strictly positive. It also explains why correlation-based models with ordinal indicators (EFA/CFA/SEM) can be distorted when Pearson correlations are used ((Flora & Curran, 2004); (Green et al., 1997)). Our contribution situates empirical practice within

these theoretical limits and provides tools to compute and visualize the bounds given fixed marginals.

This article proceeds as follows. We start by briefly elaborating the character of ordinal categorical variables (orcats), why they are generally not regarded as being metric, and why, in spite of that, they are nevertheless very frequently used in statistical procedures that assume metric level data. We then proceed in section 3 to show why correlations based on orcats tend to be constrained, which implies that their range is smaller than from -1 to +1, and why this is problematic in applied research. This raises the question how to determine the actual range of attainable values of correlations between orcats, which we address in the subsequent section. In section 5 we assess the magnitude of these constraining effects on two different sets of data. One is a typical survey dataset in which many of the variables are ordinal categorical, the oricatorcat 2019 British Election Study. The second consists of data derived from simulations covering all possible relevant conditions for pairs of ordered categorical variables. We use both sets of data to explore conditions that influence the magnitude of the constraining effect. Section 5 concludes by summarizing the findings, by elaborating the consequences for applied research, and by discussing directions for further research.

1.1 On ordinal-categorical variables (orcats), their level of measurement and their common use in quantitative analyses

Orcats are ordinal variables with a relatively small number of categories (or ranks) compared to the number of cases, thus resulting in partial orderings of cases. Although they can be found in all field of quantitative inquiry they are ubiquitous in data from standardised social science surveys. Likert items and rating scales [describe in fn Likert items and rating scales, add that there are yet other formats such as semantic-differential scales that also yield orcats] are among the most favoured ways to elicit responses in such surveys. These survey-question formats have most often between 3 and 11 ordinal categories. The numerical values assigned to the categories of orcats are most often consecutive integers, starting with either 0 or 1. [fn: explain that these values may be included in the formulation of a survey question, which is immaterial to our discussion here. The values assigned to the categories (and to the responses) reflect, in combination with the analysis procedure the analyst’s interpretation of their meaning.] . Such an assignment of values is as good as, and indeed analytically equivalent to, any other set of values that reflects the analyst’s understanding of the ordinal relations between the categories, i.e., any monotonic transformation. However, objections are often raised when orcats [fn: add references] are analysed with procedures requiring metric measurement. That requirement implies that the differences between values can validly be compared so that, e.g., the substantive difference between categories 1 and 2 is equal to that between categories 3 and 4, and half as large as the difference between categories 2 and 4, etc. Any implied interval-level interpretation of ordinal categorical variables requires therefore a compelling theoretical or empirical justification of the numerical values assigned to the categories rather than just the analyst’s fiat. In principle this is quite possible, for example by calibrating values of categories against psychophysical measurements (cf., Lodge; Tillie; others), or against the results of psychometric/linguistic analysis of verbal

response labels (cf Sönning 2024). However, such procedures are rarely integrated in applied survey research because of their high cost. In the absence of compelling justifications for values assigned to categories orcats can only be considered to provide ordinal information, not metric information. In spite of this we find that they are very commonly analysed with statistical tools that require metric measurement of variables such as product-moment correlations or procedures that build upon such correlations. We find this clearly exhibited in published articles in high-ranking academic journals [fn: explain how we did this] and in highly regarded textbooks where examples are given of the use of correlations between orcats as input for procedures such as PCA, FA or SEM [fn: examples: Kline; Byrne; xxx]. This practice is so established that it is occasionally rationalised by referring to orcats as being ‘quasi-interval level’ measures [fn: reference]. One can, however, also find more explicit justifications for this practice, which emphasise the robustness of correlation coefficients for violations of their measurement level assumptions. Labovitz (1967, 1970) in particular has compared correlations between orcats scored according to equal-distances with correlations when the variables were scored with a monotonic random procedure (which only assumes ordinality between categories). He reports that the differences are negligible, and suggests that even if the theoretical objection against unfounded interval level interpretations of orcats would be justified, that objection would nevertheless have such minor practical consequences as to make it irrelevant in most situations. Not surprisingly, Labovitz’ work has in turn be criticised itself on various grounds, most particularly of (inadvertently) mainly covering situations where this robustness holds, and overlooking other situations (cf O’Brien 1979; xxx). At this place we will not focus on discussions about the equal-distance scoring practice. Our main reason for referring to them is to demonstrate the existence of lively debates in the applied literature about potential issues when using orcats in procedures requiring metric data. The existence of these debates makes it remarkable that another important issue has –to the best of our knowledge– remained mainly overlooked in the applied literature while it has been well-known in the theoretical statistical literature. This is the problem that correlations between orcats are generally ‘constrained’, which means that they do not range between -1 and +1, but between minima and maxima that have to be calculated separately for each pair of variables. The nature of this issue is the subject of the next section.

1.2 Bounds of correlations between ordinal categorical variables (orcats)

The values that product-moment correlations between ordinal-categorical variables (orcats) can attain are generally constrained by minimum and maximum bounds of attainable values that are not -1 and +1. [fn: The maximum attainable value is +1 only when the univariate distributions of the variables are identical. But that does not imply the minimum attainable value then to be -1, as that also requires further conditions that the univariate distributions must meet (see below, in the discussion of Table 2). The minimum attainable value of a correlation is -1 when the univariate distributions are each other’s inverse, but again, that does not imply that the maximum is +1, as that requires additional conditions of the univariate distributions.]

Because the bounds of correlations between orcats are not -1 and +1 it is even more difficult

than commonly thought in applied research to interpret the strength of correlations. The well-known and often referred to criteria formulated by Cohen (1988: 77-81) for correlations in the behavioural sciences [fn: Cohen suggested, with appropriate caution, the following guidelines: $0.10 < r < 0.29$ weak; $0.30 < r < 0.49$ moderate; $0.50 < r < 1.00$ strong. Hemphill (2003) suggests on the basis of a review of several meta-analyses in psychological research that the cut-off of 0.50 for considering a correlation to be ‘strong’ may be too high in many common instances.] are explicitly based on the notion that the minimum and maximum attainable values for r are -1 and +1. But that is generally not the case and, moreover, the actual upper- and lower-bounds of r are not the same across pairs of variables, but specific to each. This state of affairs thus generates two interconnected problems when correlating orcats. First, the absence of clarity of the lower- and upper-bounds of r makes qualitative interpretations of the strength of correlations elusive. Second, because the bounds of attainable values of r are specific for each pair of variables each correlation is expressed in terms of a different, but unknown reference interval. Such correlations are therefore not comparable. To better grasp these problems, it is necessary to determine the upper- and lower bounds of correlations, given the univariate distributions of the variables involved.

Statistical scholarship has since long recognized that, in contrast to what is often thought, the upper- and lower-bounds of correlations between orcats are generally not the reference interval $[-1, +1]$. [insert references, including Shi&Huang 1992; xxxx]. Determining the actual bounds given the marginal distributions of the two variables can be done by using Whitt’s (1976) rearrangement theorem and Hardy–Littlewood–Pólya’s (1952) rearrangement inequality. For obtaining the maximum attainable value of a correlation the values of X and Y have each to be sorted independently in either ascending or descending order and then paired (comonotonic arrangement). For the minimum attainable value of the correlation (given the marginals of the two variables) the anti-comonotonic arrangement is required (sorting one of the variables in ascending and the other in descending order). A formalized elaboration is presented in Appendix 1. Moreover, the algorithm for calculating r_{min} and r_{max} has been programmed in R and is [or: will be] freely available for researchers to assess the minimum and maximum attainable values of r for pairs of variables in their own data. Details about this routine are described in Appendix 2.

A series of illustrative cases is presented in Table 2 (as in Table 1 for $n=12$ and two orcats with 4 categories each). Although the distributions of the variables given in these examples is of no particular importance, and could have been replaced by innumerable other examples, they are nevertheless helpful to arrive at some general observations about r_{min} and r_{max} :

- The values of r_{min} and r_{max} are not constant across all examples, and not even across correlations involving a specific variable (see examples (a) and (b) which involve the same X -variable). The variation in the values of the bounds of r is across pairs of variables. In other words, the bounds of r are specific for each pair.
- r_{min} can be -1, and r_{max} can be 1, but neither of these values implies the other (as seen in examples (b) and (c). The combination of $r_{min}=-1$ and $r_{max}=1$ is possible, but dependent on specific characteristics of the two marginal distributions involved (see examples (d) and (f). Both variables must not only have identical distributions, but these must also be symmetrical.

- $r_{\min}$ and $r_{\max}$ are not necessarily symmetric around 0. Such symmetry may exist, but requires specific characteristics of the pair of marginal distributions, namely that both are symmetric (see examples (e), (g) and (h)).
- The extent to which the reference interval $[-1, 1]$ is covered by the interval $[r_{\min}, r_{\max}]$ varies across pairs of variables.
- The values of $r_{\min}$ and $r_{\max}$ depend not only on the marginal distributions of the two variables, but also on the values assigned to their categories. Unless specified differently, we follow in this article the common practice of assigning consecutive integer values to the ordered categories. But any monotonic transformation of values is equally adequate to represent the ordinal character of variables. Examples (i) and (j) involve the same pair of variables, but in (i) their categories have been assigned values 1, 2, 3 and 4, while in (j) the assigned values were 1, 2.25, 2.75 and 4. The consequences of this change are visible in the values of $r_{\min}$ and $r_{\max}$. This observation implies that value assignment to categories of orcats (and the justification thereof) remains an issue of importance when using Pearson correlations on orcats. It is a different problem, however, than that of constraintment of correlations on which we focus here.

Consequences of constrained correlations in applied research

- Descriptive interpretation of correlations (Cohen’s criteria) is undermined
- It is also undermined by every correlation being expressed on its own unique scale/unique calibration
- Need a nice example of 2 correlations with $r(a, b) \leq r(b, c)$ and where $r(a, b)_{\max}$ is $< r(b, c)$: is it smaller because the linear relationship is weaker, or because the constraintment is stronger?

1.3 Background and Definitions

- Ordinal data and assumptions of interval-scale correlation
- Existing alternatives: Spearman, polychoric, polyserial; their strengths and limitations
- Coupling and rearrangement in discrete settings

2 Theoretical Framework

The problem of extremal correlations under fixed marginals connects directly to classical results in probability inequalities. Fréchet’s work on admissible joint distributions ((Fréchet, 1951)) and Hoeffding’s development of covariance bounds ((Hoeffding, 1940)) laid the groundwork. Later refinements, such as Rüschendorf’s demonstration of sharpness of Fréchet bounds ((Rüschendorf, 1981)), clarified when comonotonic and countermonotonic couplings achieve extremal values. The rearrangement inequality, and its discrete extensions, ensure that matching high values of X with high values of Y maximizes $\mathbb{E}[XY]$, while opposite pairings minimize it. This perspective is elaborated in comparison and coupling frameworks ((Müller

& Stoyan, 2002); (Thorisson, 2000)). These results provide the theoretical backbone for our derivations of $r_{\min}$ and $r_{\max}$.

2.1 Optimal Coupling and Correlation Bounds

We formalize the problem of identifying the joint distribution (coupling) of two ordinal variables with fixed marginals that maximizes or minimizes the Pearson correlation.

Proposition 2.1. *Let X and Y be discrete ordinal random variables, each with finite ordered support:*

$$X : x_1 < x_2 < \dots < x_{K_X}, \quad Y : y_1 < y_2 < \dots < y_{K_Y}$$

with marginal probability distributions $p_X(x_i) = P(X = x_i)$, $p_Y(y_j) = P(Y = y_j)$, respectively.

Then, the maximum and minimum values of the Pearson correlation coefficient r_{XY} , consistent with the fixed marginals, are obtained as follows:

- *Maximum r_{XY} : achieved by pairing largest X -values with largest Y -values (comonotonic arrangement).*
- *Minimum r_{XY} : achieved by pairing largest X -values with smallest Y -values (anti-comonotonic arrangement).*

Proof. The Pearson correlation coefficient is given by:

$$r_{XY} = \frac{E[XY] - E[X]E[Y]}{\sigma_X \sigma_Y}.$$

Given fixed marginals $p_X(x_i)$, $p_Y(y_j)$, the expectations $E[X]$, $E[Y]$ and standard deviations σ_X , σ_Y are constant. Thus, to maximize or minimize r_{XY} , we only need to maximize or minimize the expectation $E[XY]$:

$$E[XY] = \sum_{i=1}^{K_X} \sum_{j=1}^{K_Y} x_i y_j P(X = x_i, Y = y_j).$$

This proof relies on a rearrangement inequality of (Hardy et al., 1952) (Theorem 368):

For real sequences $a_1 \leq a_2 \leq \dots \leq a_n$ and $b_1 \leq b_2 \leq \dots \leq b_n$, and for any permutation π of indices $\{1, \dots, n\}$ we have the following:

$$a_1 b_1 + a_2 b_2 + \dots + a_n b_n \geq a_1 b_{\pi(1)} + a_2 b_{\pi(2)} + \dots + a_n b_{\pi(n)}$$

and

$$a_1 b_n + a_2 b_{n-1} + \dots + a_n b_1 \leq a_1 b_{\pi(1)} + a_2 b_{\pi(2)} + \dots + a_n b_{\pi(n)}$$

Equality occurs only if the permutation π results in a monotonically increasing sequence (for maximum) or monotonically decreasing sequence (for minimum).

To explicitly connect this to our problem, enumerate observations explicitly from sorted marginals:

$$X_{\downarrow} : x_{[K_X]} \geq x_{[K_X-1]} \geq \dots \geq x_{[1]}, \quad Y_{\downarrow} : y_{[K_Y]} \geq y_{[K_Y-1]} \geq \dots \geq y_{[1]}.$$

(Sorted in descending order for maximization). Next expand discrete probability distributions into explicit observation-level sequences. If $K_X \neq K_Y$, extend the shorter sequence with zero-probability categories so both sequences have equal length, which does not affect marginal probabilities or correlation.

Hence, to maximize $E[XY]$: pair the largest ranks of X with the largest ranks of Y . Explicitly, start from the top (highest values) and move downward:

- Pair as many observations of the largest X -category as possible with the largest Y -category, then next-largest, etc., until all observations are paired.
- By the Hardy–Littlewood–Pólya Rearrangement Inequality ([Hardy et al., 1952](#)), this explicitly constructed pairing yields the maximum possible sum of products, hence maximizing $E[XY]$.

Likewise, to minimize $E[XY]$ pair the largest X -categories with the smallest Y -categories, then second-largest X -categories with second-smallest Y -categories, and so forth (anti-comonotonic ordering). By the rearrangement inequality, this arrangement explicitly yields the minimal sum of products, hence minimizing $E[XY]$. ■

In words, given fixed marginals, we have shown:

- **Maximum correlation** $r_{\max}$ is achieved by comonotonic (e.g. sorting values of X and Y from largest to smallest independently and pairing) arrangement:

$$P_{\max}(X = x_{[i]}, Y = y_{[i]}) \quad \text{in either descending or ascending order.}$$

- **Minimum correlation** $r_{\min}$ is achieved by anti-comonotonic arrangement:

$$P_{\min}(X = x_{[i]}, Y = y_{[n+1-i]}) \quad \text{pairing descending } X \text{ with ascending } Y, \text{ or viceversa.}$$

With this explicit construction we can simply calculate $r_{\max}$ and $r_{\min}$ when the values of X and Y are ranks.

Corollary 1. (Analytic Forms for $r_{\min}$ and $r_{\max}$)

Let two ordinal variables X and Y have values as follows:

- X with K_X ordinal levels: $1, 2, \dots, K_X$
- Y with K_Y ordinal levels: $1, 2, \dots, K_Y$

with absolute frequencies:

- $n_X(i)$, number of observations at level i of X .
- $n_Y(j)$, number of observations at level j of Y .

Then the maximum Pearson correlation is:

$$r_{\max} = \frac{E[XY]_{\max} - \bar{X}\bar{Y}}{\sigma_X \sigma_Y}.$$

Where

$$E[XY]_{\max} = \frac{1}{N} \sum_{i=1}^{K_X} \sum_{j=1}^{K_Y} x_i y_j M_{i,j}$$

where explicitly defined frequencies $M_{i,j}$ pair observations in a comonotonic manner, calculated explicitly via the greedy iterative algorithm in as Algorithm ?? in the proof.

Then explicitly the minimal expectation is:

$$E[XY]_{\min} = \frac{1}{N} \sum_{i=1}^{K_X} \sum_{j=1}^{K_Y} x_i y_j R_{i,j}$$

Where $R_{i,j}$ is defined in the proof.

Proof. Algorithm for explicit calculation of $M_{i,j}$ (greedy method):

Initialize available frequencies explicitly: $n'_Y(j) = n_Y(j)$ for each j

- For each level x_i , from smallest $i = 1$ to largest:
 - For each level y_j , from smallest $j = 1$ to largest:
 - * Pair as many observations as possible:

$$M_{i,j} = \min \left(n_X(i) - \sum_{s=1}^{j-1} M_{i,s}, n'_Y(j) \right)$$

Next, update explicitly:

$$n'_Y(j) \leftarrow n'_Y(j) - M_{i,j}$$

Define the matrix $M_{i,j}$, the frequency with which the level x_i pairs with the level y_j . Clearly, we must have:

$$\sum_j M_{i,j} = n_X(i), \quad \sum_i M_{i,j} = n_Y(j)$$

Similarly, the minimum Pearson correlation ($r_{\min}$) achieved by anti-comonotonic alignment (pairing largest ranks of X with smallest ranks of Y) is explicitly:

$$r_{\min} = \frac{E[XY]_{\min} - \bar{X}\bar{Y}}{\sigma_X \sigma_Y}.$$

To achieve the maximum expected product $E[XY]_{\max}$, do the following: First sort ranks from largest to smallest independently for both X and Y ; then pair ranks directly from highest to lowest available observations.

Minimum Expectation

The formula for $E[XY]_{\min}$ is derived in a similar manner. Define explicitly a matrix $R_{i,j}$ giving frequencies of pairings $X = x_i$ with $Y = y_j$. For the minimal expectation, pair largest available X -values explicitly with smallest available Y -values first (greedy algorithm):

Initialize explicitly available frequencies:

$$n'_Y(j) = n_Y(j) \quad \text{for each } j$$

- For i from largest to smallest ($i = K_X$ down to 1):
 - For j from smallest to largest ($j = 1$ up to K_Y):
 - * Pair explicitly as many observations as possible:

$$R_{i,j} = \min \left(n_X(i) - \sum_{s=1}^{j-1} R_{i,s}, n'_Y(j) \right)$$

- Update explicitly available frequencies:

$$n'_Y(j) \leftarrow n'_Y(j) - R_{i,j}$$

Then explicitly the minimal expectation is:

$$E[XY]_{\min} = \frac{1}{N} \sum_{i=1}^{K_X} \sum_{j=1}^{K_Y} x_i y_j R_{i,j}$$

This explicit equation applies Whitt's rearrangement theorem , Whitt (1976) , ensuring proper maximal pairing of ranks. (Whitt (1976)) ■

Discussion

We want to construct a joint distribution table M such that:

- The **row sums** match the marginal counts for X : $\sum_j M_{i,j} = n_X(i)$
- The **column sums** match the marginal counts for Y : $\sum_i M_{i,j} = n_Y(j)$

To do this, we go **greedily**, pairing the highest available X values with the **smallest available** Y values, one cell at a time.

To avoid **exceeding** the available marginal totals in Y , we must track how much of $n_Y(j)$ is **still unassigned** at each step — and that's what $n'_Y(j)$ represents.

After assigning $M_{i,j}$ units to the cell (i, j) , we subtract that amount from the remaining pool of level y_j values, like this: $n'_Y(j) \leftarrow n'_Y(j) - M_{i,j}$. This ensures we don't **overfill** any column of the matrix.

To maximize the expectation $E[XY]$, the rearrangement inequality states we must pair values of X and Y from smallest-to-smallest, second-smallest-to-second-smallest, and so forth (comonotonic alignment).

Thus, explicitly:

- Sort X : lowest-to-highest:

$$X_{\text{sorted}} = (\underbrace{x_1, \dots, x_1}_{n_X(1)}, \underbrace{x_2, \dots, x_2}_{n_X(2)}, \dots, \underbrace{x_{K_X}, \dots, x_{K_X}}_{n_X(K_X)})$$

Sorted Y -values (ascending order for max, descending for min):

$$Y_{\text{sorted}(\text{max})} = (\underbrace{y_1, \dots, y_1}_{n_Y(1)}, \underbrace{y_2, \dots, y_2}_{n_Y(2)}, \dots, \underbrace{y_{K_Y}, \dots, y_{K_Y}}_{n_Y(K_Y)})$$

$$Y_{\text{sorted}(\text{min})} = (\underbrace{y_{K_Y}, \dots, y_{K_Y}}_{n_Y(K_Y)}, \underbrace{y_{K_Y-1}, \dots, y_{K_Y-1}}_{n_Y(K_Y-1)}, \dots, \underbrace{y_1, \dots, y_1}_{n_Y(1)})$$

The maximum and minimum expectations are simply:

Maximum expectation (pair element-by-element):

$$E[XY]_{\text{max}} = \frac{1}{N} \sum_{k=1}^N X_{\text{sorted}}(k) \cdot Y_{\text{sorted}(\text{max})}(k)$$

- **Minimum expectation:**

$$E[XY]_{\text{min}} = \frac{1}{N} \sum_{k=1}^N X_{\text{sorted}}(k) \cdot Y_{\text{sorted}(\text{min})}(k)$$

These classical bounds define feasible joint distributions given marginals and constrain the attainable values of r . There is an interesting connection to Fréchet–Hoeffding and Boole–Fréchet Inequalities, which we note but leave for future research.¹

2.1.1 Illustration

To illustrate these idea concretely, Table 1 presents an illustrative example of comonotonic and anti-comonotonic arrangements and of the resulting lower- and upper-bounds of r , for two 4-category variables and 12 cases. One can easily expand this example to pairs of variables with different numbers of categories and different numbers of cases.

¹Fréchet–Hoeffding and Boole–Fréchet Inequalities: The upper bound is always attained by r_{max} . However, the lower bound is *not* guaranteed to attain. The fundamental asymmetry arises because discrete ordinal distributions cannot smoothly bridge these gaps. Thus, the upper bound alignment is always feasible, while the lower bound alignment frequently encounters structural obstacles.

In the discrete ordinal setting, r_{max} is **always** attained by the FH upper (comonotonic) construction. (*BFbounds.md*; *Proofs-OrdinalVariables.md*; *Ordinal Variables and Correlation.md*)

The FH lower (countermonotonic) construction may be **unattainable** in discrete settings; r_{min} is then the best feasible opposite-order arrangement.

(*BFbounds.md*; *Proofs-OrdinalVariables.md*; *Ordinal Variables and Correlation.md*)

Upper bound (positive alignment):

- Sorting X and Y separately in descending order, then pairing largest-to-largest, smallest-to-smallest, **always works**.
- You never run out of “matching” observations because the descending order for both variables naturally aligns category sizes. Even if one variable has large gaps, you can still sequentially pair without difficulty.

Lower bound (negative alignment):

- You sort X descending but Y ascending. You’re pairing extremes together: highest X with lowest Y , lowest X with highest Y .
- Now you face potential “imbalance”:
 - If one category is large for X but the corresponding opposite-ranked category for Y is small, you quickly exhaust available “opposite-end” observations.
 - After exhausting the best opposites, you are forced to pair less “extreme” pairs, creating a suboptimal alignment.
- Hence, perfect negative alignment (lower bound) becomes impossible to achieve fully, leaving you short of the theoretical Fréchet–Hoeffding lower bound.

Table 1: Determining minimum and maximum attainable values of correlation between two ordered categorical variables X and Y given their marginal distributions.

(a) Observed frequency distributions

	Category 1	Category 2	Category 3	Category 4	Total
X	6	3	2	1	12
Y	3	5	2	2	12

(b) Comonotonic arrangement and pairing:

X: 1 1 1 1 1 1 2 2 2 3 3 4

Y: 1 1 1 2 2 2 2 2 3 3 4 4 $r_{\max} = 0.87831$

(c) Anti-comonotonic arrangement and pairing:

X: 1 1 1 1 1 1 2 2 2 3 3 4

Y: 4 4 3 3 2 2 2 2 2 1 1 1 $r_{\min} = -0.79466$

Table 2: Illustrative pairs of marginal distributions and their associated minimum and maximum attainable values of r ($r_{\min}$, $r_{\max}$) given these distributions. The distributions are given in counts (n=12). The categories are assigned consecutive integer values starting with 1. C1 and C2 are measures of constraint, as discussed in the text.

	Category				Total	$r_{\min}$	$r_{\max}$
	1	2	3	4			
(a) Two arbitrarily distributed ordinal categorical variables (orcats)							
X	6	3	2	1	12	-0.79466	0.87831
Y	3	5	2	2	12		
(b) One arbitrarily distributed variable, and one with same distribution inversed							
X	6	3	2	1	12	-1	0.71429
Y	1	2	3	6	12		
(c) Two variables with arbitrary identical distributions							
X	6	3	2	1	12	-0.71429	1
Y	6	3	2	1	12		
(d) Two variables with identical symmetric and unimodal distributions							
X	2	4	4	2	12	-1	1
Y	2	4	4	2	12		
(e) Two variables with different distributions, both symmetric and unimodal							
X	2	4	4	2	12	-0.91169	0.91169
Y	1	5	5	1	12		
(f) Two variables with identical symmetric bimodal distributions							
X	4	2	2	4	12	-1	1
Y	4	2	2	4	12		
(g) Two variables with different distributions, both symmetric and bimodal							
X	4	2	2	4	12	-0.95673	0.95673
Y	5	1	1	5	12		
(h) One symmetric unimodal variable, the other symmetric bimodal							
X	2	4	4	2	12	-0.89923	0.89923
Y	4	2	2	4	12		
(i) Two variables, one very left skewed, the other very right skewed							
X	1	1	1	9	12	-0.95818	0.31939
Y	8	2	1	1	12		
(j) Same as situation (i), above, but categories assigned values 1; 2.25; 2.75; 4							
X	1	1	1	9	12	-0.93321	0.33829
Y	8	2	1	1	12		

2.2 Examples

Below, “scores $0:(K - 1)$ ” means category values $[0, 1, \dots, K - 1]$. “Affine recode” means $a + b \times (\text{old score})$ with $b > 0$. “Non-affine monotone” means strictly increasing but not affine (e.g., convex spacing).

2.2.1 Notation (permutation extremes, “Lane A”)

- Build expanded vectors x and y from **counts** and **scores**.
 - $r_{\max} = \text{cor}(\text{sort}(x, \downarrow), \text{sort}(y, \downarrow))$.
 - $r_{\min} = \text{cor}(\text{sort}(x, \downarrow), \text{sort}(y, \uparrow))$.
-

2.2.2 A) Same number of categories, but $|r_{\min}| \neq r_{\max}$ (concrete numbers)

Counts (both

$$K = 5$$

):

- $$X : [100, 200, 300, 200, 200]$$
- $Y : [100, 100, 100, 300, 400]$ (skewed; not palindromic)

Scores: 0:4 for both.

- $$r_{\max} \approx 0.929398758$$
- $$r_{\min} \approx -0.881118303$$
- $$|r_{\min}| \approx 0.8811 \neq r_{\max}$$

Affine recode (10,20,30,40,50) for both:

- $r_{\max}, r_{\min}$ **unchanged** (up to rounding).

Non-affine monotone recode for Y : $[0, 1, 1.5, 3, 10]$.

- $$r_{\max} \approx 0.930261031$$
- $$r_{\min} \approx -0.863250703$$
- Values change (Pearson is **not** invariant to non-affine spacing); magnitudes still unequal.

Why unequal magnitudes? No palindromic symmetry in Y — the “palindrome-sum” pattern in y' -space isn’t zero, so $S_{\max} + S_{\min} \neq 0$.

2.2.3 B) Different numbers of categories, yet $|r_{\min}| = r_{\max}$ (concrete numbers)

Counts:

- X (4 categories): [100, 200, 300, 400]
- Y (5 categories): [100, 200, 400, 200, 100] (**palindromic**)

Scores: X : 0:3, Y : 0:4.

•

$$r_{\max} \approx 0.912870929$$

•

$$r_{\min} \approx -0.912870929$$

- $|r_{\min}| = r_{\max}$ (to machine precision)

Affine recode (e.g., 10,20,...):

- Magnitudes remain **identical**; affine transforms don't change Pearson correlation.

Non-affine monotone recode for Y : [0, 1, 1.5, 3, 10].

•

$$r_{\max} \approx 0.580072921$$

•

$$r_{\min} \approx -0.842041337$$

- **Magnitudes no longer match** because antisymmetry of the **scores** about the center is lost, even though the **counts** are palindromic.

Why equal magnitudes in the affine case? Palindromic **mass** symmetry in $Y + \text{antisymmetric scoring}$ (any affine map of $0:(K_Y - 1)$) the palindrome-sum vector $y'_i + y'_{n+1-i} \equiv 0$. Hence $S_{\max} + S_{\min} = 0$ $|r_{\min}| = r_{\max}$ for **any** X , irrespective of $K_X \neq K_Y$.

2.2.4 C) Spearman invariance (same data as in B)

Compute ranks (with average ties) **once** for x and y , then form extremes by sorting those rank vectors.

- With Y scored 0:4:
 $r_{\max}^{\text{Spearman}} \approx 0.920837215$, $r_{\min}^{\text{Spearman}} \approx -0.920837215$.
- With Y scored non-affinely [0, 1, 1.5, 3, 10]:
identical Spearman extremes: 0.920837215 and -0.920837215 .

Spearman is invariant to any strictly monotone recode because ranks depend only on order, not spacing.

2.3 what changes and what doesn't (Pearson vs Spearman)

- **Pearson:**
 - **Affine** recodes $a + b \cdot \text{score}$ (with $b > 0$) leave $r_{\max}, r_{\min}$, and any $|r_{\min}| = r_{\max}$ conclusions **unchanged**.
 - **Non-affine monotone** recodes can change $r_{\max}, r_{\min}$ and can break $|r_{\min}| = r_{\max}$ even with palindromic counts.
- **Spearman:**
 - Any strictly monotone recode leaves extremes **unchanged**.
 - Palindromic counts in Y (with the usual symmetric rank scheme) give $|r_{\min}| = r_{\max}$.

2.4 Population Bounds

In this section we further explore the theoretical properties on $r_{\min}$ and $r_{\max}$.

2.4.1 Lemma (signs of extreme correlations under monotone scoring)

Lemma 2.2. *Let X and Y be ordinal variables. Assign numeric scores to categories via ****non-decreasing**** maps s_X, s_Y that respect the category order (scores may be affine or nonlinear; ties allowed). Define two extreme couplings:*

- ****Comonotone (upper)****: pair larger scored values of X with larger scored values of Y .
- ****Antitone (lower)****: pair larger scored values of X with smaller scored values of Y .

*Let $r_{\max}$ and $r_{\min}$ be the Pearson correlations computed from these couplings, either (i) in the ****finite-sample/permutation**** sense (sort scored observations in the same vs. opposite order), or (ii) in the ****Fréchet–Hoeffding (FH)/population**** sense (pair quantiles $Q_X(u)$ with $Q_Y(u)$ vs. $Q_Y(1-u)$).*

Then:

1. $r_{\max} \geq 0$ and $r_{\min} \leq 0$.
2. Moreover, $r_{\max} > 0$ and $r_{\min} < 0$ ****iff**** both variables have ****positive variance**** under the chosen scoring (i.e., neither scored variable is almost surely constant).

(If you reverse the scoring of one variable (decreasing map), swap "upper" and "lower". For Spearman's ρ , take the scores to be midranks—monotone by construction—and the same conclusions hold.)

Proof. Proof

We give parallel proofs in the finite-sample and population settings.

A) Finite-sample (permutation) setting

Let $x_{(1)} \geq \dots \geq x_{(n)}$ and $y_{(1)} \geq \dots \geq y_{(n)}$ be the **scored** observations of X and Y sorted in nonincreasing order (monotone scoring ensures sorting by category equals sorting by score). Center them:

$$x'_i := x_{(i)} - \bar{x}, \quad y'_i := y_{(i)} - \bar{y}, \quad \bar{x} := \frac{1}{n} \sum_i x_{(i)}, \quad \bar{y} := \frac{1}{n} \sum_i y_{(i)}.$$

Define the two extreme centered sums

$$S_{\max} := \sum_{i=1}^n x'_i y'_i, \quad S_{\min} := \sum_{i=1}^n x'_i y'_{n+1-i}.$$

- **Signs.**

Chebyshev's sum inequality (rearrangement for similarly/ oppositely ordered sequences) gives

$$\frac{1}{n} \sum_i x'_i y'_i \geq \left(\frac{1}{n} \sum_i x'_i \right) \left(\frac{1}{n} \sum_i y'_i \right) = 0, \quad \Rightarrow \quad S_{\max} \geq 0,$$

and, because y'_{n+1-i} is **nondecreasing** in i ,

$$\frac{1}{n} \sum_i x'_i y'_{n+1-i} \leq \left(\frac{1}{n} \sum_i x'_i \right) \left(\frac{1}{n} \sum_i y'_{n+1-i} \right) = 0, \quad \Rightarrow \quad S_{\min} \leq 0.$$

Dividing by $\sigma_X \sigma_Y > 0$ (the standard deviations of the scored variables) yields $r_{\max} \geq 0$ and $r_{\min} \leq 0$.

- **Strictness ($\Leftrightarrow$ positive variances).**

Suppose $\sigma_X > 0$ and $\sigma_Y > 0$ (i.e., neither $\{x'_i\}$ nor $\{y'_i\}$ is identically zero). The rearrangement inequality says $S_{\max}$ is the **maximum** and $S_{\min}$ the **minimum** of $\sum_i x'_i y'_{\pi(i)}$ over all permutations π ; moreover $S_{\min} < S_{\max}$. The **average** over all permutations equals

$$\frac{1}{n!} \sum_{\pi} \sum_i x'_i y'_{\pi(i)} = \sum_i x'_i \left(\frac{1}{n!} \sum_{\pi} y'_{\pi(i)} \right) = \sum_i x'_i \cdot \frac{1}{n} \sum_j y'_j = 0.$$

Since the minimum is $<$ the maximum and the average is 0, it follows that $S_{\min} < 0 < S_{\max}$. Hence $r_{\min} < 0 < r_{\max}$.

Conversely, if $\sigma_X = 0$ or $\sigma_Y = 0$, then one centered sequence is identically zero, so $S_{\max} = S_{\min} = 0$ and the correlations are undefined or (in a limiting sense) attain 0. This matches the “only if” direction.

B) Population (FH/quantile) setting

Let $Q_X, Q_Y : [0, 1] \rightarrow \mathbb{R}$ be the quantile functions of the **scored** variables. Set

$$f(u) := Q_X(u) - \mu_X, \quad g(u) := Q_Y(u) - \mu_Y,$$

so $\int_0^1 f = \int_0^1 g = 0$. Monotone scoring preserves category order, hence Q_X, Q_Y are nondecreasing; therefore f, g are nondecreasing functions.

- **Upper/comonotone covariance:**

$$\text{Cov}_{\max} = \int_0^1 f(u) g(u) du \geq \left(\int_0^1 f \right) \left(\int_0^1 g \right) = 0$$

by Chebyshev's **integral** inequality for similarly ordered functions.

- **Lower/antitone covariance:**

$$\text{Cov}_{\min} = \int_0^1 f(u) g(1-u) du \leq \left(\int_0^1 f \right) \left(\int_0^1 g(1-u) \right) = 0,$$

since f is nondecreasing and $g(1-u)$ is **nonincreasing**.

Dividing by $\sigma_X \sigma_Y > 0$ yields $r_{\max} \geq 0$ and $r_{\min} \leq 0$.

- **Strictness.**

If both variables have positive variance, then f and g are not a.e. constant. For nontrivial nondecreasing f, g , Chebyshev's integral inequalities above are **strict** unless one function is a.e. constant. Hence $\text{Cov}_{\max} > 0$ and $\text{Cov}_{\min} < 0$, giving $r_{\max} > 0$ and $r_{\min} < 0$. Conversely, if either variance is 0, both covariances are 0.

This completes the proof. $\square$ ■

2.5 Asymetry

Symmetry and the Role of Ties

- Conditions under which $r_{\min} = -r_{\max}$

PROP: If one marginal is **palindromically symmetric** about its center rank, then $r_{\min} = -r_{\max}$.

(Cees - *r_min MAGNITUDE Asymmetry in Pearson Correlation.md*)

Necessary/sufficient symmetry condition (precisely stated)

- **Pair-specific equality** $|r_{\min}| = r_{\max} \iff \sum_i x'_i (y'_i + y'_{n+1-i}) = 0$.
(Can happen by coincidence even if Y isn't symmetric.)
- **Universal in X equality** (for **all** X):
Necessary and sufficient that **one** marginal (say Y) has

(i) **palindromic mass symmetry** $p_Y(k) = p_Y(K_Y - 1 - k)$ and

(ii) **antisymmetric scoring** $s_Y(k) + s_Y(K_Y - 1 - k) = \text{const}$ (true for any affine map of $0:(K_Y - 1)$).

This condition is **independent of K_X vs K_Y** .

2.5.1 1) setup (permutation extremes for a single sample)

Let $x_{(1)} \geq \dots \geq x_{(n)}$ and $y_{(1)} \geq \dots \geq y_{(n)}$ be the sorted sample values (or expanded from counts). Define centered sequences

$$x'_i := x_{(i)} - \bar{x}, \quad y'_i := y_{(i)} - \bar{y},$$

and the two permutation extremes

$$S_{\max} = \sum_{i=1}^n x'_i y'_i, \quad S_{\min} = \sum_{i=1}^n x'_i y'_{n+1-i},$$

with $r_{\max} = S_{\max}/(\sigma_X \sigma_Y)$, $r_{\min} = S_{\min}/(\sigma_X \sigma_Y)$.

We always have $S_{\max} \geq 0$ and $S_{\min} \leq 0$. Magnitude symmetry $|r_{\min}| = r_{\max}$ is equivalent to

$$S_{\max} + S_{\min} = 0.$$

2.5.2 2) pair-specific necessary & sufficient condition

Let

$$z_i := y'_i + y'_{n+1-i}.$$

Then

$$S_{\max} + S_{\min} = \sum_{i=1}^n x'_i z_i.$$

Hence, for this *particular* pair (x, y) ,

$$\boxed{|r_{\min}| = r_{\max} \iff \sum_i x'_i z_i = 0} \quad ()$$

(i.e., the centered X sequence is orthogonal to the “palindrome-sum” vector z).

This can hold even if Y is not symmetric—by coincidence of that inner product being zero.

2.5.3 3) universal condition (independent of X)

If you want $|r_{\min}| = r_{\max}$ to hold for **every** possible X (every centered, sorted x'), then () must hold for all x' . The only way is

$$z_i \equiv 0 \quad \text{for all } i,$$

i.e.

$$y'_i + y'_{n+1-i} \equiv 0 \quad \forall i.$$

Translating back to categories and scores, a **necessary and sufficient universal condition** is:

- **Palindromic mass symmetry for Y :** $p_Y(k) = p_Y(K_Y - 1 - k)$ for all k , and
- **Antisymmetric scoring for Y :** the score map satisfies $s_Y(k) + s_Y(K_Y - 1 - k)$ is constant (true for affine scores like $0, 1, \dots, K_Y - 1$, and for Spearman midranks under palindromic counts).

Under these, $z_i \equiv 0$ and thus $|r_{\min}| = r_{\max}$ for any X .

If either mass symmetry or score antisymmetry fails, you can generally find an X for which $|r_{\min}| \neq r_{\max}$ (often $|r_{\min}| > r_{\max}$ when Y is skewed).

- How ties in marginal distributions affect the attainability of the lower bound
- 1. $r_{\max} = \mathbf{FH \text{ upper (comonotonic), always attainable}}$ in discrete ordinal settings.
- 2. $r_{\min}$ may **not** equal FH lower in discrete settings; use best feasible opposite-order arrangement.
- 3. No universal bound on relative magnitudes: $|r_{\min}|$ may be less than, equal to, or greater than $r_{\max}$.
- 4. Palindromic symmetry of one marginal guarantees $|r_{\min}| = r_{\max}|$.

VH 3. Symmetry Analysis:

- (3a) Visualize and explore how changes from symmetric to asymmetric marginal distributions influence the joint distributions and, consequently, the correlation. Consider the potential role of entropy in measuring asymmetry.
- 4. **Investigate when the minimum and maximum correlations are asymmetric ($r_{\min} \neq -r_{\max}$):**
 - Show why and under what conditions this asymmetry occurs.
 - Visually investigate why perfect negative and positive correlations may not be symmetric in magnitude given ordinal marginals.

Special Cases and Boundary Conditions

- Equal vs. unequal numbers of categories
- Symmetric vs. asymmetric marginal distributions

Often $|r_{\min}| < r_{\max}$ for discrete asymmetrical marginals.

(*BFbounds.md*; *Ordinal Variables and Correlation.md*) - **New evidence (BES data)**: Possible to have $|r_{\min}| > r_{\max}|$.

Example: X with 4 categories, Y with 5 categories, skewed marginals;

$r_{\max} = 0.85027$, $r_{\min} = -0.89916$.

(*Cees - r_min MAGNITUDE Asymmetry in Pearson Correlation.md*)

Reason: Pearson correlation uses centered values; skewness can make negative deviations in antitone pairing larger in magnitude than positive deviations in comonotone pairing.

VH

2.6 Estiimation and Joint Uncertainty

VH

3 Measurement Issues and Metrics of Constraint

IN THIS SECTION: “How to define metrics (on *pairs* of discrete probability distributions) to *explain*:

VH

1.
$$\frac{r_{\max} - r_{\min}}{2}$$
2.
$$1 - r_{\max}$$
3.
$$| -1 - r_{\min} |$$

Or analogs for $\hat{r}^2$:

1.
$$\frac{r_{\max}^2 - r_{\min}^2}{2}$$
- 1a.
$$\frac{(r_{\max} - r_{\min})^2}{2}$$
2.
$$1 - r_{\max}^2$$
3.
$$1 - r_{\min}^2$$

As demonstrated above, the extent to which marginal distributions of orcats constrain the range of attainable value of r varies considerably. In the few illustrative examples presented in Table 2 r_{min} varied between -1 and -0.71, while r_{max} varied between 0.32 and 1. Moreover, in Table 2 r_{min} and r_{max} do not vary in lockstep but at least to some extent independent of each other.^[fn: Table 2 is a poor basis to assess the functional relationship between r_{min} and r_{max} as it contains a small number of handpicked situations. This relationship will be investigated in detail later, in section ??.] This raises the question how strong the overall constraining effect is that marginal distributions exert on the attainable values of r . It also raises the question of what factors affect the strength of this overall constraining effect. In this section we describe two ways in which the strength of constraints can be defined. Later (in section ??) we turn to the questions of the conditions that drive the strength of this effect. Perhaps the most obvious way to express the strength of the marginal constraints on the attainable values of r is by comparing the width of the interval between r_{min} and r_{max} with that of the ubiquitously used reference interval $[-1, 1]$. As the width of the latter interval is 2, it is advantageous to express the strength of the overall constraining effect –to be denoted as C_1 as

$$C_1 = \frac{|r_{min}| + |r_{max}|}{2}$$

C_1 reflects the proportion of the range of the attainable values of r compared to the theoretically maximum width of this range. Applying this measure to the illustrative cases listed in Table 2 shows that C_1 varies. Its highest value is 1, reflecting the absence of any constraining effect (see examples (d) and (f) in Table 2). The lowest value of C_1 in Table 2 is for example (i), where it is 0.639. In other words, the marginal distributions of the two variables in that example restrict the range of its attainable values to 63.9% of the traditional $[-1, 1]$ reference interval. Some might object to the C_1 measure on the grounds that it assumes that r (and thus also r_{min} and r_{max}) is expressed at interval level and that one could thus state validly that a correlation of 0.4 reflects an association twice as strong as one of 0.2. That would be so if numerical differences between correlations correspond to similar differences in observable aspects of data [insert references to correspondence NRS to ERS]. That not being the case, one could instead compare r^2 s –percentages of shared variance– rather than r s. ^[fn: This perspective is not uncontested, however, and its usefulness depends on the purpose of inspecting correlations: as reflections of the association between variables, or as reflections of associations between indicators of a latent concept (for an elaboration of this difference, cf. Ozer 1985).] This would lead to a somewhat different measure of the strength of the constraining effect of marginal distributions on the attainable values of r , as follows:

$$C_2 = \frac{r_{min}^2 + r_{max}^2}{2}$$

For the examples in Table 2, the values C_2 are, respectively, 0.701; 0.755; 0.715; 1; 0.831; 1; 0.809; 0.915 and 0.510. Obviously, C_1 and C_2 are correlated, but how strongly cannot be assessed very well from the few handpicked examples in Table 2, but it will be considered below, using more encompassing sets of data.

3.1 Possible “Metrics of Constraint”

3.1.1 Overlap-Based Measures of Asymmetry

In addition to Total Variation (TV) distance, other metrics can quantify the structural dissimilarity between a marginal distribution and its reverse. One natural approach is to assess the degree of **overlap** between the original distribution p and its reverse p^{rev} , defined by reversing the order of the categories.

We consider three complementary measures of overlap and their complements:

3.1.1.1 (i) Total Variation (TV) Distance

The TV distance, already introduced in Section 5.5, quantifies the maximum absolute discrepancy between the two distributions:

$$\text{TV}(p, p^{\text{rev}}) = \frac{1}{2} \sum_{i=1}^K |p_i - p_{K+1-i}|.$$

3.1.1.2 (ii) Bhattacharyya Coefficient (BC) and its Complement

The **Bhattacharyya Coefficient** provides a geometric measure of similarity:

$$\text{BC}(p, p^{\text{rev}}) = \sum_{i=1}^K \sqrt{p_i \cdot p_{K+1-i}}.$$

Its complement, $1 - \text{BC}$, can be interpreted as a divergence-like measure that reflects geometric differences between the original and reversed distributions. While not a metric, it is bounded between 0 and 1, with 0 indicating perfect symmetry.

3.1.1.3 (iii) Overlap Coefficient (OVL)

The **Overlap Coefficient**, sometimes referred to as the Szymkiewicz–Simpson coefficient, captures the shared probability mass:

$$\text{OVL}(p, p^{\text{rev}}) = \sum_{i=1}^K \min\{p_i, p_{K+1-i}\}.$$

The complement $1 - \text{OVL}$ thus reflects the fraction of mass that is not shared between the two distributions.

These measures differ in their interpretive emphasis: - TV focuses on the total discrepancy. - BC captures geometric congruence. - OVL captures the total shared area between the distributions.

In our simulation study (Section), we compute all three diagnostics to illustrate how marginal asymmetry correlates with asymmetry in achievable Pearson correlation bounds. These overlap-based measures are implemented in the R package `overlap` (Ridout & Linkie, 2009), which facilitates reproducible estimation and visualization.

Empirically, we find that all three metrics provide consistent, interpretable diagnostics of distributional asymmetry. Using them jointly provides a robust triangulation of symmetry breaking, especially in applied contexts where ordinal variables are discretized from latent traits or ratings.

4 Implications, Issues, and Questions

•

4.1 Interpretation of ‘Effect Size’

- Revisiting Cohen’s benchmarks in light of constrained bounds $\rightsquigarrow$ Interpretation of ‘effect size’ (Cohen et al. (2002))
- Risk of over- or under-stating association.

4.2 Comparison Across Studies / Measures

- Non-comparable bounds across datasets or variable codings.
- Need for bound-aware effect size metrics.

4.3 Hypothesis Testing

- t-test can mislead: Example from BES: (i) t-test sig; perm test not; (ii) visa versa
- Why p-values based on $[-1, 1]$ bounds may mislead $\rightsquigarrow$ need for Adjusted null distributions
- **Permutation distribution:** Generate full vectors from marginals, independently permute X and Y , compute r , repeat. Produces null distribution centered at 0. (*Proofs-OrdinalVariables.md*)

4.4 Rescaling

So, what to do? *One* idea: **linear rescaling** (preserving zero)

- Adjusted null distributions (via permutation or simulation).

4.5 Variance “Explained”

- Interpretation of r^2 is ambiguous for at least two reasons:
 1. ordcats are invariant to monotone transformation (variance is a *cardinal* property, simply not appropriate for ordinal variables)

2. is r_{max}^2 the maximal ‘amount’ of variance between the two orcats? How is this quantity to be interpreted? How is r_{min}^2 to be interpreted?

5 Monte Carlo Simulations

To investigate factors that affect the bounds of attainable values of r and thus the strength of constraints on r exerted by marginal distributions we use two different sets of data. Our first dataset is simulation generated and covers in principle all possible combinations of marginal distributions for two orcats with given numbers of categories and given n . We simulate orcats containing, respectively, 4, 5, 6, 7, 10 and 11 categories, these being by far the most common in empirical data deriving from survey studies. For each of these we generate a large number of variables, each with its own simulated univariate distribution, and thus also its own form (modality, symmetry, etc.), mean, variance, skew and kurtosis as follows. [insert description, detailing n , how many univariate distributions we generate for each number of categories; how to populate the categories, whether we impose distributional characteristics such as normality, modality, skew etc. Btw: my preference would be to impose as little as possible, and derive characteristics such as mean, variance, skew, kurtosis from the distributions generated.] We subsequently randomly select pairs of simulated variables and determine for each r_{min} , r_{max} , $C1$ and $C2$. From the resulting simulated pairs of orcats we delete the cases in which one, or both of the variables have zero variance. This results in a dataset with $n = xxx$ [insert very large number] that forms the basis of subsequent analyses. Further details are specified in Appendix 3. This simulated dataset constitutes, within the bounds of granularity of the procedure, a random sample all possible combinations of univariate distributions of pairs of orcats, and can therefore be used as an adequate basis for investigating functional relationships between sundry properties of pairs of orcats and the range of attainable values of r for those pairs. Although this simulated dataset randomly represents all possible combinations of univariate distributions of pairs of orcats, it cannot be considered to be a random sample of pairs of orcats that applied researchers would encounter in actual, empirical data such as derived from, e.g., mass surveys. This is because, when designing survey questions, researchers tend to avoid formulations that they expect to yield little variance in the responses. Therefore, we use a second dataset to investigate constraints on the attainable range of values of r , that is derived from the British Election Study 2019. [fn: insert details about BES2019]. In this study we identified xxx orcats, which define 7503 pairs, for each of which we calculated pair-wise distributional characteristics including r_{min} , r_{max} , $C1$ and $C2$. Further details on this dataset are provided in Appendix 4. Obviously, the orcats in BES2019 are not a random sample from the contents of all socio-political mass surveys in Britain in 2019, let alone of those in wider areas such as the developed western world, or in other years. But the BES can sensibly be regarded as typical for such studies, which include (national) election studies, general social surveys, and many public opinion and marketing surveys. As such, findings based on this dataset derived from the BES can sensibly be regarded as typical for what applied researchers in other countries will encounter in their survey data. These two datasets will be used to address, in the subsequent sections, questions with regard to: - Distributions of r_{min} , r_{max} and measures of overall constraint of r by marginals - Factors affecting r_{min} , r_{max} and overall constraint of r - Consequences of con-

strained correlations in applied research, including for descriptive purposes and hypothesis testing - Possibilities for reducing undesired consequences of constrained correlations

VH: (r_{min}, r_{max})

5.1 Effects of Marginal Shapes

- Shape-based elasticity of correlation bounds
- Skewness and modality heavily influence the attainable range of r .
- Marginals with extreme concentration (e.g., most responses in one or two categories) sharply restrict the possible correlation range.

5.2 Symmetry Breaking in Correlation Bounds

- When and why $r_{min} \neq -r_{max}$
- Empirical indicators of bound asymmetry
- When given marginal distributions are sufficiently asymmetric, the min and max bounds of r are not symmetric about zero.
- The shape of the null distribution of r can be strongly skewed, especially in cases with unequal marginals.

When we talk about “asymmetry” in this context, we’re referring to how different the distribution of Y is from the distribution of X . Since X is kept uniform, any deviation of Y from uniformity creates asymmetry between the distributions.

The *total variation distance* (TV) quantifies this asymmetry:

- It measures how different Y is from being symmetric around its midpoint
- When $TV = 0$: Y is symmetric (like X)
- When TV is large: Y is highly asymmetric compared to X

6 Illustration with ‘Typical’ Data: BES

We do not claim this data is representative of data-sets of ordinal variables used by social scientists. Rather, we view it as *typical* in that it is a real-world example possessing many of the features that would raise issues...

VH: (r_{min}, r_{max}) And compare with MC simulated data

7 Discussion

7.1 Comparing ‘Typical’ Data (BES) with Simulated Data

7.2 Relation to Literature

!fullcite camb76-inequalities

Full citation:

Cambanis et al. (1976)

Grether, D. M. (1976). On the Use of Ordinal Data in Correlation Analysis. *American Sociological Review*, 41(5), 908–912. <https://doi.org/10.2307/2094741>

Labovitz, S. (1967). Some observations on measurement and statistics. *Social Forces*, 46(2), 151–160. <https://doi.org/10.2307/2574595>

Labovitz, S. (1970). The assignment of numbers to rank order categories. *American Sociological Review*, 35(3), 515–524. <https://doi.org/10.2307/2092993>

Moss, J., & Grønneberg, S. (2023). Partial Identification of Latent Correlations with Ordinal Data. *Psychometrika*, 88(1), 241–252. <https://doi.org/10.1007/s11336-022-09898-y>

O’Brien, R. M. (1979). The use of Pearson’s r with ordinal data. *American Sociological Review*, 44(5), 851–857. <https://doi.org/10.2307/2094532>

Robitzsch, A. (2022). On the Bias in Confirmatory Factor Analysis When Treating Discrete Variables as Ordinal Instead of Continuous. *Axioms*, 11(4), Article 4. <https://doi.org/10.3390/axioms11040162>

Fréchet (1951)

Whitt (1976)

Hoeffding (1940)

Lehmann (1966)

Tchen (1980)

Cambanis et al. (1976)

Shih & Huang (1992)

Alwin & Krosnick (1991); Brewer & Gross (2005); Flora & Curran (2004); Gaito (1980); Green et al. (1997); Jacoby (1999); Kendall (1938); Knoke et al. (2002); Kreuter & Valliant (2007); Krosnick & Fabrigar (1997); Labovitz (1970); Olsson (1979); Spearman (1904)

7.3 Recommendations

- Bound-aware effect size interpretation.
- Bound-aware hypothesis testing.

- Possible inclusion in statistical software.

8 Conclusion

Summarize:

- Main theoretical findings.
- Key applied consequences.
- Summary: Theoretical bounds, symmetry-breaking conditions, and implications for applied research
- An R package accompanies this paper, providing tools to compute max/min bounds and perform randomization-based hypothesis tests for fixed marginals. Details are available in the appendix and online.

Impacts standard methods:

- interpretation (e.g. ‘effect size’) and comparison
- hypothesis testing

Recommendations for interpreting Pearson’s r in applied contexts

- Reporting observed r in context of its attainable bounds.
- Using our tools to test whether an observed r is unusually strong or weak given marginals.

8.1 Extensions and Open Questions

- Role of entropy and joint uncertainty in bounding behavior
- Possibility of adjusting or normalizing r
- Relation to Copula Theory (Optional)
 - “Although our setting is discrete, the problem is conceptually linked to copula-based modeling of dependence with fixed marginals. The connection is discussed as a direction for future work.”
- ‘Correction’ strategies $\rightsquigarrow$ *rescaling*
- Generalizations to higher dimensions $\rightsquigarrow$ implications for correlation matrices and Factor Analysis, SEMs, etc.
 - **PCA and Factor Analysis:** Attenuated correlations may lead to $\rightsquigarrow$ practical risks: over-factoring (more factors estimated than ‘should be’), biased structural coefficients(?)
 - * **SEM:** Biased estimation of structural paths if ordinal-level attenuation is ignored.

8.1.1 Correlation Matrices

The consequences of using Pearson correlations with ordinal data extend beyond descriptive association. In factor analysis and SEM, attenuated correlations can yield distorted loadings, inflated factor numbers, and biased structural coefficients. Empirical studies confirm this risk: (Flora & Curran, 2004) show that CFA models fitted with Pearson’s r among ordinal indicators produce biased estimates compared to models using polychoric correlations; (Green et al., 1997) demonstrate how the number of response categories alters apparent model fit. More broadly, psychometric research emphasizes that ignoring the ordinal nature of indicators can misrepresent latent structures ((Lord, 1952); (Muthén, 1984); (Bartholomew et al., 2011)). For survey-based political science and sociology, this means that correlations reported between Likert items are not only bounded by their marginals, but may also systematically bias downstream modeling.

Since the possible values r may take on depend on the marginal distributions of the two variables being correlated, and since this reference interval may – and often is – quite different for different pairs of variables, interpreting and working with correlation matrices (alternatively variance-covariance matrices) must be paid special attention. Such matrices form the basis for a number of statistical techniques, from regression to Factor Analysis or Structural Equation Modeling (CITES). In this section we present preliminary results exploring the implications of our findings above for PCA, FA, SEM.

First, some preliminaries. It is well known that any correlation matrix (using completely cases – i.e. when each pair of variables is observed for the same set of respondents), S , is positive semi-definite and symmetric. Hence, S is invertable. However, ...

Why PSD Matters for PCA:

- **Mathematically:** Eigen-decomposition works for any symmetric S , PSD or not.
- **Interpretively:** Negative eigenvalues break the “variance explained” story (variance cannot be negative).
- **Software behavior:** Most PCA routines check PD/PSD; failure can cause warnings or dropped components. (*PCA and FA PSD requirement.md*)

If Input is Symmetric but not PSD:

Example:

$$\mathbf{A} = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}, \quad \lambda = (3, -1),$$

is indefinite (one negative eigenvalue).

- PCA math still runs; eigenvalues/vectors computed.
- Interpretation fails: $\sqrt{\lambda_j}$ imaginary for $\lambda_j < 0$.
- Some software refuses input; others run but return nonsensical variance metrics. (*PCA and FA PSD requirement.md*)

PSD/PD Needs in FA:

- **ML FA:** Requires $\mathbf{S}$ be PSD; $\log \det \mathbf{S}$ undefined if indefinite.
- **Principal-axis / Image FA:** Also break with negative eigenvalues (square roots fail). (*PCA and FA PSD requirement.md*)

SEM and PSD Requirements

8.1.1.1 SEM Machinery

Observed covariance $\mathbf{S}$; implied covariance $\Sigma(\theta)$ from structural and measurement models. (*PCA and FA PSD requirement.md*)

8.1.1.2 Fit Functions and PD Needs

- **ML / GLS:** Require both $\mathbf{S}$ and $\Sigma(\theta)$ PD.
 - **WLS/DWLS:** Need PD indirectly (weight matrix invertible).
 - **ULS:** Can start from indefinite $\mathbf{S}$, but test statistics again require PD. (*PCA and FA PSD requirement.md*)
-

8.1.1.3 What Happens if Not PSD

- Software errors out (“matrix is not positive definite”).
 - Log-determinant undefined; inversion fails.
 - Even if bypassed, optimisation steps hit complex/NaN values. (*PCA and FA PSD requirement.md*)
-

8.1.1.4 Remedies in SEM

- Nearest PD smoothing (Higham 2002).
 - Ridge adjustment.
 - Analyse raw data.
 - Check rounding issues. (*PCA and FA PSD requirement.md*)
-

Core Lessons Across PCA, FA, SEM

1. **Theoretical PSD:** Covariance matrices from real data are PSD by definition.
2. **Numerical Issues:** Sampling noise, rounding, or listwise deletion can yield small negative eigenvalues.
3. **PCA:** Can compute from indefinite matrix but interpretation breaks.
4. **FA and SEM:** Require PD input; will not run on indefinite matrices.
5. **Repairs:** Adjust to nearest PSD, ridge, or recompute from raw data.

So, we consider linearly re-scaling the estimated r for each pair of variables in the correlation matrix S . Call this new rescaled correlation matrix S' . We seek to understand what the process of pair-wise linear rescaling has on the PSD-ness and invertability of S' . This is important because without an invertable correlation matrix, several analyses scholars perform on S become questionable (at best). Further, some estimators used in e.g. FA require PSD (namely, MLE).

The general result is that linear rescaling does not ‘break’ invertability (nor PSD) but does make invertability ‘worse’ vizaviz the condition number. [Discuss condition number here]

9 References

- Alwin, D. F., & Krosnick, J. A. (1991). The reliability of survey attitude measurement: The influence of question and respondent attributes. *Sociological Methods & Research*, 20(1), 139–181.
- Bartholomew, D. J., Knott, M., & Moustaki, I. (2011). *Latent Variable Models and Factor Analysis: A Unified Approach* (3rd ed.). Wiley.
- Brewer, P. R., & Gross, K. (2005). Values, framing, and citizens' thoughts about policy issues: Effects on content and quantity. *Political Psychology*, 26(6), 929–948.
- Cambanis, S., Simons, G., & Stout, W. (1976). Inequalities for $E k(X, Y)$ when the marginals are fixed. *Zeitschrift Für Wahrscheinlichkeitstheorie Und Verwandte Gebiete*, 36(4, 4), 285–294. <https://doi.org/10.1007/BF00532695>
- Cohen, J., Cohen, P., West, S. G., & Aiken, L. S. (2002). *Applied Multiple Regression/Correlation Analysis for the Behavioral Sciences* (3 edition). Routledge.
- Flora, D. B., & Curran, P. J. (2004). An empirical evaluation of alternative methods of estimation for confirmatory factor analysis with ordinal data. *Psychological Methods*, 9(4), 466–491.
- Fréchet, M. (1951). Sur les tableaux de corrélation dont les marges sont données. *Annales de l'Université de Lyon*, 14, 53–77.
- Gaito, J. (1980). Measurement scales and statistics: Resurgence of an old misconception. *Psychological Bulletin*, 87(3), 564–567.
- Green, S. B., Akey, T. M., Fleming, K. K., Hershberger, S. L., & Marquis, J. G. (1997). Effect of number of scale points on estimates of fit in confirmatory factor analysis. *Structural Equation Modeling*, 4(3), 244–260.
- Hardy, G. H., Littlewood, J. E., & Pólya, G. (1952). *Inequalities*. Cambridge University Press.
- Hoeffding, W. (1940). Masstabinvariante Korrelationstheorie. *Schriften Des Mathematischen Instituts Und Instituts Fur Angewandte Mathematik Der Universitat Berlin*, 5, 181–233.
- Jacoby, W. G. (1999). Levels of measurement and political research: An optimistic view. *American Journal of Political Science*, 43(1), 271–301.
- Jaekel, P., & Kahl, C. (2009). *Positive semi-definite correlation matrix completion*.
- Kendall, M. G. (1938). A new measure of rank correlation. *Biometrika*, 30(1/2), 81–93.
- Kinder, D. R., & Kam, C. D. (2009). *Us Against Them: Ethnocentric Foundations of American Opinion*. University of Chicago Press.
- Knoke, D., Bohrnstedt, G. W., & Mee, A. P. (2002). *Statistics for social data analysis* (4th ed.). Wadsworth.
- Kreuter, F., & Valliant, R. (2007). A survey on survey statistics: What is being used in practice? *Journal of Official Statistics*, 23(1), 1–21.
- Krosnick, J. A., & Fabrigar, L. R. (1997). Designing rating scales for effective measurement in surveys. In L. Lyberg, P. Biemer, M. Collins, E. de Leeuw, C. Dippo, N. Schwarz, & D. Trewin (Eds.), *Survey measurement and process quality* (pp. 141–164). Wiley.
- Labovitz, S. (1970). The assignment of numbers to rank order categories. *American Sociological Review*, 35(3), 515–524. <https://doi.org/10.2307/2092993>
- Lehmann, E. L. (1966). Some Concepts of Dependence. *The Annals of Mathematical Statis-*

- tics*, 37(5), 1137–1153.
- Lord, F. (1952). *A Theory of Test Scores*. Psychometric Society.
- Mishra, S. K. (2009). A note on positive semi-definiteness of some non-pearsonian correlation matrices. *Available at SSRN 1419362*.
- Müller, A., & Stoyan, D. (2002). *Comparison Methods for Stochastic Models and Risks*. Wiley.
- Muthén, B. (1984). A general structural equation model with dichotomous, ordered categorical, and continuous latent variable indicators. *Psychometrika*, 49(1), 115–132. <https://doi.org/10.1007/BF02294210>
- Olsson, U. (1979). Maximum likelihood estimation of the polychoric correlation coefficient. *Psychometrika*, 44(4), 443–460.
- Rüschendorf, L. (1981). Sharpness of Fréchet bounds. *Zeitschrift Für Wahrscheinlichkeitstheorie Und Verwandte Gebiete*, 57, 293–302.
- Shantal, M., Othman, Z., & Bakar, A. A. (2023). Impact of Missing Data on Correlation Coefficient Values: Deletion and Imputation Methods for Data Preparation. *Malaysian Journal of Fundamental and Applied Sciences*, 19(6, 6), 1052–1067. <https://doi.org/10.1113/mjfas.v19n6.3098>
- Shih, W. J., & Huang, W.-M. (1992). Evaluating Correlation with Proper Bounds. *Biometrics*, 48(4), 1207–1213. <https://doi.org/10.2307/2532712>
- Spearman, C. (1904). The proof and measurement of association between two things. *The American Journal of Psychology*, 15(1), 72–101.
- Tchen, A. H. (1980). Inequalities for Distributions with Given Marginals. *The Annals of Probability*, 8(4), 814–827.
- Thorisson, H. (2000). *Coupling, stationarity, and regeneration*. Springer-Verlag.
- Whitt, W. (1976). Bivariate Distributions with Given Marginals. *The Annals of Statistics*, 4(6), 1280–1289.

A Appendix A: Mathematical Proofs

A.1 Proof A.1

Proof of Proposition (?).

A.2 Proof A.2

Proof of Corollary 1.

B Appendix B: Software Tools and Code Listings

- Documentation and examples for the R package
- Description of randomization testing features

C Appendix C: Additional Tables and Simulations

- Supplementary empirical examples, simulation results

D Appendix D: Old Outline

1. Introduction and Conceptual Framework

- Research motivation and objectives
- Overview of limitations in current correlation testing with ordinal
- Why does this matter? $\leadsto$ consequences if these changing bounds are ignored

2. Theoretical Bounds on Correlation Coefficients

- Mathematical explanation of R-min and R-max
- Elasticity and asymmetry due to marginal distributions
 - Atainable and non-atainable bounds
- Implications for standard significance tests
- Note on Freshet bounds anomaly (appendix or highlighted sidebar)

3. Measurement Issues and Metrics of Constraint [TITLE?]

- Problems with interval assumptions in ordinal categorical data
- Alternatives to traditional scaling (psychometric, text-based)
- Concept of boundedness and its distortion effects
- Proposed *metrics of constraint*: Asymmetry; Aitchinson distance; Entropy(?); Total Variation(?)

4. Simulations and Monte Carlo Approaches

- Design of permutation and Monte Carlo simulations

- Bias and error analysis under null hypotheses
- Role of mean, skewness, variance in predicting constraint metrics

5. Empirical Illustration Using BES Data

- Justification for selection: a typical but not representative case
- Application of shrinkage and constraint metrics to BES
- Comparison to simulation outcomes

6. Implications for Applied Research and Tools

- Critique of statistical purism in significance testing
 - Critique of over-reliance on significance without consideration of boundedness or constraint metrics.
- Reframing effect size: from absolute value to bounded context
 - r^2 interpretation under bounded conditions
- RE-SCALING:
 1. Map r_{min} to -1, etc.
 2. Map $\hat{r}^2$ to $[0, 1]$, the re-scale $\hat{r}_{adj}$ by preserving sign(?)
 - Effect on PSD and invertability of adjusted correlation matrix (NOTE: only use complete cases here so as to not ‘cheat’ – Jaeckel & Kahl (2009); Mishra (2009); Shantal et al. (2023))
- Practical guidance for analysts: how to assess when elastic bounds affect inference.
- Overview of accompanying software tools or R functions developed.

7. Open Questions

8. Conclusions