

## STAT502 Lab #1

1. In this part, we will learn some of the basics of the statistical software R. We will learn how to read data in from a file, how to handle variables, how to plot and boxplot data, and how to save these plots.
  - (a) Create a directory within your PASS space (X drive) entitled “STAT502”. You may also wish to create subdirectories for separate labs, homework, etc.
  - (b) Launch R, and change the working directory to STAT502.
  - (c) Download “CH16PR07.txt” from Canvas into your STAT502 folder.
  - (d) In R, follow File → New script to create an R file to keep track of the commands you’ll use for this lab. Include the following lines of code. To run them, simply copy and paste into the R terminal window. You can switch among active windows in R with **ctrl+tab**.

```
p16.7 <- read.table("CH16PR07.txt")
p16.7
p16.7$V1
p16.7$V2
names(p16.7) <- c("improvement","rd","observation")
p16.7
p16.7$improvement
p16.7$imp #don't need to specify the whole name
p16.7$rd
plot(p16.7$imp)
plot(p16.7$rd,p16.7$imp)
boxplot(improvement~rd,data=p16.7)
p16.7$rd
p16.7$rd <- factor(p16.7$rd,labels=c("low","moderate","high"))
p16.7$rd
boxplot(improvement~rd,data=p16.7)
mean(p16.7$improvement)
sd(p16.7$improvement)
tapply(p16.7$improvement,p16.7$rd,mean)
tapply(p16.7$improvement,p16.7$rd,sd)
p16.7[2,3]; p16.7[,1]; p16.7[2,]; p16.7[,]
```

2. Download the “gpa.dat” data set from Canvas to work through the example we did in class. Use the following command to get the data into R.

```
gpa = read.table("gpa.dat",header=T)
```

- (a) Construct side-by-side boxplots for the two groups. Does it appear that there is a significant difference?
- (b) For each group, find the sample mean, variance, and sample size. Also, find the pooled sample variance.
- (c) Compute the test statistic for  $H_0 : \mu_1 - \mu_2 = 0$  versus  $H_a : \mu_1 - \mu_2 \neq 0$ :

$$t = \frac{\bar{x}_1 - \bar{x}_2}{s_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$$

- (d) Use the R functions `qt()` and `pt()` to find the critical value and  $p$ -value, respectively, for making a decision about  $H_0$  (assuming  $\alpha = .05$ ). Use the command `?qt` to see documentation for how these functions are used.
- (e) Construct a 95% confidence interval for  $\mu_1 - \mu_2$ .
- (f) Now use the R function `t.test()` to carry out the above test. If you use the options correctly (check the documentation), you should get the same answer (it also provides confidence limits).