SMR: LEICOV. Shapes, Materials and Regularity: Lecce Encounters In Calculus Of Variations

Organizers: Paolo Acampora (Università del Molise), Alessandro Carbotti, Simone Cito, Luigi Negro and Francesco Solombrino (Università del Salento)

SMR:LEICOV is a series of three meetings to bring together young researchers and experts in the field of Calculus of Variations and Partial Differential Equations. We want to give value to the intersections between shape optimization, materials science and regularity theory.

List of confirmed speakers (14)

1. Vincenzo Amato, Scuola Superiore Meridionale, Italy
   Thu 01 Oct, 16:45
2. Luciana Angiuli, Università del Salento, Italy
   Wed 30 Sep, 14:30
3. Annika Bach, Eindhoven University of Technology, Holland
   Thu 01 Oct, 14:00
4. Dorin Bucur, Université de Savoie, France
   Fri 02 Oct, 11:00
5. Maicol Caponi, Università dell'Aquila, Italy
   Thu 01 Oct, 09:00
6. Filomena De Filippis, University of Salzburg, Austria
   Wed 30 Sep, 16:00
7. Lucia De Luca, CNR, Italy
   Thu 01 Oct, 11:45
8. Manuel Friedrich, Johannes Kepler Universität, Linz, Austria
   Wed 30 Sep, 15:15
9. Ilias Ftouhi, Université de Nîmes, France
   Fri 02 Oct, 09:45
10. Alessandro Giacomini, Università di Brescia, Italy
    Fri 02 Oct, 09:00
11. Ilaria Lucardesi, Università di Pisa, Italy
    Thu 01 Oct, 16:00

12. Gloria Paoli, Università degli Studi di Napoli "Federico II", Italy
    Thu 01 Oct, 14:45
13. Marianna Porfido, TU Bergakademie Freiberg, Germany
    Thu 01 Oct, 09:45
14. Filippo Riva, Charles University, Prague, Czech Republic
    Thu 01 Oct, 11:00

See next pages for titles and abstracts

Titles and Abstracts

If it twists like an annulus, it looks like an annulus

Vincenzo Amato
Scuola Superiore Meridionale, Italy
v.amato@ssmeridionale.it
Thu 01 Oct, 16:45

Among planar domains with prescribed area and hole area, the annulus best resists torsion. We show that this optimality is stable: a domain with nearly optimal torsional rigidity is close to an annulus, and its holes are almost concentric discs. The key is tracking how asymmetry spreads through the level sets of the torsion function.

Hypercontractivity and asymptotic behaviour of generalised Mehler semigroups

Luciana Angiuli
Università del Salento, Italy
luciana.angiuli@unisalento.it
Wed 30 Sep, 14:30

A natural L^p-setting for studying semigroups associated with elliptic operators with unbounded coefficients is the one related to invariant measures. This is the case, for instance, of the classical Ornstein–Uhlenbeck semigroup $(P(t))_{t\geq 0}$, which enjoys good properties in $L^p(\gamma)$, where γ denotes the standard Gaussian measure that turns out to be the unique invariant measure associated with $(P(t))_{t\geq 0}$. One of the most relevant properties of the Ornstein–Uhlenbeck semigroup, proved by Nelson in 1973, is hypercontractivity: for any $1 < p < q < \infty$ there exists $t_0 = t_0(p,q) > 0$ such that

$$\|P(t)f\|_{L^q(\gamma)} \leq \|f\|_{L^p(\gamma)}$$

for all $f \in L^p(\gamma)$ and $t \geq t_0$. The hypercontractivity of $P(t)$ is strictly connected to the validity of the classical logarithmic Sobolev inequality with respect to the Gaussian measure and to the asymptotic behaviour of the semigroup $P(t)$ as $t \to +\infty$.

The Ornstein–Uhlenbeck semigroup is the prototype for the class of generalised Mehler semigroups arising from linear stochastic differential equations perturbed by Lévy noise. For such semigroups, the hypercontractivity property fails to hold in general. In this talk we investigate the hypercontractivity property of generalised Mehler semigroups $(T(t))_{t\geq 0}$ on the L^p-scale with respect to an invariant measure μ. When the associated invariant measure μ lacks a purely Gaussian structure, jump components may prevent the validity of Nelson's classical L^p-L^q estimates. However, a summability-improving property can be obtained in the setting of mixed-norm spaces $\mathcal{X}_{p,q}(E;\gamma,\pi)$ related to the factorization of the invariant measure $\mu = \gamma * \pi$ into a Gaussian part γ and an infinitely divisible non-Gaussian part π. As in the classical Gaussian case, some modified logarithmic Sobolev inequalities with respect to invariant measures can be inferred, as well as some results on the asymptotic behaviour of $T(t)$ as $t \to +\infty$.

Joint work with Simone Ferrari and Diego Pallara.

Monotonicity arguments for the multi-cell formula in stochastic homogenization

Annika Bach
Eindhoven University of Technology, Holland
a.bach@tue.nl
Thu 01 Oct, 14:00

Stochastic homogenization of integral functionals F_ε via Γ-convergence can be seen as a rigorous mathematical averaging procedure, where highly oscillating random integrands are in the Γ-limit replaced by a homogeneous (suitably averaged) integrand. This limiting integrand is typically determined via a multi-cell formula, a limit of minimization problems where the integral functionals F_ε are rescaled to larger and larger cubes and subjected to suitable boundary conditions. In many situations it is useful to replace those minimization problems by a family of auxiliary problems defined on a restricted but growing class of competitors, and to relate the two minimization problems via a monotonicity argument. In this talk we will present two such monotonicity arguments that we used to obtain fluctuation estimates for the multi-cell formula in the context of integral functionals defined on finite partitions and to obtain a stochastic homogenization result for integral functionals defined on manifold-valued Sobolev functions.

Joint work with Matthias Ruf (Justus-Liebig-Universität Gießen).

On a free boundary problem for the gradient magnitude of the torsion function

Dorin Bucur
Université de Savoie, France
dorin.bucur@univ-savoie.fr
Fri 02 Oct, 11:00

A recent result by Burdzy, Ftouhi, and Mariano establishes a series of geometric properties of optimizers among convex sets for the maximum of the gradient of the torsion function. In this talk, I will discuss a related free boundary problem. We prove a general result concerning the presence of flat parts in the free boundary, together with an estimate of their Hausdorff measure. We will also discuss applications to shape optimization problems.

H-convergence and Γ-convergence in the Riesz fractional setting: the nonlinear case

Maicol Caponi
Università dell'Aquila, Italy
maicol.caponi@univaq.it
Thu 01 Oct, 09:00

This talk concerns the H-convergence of nonlinear nonlocal monotone operators defined through the Riesz fractional gradient and divergence. We show that the H-convergence in this nonlocal framework is equivalent to the H-convergence of the corresponding local one. As a consequence, we obtain a H-compactness result for a suitable class of nonlocal monotone operators. We then study the Γ-convergence of nonlocal energy functionals associated with the subclass of conservative monotone operators, proving that it is equivalent to the Γ-convergence of the corresponding local energies. A key ingredient is a new uniqueness result for the integral representation of both local and nonlocal functionals. As a by-product, we obtain the Γ-compactness of the class of nonlocal energies under consideration. Finally, we show the equivalence between the H-convergence of nonlocal conservative monotone operators and the Γ-convergence of the associated energy functionals.

Joint work with G. C. Brusca (SISSA), A. Carbotti (Università del Salento), A. Maione (Politecnico di Milano) and F. Paronetto (Università di Padova).

Intrinsic Schauder estimates at nearly linear growth

Filomena De Filippis
University of Salzburg, Austria
filomena.defilippis@plus.ac.at
Wed 30 Sep, 16:00

Variational integrals at nearly linear growth appear in the theory of plasticity with logarithmic hardening, that is the borderline configuration between plasticity with power hardening and perfect plasticity. The related (very challenging) regularity theory for minima has been intensively developed over the last 25 years, see e.g. the work of Frehse & Seregin '99, Fuchs & Mingione '00, Bildhauer '03, Beck & Bulíček & Gmeineder '20, Di Marco & Marcellini '20, Gmeineder & Kristensen '22, De Filippis & Mingione '23.

In this talk, we will discuss a nonlinear potential theoretic framework for Schauder estimates for vector valued solutions of a broad class of nonautonomous variational problems at nearly linear growth. Our approach naturally embraces the variable exponent as well as the Double and Multi phase setting, yielding new regularity results in basic models and recovering optimal regularity recently established in specific cases.

Joint work with Cristiana De Filippis (Parma) and Peter Hästö (Helsinki).

Asymptotic expansion of the relative s-fractional perimeter as s tends to 0

Lucia De Luca
CNR, Italy
l.deluca@iac.cnr.it
Thu 01 Oct, 11:45

We prove a first-order Γ-convergence result (as s tends to 0) for the s-fractional perimeter relative to a given bounded set and with prescribed exterior datum. We investigate the robustness of our approach under perturbations of the exterior datum and in presence of volume constraints. We then use the Γ-convergence expansion to study the asymptotic behavior of s-minimal sets as s tends to 0.

Joint work with G. Palatucci, M. Piccinini and M. Ponsiglione.

Quasistatic evolution of cohesive-type fracture

Manuel Friedrich
Johannes Kepler Universität, Linz, Austria
manuel.friedrich@jku.at
Wed 30 Sep, 15:15

In this talk, I present an existence result for a variational model of quasistatic crack growth for a class of cohesive fracture energies with an activation threshold. The evolution is characterized by an irreversibility condition in terms of a maximal opening variable, a global stability, and an energy balance.

Joint work with Vito Crismale.

About the maximization of the gradient of the torsion function

Ilias Ftouhi
Université de Nîmes, France
ilias.ftouhi@unimes.fr
Fri 02 Oct, 09:45

In this talk, we address the problem of maximizing the supremum norm of the gradient of the torsion function over planar convex domains with fixed measure (or perimeter). We prove that an optimal shape exists, its boundary possesses C^1 regularity, and it features at least one straight segment. The proofs leverage probabilistic methods along with a novel formulation of the boundary Harnack principle for the torsion function.

Joint work with Krzysztof Burdzy (University of Washington) and Phanuel Mariano (Union College).

Periodic homogenization in scalar Hencky plasticity

Alessandro Giacomini
Università di Brescia, Italy
alessandro.giacomini@unibs.it
Fri 02 Oct, 09:00

In this talk I will deal with periodic homogenization for a two-phase elasto-plastic material in the framework of Γ-convergence under two-scale convergence. The elasto-plastic character of the homogenized problem is unclear, and will be discussed by resorting to a one field description of the associated energy. A positive answer will be given in one dimension.

Joint work with G. Francfort.

On the shape optimization of the torsional rigidity under thickness constraint among triangles

Ilaria Lucardesi
Università di Pisa, Italy
ilaria.lucardesi@unipi.it
Thu 01 Oct, 16:00

In this talk I will present some recent results about the optimization, both minimization and maximization, of a shape functional which depends on the torsional rigidity and the thickness, in the class of triangles. Besides the detection of optimal triangles, the study allows to obtain new geometric estimates on the torsional rigidity for triangles.

Joint work with Davide Zucco (Università di Torino).

A reverse Faber–Krahn inequality for the Robin Laplacian with negative boundary parameter: small coupling in all dimensions

Gloria Paoli
Università degli Studi di Napoli "Federico II", Italy
gloria.paoli@unina.it
Thu 01 Oct, 14:45

We establish Bareket's conjecture from 1977 for convex domains in all dimensions in the regime of weak boundary coupling. In other words, we consider the Laplace operator, subject to negative boundary conditions, and show that the ball maximises the first eigenvalue among all bounded convex domains of fixed volume, provided that the boundary parameter is sufficiently close to zero. The smallness depends on the volume and dimension only. The proof relies on a comparison with spherical shells with combined Neumann–Robin boundary conditions obtained via the method of parallel coordinates, which we manage to extend to all dimensions, and on a careful analysis of the corresponding radial problem.

Joint work with Nunzia Gavitone and David Krejčiřík.

The Free Lunch Theorem of homogenisation

Marianna Porfido
TU Bergakademie Freiberg, Germany
Marianna.Porfido@math.tu-freiberg.de
Thu 01 Oct, 09:45

In this talk, we revisit the classical elliptic homogenisation theory for local coefficients, i.e., L_∞-multiplication operators (H-convergence as envisioned by Murat and Tartar in the 1970's). Then, we compare it with an abstract operator-theory approach that allows for nonlocal coefficients, i.e., general bounded operators (nonlocal H-convergence introduced in [1]). With these concepts in hand, we show that H-convergence for multiplication type operators always implies nonlocal H-convergence. As a consequence, H-convergence for multiplication operators in divergence form problems will always imply H-type convergence for a different variational problem for free.

Joint work with Andreas Buchinger and Marcus Waurick.

References

[1] M. Waurick, Nonlocal H-Convergence, Calc. Var. Partial Differential Equations 57(6) (2018).
[2] A. Buchinger, M. Porfido, M. Waurick, The Free Lunch Theorem of Homogenisation, preprint (2026), doi: https://arxiv.org/abs/2606.04498

A free boundary approach to quasistatic debonding

Filippo Riva
Charles University, Prague, Czech Republic
filippo.riva@matfyz.cuni.cz
Thu 01 Oct, 11:00

Debonding models describe the evolution of an adhesive elastic membrane peeled away from a planar rigid substrate. When the process is sufficiently slow, inertial and viscous effects can be neglected, leading to a quasistatic setting. In this framework, debonding can be formulated as a variational evolution of sets. In the talk, I will present how to prove existence of so-called energetic solutions for this formulation — based on global minimizers of a suitable functional together with an energy balance — by exploiting an equivalent rewriting of the model in terms of the celebrated one-phase Bernoulli free boundary problem.

Joint work with E. Maggiorelli and E. Tolotti.