

Characterizing the γ -ray sky using differentiable probabilistic programming

Anonymous Authors¹

Abstract

We motivate the use of differentiable probabilistic programming techniques in order to account for the large model-space inherent to astrophysical γ -ray analyses. Targeting the longstanding Galactic Center γ -ray Excess (GCE) puzzle, we construct a differentiable forward model and likelihood that makes liberal use of GPU acceleration and vectorization in order to simultaneously account for a continuum of possible spatial morphologies consistent with the Excess emission in a fully probabilistic manner. Our setup allows for efficient inference over the large model space using variational methods. Beyond application to γ -ray data, a goal of this work is to showcase how differentiable probabilistic can be used as a tool to enable flexible analyses of astrophysical datasets.

1. Introduction

Analysis of γ -ray data towards the Galactic Center involves a subtle interplay between observations and theoretical models. The complexity of cosmic-ray transport processes in the region, combined with the fact that we only have access to a two-dimensional projection of this inherently three-dimensional process, means that we typically cannot make precise predictions about emission from crucial sources. Consequently, scientific conclusions drawn from γ -ray analyses can depend strongly on specific assumptions made about the modeling of the signal and background, as well as manual choices involved in the data analysis pipeline.

The γ -ray Galactic Center Excess (GCE), first identified over a decade ago (Goodenough & Hooper, 2009) using public data from the *Fermi*-LAT telescope (Atwood et al., 2009), is an excess of photons in the Inner Galaxy whose physical origin is currently unknown. Although the GCE has properties broadly compatible with those expected from annihilating dark matter (DM) (Goodenough & Hooper, 2009;

Hooper & Goodenough, 2011; Daylan et al., 2016; Calore et al., 2015; Ajello et al., 2016), competing explanations in terms of a population of unresolved astrophysical point sources (PSs), in particular millisecond pulsars (MSPs), remain viable (Abazajian et al., 2014; Abazajian, 2011). Significant effort has gone into characterizing the GCE and understanding its origin, including through inferring the properties of unresolved PSs in the Inner Galaxy region. A particularly successful method in this direction, Non-Poissonian Template Fitting (NPTF), aims to characterize different populations of unresolved PSs using their photon statistics, via a likelihood-based analysis (Lee et al., 2015; 2016). In the context of the GCE, initial applications of this method showed overwhelming support for the hypothesis of the GCE being sourced by an unresolved population of PSs spatially correlated with the observed GCE morphology (Lee et al., 2016). However, more recent studies showed the method to be susceptible to systematic biases (Leane & Slatyer, 2019; 2020a,b; Chang et al., 2020) partly stemming from degeneracies between different emission components in the region (e.g., those spatially correlated with the GCE, centrally-concentrated stellar populations, and the Galactic disk), with results depending strongly on specific assumptions made about the spatial morphologies of components. *All NPTF analyses so far have assumed inflexible, spatially rigid templates, preventing a rigorous expression and exploration of this degeneracy.*

In this work, we use differentiable probabilistic programming to alleviate this shortcoming of traditional likelihood-based fitting techniques for analyzing γ -ray data. Using GPU-native acceleration and vectorization, we construct a forward model for γ -ray emission in the Galactic Center region that can generate high-resolution spatial templates for various emission components, including those associated with the Galactic Center excess signal, on the fly. This allows for analyses that are robust to specific analysis choices, and opens up the possibility of performing model-independent analyses using the framework. In the broader context of astrophysics, this work aims to motivate the use of differentiable probabilistic programming as a way to enable high-dimensional Bayesian inference for complex descriptions of diverse astrophysical data.

¹Anonymous Institution, Anonymous City, Anonymous Region, Anonymous Country. Correspondence to: Anonymous Author <anon.email@domain.com>.

2. Methodology

We describe our forward model and likelihood, its implementation as a differentiable probabilistic program, as well as the inference strategies used to perform Bayesian posterior inference over the model parameters characterizing various emission mechanisms for Galactic Center γ -ray data.

2.1. The likelihood and forward model

In traditional spatial template fitting, photon counts x^p in different pixels p are modeled as Poisson random variables, with mean given by a linear combination of templates T_i^p , indexed i ; $x^p \sim \text{Pois}(x^p \mid \sum_i A_i T_i^p)$. A_i are the template normalizations and represent the parameters of interest. The full map-levels likelihood factorizes into a product of pixel-wise likelihoods, $p(x \mid \{A_i\}) = \prod_p p(x_p \mid \{A_i\})$.

In the presence of populations of unresolved point sources, the likelihood can be augmented to take into account the probability that a given pixel contains a point source from a given population, and the probability of that point source emitting a certain mean number of contributing photons. Point source populations can be independently characterized by a spatial distribution, specified by a template $T_{\text{PS},i}^p$, and a source-count distribution dN/dS , which jointly describes the distribution of photon counts S from individual PSs, $p(S \mid \theta_{\text{PS}})$, and their mean abundance, $\bar{n}_{\text{PS}}$; $dN/dS = \bar{n}_{\text{PS}} p(S \mid \theta_{\text{PS}})$.

Unfortunately, the inclusion of point source populations renders the likelihood intractable, since computing it would involve marginalizing over positions, counts, and cardinality of point sources from each population. The Non-Poissonian Template Fitting (NPTF) method (Lee et al., 2015; 2016) circumvents this issue by instead computing the pixel-wise likelihood, and approximating the full likelihood in a factorized form; $p(x \mid \bar{n}_{\text{PS}}, \theta_{\text{PS}}) = \prod_p p(x^p \mid \bar{n}_{\text{PS}}, \theta_{\text{PS}})$. See Mishra-Sharma et al. (2017) for further details on the method. We implement a differentiable version of the NPTF likelihood in the Jax framework (Bradbury et al., 2018), making liberal use of vectorization and automatic batching to efficiently compute the likelihood for multiple point source populations.

2.2. Flexible specification of point source populations

Previous analyses based on the NPTF framework assumed rigidly specified spatial templates for the distribution of point source populations; this is because generating new morphologies adds to considerable overhead in the likelihood calculation, rendering a traditional Monte Carlo sampling-based analysis difficult. In particular, given a morphology for the PS template, $\rho(R, z)$ (assumed cylindrically symmetric), computing the template expectation in a

give pixel p involves performing a line-of-sight integral in the pixel direction, $T_{\text{PS}}^p = \int dl \rho(R(l, p), z(l, p))$. In Jax, the density over the line of sight as well across the pixel index can be efficiently vectorized, and the computation is just-in-time (JIT) compiled for efficient execution on a GPU.

This allows us to include parametric variations on different spatial templates that have been previously included in NPTF analyses, including templates for PSs correlated with the Galactic disk and the GCE, directly as part of the inference pipeline. We model a PS population with a generalized Navarro-Frenk-White (NFW) profile, motivated by the DM expectation: $\rho_{\text{gNFW}}(r) \propto (r/r_s)^{-\gamma} (1 + r/r_s)^{-3+\gamma}$ with the inner slope γ a free parameter to be inferred during the analysis. We also model a population of PSs spatially correlated with the Galactic disk, modeled using a doubly exponential profile, $\rho_{\text{disk}}(R, z) \propto (R/R_\odot)^B \exp[-C(R - R_\odot)/R_\odot] \exp(-|z|/z_s)$ with the parameter B , C , and the scale height z_s characterizing the disk morphology and to be inferred during the analysis.

The possibility of a population of PSs correlated with the stellar density in the Galactic bulge has been extensively tested in the literature (Macias et al., 2018; 2019; Ploeg et al., 2020), and either substantiated or disfavored depending on the analysis. We allow for this possibility in our analysis by including multiple publicly-available spatial templates following stellar bulge populations (Macias et al., 2018; 2019). For both the PS as well as smooth (Poissonian) components describing the GCE, we model the spatial distribution as a hybrid template allowing for varying fractions of DM-like and bulge-like morphologies, $T_{\text{GCE}}^p = (1 - f_{\text{bulge}})T_{\text{DM}}^p + f_{\text{bulge}}T_{\text{bulge}}^p$. The bulge template itself is modeled as a linear combination of publicly available bulge morphologies, $T_{\text{bulge}}^p = \sum_i \alpha_i T_{\text{bulge},i}^p$ with α_i modeled with a symmetric Dirichlet prior, enforcing $\sum_i \alpha_i = 1$. The templates are appropriately normalized such that an un-scaled template would produce a single count-per-pixel, allowing for a physically-meaningful interpretation of the bulge fraction f_{bulge} parameter as the relative contribution to the GCE PS or smooth component from the composite bulge component.

2.3. Parameter posterior inference

Since the entire pipeline from template generation to likelihood evaluation is end-to-end differentiable, we can evaluate gradients of the log-likelihood function with respect to the parameters of interest θ characterizing the smooth as well as PS components, $\nabla_\theta \log \mathcal{L}(\theta)$, with minimal overhead. This opens up the possibility of using highly efficient gradient-based posterior inference techniques like Stochastic Variational Inference (SVI) (Hoffman et al., 2012) and Hamiltonian Monte Carlo (HMC) (Hoffman & Gelman,




Figure 1. Summaries of the inferred posteriors for disk-correlated PS, GCE-correlated PS, and GCE-correlated smooth emission morphologies. The top row shows the median inferred pixel-wise morphologies, which the bottom row shows the relative middle-68% deviation of the posterior from the median. For comparison, contours for the best-fitting bulge description are shown for the GCE PS and smooth components.

2011). Unlike traditional sampling techniques employed in the context of *Fermi* GCE analyses so far, these methods can easily scale up to high-dimensional parameter spaces. In particular, variational methods turn posterior inference into an optimization problem by fitting for a flexible functional ansatz on the posterior density and can easily deal with hundreds of parameters. This is in contrast to current public implementations of NPTF, which cannot efficiently include real-time generated templates at inference and are restricted to a modest number of parameters with traditional Monte Carlo sampling techniques. The constructed likelihood is wrapped using the probabilistic programming framework NumPyro (Phan et al., 2019), which allows for flexible specification of complex forward models as well as performing inference on them using gradient-based techniques.

In stochastic variational inference (SVI) an approximation for the posterior distribution, parameterized by φ and denoted $q_\varphi(\theta)$, is obtained by minimizing the reverse KL-divergence between the true and variational posterior distributions, $D_{\text{KL}}(q_\varphi \parallel p(\theta \mid x))$. This is done by maximizing the model log-evidence $\log p(x)$ or, in practice using the tractable evidence lower bound (ELBO), $\text{ELBO} \equiv \mathbb{E}_{\theta \sim q_\varphi(\theta)} [\log p(x, \theta) - \log q_\varphi(\theta)]$, as the optimization objective since $\log p(x) - \text{ELBO} = D_{\text{KL}}(q_\varphi \parallel p(\theta \mid x))$. The expectation in the ELBO is taken through Monte Carlo sampling from the variational distribution at each optimization step.

Since we’d like to be able to model arbitrarily flexible non-Gaussian posterior distributions, the variational family q_φ is modeled using an inverse autoregressive normalizing flow (IAF). The variational parameters—the parameters of the normalizing flow transformation—are optimized using Optax (Hessel et al., 2020) with the AdamW optimizer (Kingma & Ba, 2014; Loshchilov & Hutter, 2017) over 10,000 steps with learning rate 10^{-3} . Optimization typically takes ~ 10 minutes on a single Nvidia A100 GPU.

3. Experiments and results

We apply our pipeline to 573 weeks of *Fermi*-LAT data in the 2–20 GeV energy range. Smooth templates corresponding to isotropically-uniform emission, emission from (revolved) point sources in the 3FGL catalog, and the *Fermi* bubbles are included. Three different diffuse models – Models ‘A’ and ‘F’ from Calore et al. (2015) and Model ‘O’ from Macias et al. (2019) – are used to model the Galactic diffuse foreground emission, with the gas-correlated emission and Galactic charge density-correlated emission separately allowed be linear combinations of the corresponding components of the three templates.

Summaries of the inferred posteriors for disk-correlated PS, GCE-correlated PS, and GCE-correlated smooth emission morphologies are shown in Fig. 1. The top row shows the median inferred pixel-wise morphologies, which the bottom row shows the relative middle-68% deviation of the




Figure 2. Posteriors on select parameters characterizing the GCE and disk PS morphologies obtained by analyzing *Fermi*-LAT data in the Inner Galaxy. The bulge fraction f_{bulge} , distributions on the NFW inner slopes γ for the PS and smooth NFW profiles, as well as select parameters characterizing the disk morphology are shown.

posterior from the median (heuristically, the ‘1- σ ’ range of the posterior distribution). Contours for the best-fitting bulge-correlated component are shown using dotted red lines. It can be seen that, for both PS-like as well as smooth emission, the inferred posterior median is fairly spherically symmetric when contrasted to the bulge profile expectation; this reflects a general preference for DM-like morphologies in our analyses. We emphasize that we obtain a full posterior distribution over preferred signal morphologies, departing significantly from the rigid parametric descriptions used in previous works.

Figure 2 shows posteriors on select parameters characterizing the GCE and disk PS morphologies. The bulge fraction $f_{\text{bulge}} \sim 0.3$ quantitatively substantiates a not-insignificant preference for a DM-like morphology in the GCE emission. Posteriors on the NFW inner slopes γ for the PS and smooth NFW profiles, as well as select parameters characterizing the disk morphology are shown.

4. Discussion and outlook

We have introduced a scalable and flexible pipeline for disentangling the contribution of various emission components to the observed γ -ray sky using differentiable probabilis-

tic programming. The pipeline makes extensive use of the seamless vectorization, automatic differentiation, just-in-time compilation, and GPU acceleration enabled by the Jax framework as well as the flexible model specification and inference capabilities of the NumPyro framework. In a preliminary analysis, we applied our pipeline to Inner Galaxy *Fermi*-LAT γ -ray data with goal of characterizing the long-standing Galactic Center Excess emission while allowing for a richer description of possible signal morphologies and Galactic foreground model descriptions.

We outline a few possible extensions to the pipeline. While we have used parametric and physically-motivated morphologies for the spatial distribution of various point source populations, using our pipeline we will be able to characterize Galactic Center in a fully signal model-independent manner by scanning over a space of spatial functions on the sky map e.g., using Gaussian processes (GPs) to define the function prior. A similar GP-based technique was implemented in Mishra-Sharma & Cranmer (2020) to augment the morphology of the diffuse foreground template. Finally, since our pipeline is implemented using differential programming, it can admit arbitrarily flexible descriptions of the Galactic foreground emission, e.g. as a latent-variable generative model that spans a distribution of possible foreground models. We leave a study of these extensions to future work.

References

- Abazajian, K. N. The Consistency of *Fermi*-LAT Observations of the Galactic Center with a Millisecond Pulsar Population in the Central Stellar Cluster. *JCAP*, 03:010, 2011. doi: 10.1088/1475-7516/2011/03/010.
- Abazajian, K. N., Canac, N., Horiuchi, S., and Kaplinghat, M. Astrophysical and Dark Matter Interpretations of Extended Gamma-Ray Emission from the Galactic Center. *Phys. Rev. D*, 90(2):023526, 2014. doi: 10.1103/PhysRevD.90.023526.
- Ajello, M. et al. *Fermi*-LAT Observations of High-Energy γ -Ray Emission Toward the Galactic Center. *Astrophys. J.*, 819(1):44, 2016. doi: 10.3847/0004-637X/819/1/44.
- Atwood, W. B. et al. The Large Area Telescope on the *Fermi* Gamma-ray Space Telescope Mission. *Astrophys. J.*, 697: 1071–1102, 2009. doi: 10.1088/0004-637X/697/2/1071.
- Bradbury, J., Frostig, R., Hawkins, P., Johnson, M. J., Leary, C., Maclaurin, D., Necula, G., Paszke, A., VanderPlas, J., Wanderman-Milne, S., and Zhang, Q. JAX: composable transformations of Python+NumPy programs, 2018. URL <http://github.com/google/jax>.
- Calore, F., Cholis, I., and Weniger, C. Background Model

- Systematics for the Fermi GeV Excess. *JCAP*, 03:038, 2015. doi: 10.1088/1475-7516/2015/03/038.
- Chang, L. J., Mishra-Sharma, S., Lisanti, M., Buschmann, M., Rodd, N. L., and Safdi, B. R. Characterizing the nature of the unresolved point sources in the Galactic Center: An assessment of systematic uncertainties. *Phys. Rev. D*, 101(2):023014, 2020. doi: 10.1103/PhysRevD.101.023014.
- Daylan, T., Finkbeiner, D. P., Hooper, D., Linden, T., Portillo, S. K. N., Rodd, N. L., and Slatyer, T. R. The characterization of the gamma-ray signal from the central Milky Way: A case for annihilating dark matter. *Phys. Dark Univ.*, 12:1–23, 2016. doi: 10.1016/j.dark.2015.12.005.
- Goodenough, L. and Hooper, D. Possible Evidence For Dark Matter Annihilation In The Inner Milky Way From The Fermi Gamma Ray Space Telescope. 10 2009.
- Hessel, M., Budden, D., Viola, F., Rosca, M., Sezener, E., and Hennigan, T. Optax: composable gradient transformation and optimisation, in *jax!*, 2020. URL <http://github.com/deepmind/optax>.
- Hoffman, M., Blei, D. M., Wang, C., and Paisley, J. Stochastic Variational Inference. *arXiv e-prints*, art. arXiv:1206.7051, June 2012. doi: 10.48550/arXiv.1206.7051.
- Hoffman, M. D. and Gelman, A. The No-U-Turn Sampler: Adaptively Setting Path Lengths in Hamiltonian Monte Carlo. *arXiv e-prints*, art. arXiv:1111.4246, November 2011. doi: 10.48550/arXiv.1111.4246.
- Hooper, D. and Goodenough, L. Dark Matter Annihilation in The Galactic Center As Seen by the Fermi Gamma Ray Space Telescope. *Phys. Lett. B*, 697:412–428, 2011. doi: 10.1016/j.physletb.2011.02.029.
- Kingma, D. P. and Ba, J. Adam: A method for stochastic optimization. *arXiv preprint arXiv:1412.6980*, 2014.
- Leane, R. K. and Slatyer, T. R. Revival of the Dark Matter Hypothesis for the Galactic Center Gamma-Ray Excess. *Phys. Rev. Lett.*, 123(24):241101, 2019. doi: 10.1103/PhysRevLett.123.241101.
- Leane, R. K. and Slatyer, T. R. Spurious Point Source Signals in the Galactic Center Excess. *Phys. Rev. Lett.*, 125(12):121105, 2020a. doi: 10.1103/PhysRevLett.125.121105.
- Leane, R. K. and Slatyer, T. R. The enigmatic Galactic Center excess: Spurious point sources and signal mis-modeling. *Phys. Rev. D*, 102(6):063019, 2020b. doi: 10.1103/PhysRevD.102.063019.
- Lee, S. K., Lisanti, M., and Safdi, B. R. Distinguishing Dark Matter from Unresolved Point Sources in the Inner Galaxy with Photon Statistics. *JCAP*, 05:056, 2015. doi: 10.1088/1475-7516/2015/05/056.
- Lee, S. K., Lisanti, M., Safdi, B. R., Slatyer, T. R., and Xue, W. Evidence for Unresolved γ -Ray Point Sources in the Inner Galaxy. *Phys. Rev. Lett.*, 116(5):051103, 2016. doi: 10.1103/PhysRevLett.116.051103.
- Loshchilov, I. and Hutter, F. Decoupled weight decay regularization. *arXiv preprint arXiv:1711.05101*, 2017.
- Macias, O., Gordon, C., Crocker, R. M., Coleman, B., Patterson, D., Horiuchi, S., and Pohl, M. Galactic bulge preferred over dark matter for the Galactic centre gamma-ray excess. *Nature Astron.*, 2(5):387–392, 2018. doi: 10.1038/s41550-018-0414-3.
- Macias, O., Horiuchi, S., Kaplinghat, M., Gordon, C., Crocker, R. M., and Nataf, D. M. Strong Evidence that the Galactic Bulge is Shining in Gamma Rays. *JCAP*, 09:042, 2019. doi: 10.1088/1475-7516/2019/09/042.
- Mishra-Sharma, S. and Cranmer, K. Semi-parametric γ -ray modeling with Gaussian processes and variational inference. 10 2020.
- Mishra-Sharma, S., Rodd, N. L., and Safdi, B. R. NPTFit: A code package for Non-Poissonian Template Fitting. *Astron. J.*, 153(6):253, 2017. doi: 10.3847/1538-3881/aa6d5f.
- Phan, D., Pradhan, N., and Jankowiak, M. Composable Effects for Flexible and Accelerated Probabilistic Programming in NumPyro. *arXiv e-prints*, art. arXiv:1912.11554, December 2019. doi: 10.48550/arXiv.1912.11554.
- Ploeg, H., Gordon, C., Crocker, R., and Macias, O. Comparing the Galactic Bulge and Galactic Disk Millisecond Pulsars. *JCAP*, 12:035, 2020. doi: 10.1088/1475-7516/2020/12/035.