



Siddhant Mishra-Sharma (MIT/AI FI) | AI FI Summer School



170

18

Latent-variable modeling

Maximum-likelihood training?

$$\vartheta^* = \arg \max_{\vartheta} p_{\vartheta}(x)$$

$$= \arg \max_{\vartheta} \int p_{\vartheta}(x \mid z) p(z) \, \mathrm{d}z$$

$$= \arg \max_{\vartheta} \left\langle p_{\vartheta}(x \mid z) \right\rangle_{p(z)}$$



*Difficult to build a good estimator!*

*Dealing with  
high-dim data*



*Curse of  
dimensionality*



The intractability of  $p(x)$  is closely related to the intractability of the *posterior*  $p(z \mid x)$

$$p(z \mid x) = \frac{p(x, z)}{p(x)} \quad (\text{Bayes' theorem})$$

# Latent-variable modeling

## Maximum-likelihood training?

$$\begin{aligned}\vartheta^* &= \arg \max_{\vartheta} p_{\vartheta}(x) \\ &= \arg \max_{\vartheta} \int p_{\vartheta}(x \mid z) p(z) \, dz \\ &= \arg \max_{\vartheta} \left\langle p_{\vartheta}(x \mid z) \right\rangle_{p(z)}\end{aligned}$$

*Difficult to build a good estimator!*

The intractability of  $p(x)$  is closely related to the intractability of the *posterior*  $p(z \mid x)$

$$p(z \mid x) = \frac{p(x, z)}{p(x)} \quad (\text{Bayes' theorem})$$

# Kullback–Leibler (KL) divergence

A measure of similarity between two probability distributions

$$D_{\text{KL}}(P\|Q) = \int_{-\infty}^{\infty} dx p(x) \log \left( \frac{p(x)}{q(x)} \right)$$
$$= \left\langle \log \frac{p(x)}{q(x)} \right\rangle_{x \sim p(x)}$$

