

70. Climbing Stairs:

You are climbing a staircase. It takes n steps to reach the top.

Each time you can either climb 1 or 2 steps. In how many distinct ways can you climb to the top?

Example 1:

Input: $n = 2$

Output: 2

Explanation: There are two ways to climb to the top.

1. 1 step + 1 step

2. 2 steps

Example 2:

Input: $n = 3$

Output: 3

Explanation: There are three ways to climb to the top.

1. 1 step + 1 step + 1 step

2. 1 step + 2 steps

3. 2 steps + 1 step

Constraints:

$1 \leq n \leq 45$

53. Maximum Subarray

Given an integer array `nums`, find the subarray with the largest sum, and return its sum.

Example 1:

Input: `nums = [-2,1,-3,4,-1,2,1,-5,4]`

Output: 6

Explanation: The subarray `[4,-1,2,1]` has the largest sum 6.

Example 2:

Input: `nums = [1]`

Output: 1

Explanation: The subarray `[1]` has the largest sum 1.

Example 3:

Input: `nums = [5,4,-1,7,8]`

Output: 23

Explanation: The subarray `[5,4,-1,7,8]` has the largest sum 23.

300. Longest Increasing Subsequence

Given an integer array `nums`, return the length of the longest strictly increasing subsequence.

Example 1:

Input: `nums = [10,9,2,5,3,7,101,18]`

Output: 4

Explanation: The longest increasing subsequence is `[2,3,7,101]`, therefore the length is 4.

Example 2:

Input: `nums = [0,1,0,3,2,3]`

Output: 4

Example 3:

Input: `nums = [7,7,7,7,7,7,7]`

Output: 1

416. Partition Equal Subset Sum

Given an integer array `nums`, return `true` if you can partition the array into two subsets such that the sum of the elements in both subsets is equal or `false` otherwise.

Example 1:

Input: `nums = [1,5,11,5]`

Output: `true`

Explanation: The array can be partitioned as `[1, 5, 5]` and `[11]`.

Example 2:

Input: `nums = [1,2,3,5]`

Output: `false`

Explanation: The array cannot be partitioned into equal sum subsets.