

Submission Assignment #1 Solution

Instructor: Beilun Wang

Name: No Name, ID: 12345678

Problem 1: Basic Vector Operations

(points)

$$(1) \|\mathbf{a}\|_2 = \sqrt{1^2 + 2^2 + 3^2} = \sqrt{14}$$

$$\|\mathbf{b}\|_2 = \sqrt{8^2 + 1^2 + 2^2} = \sqrt{69}$$

$$(2) \|\mathbf{a} - \mathbf{b}\|_2 = \sqrt{9^2 + 1^2 + 1^2} = \sqrt{83}$$

$$(3) \because \mathbf{a}^T \mathbf{b} = -8 + 2 + 6 = 0$$

$$\therefore \mathbf{a} \text{ and } \mathbf{b} \text{ are orthogonal}$$

Problem 2: Basic Matrix Operations

(points)

(1) When I calculate the inverse matrix of \mathbf{A} , I use the row transformation. Then the result is as following.

$$\mathbf{A}^{-1} = \begin{pmatrix} -\frac{1}{8} & -\frac{3}{8} & \frac{3}{8} \\ \frac{3}{4} & -\frac{5}{4} & \frac{3}{4} \\ \frac{3}{4} & -\frac{3}{4} & \frac{1}{4} \end{pmatrix}$$

$$|\mathbf{A}| = 16$$

$$(2) \because |\mathbf{A}| \neq 0, \therefore r(\mathbf{A}) = 3$$

$$(3) \text{tr}(\mathbf{A}) = 1 + (-5) + 4 = 0$$

$$(4) \mathbf{A} + \mathbf{A}^T = \begin{pmatrix} 1 & -3 & 3 \\ 3 & -5 & 3 \\ 6 & -6 & 4 \end{pmatrix} + \begin{pmatrix} 1 & 3 & 6 \\ -3 & -5 & -6 \\ 6 & -6 & 4 \end{pmatrix} = \begin{pmatrix} 2 & 0 & 9 \\ 0 & -10 & -3 \\ 9 & -3 & 8 \end{pmatrix}$$

$$(5) \text{Matrix } \mathbf{A} \text{ is not orthogonal because } (1, -3, 3)^T \cdot (3, -5, 3) \neq 0$$

$$(6) |\lambda \mathbf{E} - \mathbf{A}| = \begin{vmatrix} \lambda - 1 & 3 & -3 \\ -3 & \lambda + 5 & -3 \\ -6 & 6 & \lambda - 4 \end{vmatrix}$$

After simplification, we can find $(\lambda - 4)(\lambda + 2)^2 = 0$. So we can find the three eigenvalues is: $\lambda_1 = 4$, $\lambda_2 = \lambda_3 = -2$.

When $\lambda_1 = 4$, there is one eigen vectors: $\mathbf{v}_1 = \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \\ 1 \end{pmatrix}$. When $\lambda_2 = \lambda_3 = -2$, we can get two eigen vectors:

$$\mathbf{v}_2 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \mathbf{v}_3 = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$$

$$(7) \mathbf{P} = (\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3) = \begin{pmatrix} \frac{1}{2} & 1 & -1 \\ \frac{1}{2} & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}, \text{ then we can calculate } \mathbf{P}^{-1} = \begin{pmatrix} 1 & -1 & 1 \\ -\frac{1}{2} & \frac{3}{2} & -\frac{1}{2} \\ -1 & 1 & 0 \end{pmatrix}, \Lambda = \text{diag}(4, -2, -2)$$

so $\mathbf{A} = \mathbf{P}^{-1} \Lambda \mathbf{P}$

$$(8) \|\mathbf{A}\|_F = \sqrt{1 + 9 + 9 + 9 + 25 + 9 + 36 + 36 + 16} = 5\sqrt{6}$$

$$\|\mathbf{A}\|_{l_{21}} = \sqrt{1 + 9 + 9} + \sqrt{9 + 25 + 9} + \sqrt{36 + 36 + 16} = \sqrt{19} + \sqrt{43} + 2\sqrt{22}$$

(9) $\|\mathbf{A}\|_* =$
 $\|\mathbf{A}\|_2 =$

Problem 3: Linear Equations

(points)

(1) Through the equations, we can get the augmented matrix: $\begin{pmatrix} 2 & 2 & 3 & 1 \\ 1 & -1 & 0 & -1 \\ -1 & 2 & 1 & 2 \end{pmatrix}$

Through the row transformation of the matrix, we can find: $\begin{pmatrix} 1 & 0 & 0 & -1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}$, so we can get the solution

of the equations is: $\begin{cases} x_1 = -1 \\ x_2 = 0 \\ x_3 = 1 \end{cases}$

(2) $\begin{pmatrix} 2 & 2 & 3 \\ 1 & -1 & 0 \\ -1 & 2 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$

(3) $r(A) = 3$

(4) $\begin{pmatrix} 2 & 2 & 3 & 1 & 0 & 0 \\ 1 & -1 & 0 & 0 & 1 & 0 \\ -1 & 2 & 1 & 0 & 0 & 1 \end{pmatrix}$

Then we can through row transformation to find out the inverse of A:

$$\mathbf{A}^{-1} = \begin{pmatrix} 1 & -4 & -3 \\ 1 & -5 & -3 \\ -1 & 6 & 4 \end{pmatrix}$$

$$|\mathbf{A}| = \begin{vmatrix} 2 & 2 & 3 \\ 1 & -1 & 0 \\ -1 & 2 & 1 \end{vmatrix} = \begin{vmatrix} 1 & -1 & 0 \\ 0 & 1 & 1 \\ 0 & 4 & 3 \end{vmatrix} = -1$$

(5) $\mathbf{x} = \mathbf{A}^{-1}\mathbf{b} = \begin{pmatrix} 1 & -4 & -3 \\ 1 & -5 & -3 \\ -1 & 6 & 4 \end{pmatrix} \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$

(6) $\langle \mathbf{x}, \mathbf{b} \rangle = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}^T \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} = -1 + 0 + 2 = 1$

$$\mathbf{x} \otimes \mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 0 & 1 \\ 1 & -1 & 2 \end{vmatrix} = (1, 3, 1)$$

(7) $\|\mathbf{b}\|_{l_1} = 2 \quad \|\mathbf{b}\|_{l_2} = \sqrt{2} \quad \|\mathbf{b}\|_{l_\infty} = 2$

(8) $\mathbf{y}^T \mathbf{A} \mathbf{y} = (y_1, y_2, y_3) \begin{pmatrix} 2 & 2 & 3 \\ 1 & -1 & 0 \\ -1 & 2 & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = 2y_1^2 - y_2^2 + y_3^2 + 3y_1y_2 + 2y_1y_3 + 2y_2y_3$

$$\nabla_{\mathbf{y}} \mathbf{y}^T \mathbf{A} \mathbf{y} = (4y_1 + 3y_2 + 2y_3, 3y_1 - 2y_2 + 2y_3, 2y_1 + 2y_2 + 2y_3)$$

(9) $\begin{pmatrix} 2 & 2 & 3 \\ 1 & -1 & 0 \\ -1 & 2 & 1 \\ -1 & 2 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$

(10) $\because r(\mathbf{A}_1) \leq 3, \text{ and } r(\mathbf{A}_1) \geq r(\mathbf{A}), \therefore r(\mathbf{A}_1) = 3$

$$(11) (\mathbf{A}_1, \mathbf{b}) = \begin{pmatrix} 2 & 2 & 3 & 1 \\ 1 & -1 & 0 & -1 \\ -1 & 2 & 1 & 2 \\ -1 & 2 & 1 & 2 \end{pmatrix} \Rightarrow \begin{pmatrix} 2 & 2 & 3 & 1 \\ 1 & -1 & 0 & -1 \\ -1 & 2 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{pmatrix} \Rightarrow r(\mathbf{A}_1, \mathbf{b}) = r(\mathbf{A}, \mathbf{b}), \because \mathbf{Ax} = \mathbf{b} \text{ have the only one solution, } r(\mathbf{A}) = 3. \therefore r(\mathbf{A}_1, \mathbf{b}) = 3 = r(\mathbf{A}_1)$$