



ISTA 421 + INFO 521

Machine Learning

Probability Review

Clay Morrison

claytonm@email.arizona.edu

Harvill 437A, 621-6609

References for probability

Recommend:

(lvl 1) Doing Bayesian Data Analysis (**DBDA**)

Ch 2, 4, 5

(lvl 2) First Course in Machine Learning (**FCML**)

Ch 2.2 (foundations),

Ch 2.3 (Discrete),

Ch 2.4-2.5 (Continuous)

Ch 2.6-2.7 (Expectation and Maximum Likelihood)

Ch 3 (Bayesian)

(lvl 3) Pattern Recognition and Machine Learning (**PRML**)

Ch 1.2 (foundations),

Ch 2.1-2.2 (Discrete),

Ch 2.3 (Continuous)

Google (and WikiPedia) for unfamiliar terms and alternative explanations.

Wisdom from tea dipper handle

**Good
Earth®**

In mathematics you
don't understand
things. You just get
used to them.

Johann von Neumann
(1903 - 1957)

Probability semantics

Two broad interpretations of probability
(variants exist for both)

- 1) Representation of expected frequency (“frequentist”)
- 2) Degree of belief (“Bayesian”)

There is a 20% chance of rain tomorrow.



Basic terminology and rules

Sample Space of *outcomes* (often denoted by Ω)

$\{H, T\}$

$\{1, 2, 3, 4, 5, 6\}$

An outcome is just ONE element of the sample space

A “generic” outcome is often denoted by ω

and we can say things like, e.g., “for each $\omega \in \Omega \dots$ ”



Basic terminology and rules

Sample Space of *outcomes* (often denoted by Ω)

$\{H, T\}$

$\{1, 2, 3, 4, 5, 6\}$

An outcome is just ONE element of the sample space

A “generic” outcome is often denoted by ω

and we can say things like, e.g., “for each $\omega \in \Omega \dots$ ”

Event (subset of Ω) ...does or does not contain (is true or false for) a particular outcome

odd $\{1, 3, 5\}$, even $\{2, 4, 6\}$, prime $\{2, 3, 5\}$



Basic terminology and rules

Sample Space of *outcomes* (often denoted by Ω)

$\{H, T\}$

$\{1, 2, 3, 4, 5, 6\}$

An outcome is just ONE element of the sample space

A “generic” outcome is often denoted by ω

and we can say things like, e.g., “for each $\omega \in \Omega$...”

Event (subset of Ω) ...does or does not contain (is true or false for) a particular outcome

odd $\{1, 3, 5\}$, even $\{2, 4, 6\}$, prime $\{2, 3, 5\}$

Semantics of Set Operations

Equivalence between “set” and “proposition” representations.

1. Set E : outcomes s.t. proposition E is true.
2. Union, $E \cup F$: logical OR between propositions E and F .
3. Intersection, $E \cap F$: logical AND
4. Complement, E^C : logical negation



Basic terminology and rules

Sample Space of *outcomes* (often denoted by Ω)

$\{H, T\}$

$\{1, 2, 3, 4, 5, 6\}$

An outcome is just ONE element of the sample space

A “generic” outcome is often denoted by ω

and we can say things like, e.g., “for each $\omega \in \Omega$...”

Event (subset of Ω) ...does or does not contain (is true or false for) a particular outcome

odd $\{1, 3, 5\}$, even $\{2, 4, 6\}$, prime $\{2, 3, 5\}$

Denote the **collection of measurable events**
(ones we want to assign probabilities to) by S .

S must include \emptyset and Ω

These special events represent the cases where
“nothing” among all the choices happens (impossible),
and “something” happens (certain).

Reason for being technical: It is important to be tuned
into **what** a particular probability is **about** (precisely!).



Basic terminology and rules

Sample Space of *outcomes* (often denoted by Ω)

$\{H, T\}$

$\{1, 2, 3, 4, 5, 6\}$

An outcome is just ONE element of the sample space

A “generic” outcome is often denoted by ω

and we can say things like, e.g., “for each $\omega \in \Omega$...”

Event (subset of Ω) ...does or does not contain (is true or false for) a particular outcome

odd $\{1, 3, 5\}$, even $\{2, 4, 6\}$, prime $\{2, 3, 5\}$

Denote the **collection of measurable events**
(ones we want to assign probabilities to) by S .

S must include \emptyset and Ω

S is *closed* under set operations

...aka: σ -algebra

$\alpha, \beta \in S \Rightarrow \alpha \cup \beta \in S, \alpha \cap \beta \in S, \alpha^c = \Omega - \alpha \in S$, etc.

Translation: We need to be able to deal with concepts such as “either A or B” happens, or “both A and B” happen.

E.g., I’ll accept either an even or prime number

E.g., If I roll a 3, it is both odd and prime



Basic terminology and rules

Probability Space

A **probability space** is a sample space augmented with a function, P , that assigns a **probability** to each event, $E \subset S$.

Kolmogorov Axioms

1. $0 \leq P(E) \leq 1$ for all $E \subset S$.
2. $P(\Omega) = 1$.
3. If $E \cap F = \emptyset$ then $P(E \cup F) = P(E) + P(F)$.

Important Consequences

1. $P(\emptyset) = 0$.
2. $P(E^C) = 1 - P(E)$
3. In general, $P(E \cup F) = P(E) + P(F) - P(E \cap F)$.



Random Variables

Random variables

Defined by **functions** mapping **outcomes** (ω) to **values**

A random variable is a way of reporting an attribute of an outcome

Typically r.v. are denoted by uppercase letters (e.g., X)

Generic values are corresponding lower case letters (e.g., x)

Shorthand: $P(x) = P(X=x)$

Value “type” is arbitrary (typically categorical or real)

Example (from K&F)

Outcomes are student grades (A,B,C)

Random variable $G = f_{\text{GRADE}}(\text{student})$

$$P('A') = P(G = 'A') = P(\{ \omega \in \Omega : f_{\text{GRADE}}(\omega) = 'A' \})$$

We sometimes use sets, but usually R.Vs.: $P(\overbrace{A \cap B \cap C}^{\text{Sets}}) \equiv P(\overbrace{A, B, C}^{\text{R. Vs.}})$



Random Variables

Random Variable

- ▶ Formally, a **random variable** is a function, X that assigns a number to each outcome in S (e.g., dead $\rightarrow 0$, alive $\rightarrow 1$).
- ▶ Key consequence: a random variable divides the sample space into **equivalence classes**: sets of outcomes that share some property (differ only in ways irrelevant to X)

Example

- ▶ Let S = all sequences of 3 coin tosses.
- ▶ We can define a r.v. X that counts number of heads.
- ▶ Then HHT and HTH are equivalent in the eyes of X :

$$X(HHT) = X(HTH) = 2$$



Random Variables

Distribution of a Random Variable

- ▶ The expression $P(X = x)$ refers to the probability of the event $E = \{\omega \in S : X(\omega) = x\}$.
- ▶ Sometimes we can obtain it by breaking it down into simpler, mutually exclusive events and adding their probabilities (Kolmogorov axiom 3)

Example

- ▶ $S =$ all sequences of 3 coin tosses.
- ▶ $X(\omega) =$ # of heads in ω .

$$\begin{aligned}\{X = 2\} &= \{HHT\} \cup \{HTH\} \cup \{THH\} \\ P(X = 2) &= P(HHT) + P(HTH) + P(THH) \\ &= \frac{1}{8} + \frac{1}{8} + \frac{1}{8}\end{aligned}$$



Random Variables

Distribution of a Random Variable

- ▶ Similarly, $P(X < x)$ is the probability of the event $E = \{\omega \in S : X(\omega) < x\}$.
- ▶ Can sometimes obtain it the same way as we did above.

Example

- ▶ $S =$ all sequences of 3 coin tosses.
- ▶ $X(\omega) =$ # of heads in ω .

$$\begin{aligned}\{X < 2\} &= \{TTT\} \cup \{TTH\} \cup \{THT\} \cup \{HTT\} \\ P(X < 2) &= P(TTT) + P(TTH) + P(THT) + P(HTT) \\ &= \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8}\end{aligned}$$



Random Variables

Distribution of a Random Variable

Example, continued

- Notice that in this example we could also have written

$$\begin{aligned}\{X < 2\} &= \{X = 0\} \cup \{X = 1\} \\ P(X < 2) &= P(X = 0) + P(X = 1)\end{aligned}$$

which is useful if we have already calculated $P(X = x)$ for each value of x .

- This always works if X is always an integer.

Joint Probability

Joint Probability

- ▶ We have already seen the concept of *intersecting events*: $A \cap B$ is the event that occurs when *both* A and B are true *at the same time*.
- ▶ $P(A \cap B)$ is called the **joint probability** of A and B .
- ▶ If A is $\{X = x\}$ and B is $\{Y = y\}$, then $A \cap B$ means $X = x$ and $Y = y$ *at the same time*.
- ▶ If X and Y are discrete, $P(X = x, Y = y)$, for different combinations of x and y , characterize the **joint distribution** of X and Y .

We write $P(x, y)$ for $P(\{w \in \Omega : X(w) = x \text{ and } Y(w) = y\})$

Alternatively, $P((X = x) \cap (Y = y))$


Note that the comma in the usual form, $P(x, y)$, is read as "and".

Here events are being defined by assignments of random variables

Joint Probability

$$P(A) \left\{ \begin{array}{c} a_1 \\ a_2 \\ \\ a_n \end{array} \right. \begin{array}{c} P(a_1) \\ \\ \\ \end{array}$$

Joint Probability



Joint Probability


Joint Probability

		$P(B)$				
		b_1	b_2	b_m		
			$P(b_2)$			
$P(A)$	a_1	$P(a_1)$	$P(a_1, b_2)$			
	a_2					
	a_n					

$P(A, B)$

Joint Probability

Joint Probability



Marginalization: $P(A) = \sum_{b \in B} P(A, B)$ *

Formulas that you should be comfortable with are marked by * .


Conditional Probability

“probability in context”

Conditional probability (definition)

$$P(A|B) \equiv \frac{P(A \cap B)}{P(B)}$$

*



Conditional Probability

“probability in context”

Conditional probability (definition)

$$P(A|B) \equiv \frac{P(A \cap B)}{P(B)} \quad *$$

Example: what is the probability that you roll 2 (on a six sided die), given that you know you have rolled a prime number?

		$P(B)$			
		b_1	b_2	b_m	
			$P(b_2)$		
$P(A)$	a_1	$P(a_1)$	$P(a_1 b_2)$		
	a_2				
	a_n				

$P(A, B)$

Product Rule

“probability in context”

Conditional probability (definition)

$$P(A|B) \equiv \frac{P(A \cap B)}{P(B)} \quad *$$

Applying a bit of algebra,

$$P(A \cap B) = P(B)P(A|B)$$

Chain (Product) Rule

“probability in context”

Conditional probability (definition)

$$P(A|B) \equiv \frac{P(A \cap B)}{P(B)} \quad *$$

Applying a bit of algebra,

$$P(A \cap B) = P(B)P(A|B)$$

In general, we have the **chain (product)** rule:

Product	$P(A_1 \cap A_2) = P(A_1)P(A_2 A_1)$
Chain	$P(A_1 \cap A_2 \cap \dots A_N) = P(A_1)P(A_2 A_1)P(A_3 A_1 \cap A_2) \dots P(A_N A_1 \cap A_2 \cap \dots A_{N-1}) \quad *$

Bayes Rule

Going back to the definition of conditional probability

$$P(A|B) \equiv \frac{P(A \cap B)}{P(B)}$$

Applying a little bit more algebra,

$$P(A \cap B) = P(A)P(B|A)$$

$$\text{and } P(A \cap B) = P(B)P(A|B)$$

$$\text{and thus } P(B)P(A|B) = P(A)P(B|A)$$

$$\text{and we get } P(A|B) = \frac{P(A)P(B|A)}{P(B)}$$

Bayes rule *

Bayes Rule

Going back to the definition of conditional probability

$$P(A|B) \equiv \frac{P(A \cap B)}{P(B)}$$

Applying a little bit more algebra,

$$P(A \cap B) = P(A)P(B|A)$$

$$\text{and } P(A \cap B) = P(B)P(A|B)$$

$$\text{and thus } P(B)P(A|B) = P(A)P(B|A)$$

$$\text{and we get } P(A|B) = \frac{P(A)P(B|A)}{P(B)}$$

Pro tip!: Common to represent denominator as marginalization of numerator:

$$\begin{aligned} P(B) &= \sum_{a \in A} P(A, B) \\ &= \sum_{a \in A} P(A)P(B|A) \end{aligned}$$

Bayes rule *